



## The Continuous Dependence and Numerical Approximation of the Solution of the Quasilinear Pseudo-Parabolic Problem with Periodic Boundary Condition

Irem Baglan, Fatma Kanca,

Department of Mathematics, Kocaeli University, Kocaeli 41380, Turkey  
isakinc@kocaeli.edu.tr

Department of Management Information Systems, Kadir Has University,  
34083, Istanbul, Turkey

fatma.kanca@khas.edu.tr

### ABSTRACT

In this paper we consider a pseudo- parabolic equation with a periodic boundary condition and we prove the stability of a solution on the data. We give a numerical example for the stability of the solution on the data.

### Keywords:

Psedu-parabolic quasilinear equation; periodic-boundary condition; finite difference method.



## Council for Innovative Research

Peer Review Research Publishing System

**Journal:** Journal of Advances in Mathematics

Vol 5, No. 3

[editor@cirworld.com](mailto:editor@cirworld.com)

[www.cirworld.com](http://www.cirworld.com), [member.cirworld.com](http://member.cirworld.com)



## INTRODUCTION

Consider the following mixed problem

$$u_t = u_{xx} - \epsilon u_{xxt} + f(x, t, u), (x, t) \in D := \{0 < x < \pi, 0 < t < T\} \tag{1}$$

$$u(0, t) = u(\pi, t), t \in [0, T] \tag{2}$$

$$u_x(0, t) = u_x(\pi, t), t \in [0, T] \tag{3}$$

$$u(x, 0) = \varphi(x), x \in [0, \pi] \tag{4}$$

for a quasilinear parabolic equation with the nonlinear source term  $f(x, t, u)$ . The functions  $\varphi(x)$  and  $f(x, t, u)$  are given functions on  $[0, \pi]$  and  $D \times (-\infty, \infty)$  respectively. Let  $\epsilon > 0$  is small parameter.

Denote the solution of the problem (1)-(4) by  $u = u(x, t, \epsilon)$ .

The existence, uniqueness and convergence of the weak generalized solution the problem (1)-(4) are considered in [1]. The numerical solution of parabolic problem is considered in [3].

In this study we prove the continuous dependence of the solution  $u = u(x, t, \epsilon)$  upon the data  $\varphi(x)$  and  $f(x, t, u)$ . This kind of conditions arise from many important applications in heat transfer, life sciences, etc. For example the periodic conditions are used on lunar theory and in the system of rocket firing in [2]. Then we give a numerical example using the method for the stability.

## CONTINUOUS DEPENDENCE UPON THE DATA

In this section, we shall prove the continuous dependence of the solution  $u = u(x, t, \epsilon)$  using an iteration method. The continuous dependence upon the data for linear problems by different methods are shown.

**Theorem:** Under the following assumptions, the solution  $u = u(x, t, \epsilon)$  depends continuously upon the data.

(A<sub>1</sub>) Let the function  $f(x, t, u)$  is continuous with respect to all arguments in  $D \times (-\infty, \infty)$  and satisfies the following condition

$$|f(t, x, u) - f(t, x, v)| \leq b(x, t)|u - v|,$$

Where  $b(x, t) \in L^2(D), b(x, t) \geq 0$ ,

$$(A_2) f(x, t, 0) \in C^2[0, \pi], t \in [0, \pi]$$

$$(A_3) \varphi(x) \in C^2[0, \pi].$$

**Proof:** Let  $\phi = \{\varphi, f\}$  and  $\bar{\phi} = \{\bar{\varphi}, \bar{f}\}$  be two sets of data which satisfy the conditions (A<sub>1</sub>)-(A<sub>3</sub>)

Let  $u = u(x, t, \epsilon)$  and  $v = v(x, t, \epsilon)$  the solutions of the problem (1)-(4) corresponding to the data  $\phi$  and  $\bar{\phi}$  respectively and

$$|f(t, x, 0) - \bar{f}(t, x, 0)| \leq \epsilon, \text{ for } \epsilon \geq 0.$$

The solutions of (1)-(4)  $u = u(x, t, \epsilon)$  and  $v = v(x, t, \epsilon)$  in the following form, respectively,

$$u_0(t, \epsilon) = \varphi_0 + \frac{2}{\pi} \int_0^t \int_0^\pi f(\xi, \tau, u(\xi, \tau, \epsilon)) d\xi d\tau,$$

$$u_{ck}(t, \epsilon) = \varphi_{ck} e^{\frac{-(2k)^2 t}{1+\epsilon(2k)^2}} + \frac{1}{1+\epsilon(2k)^2} \int_0^t \int_0^\pi f(\xi, \tau, u(\xi, \tau, \epsilon)) e^{\frac{-(2k)^2(t-\tau)}{1+\epsilon(2k)^2}} \cos 2k\xi d\xi d\tau,$$

$$u_{sk}(t, \epsilon) = \varphi_{sk} e^{\frac{-(2k)^2 t}{1+\epsilon(2k)^2}} + \frac{1}{1+\epsilon(2k)^2} \int_0^t \int_0^\pi f(\xi, \tau, u(\xi, \tau, \epsilon)) e^{\frac{-(2k)^2(t-\tau)}{1+\epsilon(2k)^2}} \sin 2k\xi d\xi d\tau,$$

$$v_0(t, \epsilon) = \bar{\varphi}_0 + \frac{2}{\pi} \int_0^t \int_0^\pi \bar{f}(\xi, \tau, v(\xi, \tau, \epsilon)) d\xi d\tau,$$

$$v_{ck}(t, \epsilon) = \bar{\varphi}_{ck} e^{\frac{-(2k)^2 t}{1+\epsilon(2k)^2}} + \frac{1}{1+\epsilon(2k)^2} \int_0^t \int_0^\pi \bar{f}(\xi, \tau, v(\xi, \tau, \epsilon)) e^{\frac{-(2k)^2(t-\tau)}{1+\epsilon(2k)^2}} \cos 2k\xi d\xi d\tau,$$

$$v_{sk}(t, \epsilon) = \bar{\varphi}_{sk} e^{\frac{-(2k)^2 t}{1+\epsilon(2k)^2}} + \frac{1}{1+\epsilon(2k)^2} \int_0^t \int_0^\pi \bar{f}(\xi, \tau, v(\xi, \tau, \epsilon)) e^{\frac{-(2k)^2(t-\tau)}{1+\epsilon(2k)^2}} \sin 2k\xi d\xi d\tau,$$



From the condition of the theorem we have  $u^{(0)}(t, \varepsilon)$  and  $v^{(0)}(t, \varepsilon) \in B$ . We will prove that the other sequentially approximations satisfy this condition.

$$u_0^{(N+1)}(t, \varepsilon) = u_0^{(0)}(t, \varepsilon) + \frac{2}{\pi} \int_0^t \int_0^\pi f(\xi, \tau, u^{(N)}(\xi, \tau, \varepsilon)) d\xi d\tau,$$

$$u_{ck}^{(N+1)}(t, \varepsilon) = u_{ck}^{(0)}(t, \varepsilon) + \frac{1}{1+\varepsilon(2k)^2} \int_0^t \int_0^\pi f(\xi, \tau, u^{(N)}(\xi, \tau, \varepsilon)) e^{\frac{-(2k)^2(t-\tau)}{1+\varepsilon(2k)^2}} \cos 2k\xi d\xi d\tau, \quad (5)$$

$$u_{sk}^{(N+1)}(t, \varepsilon) = u_{sk}^{(0)}(t, \varepsilon) + \frac{1}{1+\varepsilon(2k)^2} \int_0^t \int_0^\pi f(\xi, \tau, u^{(N)}(\xi, \tau, \varepsilon)) e^{\frac{-(2k)^2(t-\tau)}{1+\varepsilon(2k)^2}} \sin 2k\xi d\xi d\tau,$$

$$v_0^{(N+1)}(t, \varepsilon) = v_0^{(0)}(t, \varepsilon) + \frac{2}{\pi} \int_0^t \int_0^\pi \bar{f}(\xi, \tau, v^{(N)}(\xi, \tau, \varepsilon)) d\xi d\tau,$$

$$v_{ck}^{(N+1)}(t, \varepsilon) = v_{ck}^{(0)}(t, \varepsilon) + \frac{1}{1+\varepsilon(2k)^2} \int_0^t \int_0^\pi \bar{f}(\xi, \tau, v^{(N)}(\xi, \tau, \varepsilon)) e^{\frac{-(2k)^2(t-\tau)}{1+\varepsilon(2k)^2}} \cos 2k\xi d\xi d\tau, \quad (6)$$

$$v_{sk}^{(N+1)}(t, \varepsilon) = v_{sk}^{(0)}(t, \varepsilon) + \frac{1}{1+\varepsilon(2k)^2} \int_0^t \int_0^\pi \bar{f}(\xi, \tau, v^{(N)}(\xi, \tau, \varepsilon)) e^{\frac{-(2k)^2(t-\tau)}{1+\varepsilon(2k)^2}} \sin 2k\xi d\xi d\tau,$$

where  $u_0^{(0)}(t, \varepsilon) = \varphi_0, u_{ck}^{(0)}(t, \varepsilon) = \varphi_{ck} e^{\frac{-(2k)^2 t}{1+\varepsilon(2k)^2}}, u_{sk}^{(0)}(t, \varepsilon) = \varphi_{sk} e^{\frac{-(2k)^2 t}{1+\varepsilon(2k)^2}}$ , and  $v_0^{(0)}(t, \varepsilon) = \bar{\varphi}_0, v_{ck}^{(0)}(t, \varepsilon) = \bar{\varphi}_{ck} e^{\frac{-(2k)^2 t}{1+\varepsilon(2k)^2}}, v_{sk}^{(0)}(t, \varepsilon) = \bar{\varphi}_{sk} e^{\frac{-(2k)^2 t}{1+\varepsilon(2k)^2}}$ .

First of all, we write  $N = 0$  in (5)-(6). we consider  $u^{(1)}(t, \varepsilon) - v^{(1)}(t, \varepsilon)$

$$\begin{aligned} u^{(1)}(t, \varepsilon) - v^{(1)}(t, \varepsilon) &= \frac{u_0^{(1)}(t, \varepsilon) - v_0^{(1)}(t, \varepsilon)}{2} + \sum_{k=1}^\infty \left[ (u_{ck}^{(1)}(t, \varepsilon) - v_{ck}^{(1)}(t, \varepsilon)) \cos 2kx + (u_{sk}^{(1)}(t, \varepsilon) - v_{sk}^{(1)}(t, \varepsilon)) \sin 2kx \right] \\ &= (\varphi_0 - \bar{\varphi}_0) + \frac{2}{\pi} \int_0^t \int_0^\pi [f(\xi, \tau, u^{(0)}(\xi, \tau, \varepsilon)) - \bar{f}(\xi, \tau, v^{(0)}(\xi, \tau, \varepsilon))] d\xi d\tau + (\varphi_{ck} - \bar{\varphi}_{ck}) e^{\frac{-(2k)^2 t}{1+\varepsilon(2k)^2}} \\ &+ \frac{1}{1+\varepsilon(2k)^2} \int_0^t \int_0^\pi [f(\xi, \tau, u^{(0)}(\xi, \tau, \varepsilon)) - \bar{f}(\xi, \tau, v^{(0)}(\xi, \tau, \varepsilon))] e^{\frac{-(2k)^2(t-\tau)}{1+\varepsilon(2k)^2}} \cos 2k\xi d\xi d\tau \quad (7) \\ &+ (\varphi_{sk} - \bar{\varphi}_{sk}) e^{\frac{-(2k)^2 t}{1+\varepsilon(2k)^2}} + \frac{1}{1+\varepsilon(2k)^2} \int_0^t \int_0^\pi [f(\xi, \tau, u^{(0)}(\xi, \tau, \varepsilon)) - \bar{f}(\xi, \tau, v^{(0)}(\xi, \tau, \varepsilon))] e^{\frac{-(2k)^2(t-\tau)}{1+\varepsilon(2k)^2}} \sin 2k\xi d\xi d\tau \end{aligned}$$

Adding and subtracting

$$\frac{2}{\pi} \int_0^t \int_0^\pi f(\xi, \tau, 0) d\xi d\tau, \frac{1}{1+\varepsilon(2k)^2} \int_0^t \int_0^\pi f(\xi, \tau, 0) e^{\frac{-(2k)^2(t-\tau)}{1+\varepsilon(2k)^2}} \cos 2k\xi d\xi d\tau, \frac{1}{1+\varepsilon(2k)^2} \int_0^t \int_0^\pi f(\xi, \tau, 0) e^{\frac{-(2k)^2(t-\tau)}{1+\varepsilon(2k)^2}} \sin 2k\xi d\xi d\tau,$$

to both of side and applying Cauchy Inequality, Hölder Inequality, Lipschitz Condition and Bessel Inequality to the right side of (7) respectively, we obtain:

$$|u^{(1)}(t, \varepsilon) - v^{(1)}(t, \varepsilon)| \leq \frac{|u_0^{(1)}(t, \varepsilon) - v_0^{(1)}(t, \varepsilon)|}{2} + \sum_{k=1}^\infty [ |u_{ck}^{(1)}(t, \varepsilon) - v_{ck}^{(1)}(t, \varepsilon)| + |u_{sk}^{(1)}(t, \varepsilon) - v_{sk}^{(1)}(t, \varepsilon)| ] \leq \|\varphi - \bar{\varphi}\| + 3T + \pi 6\pi 0t 0\pi b 2\xi, \tau d\xi d\tau 12u 0t, \varepsilon + 3T + \pi 6\pi 0t 0\pi b 2\xi, \tau d\xi d\tau 12v 0t, \varepsilon$$

$$+ \frac{\sqrt{3T} + \pi}{\sqrt{6\pi}} \left( \int_0^t \int_0^\pi [f^2(\xi, \tau, 0) - \bar{f}^2(\xi, \tau, 0)] d\xi d\tau \right)^{1/2},$$

where

$$A_T = \|\varphi - \bar{\varphi}\| + \frac{\sqrt{3T} + \pi}{\sqrt{6\pi}} \|b(x, t)\| \|u^{(0)}(t, \varepsilon)\| + \frac{\sqrt{3T} + \pi}{\sqrt{6\pi}} \|\bar{b}(x, t)\| \|v^{(0)}(t, \varepsilon)\| + \frac{\sqrt{3T} + \pi}{\sqrt{6\pi}} \|f - \bar{f}\|$$

$$\|\varphi - \bar{\varphi}\| = \max \frac{|\varphi_0 - \bar{\varphi}_0|}{2} + \sum_{k=1}^\infty |\varphi_{ck} - \bar{\varphi}_{ck}| + |\varphi_{sk} - \bar{\varphi}_{sk}|.$$

For  $N = 1$ ,



$$|u^{(2)}(t, \varepsilon) - v^{(2)}(t, \varepsilon)| \leq \frac{|u^{(2)}(t, \varepsilon) - v^{(2)}(t, \varepsilon)|}{2} + \sum_{k=1}^{\infty} [|u_{ck}^{(2)}(t, \varepsilon) - v_{ck}^{(2)}(t, \varepsilon)| + |u_{sk}^{(2)}(t, \varepsilon) - v_{sk}^{(2)}(t, \varepsilon)|] \leq \frac{\sqrt{3T+\pi}}{\sqrt{6\pi}} \left( \int_0^t \int_0^\pi b^2(\xi, \tau) d\xi d\tau \right)^{\frac{1}{2}} A_T + \frac{\sqrt{3T+\pi}}{\sqrt{6\pi}} \left( \int_0^t \int_0^\pi \bar{b}^2(\xi, \tau) d\xi d\tau \right)^{\frac{1}{2}} A_T.$$

For  $N = 2$ ,

$$|u^{(3)}(t, \varepsilon) - v^{(3)}(t, \varepsilon)| \leq \frac{|u^{(3)}(t, \varepsilon) - v^{(3)}(t, \varepsilon)|}{2} + \sum_{k=1}^{\infty} [|u_{ck}^{(3)}(t, \varepsilon) - v_{ck}^{(3)}(t, \varepsilon)| + |u_{sk}^{(3)}(t, \varepsilon) - v_{sk}^{(3)}(t, \varepsilon)|] \leq \frac{\sqrt{3T+\pi}}{\sqrt{6\pi}} \left( \int_0^t \int_0^\pi b^2(\xi, \tau) |u^{(2)}(t, \varepsilon) - v^{(2)}(t, \varepsilon)|^2 d\xi d\tau \right)^{\frac{1}{2}} + \frac{\sqrt{3T+\pi}}{\sqrt{6\pi}} \left( \int_0^t \int_0^\pi \bar{b}^2(\xi, \tau) |u^{(2)}(t, \varepsilon) - v^{(2)}(t, \varepsilon)|^2 d\xi d\tau \right)^{\frac{1}{2}} \leq \frac{\sqrt{3T+\pi}}{\sqrt{6\pi}} A_T \left( \int_0^t \int_0^\pi b^2(\xi, \tau) \left( \int_0^\tau \int_0^\pi b^2(\xi_1, \tau_1) d\xi_1 d\tau_1 \right) d\xi d\tau \right)^{\frac{1}{2}} + \frac{\sqrt{3T+\pi}}{\sqrt{6\pi}} A_T \left( \int_0^t \int_0^\pi \bar{b}^2(\xi, \tau) \left( \int_0^\tau \int_0^\pi \bar{b}^2(\xi_1, \tau_1) d\xi_1 d\tau_1 \right) d\xi d\tau \right)^{\frac{1}{2}} \leq \left( \frac{\sqrt{3T+\pi}}{\sqrt{6\pi}} \right)^2 A_T \frac{1}{\sqrt{2}} \left[ \left( \int_0^t \int_0^\pi b^2(\xi, \tau) d\xi d\tau \right)^2 \right]^{\frac{1}{2}} + \left( \frac{\sqrt{3T+\pi}}{\sqrt{6\pi}} \right)^2 A_T \frac{1}{\sqrt{2}} \left[ \left( \int_0^t \int_0^\pi \bar{b}^2(\xi, \tau) d\xi d\tau \right)^2 \right]^{\frac{1}{2}}$$

In the same way, for a general value of  $N$  we have

$$|u^{(N+1)}(t, \varepsilon) - v^{(N+1)}(t, \varepsilon)| \leq \frac{|u^{(N+1)}(t, \varepsilon) - v^{(N+1)}(t, \varepsilon)|}{2} + \sum_{k=1}^{\infty} [|u_{ck}^{(N+1)}(t, \varepsilon) - v_{ck}^{(N+1)}(t, \varepsilon)| + |u_{sk}^{(N+1)}(t, \varepsilon) - v_{sk}^{(N+1)}(t, \varepsilon)|] \leq A_T a_N \leq a_N (\|\varphi - \bar{\varphi}\| + C(t) + M_1 \|f - \bar{f}\|)$$

where

$$a_N = \left( \frac{\sqrt{3T+\pi}}{\sqrt{6\pi}} \right)^N \frac{1}{\sqrt{N!}} \left[ \left( \int_0^t \int_0^\pi b^2(\xi, \tau) d\xi d\tau \right)^2 \right]^{\frac{N}{2}} + \left( \frac{\sqrt{3T+\pi}}{\sqrt{6\pi}} \right)^N \frac{1}{\sqrt{N!}} \left[ \left( \int_0^t \int_0^\pi \bar{b}^2(\xi, \tau) d\xi d\tau \right)^2 \right]^{\frac{N}{2}},$$

$$M_1 = \left( \frac{\sqrt{3T+\pi}}{\sqrt{6\pi}} \right)^N.$$

( $M$  = The sequence  $a_N$  is convergent then we can write  $a_N \leq N, \forall N$ )

It follows from the estimation ([3]) that  $\lim_{N \rightarrow \infty} u^{(N+1)}(t) = u(t)$ ,

then let  $N \rightarrow \infty$  for last equation

$$|u(t) - v(t)| \leq M \|\varphi - \bar{\varphi}\| + M_2 \|f - \bar{f}\|$$

where  $M_2 = M \cdot M_1$ .

If  $\|f - \bar{f}\| \leq \varepsilon, \|\varphi - \bar{\varphi}\| \leq \varepsilon$  then  $|u(t) - v(t)| \leq \varepsilon$ .

### NUMERICAL PROCEDURE FOR THE NONLINEAR PROBLEM (1)-(4)

We construct an iteration algorithm for the linearization of the problem (1)-(4):

$$\left( \frac{\partial u^{(n)}}{\partial t} \right) = \left( \frac{\partial^2 u^{(n)}}{\partial x^2} \right) - \varepsilon \left( \frac{\partial^3 u^{(n)}}{\partial x^2 \partial t} \right) + f(x, t, u^{(n-1)}), \quad (x, t) \in D \tag{8}$$

$$u^{(n)}(0, t) = u^{(n)}(\pi, t), \quad t \in [0, T] \tag{9}$$

$$u_x^{(n)}(0, t) = u_x^{(n)}(\pi, t), \quad t \in [0, T] \tag{10}$$

$$u^{(n)}(x, 0) = \varphi(x), \quad x \in [0, \pi]. \tag{11}$$

Let  $u^{(n)}(x, t) = v(x, t)$  and  $f(x, t, u^{(n-1)}) = \tilde{f}(x, t)$ . Then the problem (8)-(11) can be written as a linear problem:

$$\left( \frac{\partial v}{\partial t} \right) = \left( \frac{\partial^2 v}{\partial x^2} \right) - \varepsilon \left( \frac{\partial^3 v}{\partial x^2 \partial t} \right) + \tilde{f}(x, t) \quad (x, t) \in D \tag{12}$$

$$v(0, t) = v(\pi, t), \quad t \in [0, T] \tag{13}$$

$$v_x(0, t) = v_x(\pi, t), \quad t \in [0, T] \tag{14}$$

$$v(x, 0) = \varphi(x), \quad x \in [0, \pi]. \tag{15}$$

We use the finite difference method to solve (12)-(15).

We subdivide the intervals  $[0, \pi]$  and  $[0, T]$  into  $M$  and  $N$  subintervals of equal lengths  $h = (\pi/M)$  and  $\tau = (T/N)$ , respectively.

Then, we add two lines  $x = 0$  and  $x = (M + 1)\pi h$  to generate the fictitious points needed for dealing with the secondary boundary condition.



We choose the implicit scheme, which is absolutely stable and has a second order accuracy in  $h$  and a first order accuracy in  $\tau$ .

The implicit monotone difference scheme for (12)-(15) is as follows:

$$\frac{(v_i^{j+1} - v_i^j)}{\tau} = \frac{1}{h^2} (v_{i-1}^{j+1} - 2v_i^{j+1} + v_{i+1}^{j+1}) - \frac{\varepsilon}{h^2\tau} (v_{i-1}^{j+1} - 2v_i^{j+1} + v_{i+1}^{j+1}) - (v_{i-1}^j - 2v_i^j + v_{i+1}^j) + \tilde{f}_i^{j+1}$$

$$v_i^0 = \phi_i, v_0^j = v_M^j, v_{x,0}^j = v_{x,M}^j$$

where  $0 \leq i \leq M$  and  $1 \leq j \leq N$  are the indices for the spatial and time steps, respectively,  $v_i^j$  is the approximation to  $v(x_i, t_j)$ ,  $\tilde{f}_{i,j} = \tilde{f}(x_i, t_j)$ ,  $\phi_i = \varphi(x_i)$ ,  $x_i = ih$ ,  $t_j = j\tau$ .

At the  $t = 0$  level, adjustment should be made according to the initial condition and the compatibility requirements.

### Numerical Example

In this section, we will consider an example of numerical solution of the problem (1)-(3).

This problem was solved by applying the iteration scheme and the finite difference scheme which were explained in the Section 2. The condition

$$error(i, j) := \|u_i^{j(n+1)} - u_i^{j(n)}\|_\infty$$

with

$$error(i, j) := 10^{-3}$$

was used as a stopping criteria for the iteration process.

**Example:** Consider the problem

$$\left(\frac{\partial u}{\partial t}\right) - \left(\frac{\partial^2 u}{\partial x^2}\right) - \varepsilon \left(\frac{\partial^3 u}{\partial x^2 \partial t}\right) = 4(-\cos 2x + (\sin 2x)^2)(1 - \varepsilon)u, (x, t) \in D$$


$$u(x, 0) = \exp(\cos 2x), x \in [0, \pi],$$

$$u(0, t) = u(\pi, t), t \in [0, T], u_x(0, t) = u_x(\pi, t), t \in [0, T].$$

It is easy to see that the analytical solution of this problem is


$$u(x, t) = \exp(t + \cos 2x)$$

The comparisons between the analytical solution and the numerical finite difference solution for different  $\varepsilon$  values when  $T = 1$  are shown in Figure 1 and 2.



**Figure 1 - The exact and numerical solutions of  $u(x,1)$**


The exact and numerical solutions of  $u(x,1)$  for  $\varepsilon=0$ , the exact solution is shown with dashes line.



**Figure 2 -The exact and numerical solutions of  $u(x,1)$**

The exact and numerical solutions of  $u(x,1)$  for  $\epsilon=0.05$ , the exact solution is shown with dashes line.

In Figure3 we show that the analytical solution for  $\epsilon = 0$  and the numerical solution for  $\epsilon = 0, \epsilon = 0.1, \epsilon = 0.05$ .



**Figure 3 - The exact and numerical solutions of  $u(x,1)$**

The exact and numerical solutions of  $u(x,1)$ ,  $(-\Delta)$  for  $\epsilon=0$ ,  $(-)$  for  $\epsilon=0.05$ ,  $(-*)$  for  $\epsilon=0.1$ , the exact solution is shown with dashes line.

Relative errors obtained on different grids for different  $\epsilon$  are shown on Table 1.

**Table 1. The relative errors for different grids for  $\epsilon = 0, \epsilon = 0.05$  and  $\epsilon = 0.1$ .**

$h$	$\tau$	$\epsilon = 0$	$\epsilon = 0.05$	$\epsilon = 0.1$
0.1571	0.025	0.0142	0.0250	0.0553
0.1571	0.0125	0.0151	0.0269	0.0624
0.0785	0.025	0.0160	0.0334	0.0642
0.0785	0.0125	0.0179	0.0373	0.0720



From Figure 3 and Table 1,2 and 3 when  $\varepsilon$  approximates to zero the numerical solutions converge to the exact solution for  $\varepsilon = 0$  like theoretical results .

## REFERENCES

- [1] Halilov H.,Ciftci(Baglan) I. 2009.Fourier Method for a Qusilinear Pseudo-Parabolic Equation with Periodic Boundary Condition.Hacettepe International Journal of Pure and Applied Mathematics. 52, 717-727.
- [2] Hill GW. 1886 On the part of the motion of the lunar perigee which is a function ofthe mean motions of the sun and moon. ActaMathematica .8, 1-36.
- [3] Kanca F., Baglan I. 2013 Continuous dependence on data for a solution of the quasilinear parabolic equation with a periodic boundary condition.Boundary Value Problems.28.
- [4] Sakınc(Baglan) I.2010 Numerical Solution of a Qusilinear Parabolic Problem with Periodic Boundary Condition. HacettepeJournal of Mathematics and Statistics.39(2), 183-189.

