



Moments of order statistics from nonidentically Distributed Lomax, exponential Lomax and exponential Pareto Variables

Nasr Ibrahim Rashwan
Faculty of Commerce – Tanta University, Egypt
nasrrashwan@yahoo.com

Abstract

In this paper, the probability density function and the cumulative distribution function of the r^{th} order statistic arising from independent nonidentically distributed (INID) Lomax, exponential Lomax and exponential Pareto variables are presented. The moments of order statistics from INID Lomax, exponential Lomax and exponential Pareto were derived using the technique established by Barakat and Abdelkader. Also, numerical examples are given.

Keywords

Moments of order statistics, nonidentically distributed order statistics, Lomax distribution, exponential Lomax distribution, exponential Pareto distribution..

1. Introduction

Three techniques have been established in literature to compute moments of order statistics of independent nonidentically distributed random variables. The first technique is created by Balakrishnan (1994). This technique requires a basic relation between to probability density function (pdf) and the cumulative distribution function (cdf) and referred to as differential equation technique (DET). Many authors used to this technique to derive the moments of INID order statistics for continuous distributions (See, Childs and Balakrishnan (2006) and Mohie Elidin et al (2007)).

Barakat and Abdelkader (2000) established the second technique to compute the moments of order statistics from nonidentically distributed weibull variables and referred to as (BAT). This technique that the cdf of the distribution can be written in the form $F(x) = 1 - \theta(x)$, which is satisfied in this distribution. Many authors used to BAT technique to compute the moments of INID order statistics for several continuous distributions (more details, see Abdelkader (2004a) Abdelkader (2004b), Jamjoom (2006), Abdelkader (2010), Jamjoom and Al-Saiary (2010) and Jamjoom and Al-Saiary (2013)).

The third technique was developed by Jamjoom and Al-Saiary (2011) which is the moment generating function technique it depends on (BAT) and referred to as (M.G.F. BAT). The third technique was used by Al-Saiary (2015) to compute the moments of INID order statistics for standard type II generalized logistic variables.

In this paper the pdf and the cdf of the r^{th} order statistic arising from INID Lomax, exponential Lomax and exponential Pareto distributions are given in section 2. In section 3, we drive the moments of the r^{th} order statistics of INID random variables arising from Lomax, exponential Lomax and exponential Pareto using (BAT). Finally, some conclusions are addressed in section 4.

2. Nonidentical order statistics from Lomax, exponential Lomax and exponential Pareto distributions

Let X_1, X_2, \dots, X_n be independent random variables having cumulative distribution functions (cdf) $F_1(x), \dots, F_n(x)$ and probability density functions (pdfs) $f_1(x), f_2(x), \dots, f_n(x)$, respectively. Let $x_{1:n} \leq \dots \leq x_{n:n}$ denote the order statistics obtained by arranging the n in increasing order of magnitude. Then the pdf of the r^{th} order statistic $X_{r:n}$, ($1 \leq r \leq n$), can be written as:

$$f_{r:n}(x) = \frac{1}{(r-1)!(n-r)!} \sum_P \prod_{a=1}^{r-1} F_{i_a}(x) f_{i_r}(x) \prod_{c=r+1}^n [1-F_{i_c}(x)] \quad (1)$$

Where \sum_P denotes the summation overall $n!$ permutations (i_1, i_2, \dots, i_n) of $(1, 2, \dots, n)$ (See Bapat and Beg (1989)). Put the previous pdf of the r^{th} order statistic $X_{r:n}$ in the form of permanent as:

$$f_{r:n}(x) = \frac{1}{(r-1)!(n-r)!} \text{Per} \begin{bmatrix} \underbrace{F(x)}_{r-1} & \underbrace{F(x)}_1 & \underbrace{(1-F(x))}_{n-r} \end{bmatrix}$$



$$= \frac{1}{(r-1)! (n-r)!} \begin{matrix} \left. \begin{matrix} F_1(x) & F_2(x) & \dots & F_n(x) \\ \vdots & \vdots & & \vdots \\ F_1(x) & F_2(x) & \dots & F_n(x) \\ f_1(x) & f_2(x) & \dots & f_n(x) \\ 1-F_1(x) & 1-F_2(x) & \dots & 1-F_n(x) \\ \vdots & \vdots & & \vdots \\ 1-F_1(x) & 1-F_2(x) & \dots & 1-F_n(x) \end{matrix} \right\} \begin{matrix} (r-1) \text{ rows} \\ 1 \text{ row} \\ (n-r) \text{ rows} \end{matrix} \end{matrix} \quad (2)$$

and the cdf of the r^{th} order statistic $X_{r:n}$ can be written as:

$$F_{r:n}(x) = \sum_{i=r}^n \sum_{P_j} \prod_{a=1}^i F_{i_a}(x) \prod_{a=j+1}^n [1-F_{i_a}(x)] \quad (3)$$

Where P_j is all permutations of i_1, i_2, \dots, i_n for $1, 2, \dots, n$ which satisfy $i_1 < i_2 < \dots < i_j$ and $i_{j+1} < i_{j+2} < \dots < i_n$ and it is conveniently expressed in terms of permanent as

$$F_{r:n}(x) = \sum_{i=r}^n \frac{1}{i! (n-i)!} \text{Per} \left[\underbrace{F(x)}_i \quad \underbrace{(1-F(x))}_{n-i} \right], \quad -\infty < x < \infty \quad (4)$$

Where $F(x)$ and $[1 - F(x)]$ denote the column vectors $[F_1(x), F_2(x), \dots, F_n(x)]'$ and $[1 - F_1(x), 1 - F_2(x), \dots, 1 - F_n(x)]'$ respectively. So the $F_{r:n}(x)$ can be rewritten as

$$F_{r:n}(x) = \sum_{i=r}^n \frac{1}{i! (n-i)!} \text{Per} \left[\begin{matrix} F_1(x) & 1-F_1(x) \\ F_2(x) & 1-F_2(x) \\ \vdots & \vdots \\ F_n(x) & 1-F_n(x) \end{matrix} \right], \quad -\infty < x < \infty \quad (5)$$

The cdf of the smallest order statistic ($r = 1$) $X_{1:n}$ is given by:

$$F_{L:n}(x) = \sum_{i=1}^n \frac{1}{i! (n-i)!} \text{Per} \left[\begin{matrix} F_1(x) & 1-F_1(x) \\ F_2(x) & 1-F_2(x) \\ \vdots & \vdots \\ F_n(x) & 1-F_n(x) \end{matrix} \right], \quad -\infty < x < \infty$$

and the cdf of the largest order statistic ($r = n$) $X_{n:n}(x)$ is given by:

$$F_{n:n}(x) = \frac{1}{n!} \text{Per} \left[\begin{matrix} F_1(x) \\ \vdots \\ F_n(x) \end{matrix} \right], \quad -\infty < x < \infty$$

Note that permanent is a square matrix, which is defined similar to the determinants except that all elements in the expansion have a positive sign (Minc (1987) and Balakrishnan (1994)).

In this paper, we consider the case where the random variables $X_i, i = 1, 2, \dots, n$ are independent and nonidentical having Lomax (L), exponential Lomax (EL) and exponential Pareto (EP) distributions with cdfs, $F_L(x), F_{EL}(x)$ and $F_{EP}(x)$ and pdfs $f_L(x), f_{EL}(x)$ and $f_{EP}(x)$ respectively (more details see, Lemonte and Corderio (2013), Abdel Al-Kadim and Boshi (2013) and El-Bassiouny et al (2015)).



Where

$$F_{Li}(x) = 1 - (1 + \beta x)^{-\alpha_i}, f_{Li}(x) = \beta \alpha_i (1 + \beta x)^{-(\alpha_i+1)}, x > 0, \alpha_i, \beta > 0 \quad (6)$$

$$F_{ELi}(x) = 1 - e^{-\lambda_i(\frac{\beta}{x+\beta})^{-\alpha}}, f_{ELi}(x) = \frac{\alpha \lambda_i}{\beta} (\frac{\beta}{x+\beta})^{-(\alpha+1)} e^{-\lambda_i(\frac{\beta}{x+\beta})^{-\alpha}}, x > -\beta, \alpha, \beta, \lambda_i > 0 \quad (7)$$

$$F_{EPi}(x) = 1 - e^{-\lambda_i(\frac{x}{p})^\theta}, f_{EPi}(x) = \frac{\lambda_i \theta}{p} (\frac{x}{p})^{\theta-1} e^{-\lambda_i(\frac{x}{p})^\theta}, x > 0, \theta, p, \lambda_i > 0 \quad (8)$$

3. The moment of the r^{th} order statistics arising from INID Lomax, exponential lomax and exponential Pareto random variables

In this section, we drive the moments of order statistics from INID random variables arising from L, EL and EP. We need to following theorem which is established by Barakat and Abdelkader (2003).

Theorem 1: Let X_1, X_2, \dots, X_n be independent nonidentically distribution random variables, The k^{th} moment of order statistics $\mu_{r:n}^{(k)}$ for $1 \leq r \leq n$ and $k = 1, 2, \dots$ is given by:

$$\mu_{r:n}^{(k)} = \sum_{j=n-r+1}^n (-1)^{j-(n-r+1)} \binom{j-1}{n-r} I_j(k) \quad (9)$$

Where

$$I_j(k) = \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} \dots \sum_{k \int_0^\infty x^{k-1} \prod_{t=1}^j G_{i_t}(x) dx}, j=1,2,\dots,n, \quad (10)$$

$G_{i_t}(x) = 1 - F_{i_t}(x)$ with (i_1, i_2, \dots, i_n) are permutations of $(1, 2, \dots, n)$ for which $i_1 < i_2 < \dots < i_n$. The proof of this Theorem 1 see to Barakat and Abdelkader (2003).

3.1. Moments of order statistics from INID Lomax random variables

The following Theorem gives an explicit expression for $I_j(k)$ when X_1, X_2, \dots, X_n are INID Lomax random variables.

Theorem 2: for $1 \leq r \leq n$ and $k = 1, 2, \dots$

$$I_j(k) = \frac{k}{\beta^{k-1}} \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} \dots \sum_{B(k, \sum_{t=1}^j \alpha_{i_t} - k)} \quad (11)$$

Where $B(., .)$ is the complete beta function

Proof: on applying Theorem 1 and using eq. (10), we get

$$\begin{aligned} I_j(k) &= \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} \dots \sum_{k \int_0^\infty x^{k-1} \prod_{t=1}^j [1 - F_{Li}(x)]} \\ I_j(k) &= \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} \dots \sum_{k \int_0^\infty x^{k-1} \prod_{t=1}^j (1 + \beta x)^{-\alpha_{i_t}}} \\ &= \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} \dots \sum_{k \int_0^\infty \frac{x^{k-1}}{(1 + \beta x)^{\sum_{t=1}^j \alpha_{i_t}}} \\ &= \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} \dots \sum_{\frac{k}{\beta^{k-1}} \int_0^\infty \frac{(\beta x)^{k-1}}{(1 + \beta x)^{\sum_{t=1}^j \alpha_{i_t}}} dx} \quad (12) \end{aligned}$$



By using the known relation $B(p, q) = \int_0^{\infty} \frac{x^{p-1}}{(1+x)^{p+q}} dx$, then

$$\int_0^{\infty} \frac{(\beta x)^{k-1}}{(1+\beta x)^{\sum_{t=1}^j \alpha_{i_t}}} dx = B\left[k, \sum_{t=1}^j \alpha_{i_t} - k\right] \quad (13)$$

By substituting eq. (13) in eq. (12), we get Theorem 2.

Result 1: Substituting eq. (11) in eq. (9), The k^{th} moments of the r^{th} order statistics from INID Lomax distribution can be written as:

$$\mu_{r:n}^{(k)} = \sum_{j=n-r+1}^n (-1)^{j-(n-r+1)} \binom{j-1}{n-r} \frac{k}{\beta^{k-1}} \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} B\left[k, \sum_{t=1}^j \alpha_{i_t} - k\right] \quad (14)$$

Also, we can obtain the k^{th} moment of the smallest order statistic $x_{1:n}$ and the largest order statistic $x_{n:n}$ from INID Lomax as follows:

$$\mu_{1:n}^{(k)} = \frac{k}{\beta^{k-1}} B\left[k, \sum_{i=1}^n \alpha_{i_t} - k\right] \quad (15)$$

and

$$\mu_{n:n}^{(k)} = \sum_{j=1}^n (-1)^{j-1} \frac{k}{\beta^{k-1}} \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} B\left[k, \sum_{t=1}^j \alpha_{i_t} - k\right] \quad (16)$$

Example (1)

Let $n = 3$, $X_1 \sim \text{Lomax}(\beta = 2, \alpha_1 = 3)$, $X_2 \sim \text{Lomax}(\beta = 2, \alpha_2 = 3)$ and $X_3 \sim \text{Lomax}(\beta = 2, \alpha_3 = 4)$. When $k = 1$, then

Using eq. (15)

$$\mu_{1:3} = B\left(1, \sum_{i=1}^n \alpha_i - 1\right) = B(1, \alpha_1 + \alpha_2 + \alpha_3 - 1) = 0.1111$$

Using eq. (14)

$$\begin{aligned} \mu_{2:3} &= \sum_{j=2}^3 (-1)^{j-2} \binom{j-1}{1} I_j(1) = I_2(1) - 2 I_3(1) \\ &= [B(1, \alpha_1 + \alpha_2 - 1) + B(1, \alpha_1 + \alpha_3 - 1) + B(1, \alpha_2 + \alpha_3 - 1)] - 2 B(1, \alpha_1 + \alpha_2 + \alpha_3 - 1) = 0.3111 \end{aligned}$$

and using eq. (16)

$$\mu_{3:3} = \sum_{j=1}^3 (-1)^{j-1} I_j(1) = I_1(1) - I_2(1) + I_3(1) = 0.9111$$

Where $I_1(1) = B(1, \alpha_1 - 1) + B(1, \alpha_2 - 1) + B(1, \alpha_3 - 1)$

3.2. Moments of order statistic from INID exponential Lomax random variables

Theorem 3: for $1 \leq r \leq n$ and $k = 1, 2, \dots$,

$$I_j^{(k)} = \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} \frac{k \beta^k}{\alpha} \sum_{L=0}^{k-1} (-1)^L \binom{k-1}{L} \frac{\Gamma\left(\frac{k-L}{\alpha}\right)}{\left(\sum_{t=1}^j \lambda_{i_t}\right)^{\frac{k-L}{\alpha}}} \quad (17)$$

Proof: By using Theorem 1 and eq. (10), then

$$\begin{aligned} I_j^{(k)} &= \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} k \int_{-\beta}^{\infty} x^{k-1} \prod_{t=1}^j [1 - F_{EL_i}(x)] dx \\ &= \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} k \int_{-\beta}^{\infty} x^{k-1} e^{-\left(\frac{\beta}{x+\beta}\right)^{\alpha} \sum_{t=1}^j \lambda_{i_t}} dx \end{aligned}$$



Let, $y = \left(\frac{\beta}{x+\beta}\right)^{-\alpha}$, $y(-\beta) = 0$, $y(\infty) = \infty$, $x = \beta(y^{\frac{1}{\alpha}} - 1)$ and $dx = \frac{\beta}{\alpha} y^{\frac{1}{\alpha}-1} dy$, then

$$\begin{aligned} I_j(k) &= \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} \dots \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} \frac{k \beta^k}{\alpha} \int_0^\infty y^{\frac{k-1}{\alpha}} (1-y^{-1/\alpha})^{k-1} e^{-y \sum_{t=1}^j \lambda_{i_t}} y^{\frac{1}{\alpha}-1} dy \\ &= \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} \dots \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} \frac{k \beta^k}{\alpha} \sum_{L=0}^{k-1} (-1)^L \binom{k-1}{L} \int_0^\infty y^{\frac{(k-L-1)}{\alpha}} e^{-y \sum_{t=1}^j \lambda_{i_t}} dy \\ &= \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} \dots \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} \frac{k \beta^k}{\alpha} \sum_{L=0}^{k-1} (-1)^L \binom{k-1}{L} \frac{\Gamma\left(\frac{k-L}{\alpha}\right)}{\left(\sum_{t=1}^j \lambda_{i_t}\right)^{\frac{(k-L)}{\alpha}}} \end{aligned}$$

Result 2: Substituting eq.(17) in eq.(9), the k^{th} moments of the r^{th} order statistics from INID exponential Lomax can be written as:

$$\begin{aligned} \mu_{r:n}^{(k)} &= \sum_{j=n-r+1}^n (-1)^{j-(n-r+1)} \binom{j-1}{n-r} \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} \dots \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} \frac{k \beta^k}{\alpha} \sum_{L=0}^{k-1} (-1)^L \\ &\quad \times \frac{\Gamma\left(\frac{k-L}{\alpha}\right)}{\left(\sum_{t=1}^j \lambda_{i_t}\right)^{\frac{(k-L)}{\alpha}}} \end{aligned} \quad (18)$$

Also, the k^{th} moment of the smallest order statistic $x_{1:n}$ from INID exponential Lomax random variables can be written as:

$$\mu_{1:n}^{(k)} = \frac{k \beta^k}{\alpha} \sum_{L=0}^{k-1} (-1)^L \binom{k-1}{L} \frac{\Gamma\left(\frac{k-L}{\alpha}\right)}{\left(\sum_{i=1}^n \lambda_i\right)^{\frac{(k-L)}{\alpha}}} \quad (19)$$

and the k^{th} moment of the largest order statistic $x_{n:n}$ from INID exponential Lomax random variables can be written as:

$$\mu_{n:n}^{(k)} = \sum_{j=1}^n (-1)^{j-1} \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} \dots \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} \frac{k \beta^k}{\alpha} \sum_{L=0}^{k-1} (-1)^L \binom{k-1}{L} \frac{\Gamma\left(\frac{k-L}{\alpha}\right)}{\left(\sum_{t=1}^j \lambda_{i_t}\right)^{\frac{(k-L)}{\alpha}}} \quad (20)$$

Example (2)

Let $n = 3$, $X_1 \sim \text{EL}(\beta = 3, \alpha = 2, \lambda_1 = 1)$, $X_2 \sim \text{EL}(\beta = 3, \alpha = 2, \lambda_2 = 2)$ and $X_3 \sim \text{EL}(\beta = 3, \alpha = 2, \lambda_3 = 3)$. when $k = 1$, then

Using eq. (19)

$$\mu_{1:3} = \frac{2}{3} \cdot \frac{\Gamma(0.5)}{\left(\sum_{i=1}^3 \lambda_i\right)^{0.5}} = 1.0853$$

$$\text{Using eq. (18),} \quad \mu_{2:3} = I_2(1) - 2 I_3(1) = 1.8817$$

Where

$$I_2(1) = \frac{2}{3} \Gamma(.5) \left[\frac{1}{(\lambda_1 + \lambda_2)^{0.5}} + \frac{1}{(\lambda_1 + \lambda_3)^{0.5}} + \frac{1}{(\lambda_2 + \lambda_3)^{0.5}} \right] = 4.0524 \text{ and}$$

$$I_3(1) = \frac{2}{3} \frac{\Gamma(.5)}{(\lambda_1 + \lambda_2 + \lambda_3)^{0.5}} = 1.0853$$



and using eq. (20), $\mu_{3:3} = I_1(1) - I_2(1) + I_3(1) = 3.1053$
Where

$$I_1(1) = \frac{3}{2} \Gamma(.5) \left[\frac{1}{\lambda_1^{0.5}} + \frac{1}{\lambda_2^{0.5}} + \frac{1}{\lambda_3^{0.5}} \right] = 6.0724$$

3.3. Moments of order statistics from INID exponential Pareto random variables

Theorem 4: for $1 \leq r \leq n$ and $k = 1, 2, \dots$,

$$I_j(k) = \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} \dots \sum_{k} \frac{k p^k}{\theta} \frac{\Gamma\left(\frac{k}{\theta}\right)}{\left(\sum_{t=1}^j \lambda_{i_t}\right)^{\frac{k}{\theta}}} \quad (21)$$

Proof: on applying Theorem 1 and using eq.(10), Then

$$\begin{aligned} I_j(k) &= \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} \dots \sum_{k} \int_0^{\infty} x^{k-1} \prod_{t=1}^j [1 - F_{EP_i}(x)] dx \\ &= \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} \dots \sum_{k} \int_0^{\infty} x^{k-1} e^{-\frac{x}{p} \sum_{t=1}^j \lambda_{i_t}} dx \end{aligned}$$

Let $y = \left(\frac{x}{p}\right)^{\theta}$, $y(0) = 0$, $y(\infty) = \infty$, $x = p y^{\frac{1}{\theta}}$ and $dx = \frac{p}{\theta} y^{\frac{1}{\theta}-1} dy$, then

$$\begin{aligned} I_j(k) &= \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} \dots \sum_{k} \frac{k p^k}{\alpha} \int_0^{\infty} y^{k-1} e^{-y \sum_{t=1}^j \lambda_{i_t}} dy \\ &= \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} \dots \sum_{k} \frac{k p^k}{\theta} \cdot \frac{\Gamma\left(\frac{k}{\theta}\right)}{\left(\sum_{t=1}^j \lambda_{i_t}\right)^{\frac{k}{\theta}}} \end{aligned}$$

Result 3: Substituting eq.(21) in eq.(9), the k^{th} moments of the r^{th} order statistic from INID exponential Pareto random variables can be written as:

$$\mu_{r:n}^{(k)} = \sum_{j=n-r+1}^n (-1)^{j-(n-r+1)} \binom{j-1}{n-r} \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} \dots \sum_{k} \frac{k p^k}{\theta} \cdot \frac{\Gamma\left(\frac{k}{\theta}\right)}{\left(\sum_{t=1}^j \lambda_{i_t}\right)^{\frac{k}{\theta}}} \quad (22)$$

Also, we can obtain the k^{th} moment of $x_{1:n}$ and $x_{n:n}$ from INID exponential Pareto as follows:

$$\mu_{1:n}^{(k)} = \frac{k p^k}{\theta} \cdot \frac{\Gamma\left(\frac{k}{\theta}\right)}{\left(\sum_{i=1}^n \lambda_i\right)^{\frac{k}{\theta}}} \quad (23)$$

and

$$\mu_{n:n}^{(k)} = \sum_{j=1}^n (-1)^{j-1} \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} \dots \sum_{k} \frac{k p^k}{\theta} \cdot \frac{\Gamma\left(\frac{k}{\theta}\right)}{\left(\sum_{t=1}^j \lambda_{i_t}\right)^{\frac{k}{\theta}}} \quad (24)$$



Example (3)

Let $n = 5$, $X_1 \sim EP (P = 2, \theta = 1, \lambda_1 = 1)$, $X_2 \sim EP (P = 2, \theta = 1, \lambda_2 = 2)$,

$X_3 \sim EP (P = 2, \theta = 1, \lambda_3 = 3)$, $X_4 \sim EP (P = 2, \theta = 1, \lambda_4 = 4)$ and

$X_5 \sim EP (P = 2, \theta = 1, \lambda_5 = 5)$. when $k = 1$, then

Using eq. (23)

$$\mu_{1:5} = 2 \cdot \frac{\Gamma(1)}{\left(\sum_{i=1}^5 \lambda_i \right)} = 0.1333$$

Using eq. (22)

$$\mu_{2:5} = \sum_{j=4}^5 (-1)^{j-4} \binom{j-1}{3} I_j(1) = I_4(1) - 4 I_5(1) = 0.3119$$

Where

$$I_4(1) = 2 \left[\frac{\Gamma(1)}{\lambda_1 + \lambda_2 + \lambda_3 + \lambda_4} + \frac{\Gamma(1)}{\lambda_1 + \lambda_2 + \lambda_3 + \lambda_5} + \frac{\Gamma(1)}{\lambda_1 + \lambda_2 + \lambda_4 + \lambda_5} + \frac{\Gamma(1)}{\lambda_1 + \lambda_3 + \lambda_4 + \lambda_5} + \frac{\Gamma(1)}{\lambda_2 + \lambda_3 + \lambda_4 + \lambda_5} \right]$$

$$= 0.8451 \text{ and}$$

$$I_5(1) = 2 \cdot \frac{\Gamma(1)}{\lambda_1 + \lambda_2 + \lambda_3 + \lambda_4 + \lambda_5} = 0.1333$$

and using eq. (24)

$$\mu_{5:5} = \sum_{j=1}^5 (1)^{j-1} I_j(1) = I_1(1) - I_2(1) + I_3(1) - I_4(1) + I_5(1) = 0.6969$$

Where for example. When $n = 5$, then

$$I_2(k) = \frac{kp^k}{\theta} \sum_{1 \leq i_1 < i_2 \leq 5} \dots \sum \frac{\Gamma\left(\frac{k}{\theta}\right)}{\left(\sum_{t=1}^2 \lambda_{i_t} \right)^{(k/\theta)}$$

$$= \frac{kp^k}{\theta} \Gamma\left(\frac{k}{\theta}\right) \left[\frac{1}{(\lambda_1 + \lambda_2)^{(k/\theta)}} + \frac{1}{(\lambda_1 + \lambda_3)^{(k/\theta)}} + \frac{1}{(\lambda_1 + \lambda_4)^{(k/\theta)}} + \frac{1}{(\lambda_1 + \lambda_5)^{(k/\theta)}} + \frac{1}{(\lambda_2 + \lambda_3)^{(k/\theta)}} \right.$$

$$\left. + \frac{1}{(\lambda_2 + \lambda_4)^{(k/\theta)}} + \frac{1}{(\lambda_2 + \lambda_5)^{(k/\theta)}} + \frac{1}{(\lambda_3 + \lambda_4)^{(k/\theta)}} + \frac{1}{(\lambda_3 + \lambda_5)^{(k/\theta)}} + \frac{1}{(\lambda_4 + \lambda_5)^{(k/\theta)}} \right]$$

4. Conclusion

In this paper, exact moments of the r^{th} order statistic from independent and nonidentically distributed random variables for the Lomax, exponential Lomax and exponential Pareto distribution are derived using the BAT technique. Some numerical examples are presented to illustrate the theorems and results of this study.

References

- [1] Abdelkader, Y.H., (2004a), Computing the moments of order statistics from nonidentically distributed erlang variables, Statistical Paper, 45, 563-570.



- [2] Abdelkader, Y.H., (2004b), Computing the moments of order statistics from nonidentically distributed gamma variables with applications, *Int. J. Math. Game Theo. Algebra*, 14, 1 – 8.
- [3] Abdelkader, Y.H., (2010), Computing the moments of order statistics from independent nonidentically distributed beta random variables, *Stat. Papers*, 51, 307-313.
- [4] Abed Al-Kadim, K. and Boshi, M.A., (2013), Exponential Pareto distribution, *Mathematical Theory and Modeling*, 3, 135-146.
- [5] Al-Saiary, Z.A., (2015), Order statistics from nonidentical standard type generalized logistic variables and applications at moments, *American Journal of Theoretical and applied statistics*, 4, 1 – 5.
- [6] Balakrishnan, N., (1994), Order statistics from nonidentically exponential random variables and some applications, *comput. in statistics Data-Anal.*, 18, 203-253.
- [7] Bapat, R.B. and Beg, M.I., (1989), order statistics from nonidentically distributed variables and permanents, *Sankhya*, A, 51, 79-93.
- [8] Barakat, H.M. and Abdelkader, Y.H., (2000), Computing the moments of order statistics from nonidentically distributed weibull, *Journal of Computation and applied mathematics*, 117, 85 – 90.
- [9] Barakat, H.M. and Abdelkader, Y.H., (2003), Computing the moments of order statistics from nonidentical random variables, *Statistical Methods and applications*, 13, 15 – 26.
- [10] Childs, A. and Balakrishnan, N., (2006), Relations for order statistics from nonidentical Logistic random variables and assessment of the effect of multiple outliers on bias of linear estimators, *J. Stat. Plann. Inference*, 136, 2227 – 2253.
- [11] El-Bassiouny, A.H., Abdo, N. and Shahen, H.S., (2015), Exponential Lomax distribution, *International Journal of Computer applications*, 121, 24– 29.
- [12] Jamjoom, A.A. (2006), Computing the moments of order statistics from nonidentically distributed Burr XII random variables, *Journal of Mathematics and Statistics*, 2, 432– 438.
- [13] Jamjoom, A.A. and Al-Saiary, Z. A. (2010), Moments of order statistics from nonidentically distributed three parameters beta I and erlang truncated exponential variables, *Journal of Mathematics and statistics*, 6, 442-448.
- [14] Jamjoom, A.A., and Al-Saiary, Z. A., (2011), Moment generating function technique for moments of order statistics from nonidentically distributed random variables, *International Journal of Statistics and System*, 6, 177-188.
- [15] Jamjoom, A.A. and Al-Saiary, Z. A., (2013), Moments of nonidentical order statistics from Burr XII distribution with gamma and normal outliers, *Journal of Mathematics and statistics*, 9, 51- 61.
- [16] Lemonte, A.J. and Cordeiro, G. M., (2013), An extended Lomax distribution, *Statistics*, 47, 800 – 816.
- [17] Minc, H., (1987), Theory of Permanents 1981 – 1985, *Linear and multi linear Algebra*, 21, 109 – 198.
- [18] Mohie Elidin, M., Mahmoud, M., Moshref, M. and Mohamed, M. (2007), on independent and nonidentical order statistics and associated inference, Department of mathematics, Cairo, Al-Azhar University.