



Hermite collocation method for solving Hammerstein integral equations

Y. A. Amer¹, A. M. S. Mahdy^{1, 2} and H. A. R. Namoos³

¹ Department of Mathematics, Faculty of Science, ZagazigUniversity, Zagazig, Egypt
yaser31270@yahoo.com

²Department of Mathematics, Faculty of Science, Taif University, Saudi Arabia
amr_mahdy85@yahoo.com

³Department of Mathematics, Faculty of Education Ibn-Al-Haitham, Baghdad University, Baghdad- Iraq
abdulah123s@yahoo.com

ABSTRACT

In this paper, we are presenting Hermite collocation method to solve numerically the Fredholm-Volterra-Hammerstein integral equations. We have clearly presented a theory to find ordinary derivatives. This method is based on replacement of the unknown function by truncated series of well known Hermite expansion of functions. The proposed method converts the equation to matrix equation which corresponding to system of algebraic equations with Hermite coefficients. Thus, by solving the matrix equation, Hermite coefficients are obtained. Some numerical examples are included to demonstrate the validity and applicability of the proposed technique.

Keywords

Fredholm-Hammerstein integral equations, Hermite polynomials, Volterra integral equation.

1. INTRODUCTION

The linear and non linear Fredholm and Volterra integral equation have been a growing interest in recent years ([21], [22]). This are an important branch of modern mathematics and arise frequently in many applied areas which include engineering, mechanics, physics, chemistry, astronomy and biology ([1], [5]). There are several methods for approximating the solution of linear, non-linear integral equations ([6]-[10], [13]). and solving fractional integro-differential equations ([11],[12]).

We consider the Hammerstein integral equations in the forms ([21],[22]):-

$$z(y) = f(y) + \lambda_1 \int_0^1 K_1(y, r)F(z(r))dr + \lambda_2 \int_0^y K_2(y, r)G(z(r))dr. (1)$$

Where $f(y)$, $K_1(y, r)$ and $K_2(y, r)$ are given functions, $0 \leq y, r \leq 1$, and λ_1, λ_2 are arbitrary constants.

Hermite polynomials are widely used in numerical computation. One of the advantages of using Hermite polynomials as a tool for expansion functions is the good representation of smooth functions by finite Hermite expansion provided that the function $z(y)$ is infinitely differentiable ([2], [15], [18], [20]). The Hermite collocation method in [16] solving convection diffusion equation, in [17] solving linear differential equations and in [19] solving linear complex differential equation.

The paper is organized as follows: Section 2, we will study some properties of the Hermite polynomials. In Section 3, we take his idea about an approximate formula of the integral derivative. In Section 4, procedure solution using the proposed numerical method. In Section 5, we give numerical implementation. In Section 6, the paper ends with a brief conclusion.

2. Some properties of the Hermite polynomials

Definition:-The Hermite polynomials are given by ([2], [20]):-

$$H_n(y) = (-1)^n e^{y^2} \frac{d^n}{dz^n} e^{-y^2}.$$

Some main properties of these polynomials are :

The Hermite polynomials evaluated at zero argument $H_n(0)$ and are called Hermite number as follows ([2], [20]):-

$$H_n(0) = \begin{cases} 0, & \text{if } n \text{ is odd,} \\ (-1)^{\frac{n}{2}} 2^{\frac{n}{2}} (n-1)!, & \text{if } n \text{ is even,} \end{cases} (2)$$

Where $(n-1)!$ is the factorial.

The polynomials $H_n(y)$ are orthogonal with respect to the weight function $\omega(y) = e^{-y^2}$ with the following condition [2]:-



$$\int_{-\infty}^{\infty} H_n(y)H_m(y)\omega(y)dy = \sqrt{\pi}2^n n! \delta_{nm}.$$

3. An approximate formula of the fractional derivative

The Hermite polynomials are defined on \mathbb{R} and can be determined with the aid of the following recurrence formula ([2], [14]):-

$$H_{n+1}(y) = 2yH_n(y) - 2nH_{n-1}(y), \quad H_0(y) = 1, \quad H_1(y) = 2y, \quad n = 1, 2, \dots$$

The analytic form of the Hermite polynomials of degree n is given by [2]

$$H_n(y) = n! \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \frac{(-1)^k 2^{n-2k}}{(k)!(n-2k)!} y^{n-2k}. \quad (3)$$

In consequence, for the p -th derivatives of Hermite polynomials the following relation hold:

$$H_n^{(p)}(y) = 2^p \frac{n!}{(n-p)!} H_{n-p}(y) = v_{n,p} H_{n-p}(y), \quad v_{n,p} = 2^p \frac{n!}{(n-p)!}. \quad (4)$$

The function $z(y) \in L^2_{\omega(y)}(\mathbb{R})$, may be expressed in terms of Hermite polynomials as follows

$$z(y) = \sum_{k=0}^{\infty} c_k H_k(y) \quad (5)$$

where the coefficients c_n are given by

$$c_n = \frac{1}{\sqrt{\pi}2^n n!} \int_{-\infty}^{\infty} z(y) H_n(y) \omega(y) dy, \quad n = 0, 1, \dots \quad (6)$$

In practice, only the first $(m + 1)$ –terms of Hermite polynomials are considered. Then we have [2]

$$z_m(y) = \sum_n^m c_n H_n(y). \quad (7)$$

The main approximate formula of the ordinary (integral) derivative is given in the following theorem.

Theorem 1. Let $z(y)$ be approximated by Hermite polynomials as (7) and also suppose $q > 0$, then

$$D^q(z_m(y)) \cong \sum_{n=q}^m \left[n! \sum_{k=q}^{\ell} c_n B_{n,k}^{(q)} y^{n-2k-q} \right], \quad (8)$$

where $\ell = \frac{n-q}{2}$ and $B_{n,k}^{(q)}$ is given by

$$B_{n,k}^{(q)} = \frac{(-1)^k 2^{n-2k}}{(k)!(n-2k-q)!}$$

Proof: since the non linear operation we have

$$D^q(z_m(y)) = \sum_{n=0}^m c_n D^q(H_n(y)). \quad (9)$$

It is clear that $D^q H_n(y) = 0$, $n = 0, 1, \dots, q - 1$, $q > 0$ Therefore, for $N = q, q + 1, \dots, m$.

Substituting equation (3) in (9) we have

$$D^q(z_m(y)) = \sum_{n=0}^m n! c_n \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \frac{(-1)^k 2^{n-2k}}{(k)!(n-2k)!} D^q y^{n-2k}.$$

$$D^q(z_m(y)) = \sum_{n=q}^m n! \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} c_n \frac{(-1)^k 2^{n-2k}}{(k)!(n-2k-q)!} y^{n-2k-q}. \quad (10)$$

4. Procedure solution using the proposed numerical method

We consider the Fredholm-Volterra integral equation (1), ([21], [22]).

We substitute the Eq. (7) into Eq. (1) we get:-



$$\sum_{n=0}^m c_n H_n(y) = f(y) + \lambda_1 \int_0^1 K_1(y, r) F(\sum_{n=0}^m c_n H_n(r)) dr + \lambda_2 \int_0^y K_2(y, r) G(\sum_{n=0}^m c_n H_n(r)) dr. \quad (11)$$

On the Hermite collocation points depended by:-

$$y_i = -1 + \frac{2}{m} i, \quad i = 0, 1, \dots, m \quad (12)$$

we collocate Eq.(11) with the points (12) to obtain

$$\sum_{n=0}^m c_n H_n(y_p) = f(y_j) + \lambda_1 \int_0^1 K_1(y_p, r) F(\sum_{n=0}^m c_n H_n(r)) dr + \lambda_2 \int_0^{y_p} K_2(y_p, r) G(\sum_{n=0}^m c_n H_n(r)) dr. \quad (13)$$

The integral terms in Eq. (13) can be found using composite trapezoidal integration technique as:

$$\int_0^1 K_1(y_p, r) F(z(r)) dr \cong \frac{h}{2} (\Omega_1(r_0) + \Omega_1(r_m) + 2 \sum_{k=1}^{m-1} \Omega_1(r_k)). \quad (14)$$

Where $\Omega_1(r) = K_1(y_j, r) F(z(r))$, $h = \frac{1}{m}$, for an arbitrary integer m , $r_i = ih$, $i = 0, 1, \dots, m$

And

$$\int_0^{y_p} K_2(y_p, r) G(z(r)) dr \cong \frac{h_p}{2} (\Omega_2(\bar{r}_0) + \Omega_2(\bar{r}_m) + 2 \sum_{k=1}^{m-1} \Omega_2(\bar{r}_k)). \quad (15)$$

Where $\Omega_2(r) = K_2(y_p, r) G(z(r))$, $h_p = \frac{y_p}{m}$, for an arbitrary integer m , $\bar{r}_i = ih$.

Eq. (13) gives $(m + 1)$ system of linear or non-linear algebraic equations, which can be solved for c_k , $k = 0, 1, \dots, m$. So the unknown function $z(y)$ can be found.

5. Numerical Implementation

In this section, to a chive the validity, the accuracy and support our theoretical discussion of the proposed method, we give some computational results of numerical examples.

Example1. Consider Eq. (1) with the following functions and coefficients ([21], [22]).

$$f(y) = y^3 - (6 - 2e)e^y, \quad \lambda_1 = 1, \quad \lambda_2 = 1, \\ K_1(y, r) = e^{(y+r)}, \quad K_2(y, r) = 0, \quad F(z(r)) = z(r), \quad G(z(r)) = 0.$$

Eq. (1) takes following the form

$$z(y) = y^3 - (6 - 2e)e^y + \int_0^1 e^{(y+r)} z(r) dr. \quad (16)$$

We apply the suggested method with $m = 3$, and approximate the solution $z(y)$ as follows

$$z_m(y) = \sum_{n=0}^3 c_n H_n(y). \quad (17)$$

By the same procedure in the previous section and using Eq. (13) we have

$$\sum_{n=0}^m c_n H_n(y) = y^3 - (6 - 2e)e^y + \frac{h}{2} \left(\Omega_1(r_0) + \Omega_1(r_m) + 2 \sum_{k=1}^{m-1} \Omega_1(r_k) \right), \\ \sum_{n=0}^3 c_n H_n(y_p) - (y_p^3 - (6 - 2e)e^{y_p}) - \frac{h}{2} (\Omega_1(r_0) + \Omega_1(r_3) + 2 \sum_{k=1}^2 \Omega_1(r_k)) = 0, \quad (18) \\ p = 0, 1, 2.$$

Where $\Omega_1(r) = e^{(r+y_p)} \sum_{n=0}^3 c_n H_n(y_p)$ and the nodes $r_{\ell+1} = r_\ell + h$, $\ell = 0, 1, \dots, m$. $r_0 = 0$ and $h = \frac{1}{m}$.

Eq. (18) represents linear system of $m + 1$ algebraic equations in the coefficients c_i , by solving it using the conjugate gradient method.

The exact solution of this example is $z(y) = y^3$ ([21], [22]).




Fig. 1. The behavior of the exact solution and the approximate solution at $m = 3$.

The behavior of the approximate solution using the proposed method with $m = 3$ and the exact solution are presented in Figures, 1-4. From this Figures 1-4, it is clear that the proposed method can be considered as an efficient method to solve the linear integral equations ([21], [22]).

Example 2. Consider Eq. (1) with the following functions and coefficients ([21], [22]).

$$f(y) = 2ye^y - e^y + 1, \quad \lambda_1 = 1, \quad \lambda_2 = -1,$$

$$K_1(y, r) = 0, \quad K_2(y, r) = (r + y), \quad F(z(r)) = 0, \quad G(z(r)) = e^{z(r)}.$$

Eq. (1) takes following the form

$$z(y) = 2ye^y - e^y + 1 - \int_0^y (r + y)e^{z(r)} dr. \quad (19)$$

We apply the suggested method with $m = 3$, and approximate the solution $z(y)$ as follows

$$z_m(y) = \sum_{n=0}^3 c_n H_n(y). \quad (20)$$

By the same procedure in the previous section and using Eq. (13) we have

$$\sum_{n=0}^m c_n H_n(y) = 2ye^y - e^y + 1 - \frac{h}{2} \left(\Omega_2(\bar{r}_0) + \Omega_2(\bar{r}_m) + 2 \sum_{k=1}^{m-1} \Omega_2(\bar{r}_k) \right),$$

$$\sum_{n=0}^3 c_n H_n(y_p) - (2y_p e^{y_p} - e^{y_p} + 1) + \frac{h}{2} (\Omega_2(\bar{r}_0) + \Omega_2(\bar{r}_3) + 2 \sum_{k=1}^2 \Omega_2(\bar{r}_k)) = 0, \quad (21)$$

$$p = 0, 1, 2.$$

Where the nodes $r_{\ell+1} = r_\ell + h, \ell = 0, 1, \dots, m, r_0 = 0$ and $h_p = \frac{y_p}{m}, \Omega_2(r)(r + y_p)e^{z(r)}$.

Eq. (21) presents non-linear system of $m + 1$ algebraic equations in the coefficients c_i . by solving it by using the Newton iteration method with suitable initial solution.

The exact solution of this problem is $z(y) = y$ ([21], [22]).




Fig. 2. The behavior of the exact solution and the approximate solution at $m = 3$.

Example 3. Consider Eq. (1) with the following functions and coefficients ([21], [22]).

$$f(y) = y^e + 1, \quad \lambda_1 = -1, \quad \lambda_2 = 1,$$

$$K_1(y, r) = r + y, \quad K_2(y, r) = 0, \quad F(z(r)) = e^{z(r)}, \quad G(z(r)) = 0.$$

Eq. (1) takes following the form

$$z(y) = y^e + 1 - \int_0^1 (r + y) e^{z(r)} dr. \quad (22)$$

We apply the suggested method with $m = 3$, and approximate the solution $z(y)$ as follows

$$z_m(y) = \sum_{n=0}^3 c_n H_n(y). \quad (23)$$

By the same procedure in the previous section and using Eq. (13) we have

$$\sum_{n=0}^m c_n H_n(y) = y^e + 1 - \frac{h}{2} \left(\Omega_1(r_0) + \Omega_1(r_m) + 2 \sum_{k=1}^{m-1} \Omega_1(r_k) \right),$$

$$\sum_{n=0}^3 c_n H_n(y_p) - (y^e + 1) + \frac{h}{2} (\Omega_1(r_0) + \Omega_1(r_3) + 2 \sum_{k=1}^2 \Omega_1(r_k)) = 0, \quad (24)$$

$$p = 0, 1, 2.$$

Where the nodes $r_{\ell+1} = r_\ell + h, \ell = 0, 1, \dots, m, r_0 = 0$ and $h = \frac{1}{m}, \Omega_1(r) = (r + y) e^{z(r)}$.

Eq. (24) presents non-linear system of algebraic equations by solving it using the well known Newton iteration method with suitable initial solution.

The exact solution of this problem is $z(y) = y$ ([21], [22]).




Fig. 3. The behavior of the exact solution and the approximate solution at $m = 3$.

Example 4. Consider Eq. (1) with the following functions and coefficients ([21], [22]).

$$f(y) = \frac{y}{2} - \frac{y^4}{12} - \frac{1}{3}, \quad \lambda_1 = 1, \quad \lambda_2 = -1,$$

$$K_1(y, r) = r + y, \quad K_2(y, r) = r - y, \quad F(z(r)) = z(r), \quad G(z(r)) = z^2(r).$$

Eq. (1) takes the following form

$$z(y) = \frac{y}{2} - \frac{y^4}{12} - \frac{1}{3} + \int_0^1 (r + y)z(r) dr + \int_0^y (r - y)z^2(r)dr. \quad (25)$$

We apply the suggested method with $m = 3$, and approximate the solution $z(y)$ as follows

$$z_m(y) = \sum_{n=0}^3 c_n H_n(y). \quad (26)$$

By the same procedure in the previous section and using Eq. (13) we have

$$\begin{aligned} \sum_{n=0}^m c_n H_n(y) &= \frac{y}{2} - \frac{y^4}{12} - \frac{1}{3} + \frac{h}{2} \left(\Omega_1(r_0) + \Omega_1(r_m) + 2 \sum_{k=1}^{m-1} \Omega_1(r_k) \right) \\ &\quad - \frac{h}{2} \left(\Omega_2(\bar{r}_0) + \Omega_2(\bar{r}_m) + 2 \sum_{k=1}^{m-1} \Omega_2(\bar{r}_k) \right), \\ \sum_{n=0}^3 c_n H_n(y_p) - \left(\frac{y}{2} - \frac{y^4}{12} - \frac{1}{3} \right) - \frac{h}{2} (\Omega_1(r_0) + \Omega_1(r_3) + 2 \sum_{k=1}^2 \Omega_1(r_k)) \\ &\quad + \frac{h}{2} (\Omega_2(\bar{r}_0) + \Omega_2(\bar{r}_3) + 2 \sum_{k=1}^2 \Omega_2(\bar{r}_k)) = 0, \quad p = 0, 1, 2. \end{aligned} \quad (27)$$

Where the nodes $r_{\ell+1} = r_\ell + h$, $\bar{r}_{\ell+1} = \bar{r}_\ell + h$, $\ell = 0, 1, \dots, m$, $r_0 = \bar{r}_0 = 0$ and $h = \frac{1}{m}$, $h_p = \frac{y_p}{m}$ and $\Omega_1(r) = (r + y)z(r)$, $\Omega_2(r) = (y_p - r)z^2(r)$.

Eq. (27) presents non-linear system of $m + 1$ algebraic equations by solving it using Newton iteration method.

The exact solution of this problem is $z(y) = y$ ([21], [22]).




Fig. 4. The behavior of the exact solution and the approximate solution at $m = 3$.

6. Conclusion

In this articles, we approximate method for the solution of linear and non-linear Fredholm-Volterra integral equations using Hermite collocation method. A comparison of the exact solution reveals that the presented method is very effective and convenient. The numerical results show that the accuracy improves with increasing m , hence for better results, using number m is recommended. Also, from the obtained approximate solution, we can conclude that the proposed method gives the solution in an excellent agreement with the exact solution. All computations are done using Maple programming.

References

- [1] R. P. Agarwal, Boundary value problems for higher order integro-differential equations, *Nonlinear Anal. Theory Methods Appl*, 9 (1983) 259-270.
- [2] L.C. Andrews, *Special Functions For Engineers and Applied Mathematicians*, Macmillan publishing company, New York, (1985).
- [3] E. Babolian, F. Fattahzadeh and E. GolparRaboky, A Chebyshev approximation for solving nonlinear integral equations of Hammerstein type, *Applied Mathematics and Computation*, 189 (2007) 641-46.
- [4] A. K. Borzabadi, A.V. Kamyad and H. H. Mehne, A different approach for solving the nonlinear Fredholm integral equations of the second kind, *Applied Mathematics and Computation*, 173 (2006) 724-735.
- [5] L.M. Delves and J.L. Mohamad, *Computational Methods for Integral Equations*, Cambridge University press, (1985).
- [6] M. M. Khader, Introducing an efficient modification of the homotopy perturbation method by using Chebyshev polynomials, *Arab Journal of Mathematical Sciences*, 18 (2012) 61-71.
- [7] S.T. Mohamed and M. M. Khader, Numerical solutions to the second order Fredholm integro-differential equations using the spline functions expansion, *Global Journal of Pure and Applied Mathematics*, 34 (2011) 21-29.
- [8] N. Kurt and M. Sezer, Polynomial solution of high-order linear Fredholm integro-differential equations with constant coefficients, *Journal of the Franklin Institute*, 345 (2008) 839-850.
- [9] M. Shahrezaee, Solving an integro-differential equation by Legendre polynomials and Block-pulse functions, *Dynamical Systems and Applications*, (2004) 642-647.
- [10] N. H. Sweilam, Fourth order integro-differential equations using variational iteration method, *Comput. Maths. Appl.*, 54 (2007) 1086-1091.
- [11] N. H. Sweilam, M. M. Khader and R. F. Al-Bar, Homotopy perturbation method for linear and nonlinear system of fractional integro-differential equations, *International Journal of Computational Mathematics and Numerical Simulation*, 1(2008) 73-87.
- [12] N. H. Sweilam and M. M. Khader, A Chebyshev pseudo-spectral method for solving fractional order integro-differential equations, *ANZIAM*, 51 (2010) 464-475
- [13] S. Yousefi and M. Razzaghi, Legendre wavelet method for the nonlinear Volterra-Fredholm integral equations,



Math. Comp. Simul., 70 (2005) 1-8.

- [14] D. Funaro, Polynomial Approximations of Differential Equation, Springer-Verlag, 1992.
- [15] W. R. Dyksen and R. E. Lynch, A new decoupling technique for the Hermite cubic collocation equations arising from boundary value problems, Math. Comput. Simul. 54, p. (359-372), 2000.
- [16] S. H. Brill, Analytic solution of Hermite collocation discretization of the steady state convection diffusion equation, Inter J. Differential Equation and Application, 4, p. (141-155), 2002.
- [17] Z. K. Bojdi, S. Ahmadi-Asl and A. Aminataei, Operational matrices with respect to Hermite polynomials and their applications in solving linear differential equations with variable coefficients, Journal of Linear and Topological Algebra, 2(2) p.(91-103), 2013.
- [18] B. Bialecki, A fast domain decomposition poisson solver on a rectangle for Hermitebicubic orthogonal spline collocation, Siam Journal Numerical Analysis, 30, p.(425-434), 1993.
- [19] M. Bagherpoorfard and F. A. Ghassabzade, Hermite matrix polynomial collocation method for linear complex differential equation and some comparisons, Journal of Applied Mathematics and Physics, 1, p.(58-64), 2013.
- [20] E. M. Solouma and M. M. Khader Analytical and numerical simulation for solving the system of non-linear fractional dynamical model of marriage International Mathematical Forum January 2016.
- [21] N. H. Sweilam, M. M. Khader, and W. Y. Kota On the Numerical Solution of Hammerstein Integral Equations using Legendre Approximation International Journal of Applied Mathematical Research, 1(1) (212) 65-76.
- [22] A. M. S. Mahdy, E. M. H. Mohamed and G. M. A. Marai, Numerical solution of fractional integro-differential equation by least squares method and shifted chebyshev polynomials of the third Kind method, The oretical Mathematics and Applications, 6(4), (2016), 87-101.