



Attracting sets of nonlinear difference equations with time-varying delays

Danhua He

Department of Mathematics, Zhejiang International Studies University, Hangzhou, 310023, PR China

danhuahe@126.com

Abstract

In this paper, a class of nonlinear difference equations with time-varying delays is considered. Based on a generalized discrete Halanay inequality, some sufficient conditions for the attracting set and the global asymptotic stability of the nonlinear difference equations with time-varying delays are obtained.

Keywords: Halanay inequality, Nonlinear difference equations, Attracting set, Global asymptotic stability.

Subject Classification: Mathematics (2010):39A22, 39A30.

Subject Classification: Image Processing

Language: English

Date of Publication: 20-09-2018

DOI: 10.24297/jam.v14i2.7668

ISSN: 2347 - 1921

Volume: 14 Issue: 02

Journal: Journal Of Advances In Mathematics

Website: <https://cirworld.com>



This work is licensed under a Creative Commons Attribution 4.0 International License.



1 Introduction

Delay difference equations are an important class of discrete-time dynamical systems whose future states depend on not only the present states, but also the past states. In the past few decades, delay difference equations have attracted considerable research interest because these equations play an essential role in discrete analogues and numerical solutions of delay differential equations. A massive literature on the stability analysis of delay difference equations is available [1-12]. It should be noticed that the equilibrium point sometimes does not exist in many real physical systems, especially in nonlinear delay difference equations. Therefore, it is more interesting for nonlinear delay difference equations to study the attracting set than to study the stability. However, not much has been developed in the study of attracting sets for the nonlinear difference equations with time-varying delays. Motivated by the above discussions, the main aim of this paper is to study the attracting set of the nonlinear difference equations with time-varying delays. Based on a generalized discrete Halanay inequality, some sufficient conditions for the attracting set and the global asymptotic stability of the nonlinear difference equations with time-varying delays are obtained, with no subheadings.

2 Model description and preliminaries

Let \mathbb{R} , \mathbb{R}^+ and \mathbb{R}_0^+ denote the set of real numbers, positive real numbers, nonnegative real numbers, respectively. Let $\|\bullet\|$ denote the Euclidean norm, \mathbb{Z} and \mathbb{Z}^+ represent the set of integers and positive integers, respectively. $\mathbb{Z}^r = \{z \in \mathbb{Z} : z \geq -r\}$. For simplicity, we denote $x(m)$ and $x_j(m)$ by x_m and $x_{j,m}$, respectively. For a sequence of real number $\{x_m\}$, the difference operator Δ on x_m is defined as $\Delta x_m = x_{m+1} - x_m$.

Consider the following nonlinear difference equations with time-varying delays:

$$\begin{cases} x_i(m+1) = c_i x_i(m) + \sum_{j=1}^n a_{ij} f_j(x_j(m)) + \sum_{j=1}^n b_{ij} f_j(m, x_{j,m}, x_{j,m-h_1}, \dots, x_{j,m-h_r}) + I_i, m \in \mathbb{Z}^+ \\ x_i(m) = \phi_i(m), m = -h_r, -h_{r+1}, \dots, 0, i = 1, 2, \dots, n, \end{cases} \quad (1)$$

where c_i, a_{ij}, b_{ij}, I_i ($i, j = 1, 2, \dots, n$) are real constants, $0 = h_0 < h_1 < \dots < h_r$.

For convenience, we shall rewrite (1) in the vector form:

$$\begin{cases} x_{m+1} = Cx_m + Af(x_m) + Bg(m, x_m, x_{m-h_1}, \dots, x_{m-h_r}) + I, m \in \mathbb{Z}^+ \\ x_m = \phi_m, m = -h_r, -h_{r+1}, \dots, 0, i = 1, 2, \dots, n, \end{cases} \quad (2)$$

where $x_m = (x_{1,m}, x_{2,m}, \dots, x_{n,m})^T$, $C = \text{diag}\{c_1, \dots, c_n\}$, $A = (a_{ij})_{n \times n}$, $B = (b_{ij})_{n \times n}$, $I = (I_1, I_2, \dots, I_n)^T$, $\phi_m = (\phi_{1,m}, \dots, \phi_{n,m})^T$. Throughout this paper, we assume that Eq. (2) has at least one solution x_m for any given initial function ϕ .

Definition 2.1. The set $S \subset \mathbb{R}^n$ is called a global attracting set of (2), if for any initial function ϕ , the solution x_m satisfies $\text{dist}(x_m, S) \rightarrow 0$ as $m \rightarrow \infty$, where $\text{dist}(\phi, S) = \inf_{\varphi \in S} \rho(\phi, \varphi)$ for $\varphi \in \mathbb{R}^n$, $\rho(\bullet, \bullet)$ is any distance in \mathbb{R}^n .



Lemma 2.1. [13] For any $\varepsilon > 0$, $x, y \in \mathbb{R}^n$,

$$2X^T Y \leq \varepsilon X^T X + \varepsilon^{-1} Y^T Y.$$

Lemma 2.2. [8] Let $q_i, \gamma \in \mathbb{R}_0^+$, $h_i \in \mathbb{Z}^+$, $i = 1, \dots, r$; where $0 = h_0 < h_1 < \dots < h_r$ and $\sum_{i=0}^r q_i < p \leq 1$, and let

$\{x_n\}_{n \in \mathbb{Z}^{-h_r}}$ be a sequence of real numbers satisfying the inequality

$$\Delta x_n \leq -p x_n + \sum_{i=0}^r q_i x_{n-h_i} + \gamma, \quad n \in \mathbb{Z}^0. \quad (3)$$

Then there exists $\lambda_0 \in (0, 1)$ such that

$$x_n \leq \max \{0, x_0, x_{-h_1}, \dots, x_{-h_r}\} \lambda_0^n + \Lambda, \quad n \in \mathbb{Z}^0, \quad (4)$$

where $\Lambda = (p - \sum_{i=0}^r q_i)^{-1} \gamma > 0$. Moreover, λ_0 may be chosen as the smallest root of the polynomial

$$P(\lambda) = \lambda^{h_{r+1}} - (1 - p + q_0) \lambda^{h_r} - q_1 \lambda^{h_r - h_1} - \dots - q_{r-1} \lambda^{h_r - h_{r-1}} - q_r, \quad (5)$$

which lies in the interval $(0, 1)$.

3 Main results

In this section, we will obtain several sufficient conditions for the attracting set of (2) by Lemma 2.2.

The following assumptions are needed for our discussion.

(A₁) For any $x_m \in \mathbb{R}^n$, there exists positive definite symmetric matrix J such that

$$f^T(x_m) f(x_m) \leq x_m^T J x_m. \quad (6)$$

(A₂) There exist positive definite symmetric matrices L_i such that

$$g^T(m, x_m, x_{m-h_1}, \dots, x_{m-h_r}) g(m, x_m, x_{m-h_1}, \dots, x_{m-h_r}) \leq \sum_{i=0}^r x_{m-h_i}^T L_i x_{m-h_i}, \quad (7)$$

for all $(m, x_m, x_{m-h_1}, \dots, x_{m-h_r}) \in \mathbb{Z}^0 \times \mathbb{R}^{r+1}$.

Theorem 3.1 Suppose that (A₁) and (A₂) hold. If there exists a positive definite symmetric matrix P such that $\sum_{i=0}^r \beta_i < -(\gamma_1 + \gamma_2) \leq 1$, then every solution $x(m)$ of (2) satisfies

$$\|x_m\| \leq \max \{0, \|x_0\|, \|x_{-h_1}\|, \dots, \|x_{-h_r}\|\} (\sqrt{\lambda_0})^n + ((-\gamma_1 + \gamma_2) - \sum_{i=0}^r \beta_i)^{-1} \gamma_3)^{\frac{1}{2}},$$



where

$$\gamma_1 = \frac{\lambda_{\max}(C^T PC + \varepsilon_1^{-1} C^T PAA^T PC + \varepsilon_2^{-1} C^T PBB^T PC + \varepsilon_3^{-1} C^T PPC + \varepsilon_1 J)}{\lambda_{\min}(P)}$$

$$\gamma_2 = \frac{\lambda_{\max}(A^T PA + \varepsilon_4^{-1} A^T PPA + \varepsilon_5^{-1} A^T PBB^T PA) \lambda_{\max}(J) - \lambda_{\min}(P)}{\lambda_{\min}(P)}$$

$$\beta_i = \frac{[\varepsilon_2 + \varepsilon_5 + \lambda_{\max}(B^T PB + \varepsilon_6^{-1} B^T PPB)] \lambda_{\max}(L_i)}{\lambda_{\min}(P)}$$

$$\gamma_3 = \frac{(\varepsilon_3 + \varepsilon_4 + \varepsilon_6) I^T I + I^T P I}{\lambda_{\min}(P)}$$

and the constant $\lambda_0 \in (0,1)$ may be chosen as the smallest root of the polynomial

$$P(\lambda) = \lambda^{h_r+1} - (1 + (\gamma_1 + \gamma_2) + \beta_0) \lambda^{h_r} - \beta_1 \lambda^{h_r-h_1} - \dots - \beta_{r-1} \lambda^{h_r-h_{r-1}} - \beta_r. \quad (8)$$

As a consequence,

$$S = \left\{ \phi \in \mathbb{R}^n \mid \|\phi\| \leq \left((-\gamma_1 - \gamma_2) - \sum_{i=0}^r \beta_i \right)^{-1} \gamma_3 \right\}^{\frac{1}{2}} \quad (9)$$

is a positive attracting set of (2).

Proof. Choosing the Lyapunov functional candidate

$$V(x) \leq x^T P x.$$

Obviously, we have

$$\lambda_{\min}(P) \|x\| \leq V(x) \leq \lambda_{\max}(P) \|x\|. \quad (10)$$

Then, it follows that

$$\begin{aligned} V(x_{m+1}) &= x_{m+1}^T P x_{m+1} \\ &= [C x_m + A f(x_m) + B g(m, x_m, x_{m-h_1}, \dots, x_{m-h_r}) + I]^T P [C x_m \\ &\quad + A f(x_m) + B g(m, x_m, x_{m-h_1}, \dots, x_{m-h_r}) + I] \\ &= x_m^T C^T P C x_m + x_m^T C^T P A f(x_m \end{aligned}$$



$$\begin{aligned}
& + x_m^T C^T P B g(m, x_m, x_{m-h_1}, \dots, x_{m-h_r}) + x_m^T C^T P I + f^T(x_m) A^T P C x_m \\
& + f^T(x_m) A^T P A f(x_m) + f^T(x_m) A^T P B g(m, x_m, x_{m-h_1}, \dots, x_{m-h_r}) \\
& + f^T(x_m) A^T P I + g^T(m, x_m, x_{m-h_1}, \dots, x_{m-h_r}) B^T P C x_m \\
& + g^T(m, x_m, x_{m-h_1}, \dots, x_{m-h_r}) B^T P A f(x_m) \\
& + g^T(m, x_m, x_{m-h_1}, \dots, x_{m-h_r}) B^T P B g(m, x_m, x_{m-h_1}, \dots, x_{m-h_r}) \\
& + g^T(m, x_m, x_{m-h_1}, \dots, x_{m-h_r}) B^T P I + I^T P C x_m + I^T P A f(x_m) \\
& + I^T P B g(m, x_m, x_{m-h_1}, \dots, x_{m-h_r}) + I^T P I.
\end{aligned} \tag{11}$$

From (11) and Lemma 2.1, we have

$$\begin{aligned}
V(x_{m+1}) & = x_m^T [C^T P C + \varepsilon_1^{-1} C^T P A A^T P C + \varepsilon_2^{-1} C^T P B B^T P C + \varepsilon_3^{-1} C^T P P C] x_m \\
& + \varepsilon_1 f^T(x_m) f(x_m) + \varepsilon_2 g^T(m, x_m, x_{m-h_1}, \dots, x_{m-h_r}) g(m, x_m, x_{m-h_1}, \dots, x_{m-h_r}) \\
& + f^T(x_m) [A^T P A + \varepsilon_4^{-1} A^T P P A + \varepsilon_5^{-1} A^T P B B^T P A] f(x_m) \\
& + \varepsilon_5 g^T(m, x_m, x_{m-h_1}, \dots, x_{m-h_r}) g(m, x_m, x_{m-h_1}, \dots, x_{m-h_r}) \\
& + g^T(m, x_m, x_{m-h_1}, \dots, x_{m-h_r}) [B^T P B + \varepsilon_6^{-1} B^T P P B] g(m, x_m, x_{m-h_1}, \dots, x_{m-h_r}) \\
& + (\varepsilon_3 + \varepsilon_4 + \varepsilon_6) I^T I + I^T P I
\end{aligned} \tag{12}$$

Utilizing (12) and Assumptions (A₁) and (A₂), we obtain

$$\begin{aligned}
V(x_{m+1}) & = x_m^T [C^T P C + \varepsilon_1^{-1} C^T P A A^T P C + \varepsilon_2^{-1} C^T P B B^T P C + \varepsilon_3^{-1} C^T P P C + \varepsilon_1 J] x_m \\
& + \lambda_{\max}(A^T P A + \varepsilon_4^{-1} A^T P P A + \varepsilon_5^{-1} A^T P B B^T P A) x_m^T J x_m \\
& + [\varepsilon_2 + \varepsilon_5 + \lambda_{\max}(B^T P B + \varepsilon_6^{-1} B^T P P B)] \left(\sum_{i=0}^r x_{m-h_i}^T L_i x_{m-h_i} \right) \\
& + (\varepsilon_3 + \varepsilon_4 + \varepsilon_6) I^T I + I^T P I.
\end{aligned} \tag{13}$$

From (10) and (13) we have

$$\begin{aligned}
\lambda_{\min}(P) \|x_{m+1}\|^2 & \leq V(x_{m+1}) \\
& = \lambda_{\max}(C^T P C + \varepsilon_1^{-1} C^T P A A^T P C + \varepsilon_2^{-1} C^T P B B^T P C + \varepsilon_3^{-1} C^T P P C + \varepsilon_1 J) \|x_m\|^2 \\
& + \lambda_{\max}(A^T P A + \varepsilon_4^{-1} A^T P P A + \varepsilon_5^{-1} A^T P B B^T P A) \lambda_{\max}(J) \|x_m\|^2 \\
& + [\varepsilon_2 + \varepsilon_5 + \lambda_{\max}(B^T P B + \varepsilon_6^{-1} B^T P P B)] \left(\sum_{i=0}^r \lambda_{\max}(L_i) \|x_{m-h_i}\|^2 \right) \\
& + (\varepsilon_3 + \varepsilon_4 + \varepsilon_6) I^T I + I^T P I.
\end{aligned} \tag{14}$$

Let $y_m = \|x_m\|^2$, we have



$$\Delta y_m \leq (\gamma_1 + \gamma_2) y_m + \left(\sum_{i=0}^r \beta_i y_{m-h_i} \right) + \gamma_3. \quad (15)$$

Therefore, by Lemma 2.2, we obtain

$$y_m \leq \max \left\{ 0, y_0, y_{-h_1}, \dots, y_{-h_r} \right\} \lambda_0^n + \left(-(\gamma_1 + \gamma_2) - \sum_{i=0}^r \beta_i \right)^{-1} \gamma_3 \quad (16)$$

This means

$$\|x_m\| \leq \max \left\{ 0, \|x_0\|, \|x_{-h_1}\|, \dots, \|x_{-h_r}\| \right\} (\sqrt{\lambda_0})^n + \left(-(\gamma_1 + \gamma_2) - \sum_{i=0}^r \beta_i \right)^{-1} \gamma_3^{\frac{1}{2}}, \quad (17)$$

where $\lambda_0 \in (0,1)$ may be chosen as the smallest root of the polynomial (8). The proof is completed.

Corollary 3.1 Suppose that (A_1) and (A_2) hold. If exists a positive definite symmetric matrix P such that $\sum_{i=0}^r \beta_i < -(\gamma_1 + \gamma_2) \leq 1$, then the equation (2) with $I = 0$ is globally exponentially stable with the exponential convergence rate $r = \sqrt{\lambda_0}$, where $\lambda_0 \in (0,1)$ may be chosen as the smallest root of the polynomial (8).

Proof. Substituting $I = 0$ into γ_3 yields $\gamma_3 = 0$. Therefore, from (17) we have

$$\|x_m\| \leq \max \left\{ 0, \|x_0\|, \|x_{-h_1}\|, \dots, \|x_{-h_r}\| \right\} (\sqrt{\lambda_0})^n, \quad (18)$$

where $\lambda_0 \in (0,1)$ may be chosen as the smallest root of the polynomial (8). The proof is completed.

Let $\varepsilon_i (i = 1, 2, \dots, 6) = 1$ and P be the n -dimensional unit matrix, then from Theorem 3.1 and Corollary 3.1 we obtain the following Corollary 3.2 and Corollary 3.3, respectively.

Corollary 3.2 Suppose that (A_1) and (A_2) hold. If there exists a positive definite symmetric matrix P such that $\sum_{i=0}^r \hat{\beta}_i < -(\hat{\gamma}_1 + \hat{\gamma}_2) \leq 1$, then every solution $x(m)$ of (2) satisfies

$$\|x_m\| \leq \max \left\{ 0, \|x_0\|, \|x_{-h_1}\|, \dots, \|x_{-h_r}\| \right\} (\sqrt{\lambda_0})^n + \left(-(\hat{\gamma}_1 + \hat{\gamma}_2) - \sum_{i=0}^r \hat{\beta}_i \right)^{-1} \hat{\gamma}_3^{\frac{1}{2}},$$

where

$$\hat{\gamma}_1 = \lambda_{\max} (C^T P C + \varepsilon_1^{-1} C^T P A A^T P C + \varepsilon_2^{-1} C^T P B B^T P C + \varepsilon_3^{-1} C^T P P C + \varepsilon_1 J$$



$$\hat{\gamma}_2 = \lambda_{\max} (A^T P A + \varepsilon_4^{-1} A^T P P A + \varepsilon_5^{-1} A^T P B B^T P A) \lambda_{\max} (J) - 1$$

$$\hat{\beta}_i = [\varepsilon_2 + \varepsilon_5 + \lambda_{\max} (B^T P B + \varepsilon_6^{-1} B^T P P B)] \lambda_{\max} (L_i)$$

$$\hat{\gamma}_3 = (\varepsilon_3 + \varepsilon_4 + \varepsilon_6) I^T I + I^T P I$$

and the constant $\lambda_0 \in (0,1)$ may be chosen as the smallest root of the polynomial

$$P(\lambda) = \lambda^{h_{r+1}} - (1 + (\hat{\gamma}_1 + \hat{\gamma}_2) + \hat{\beta}_0) \lambda^{h_r} - \hat{\beta}_1 \lambda^{h_r - h_1} - \dots - \hat{\beta}_{r-1} \lambda^{h_r - h_{r-1}} - \hat{\beta}_r. \quad (19)$$

As a consequence,

$$S = \left\{ \phi \in \mathbb{R}^n \mid \|\phi\| \leq \left((-\hat{\gamma}_1 - \hat{\gamma}_2) - \sum_{i=0}^r \hat{\beta}_i \right)^{-1} \hat{\gamma}_3^{\frac{1}{2}} \right\} \quad (20)$$

is a positive attracting set of (2).

Corollary 3.3 Suppose that (A_1) and (A_2) hold. If exists a positive definite symmetric matrix P such that $\sum_{i=0}^r \hat{\beta}_i < -(\hat{\gamma}_1 + \hat{\gamma}_2) \leq 1$, then the equation (2) with $I = 0$ is globally exponentially stable with the exponential convergence rate $r = \sqrt{\lambda_0}$, where $\lambda_0 \in (0,1)$ may be chosen as the smallest root of the polynomial (19).

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Funding Statement

The work is supported by the National Natural Science Foundation of China under Grants 11501518 and the Zhejiang Natural Science Foundation of China under Grants LQ16A010004.

References

- [1] J. Cermak, J. Jansky, Stability switches in linear delay difference equations, *Applied Mathematics and Computation* 243 (2014) 755-766.
- [2] J. Cermak, J. Jansky, Explicit stability conditions for a linear trinomial delay difference equation, *Applied Mathematics Letters* 43 (2015) 56-60.
- [3] N.H. Sau, P. Niamsup, V.N. Phat, Positivity and stability analysis for linear implicit difference delay equations, *Linear Algebra and its Applications* 510 (2016) 25-41.
- [4] M.M. Kipnis, R.M. Nigmatullin, Stability of the trinomial linear difference equations with two delays, *Autom. Remote Control* 65 (11) (2004) 1710-1723.



- [5] E. Kaslik, Stability results for a class of difference systems with delay, *Adv. Differ. Equ.* 2009 (2009). Article ID 938492, 13 pp
- [6] M.M. Kipnis, R.M. Nigmatullin, Stability of some difference equations with two delays, *Autom. Remote Control* 64 (5) (2003) 782-790.
- [7] D. Melchor-Aguilar, Exponential stability of linear continuous time difference systems with multiple delays, *Systems & Control Letters* 62 (2013) 811-818.
- [8] L. Xu, Generalized discrete Halanay inequalities and the asymptotic behavior of nonlinear discrete systems, *Bulletin of the Korean Mathematical Society* 50(5)(2013)1555-1565.
- [9] R.P. Agarwal, Y.H. Kim, S.K. Sen, New discrete Halanay inequalities: stability of difference equations, *Commun. Appl. Anal.* 12 (2008) 83--90.
- [10] A. Yu. Aleksandrov, E.B. Aleksandrova, Delay-independent stability conditions for a class of nonlinear difference systems, *Journal of the Franklin Institute* 355 (2018) 3367-3380.
- [11] E. Liz, J.B. Ferreiro, A Note on the global stability of generalized difference equations, *Appl. Math. Lett.* 15 (2002) 655--659.
- [12] Z.G. Yang, D.Y. Xu, Mean square exponential stability of impulsive stochastic difference equations, *Appl. Math. Lett.* 20 (2007) 938--945.
- [13] Q. Song, Z. Zhao, Global dissipativity of neural networks with both variable and unbounded delays, *Chaos, Solitons and Fractals* 25 (2005) 393-401.
- [14] L. Huang, *Linear Algebra in Systems and Control Theory*. Beijing, China: Sci. Press, 1984.