

Malacosporean myxozoans exploit a diversity of fish hosts

Journal:	<i>Parasitology</i>
Manuscript ID	PAR-2018-0350.R1
Manuscript Type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Naldoni, Juliana; Universidade Federal de Sao Paulo - Campus Diadema, Departamento de Ecologia e Biologia Evolutiva Adriano, Edson; Universidade Federal de São Paulo, Ciências Biológicas ; Universidade Estadual de Campinas, Biologia animal Hartigan, Ashlie; Natural History Museum, Life Sciences; Kings College London, Sayer, Carl; University College London, Department of Geography Okamura, B; Natural History Museum, Zoology
Key Words:	Buddenbrockia, Tetracapsuloides, SSU rDNA sequence, ultrastructure, sporogenesis, Myxozoa, Cnidaria

SCHOLARONE™
Manuscripts

1 **Malacosporean myxozoans exploit a diversity of fish hosts**

2 Juliana Naldoni^{1,2*}, Edson A. Adriano^{1,3}, Ashlie Hartigan², Carl Sayer⁴ and Beth Okamura²

3

4 ¹Departamento de Ciências Biológicas, Universidade Federal de São Paulo (UNIFESP),

5 Diadema, SP, Brazil

6 ²Department of Life Sciences, Natural History Museum, Cromwell Road, London SW7 5BD,

7 United Kingdom

8 ³Departamento de Biologia Animal, Instituto de Biologia, Universidade Estadual de Campinas,

9 Campinas, SP, Brazil

10 ⁴Environmental Change Research Centre, Department of Geography, University College London,

11 Gower Street, London, WC1E 6BT, United Kingdom

12

13 * Corresponding author: Departamento de Ciências Biológicas, Universidade Federal de São

14 Paulo (UNIFESP), Rua Professor Arthur Riedel, 275, Jardim Eldorado, 09972-270 Diadema, SP,

15 Brazil. E-mail: jnaldoni@gmail.com

16

17

18

19

20

21

22 **Abstract**

23 Myxozoans are widespread and common endoparasites of fish with complex life cycles, infecting
24 vertebrate and invertebrate hosts. There are two classes: Myxosporidia and Malacosporea. To date
25 some 2,500 myxosporidian species have been described. By comparison, there are only five
26 described malacosporean species. Malacosporean development in the invertebrate hosts
27 (freshwater bryozoans) has been relatively well studied but is poorly known in fish hosts. Our aim
28 was to investigate the presence and development of malacosporeans infecting a diversity of fish
29 from Brazil, Europe and the USA. We examined kidney from 256 fish belonging variously to the
30 Salmonidae, Cyprinidae, Nemacheilidae, Esocidae, Percidae, Polyodontidae, Serrasalmidae,
31 Cichlidae and Pimelodidae. Malacosporean infections were detected and identified by PCR and
32 SSU rDNA sequencing, and the presence of sporogonic stages was evaluated by ultrastructural
33 examination. We found five malacosporean infections in populations of seven European fish
34 species (brown trout, rainbow trout, white fish, dace, roach, gudgeon and stone loach).
35 Ultrastructural analyses revealed sporogonic stages in kidney tubules of **three** fish species (brown
36 trout, roach and stone loach), providing evidence that fish belonging to at least three families are
37 true hosts. These results expand the range of fish hosts exploited by malacosporeans to complete
38 their life cycle.

39

40 **Key words:** *Buddenbrockia*, *Tetracapsuloides*, SSU rDNA sequence, ultrastructure, sporogenesis,
41 Myxozoa, Cnidaria.

42 **Key findings**

- 43 - Infections of five malacosporean species were detected in populations of seven European
- 44 fish species.
- 45 - Sporogonic stages were observed in kidney tubules of **three** fish species.
- 46 - Molecular and ultrastructural data provided evidence that fish belonging to at least three
- 47 families are true hosts.
- 48 - A second malacosporean life cycle is resolved.

49

50 **Introduction**

51 Myxozoans are microscopic, obligate, endoparasitic cnidarians with complex life cycles (Okamura
52 *et al.*, 2015a). Transmission from host to host is achieved by multicellular spores whose
53 morphologies have been used extensively for taxonomic purposes. However, as it became clear
54 that convergence in spore morphotypes could be problematic, researchers have increasingly
55 incorporated small subunit ribosomal DNA (SSU rDNA) sequences as additional data for the
56 reliable identification of species (Kent *et al.*, 2001; Atkinson *et al.*, 2015). Myxozoans are
57 comprised of two lineages: the speciose Myxosporidia and the species-poor Malacosporea.
58 Collectively there are some 2,500 described myxozoan species (Okamura *et al.*, 2018).
59 Myxosporidian lifecycles involve annelids as definitive hosts and vertebrates, mainly fishes, as
60 intermediate hosts. Malacosporeans use freshwater bryozoans as definitive hosts and fish as
61 intermediate hosts.

62 To date only five malacosporean species have been described (Patra *et al.*, 2016). There are two
63 malacosporean genera: *Tetracapsuloides* and *Buddenbrockia*. Species in both genera develop as
64 sac-like or vermiform (myxoworm) stages in the body cavity of their freshwater bryozoan hosts
65 (Hartikainen *et al.* 2014). Spores produced within sacs and myxoworms are infectious to fish. The
66 only malacosporean whose life cycle has been resolved and whose development in both hosts has

67 been characterized is *Tetracapsuloides bryosalmonae*, the causative agent of salmonid proliferative
68 kidney disease (PKD). *T. bryosalmonae* develops as sacs in the body cavity of freshwater
69 bryozoans (Anderson *et al.*, 1999; Canning *et al.*, 1999) and as pseudoplasmodia in kidney tubules
70 of salmonid fish (Kent and Hedrick, 1985). Spores released from bryozoans infect fish (Feist *et al.*,
71 2001) and spores passed with fish urine infect bryozoans (Hedrick *et al.*, 2004; Morris and Adams,
72 2006).

73 Recent studies provide evidence of a further twelve undescribed species of malacosporeans from
74 a diversity of bryozoan and fish hosts (Bartošová-Sojková *et al.*, 2014; Hartikainen *et al.*, 2014;
75 Patra *et al.*, 2016). These results suggest that there is substantially greater diversity of
76 malacosporeans than is currently appreciated and that further investigations may link
77 malacosporeans detected in fish with those detected in bryozoans, thereby resolving life cycles.
78 The detection of undescribed malacosporeans in fish has, however, largely been gained by PCR
79 and sequencing of material from fish kidney without ascertaining spore development in the putative
80 fish hosts. It is therefore possible that putative fish hosts may be accidental (Bartošová-Sojková *et*
81 *al.*, 2014; Hartikainen *et al.*, 2014). For example, larvae of some nematode parasites (e.g.
82 *Ancylostoma braziliense*, *Ancylostoma caninum*, *Toxocara canis* and *Gnasthostoma spinigerum*)
83 may begin their development in humans but only develop to mature adult worms in their true
84 mammal hosts (Rey, 2008). **Similarly, malacosporeans could invade fish as blood stages that are**
85 **detected by PCR of fish kidney but fish may not support subsequent spore development.** Indeed,
86 this has been observed when local *T. bryosalmonae* strains infect exotic rainbow trout in the United
87 Kingdom (Bucke *et al.*, 1991; Morris *et al.*, 1997).

88 The aim of this study was to characterize malacosporeans in a diversity of fish kidney material,
89 employing both molecular and ultrastructural methods. By using this combined approach, we are
90 able to confirm that malacosporeans exploit fish hosts belonging to at least three families. In
91 addition, we are able to resolve a second malacosporean life cycle by linking the vertebrate and

92 invertebrate hosts. Extending our knowledge of malacosporean host diversity is of general
93 importance for understanding biodiversity, ecology and co-evolutionary relationships in freshwater
94 systems and could be relevant for diagnosis and control of emerging diseases in aquaculture or wild
95 fish populations in our changing world.

96

97 **Material and methods**

98 A total of 256 fish kidney were screened for the presence of malacosporean DNA. **The material we**
99 **studied was gained by a mixture of general and targeted sampling. The former involved taking**
100 **advantage of ongoing project work sampling fish in the River Stour (electrofishing) and in**
101 **Blickling Lake (rod fishing), and screening for fish parasites in practical classes in Switzerland**
102 **(caught by net). More targeted sampling included material collected during surveys for parasites in**
103 **paddlefish in the USA and by specifically sampling fish on farms in Brazil.** Nineteen fish species
104 belonging to 9 fish families (Salmonidae, Cyprinidae, Nemacheilidae, Esocidae, Percidae,
105 Polyodontidae, Serrasalminidae, Cichlidae and Pimelodidae) were sampled from the United
106 Kingdom (UK), Switzerland, Brazil and the United States of America (USA) (Tables 1 and 2). We
107 included archived samples of brown and rainbow trout that were known to be infected by *T.*
108 *bryosalmonae* to provide comparative material because the development of *T. bryosalmonae* is
109 well known. The fish were euthanized in Brazil by benzocaine overdose, in accordance with
110 Brazilian law (Federal Law No. 11.794, dated 8 October 2008 and Federal Decree No. 6899, dated
111 15 July 2009), and in Europe by a blow to the head, followed by severance of the spinal cord.
112 Approximately 27mm³ of tissue was immediately dissected from the posterior portion of the
113 kidney. One half of the kidney material was fixed in 99% ethanol for the molecular analysis and
114 the other half in 2.5% glutaraldehyde in 0.2M sodium cacodylate buffer for the ultrastructural
115 studies described below.

116

117 *DNA extraction, PCR amplification, sequencing and species identification*

118 The DNA was extracted using a DNeasy® Blood & Tissue kit (Qiagen, USA), following the
119 manufacturer's instructions. Malacosporean specific mala-f and mala-r primers (Grabner and El-
120 Matbouli, 2010) were used in PCRs for all samples, amplifying approximately 680 bp of the SSU
121 rDNA. Malacosporean-specific budd-f and budd-r primers (Grabner and El-Matbouli, 2010) were
122 then subsequently used to amplify almost complete length SSU rDNA giving a product that is
123 approximately 1784 bp. **General myxozoan primers such as MedlinA and MedlinB (Medlin *et al.*,
124 1988) were trialed in pilot work but did not amplify any malacosporeans.**

125 Polymerase chain reactions (PCRs) were carried out in 25 µL reaction volumes using 100 ng of
126 extracted DNA, 5 ×Go Taq Flexi Buffer (Promega Madison, WI, USA), 10 mM dNTP mix, 25mM
127 MgCl₂, 10mM for each primer and 1×GoTaq G2 Flexi DNA polymerase (Promega, Madison, WI,
128 USA). The original cycling conditions were used for mala-f and mala-r primers as described by
129 Grabner and El-Matbouli (2010). For runs using budd-f and budd-r primers an initial denaturation
130 stage at 95 °C for 5 min was followed by 40 cycles of denaturation at 95 °C for 45 s, annealing at
131 61 °C for 45 s, extension at 72 °C for 105 s, finishing with an extended elongation stage at 72 °C
132 for 8 min. **The cycling conditions were modified from Grabner and El-Matbouli (2010) for budd-f
133 and budd-r primers to increase primer specificity.**

134 Ultrastructural investigation suggested the presence of sphaerosporid myxozoans in kidney of
135 white fish and dace. To identify cases presenting simultaneous infections of both sphaerosporids
136 and malacosporeans all kidney material was screened by nested PCR using the primers and
137 conditions outlined in Patra *et al.*, 2018 (Erib 1 and 10 primers for primary PCR, [95 °C for 5 min,
138 35 cycles of 94 °C for 1 min, 60°C for 1 min, extension 90 s] followed by final extension of 5 mins
139 and SphFWSSU1243F and SphFWSSU3418R for nested PCR primers [95 °C for 5 min, 35 cycles
140 of 94 °C for 1 min, 56°C for 1 min, extension 90 s] followed by final extension of 5 mins).

141 PCR products were electrophoresed in 2.0% agarose gel, stained with gel red and analysed by a
142 Syngene Transilluminator. PCR products were purified using a Gel/PCR DNA Fragment

143 Extraction Kit (Geneaid Biotech Ltd., USA) and sequenced. This work was conducted in the
144 Molecular Biology Unit of the Natural History Museum, London (NHM) using the Applied
145 Biosystems 3730xl DNA Analyser for Sanger sequencing. OTUs were compared with SSU rDNA
146 sequence data in GenBank and species identity was based on $\geq 99\%$ similarity (Bartošová-Sojková
147 *et al.*, 2014; Bartošová and Fiala, 2011; Whipps and Kent, 2006). An alignment of the **original**
148 SSU rDNA sequences obtained in this study and related species from Genbank (**see table 1 for**
149 **sequence length**), was used to produce a pairwise **dissimilarity** matrix using MEGA 6.0 (Tamura
150 *et al.*, 2011).

151

152 *Electron microscopy*

153 Pieces of kidney were fixed in 2.5% glutaraldehyde in 0.2M sodium cacodylate buffer, pH 7.4 and
154 post-fixed in 1% OsO₄ in cacodylate buffer. Matching kidney material revealed to be positive for
155 malacosporean infection (as identified by PCR and sequencing) was then dehydrated in a graded
156 series of ethanol and embedded in Agar 100 resin (Agar Scientific, Stansted, UK) via propylene
157 oxide. Semi-thin sections were stained with toluidene blue and ultrathin sections with uranyl
158 acetate and lead citrate. Material was examined using a Hitachi H-7650 transmission electron
159 microscope available at the NHM's sister institute, Jodrell Laboratory at Kew Gardens and a LEO
160 906 electron microscope available at the University of Campinas (UNICAMP), São Paulo, Brazil.
161 **We attempted to locate malacosporean infections in at least five kidneys that were identified as**
162 **positive by PCR of each fish species. The number of kidneys analysed by ultrastructure was**
163 **ultimately constrained by availability, suitability of material, and time (see Table 3).**

164

165 **Results**

166 Malacosporean infections were detected in the kidney of fishes originating from the UK and
167 Switzerland. Five species were identified: *T. bryosalmonae*; *Tetracapsuloides* sp. 4 (Bartošová-
168 Sojková *et al.*, 2014) (also referred as *Tetracapsuloides* sp. 3 [Patra *et al.*, 2016] and from here on

169 called *Tetracapsuloides* sp. 4); *Tetracapsuloides* sp. 5 (Bartošová-Sojková *et al.*, 2014) (also
170 referred as *Tetracapsuloides* sp. 2 [Patra *et al.*, 2016] and from here on called *Tetracapsuloides* sp.
171 5); *B. plumatellae*; and *Buddenbrockia* sp. 2 (Hartikainen *et al.*, 2014) (also referred as
172 *Buddenbrockia* sp. 4 [Patra *et al.*, 2016] and from here on called *Buddenbrockia* sp. 2). The
173 infection prevalences of these species ranged from 5.5% to 100% (Table 1).

174 *Tetracapsuloides bryosalmonae* was identified to infect 85.7% of white fish from Lake Lucerne
175 (n = 7), 77.7% of the brown trout specimens examined from the River Stour (n = 9), 89.0% of
176 brown trout from the River Brubach (n = 18), and 100% of rainbow trout from the River Furtbach
177 (n = 12) (Table 1). Young amorphous sporogonic stages (Fig. 1 A) and mature spores (Fig. 1 B-D)
178 of *T. bryosalmonae* were found in kidney tubules of brown trout from the River Stour. Young
179 amorphous sporogonic stages containing sporoplasmosomes with a lucent area (Fig. 1E-F) and a
180 pseudoplasmodium connected to the kidney tubule wall via pseudopodia (Fig. 1 G-H) were
181 observed in white fish from Lake Lucerne. However, because PCR indicated co-infection of this
182 material (see below), it is possible these are immature sporogonic stages of a sphaerosporid.
183 Unfortunately, although five white fish kidneys were examined by ultrastructure, the only kidney
184 that revealed developmental stages was this co-infected material (Table 3).

185 *Buddenbrockia plumatellae* infected 33.3% (n = 16) of the dace specimens from the River Stour,
186 53.8% of the roach from Blickling Lake (n = 13), and 5.5% of the roach from Lake Lucerne (n =
187 18) (Table 1). Unfortunately, ultrastructure was uninformative, being compromised by
188 degeneration of dace material.

189 *Tetracapsuloides* sp. 4 was detected in the kidney of roach sampled from the River Stour at a
190 prevalence of 6.3% (n = 16) (Table 1). Advanced developmental stages anchored to the kidney
191 tubule wall via pseudopodia showed polar capsules and sporoplasmosomes with the characteristic
192 lucent area (Fig. 2A-D).

193 *Tetracapsuloides* sp. 5 was detected in the kidney of gudgeon from the River Stour at a
194 prevalence of 42.9% (n = 21) (Table 1). Clear developmental stages of spores were not observed
195 in kidney tubules.

196 *Buddenbrockia* sp. 2 was detected in 100% (n = 20) of the stone loach specimens sampled from
197 the River Stour (Table 1). Ultrastructural analysis revealed sporogonic stages and mature spores in
198 kidney tubules (Fig. 3 A-E).

199 There was low divergence between the SSU rDNA sequences of the malacosporean species
200 found in this study and the most similar sequences available in Genbank (ranging from 0.1 to 0.3%)
201 (Table 4).

202 Sphaerosporid co-infections were identified in four individuals, two dace from Kent and two
203 white fish from Zurich. Both host species have previously been reported with sphaerosporid
204 infections in Europe (El-Matbouli and Hoffman, 1996; Patra *et al.*, 2018). Rounded sporogonic
205 stages (Fig. 4 A and D) with an electron-dense material surrounding each early developmental
206 spore stage (black arrows in Fig. 4 A, B, E, F) were observed in white fish and dace. The same
207 electron-dense material was observed forming the hard valves of a mature spore in white fish (Fig.
208 4 C).

209 **Malacosporean infections were detected by ultrastructure for material that was positive by PCR**
210 **in 4 of 33 cases that were examined (see table 3).**

211

212 **Discussion**

213 *Malacosporeans exploit a diversity of fish hosts*

214 Our results demonstrate that a range of fish hosts belonging to different families are used by the
215 two currently recognised malacosporean genera, *Tetracapsuloides* and *Buddenbrockia*. Infection

216 of trout by *T. bryosalmonae* has been known for decades (Kent and Hedrick, 1985), with many
217 studies demonstrating development in kidney tubules of brown and rainbow trout in the UK and
218 the USA (Kent and Hedrick, 1986; Clifton-Hadley and Feist, 1989; Morris *et al.*, 2000; Hedrick *et al.*
219 *al.*, 1993). *T. bryosalmonae* has been suggested to infect all salmonid species (Hedrick *et al.*, 1993)
220 but whether all species serve as effective hosts is unclear. The consistent lack of sporogony in
221 exotic rainbow trout in Europe (Grabner and El-Matbouli, 2008; Hartikainen and Okamura, 2015)
222 demonstrates that, although some salmonids are susceptible to infection, they are accidental hosts.
223 Our results suggest that white fish in Switzerland may also serve as hosts of *T. bryosalmonae* but
224 we were unable to definitively confirm spore production in white fish that were not also infected
225 with sphaerosporids. The prevalences of *T. bryosalmonae* infections were similar in white fish,
226 brown trout and rainbow trout, although it should be stressed that this observation is based on
227 relatively low sample sizes.

228 Our further studies of *Tetracapsuloides* spp. were also informative. The presence of sporogonic
229 stages including advanced developmental stages of spores of *Tetracapsuloides* sp. 4 in kidney
230 tubules in roach from the River Stour, imply that roach is a true host. The prevalence of infection
231 (6.3%) was lower than that reported in a study based on molecular analyses of roach kidney
232 material from the Czech Republic (100%; Bartošová-Sojková *et al.*, 2014). However, the high
233 prevalence reported by Bartošová-Sojková *et al.* (2014) is very likely biased by low sample size (n
234 = 2). *Tetracapsuloides* sp. 5 was detected in kidney of gudgeon from the River Stour where the
235 prevalence of infection was 43.0%. Previous molecular investigation detected this species in
236 gudgeon in the Czech Republic, with prevalences of 33.0 and 91.0% (Bartošová-Sojková *et al.*,
237 2014). We did not observe sporogonic stages in kidney tubules and thus cannot confirm the host
238 status of gudgeon. Nevertheless, recurrent detection of this species in gudgeon (Bartošová-Sojková
239 *et al.*, 2014; Patra *et al.*, 2016) often at substantial prevalences suggest that infection by
240 *Tetracapsuloides* sp. 5 is common. Further work is required to clarify the host status of gudgeon.

241 Bartošová-Sojková *et al.* (2014) found *B. plumatellae* infections in dace, roach, and bleak
242 (*Alburnus alburnus*) in the Czech Republic at 100%, 60.0%, and 46.0% infection prevalences,
243 respectively. We detected this species in dace from the River Stour (infection prevalence = 33.3%)
244 and in roach from Blickling Lake (infection prevalence = 53.8%) and Lake Lucerne (infection
245 prevalence = 5.5%). Grabner and El-Matbouli (2010) showed, in a cohabitation study, that *B.*
246 *plumatellae* was transmitted from bryozoans to carp and minnow. The collective evidence thus
247 suggests that of *B. plumatellae* is able to exploit a range of cyprinid hosts but it remains to be
248 confirmed whether roach and dace support sporogony.

249 Our detection of *Buddenbrockia* sp. 2 in stone loach is the first time this malacosporean has
250 been linked with a fish host. Hartikainen *et al.* (2014) reported infections of *Buddenbrockia* sp. 2
251 which develops as myxoworms in the bryozoan *Fredericella sultana* sampled in the UK,
252 Switzerland and Germany. Our ultrastructural analyses revealed sporogonic stages and mature
253 spores in kidney tubules, identifying stone loach as a true host. Thus, the life cycle of
254 *Buddenbrockia* sp. 2 appears to be resolved, with the parasite exploiting *F. sultana* as an
255 invertebrate host and *Barbatula barbatula* as a vertebrate host. It is of course conceivable that
256 further fish and bryozoan hosts may be used.

257 It should be noted that no signals of kidney infection were observed when fish were dissected
258 to collect material for study, an observation in keeping with the general view that many myxozoan
259 infections are innocuous and/or have little impact on fish hosts (Schulman, 1990; Lom and Dyková,
260 1992). It is also consistent with the weak or absent immune response, typically observed in natural
261 fish hosts of malacosporeans (Grabner and El-Matbouli, 2008). Such inapparent infections almost
262 certainly contribute to the general lack of investigation of malacosporean infections in fish.
263 Notably, inapparency also characterised the infections of brown trout and white fish by *T.*
264 *bryosalmonae*, suggesting that environmental conditions and/or fish health status were not

265 conducive for PKD development. However, the high infection prevalences (78.0% in the River
266 Stour and 87.0% in Lake Lucerne) suggest that many fish may have the potential to develop disease.

267 We should also note that in only some 12% of cases (4/33) where we obtained positive PCR
268 results were we able to detect malacosporean stages by ultrastructure in the paired kidney material
269 (Table 3). In some cases this was due to degraded material (e.g. dace). In other cases, this could
270 reflect little proliferation and development in kidney, which then made detection by ultrastructure
271 very difficult and eventually we ceased searching.

272

273 *The challenge of identifying fish hosts*

274 The confirmation of malacosporean fish host status is variously challenging. For example, the lack
275 of detection of malacosporeans in material from Brazil may reflect seasonality, low sample sizes,
276 lack of examination of appropriate fish age classes or absence of bryozoan hosts where the fish
277 were sampled. We anticipate that malacosporeans are present in Brazil in view of observations by
278 Marcus (1941) of *Buddenbrockia* infections in bryozoans in São Paulo State. In addition,
279 ultrastructural detection of sporogonic stages of malacosporeans in the vertebrate host is
280 complicated. It requires extensive sectioning of embedded material to search for small
281 developmental stages, and spores that may be patchily distributed in kidney tubules. The failure to
282 detect sporogonic stages in some cases may simply arise from a limited number of parasites or
283 because the infection has not yet matured. This is compounded by the inapparency of many
284 infections at the macroscopic level.

285 As shown here, a targeted approach employing associated SSU rDNA sequencing to confirm
286 infection status will at least identify what material to investigate. The alternative approach of
287 conducting transmission trials to confirm that infection is transmitted from fish to bryozoans
288 requires fish husbandry, and permits are often required for such work. We suggest a potential

289 alternative molecular approach for future identification of fish hosts by determining whether genes
290 specifically involved in polar capsule development (e.g. minicollagens and nematogalactins
291 Holland *et al.*, 2011; Shpirer *et al.*, 2014, NSPs 1-7 Shpirer *et al.*, 2018) are expressed in infected
292 kidney. The rationale is that polar capsules are only present in malacosporean spores and thus the
293 detection of such expressed genes would indicate spore development. In practical terms this would
294 involve preservation of kidney material in e.g. RNAlater and confirmation that these genes are not
295 expressed in pre-sporogonic developmental stages.

296

297 *Ultrastructural distinction of malacosporean and myxosporean sporogonic stages in fish kidney*
298 *tubules*

299 We found co-infections of *T. bryosalmonae* and a sphaerosporid species in white fish and of *B.*
300 *plumatellae* and a sphaerosporid species in dace. The most apparent morphological difference
301 between myxosporean and malacosporean spores is the composition of their valves, which are
302 hardened in myxosporeans but remain soft in malacosporeans (Anderson *et al.*, 1999; Canning and
303 Okamura, 2004). The hardening of myxosporean spore valves is achieved by secretion of chitin
304 (Munoz *et al.*, 1999; 2000; Liu *et al.*, 2011), which may also be associated with internal organelles
305 (Lukeš *et al.*, 1993; Munoz *et al.*, 2000; Redondo *et al.*, 2008). This glycoprotein likely protects
306 myxosporean spores from environmental degradation and maintains spore shape (Munoz *et al.*,
307 1999; Kaltner *et al.*, 2007; Estensoro *et al.*, 2013). Unprotected malacosporean spores degrade
308 relatively quickly upon release from fish (in <24 h) (de Kinkelin *et al.*, 2002) compared to the
309 chitin-protected spores of myxosporeans. Accordingly, electron microscopy of mature
310 myxosporean spores reveals electron dense valves, and in immature spores an accumulation of
311 electron dense material (inferred to be valve forming material) is observed in the cytoplasm of
312 valvogenic cells (Adriano *et al.*, 2009; Moreira *et al.*, 2014; Morsy *et al.*, 2016). Fibrillar electron
313 dense material reported in valvogenic cells of *Sphaerospora jiroveci* forms a continuous layer

314 enclosing the developing spore (Dyková and Lom, 1997). Further studies are required to confirm
315 whether this electron dense material is chitin and whether its presence is characteristic of
316 sphaerosporids (Fig. 4B-F).

317 At least at present, there appear to be no other reliable morphological features that could be
318 used to distinguish between malacosporean and myxosporean developmental stages. For example,
319 although the lucent area of sporoplasmosomes has been highlighted as a malacosporean feature and
320 is evident in the sporoplasmosomes in roach (Figure 2C-D) and white fish (Figure 1E-F), such
321 lucent areas are also occasionally observed in sporoplasmosomes of myxosporeans (Lom *et al.*,
322 1989; Álvarez-Pellitero *et al.*, 2002; Morris and Freeman 2009; Naldoni, pers. obs.). In addition,
323 the noted tendency for a peripheral distribution of sporoplasmosomes in primary cells of
324 malacosporeans (Canning *et al.*, 2000; Canning *et al.*, 2009) is also sometimes occasionally
325 observed in the primary cytoplasm of myxosporeans (Supamattaya *et al.*, 1993). We would note
326 that in our experience malacosporean sporogonic stages in kidney are difficult to find relative to
327 myxosporean stages. Thus, frequent detection of developmental stages in histological sections
328 could suggest the presence of myxosporeans.

329

330 *Final comments*

331 Our work has revealed five malacosporeans infecting a variety of fish hosts in the UK and
332 Switzerland, contributing to the growing evidence of a hidden diversity of vertebrate hosts that are
333 exploited by this myxozoan lineage. Further study is necessary to formally describe some of these
334 malacosporean species, to determine if gudgeon act as true hosts of *Tetracapsuloides* sp. 5, to
335 confirm that white fish are true hosts of *T. bryosalmonae*, and to ascertain whether *B. plumatellae*
336 is able to utilise both dace and roach as hosts. It is also clear that the two malacosporean genera are
337 not restricted to exploiting single fish families with *Tetracapsuloides* spp. exploiting members of

338 Salmonidae and Cyprinidae and *Buddenbrockia* spp. exploiting members of Cyprinidae and
339 Nemacheilidae (Table 1).

340 Screening both bryozoans and fish has provided vital information about malacosporean
341 diversities and distributions (Bartošová-Sojková *et al.*, 2014; Hartikainen *et al.*, 2014; Patra *et al.*,
342 2016) and may lead to the resolution of life cycles (this study). Further studies should aim to
343 determine whether malacosporeans generally show broad fish host use or whether some may be
344 specialists. In addition, resolution of accidental hosts and the consequences of dead-end infections
345 could lead to future studies on host-parasite interactions, effective immune responses and the
346 potential dilution of infectious stages when non-permissive hosts are abundant. Finally, resolving
347 malacosporean hosts may help us to understand the range of early hosts that were exploited as
348 cnidarians evolved endoparasitism in view of the primitive nature of malacosporeans (Okamura *et*
349 *al.*, 2015b).

350

351 **Acknowledgements.** The authors thank: Hanna Hartikainen (EAWAG, Zurich) and Heike
352 Schmidt-Posthaus (University of Bern) for help with access to or provision of fish kidney material
353 from the River Furtbach, Lake Lucerne, and the River Brubach, and Joe Huddart and Guy
354 Woodward (Imperial College, London) for enabling sampling kidney of fish from the River Stour,
355 UK. We also thank Pavla Bartošová-Sojková for drawing our attention to the presence of a
356 sphaerosporid infection in kidney material and advising on sphaerosporid development.

357

358 **Financial support.** This study was financed in part by the Coordenação de Aperfeiçoamento de
359 Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001 and by the grant #2013/21374-6,
360 São Paulo Research Foundation (FAPESP) to E. A. Adriano. E. A. Adriano received a research
361 productivity grant from the Brazilian Fostering Agency CNPq (grant #301886/2016-4). J. Naldoni
362 was supported by the scholarship #2016/08831-7, São Paulo Research Foundation (FAPESP).

363

364 **Reference**

365

366 **Adriano, EA, Arana, S, Carriero, MM, Naldoni, J, Ceccarelli, PS and Maia, AAM** (2009)
367 Light, electron microscopy and histopathology of *Myxobolus salminus* n. sp., a parasite of
368 *Salminus brasiliensis* from the Brazilian Pantanal. *Veterinary Parasitology* **165**, 25–29.

369 **Alvarez-Pellitero, P, Molnár, K, Sitjà-Bobadilla, A and Székely, C** (2002) Comparative
370 ultrastructure of the actinosporean stages of *Myxobolus bramae* and *M. psuedodispar*
371 (Myxozoa). *Parasitology Research* **88**, 198-207.

372 **Anderson, CL, Canning, EU and Okamura B** (1999) Molecular data implicate bryozoans as
373 hosts for PKX (Phylum Myxozoa) and identify a clade of bryozoan parasites within the
374 Myxozoa. *Parasitology* **119**, 555–561.

375 **Atkinson, SD, Bartošová-Sojková, P, Whipps, CM and Bartholomew, JL** (2015) Approaches
376 for characterising myxozoan species. In Okamura B, Gruhl A and Bartholomew JL (ed.)
377 *Myxozoan evolution, ecology and development*, Switzerland: Springer International Publishing,
378 pp. 111-123.

379 **Bartošová-Sojková, P, Hrabcová, M, Pecková, H, Patra, S, Kodádková, A, Jurajda, P, Týmł,**
380 **T and Holzer, AS** (2014) Hidden diversity and evolutionary trends in malacosporean parasites
381 (Cnidaria: Myxozoa) identified using molecular phylogenetics. *International Journal for*
382 *Parasitology* **44**, 565–577.

383 **Bartošová, P and Fiala, I** (2011) Molecular evidence for the existence of cryptic species
384 assemblages of several myxosporeans (Myxozoa). *Parasitology Research* **108**, 573–583.

- 385 **Bucke, D, Feist, SW and Clifton-Hadley, RS** (1991) The occurrence of proliferative kidney
386 disease (PKD) in cultured and wild fish: further investigations. *Journal of Fish Diseases* **14**,
387 583–588.
- 388 **Canning, EU, Curry, A and Okamura, B** (2009). Early development of the myxozoan
389 *Buddenbrockia plumatellae* in the bryozoans *Hyalinella punctata* and *Plumatella fungosa*, with
390 comments on taxonomy and systematics of the Myxozoa. *Folia parasitologica* **55**, 241-55.
- 391 **Canning, EU and Okamura, B** (2004) Biodiversity and Evolution of the Myxozoa. *Advances in*
392 *Parasitology* **56**, 43-131.
- 393 **Canning, EU, Curry, A, Feist, SW, Longshaw, M and Okamura, B** (2000) A new class and
394 order of myxozoans to accommodate parasites of bryozoans with ultrastructural observations on
395 *Tetracapsula bryosalmonae* (PKX organism). *Journal of Eukaryotic Microbiology* **47**, 456-468.
- 396 **Canning, EU, Curry, A, Feist, SW, Longshaw, M and Okamura, B** (1999) *Tetracapsula*
397 *bryosalmonae* n. sp. for PKX organism the cause of PKD in salmonid fish. *Bulletin of the*
398 *European Association of Fish Pathologists* **19**, 203–206.
- 399 **Clifton-Hadley, RS and Feist, SW** (1989) Proliferative kidney disease in brown trout *Salmo*
400 *trutta*: further evidence of a myxosporean aetiology. *Diseases of Aquatic Organisms* **6**, 99-103.
- 401 **de Kinkelin, P, Gay, M and Forman, S** (2002) The persistence of infectivity of *Tetracapsula*
402 *bryosalmonae*-infected water for rainbow trout, *Oncorhynchus mykiss* (Walbaum). *Journal of*
403 *Fish Diseases* **25**, 477–482.
- 404 **Dyková, I and Lom, L** (1997) Light and Electron Microscope Observations on *Sphaerospora*
405 *ojiroveci* n. sp. (Myxozoa) from the Kidney of *Pangasius sutchi* (Teleostei). *European Journal*
406 *of Protistology* **33**, 444-451.

- 407 **El-Matbouli, M and Hoffman, RW** (1996) Light and electron microscopic descriptions of
408 *Sphaerospora coregoni* El-Matbouli, Hoffmann and Kern, 1995 (Myxosporea:
409 Sphaerosporidae) from the kidney of Whitefish (*Coregonus lavaretus*). *European Journal for*
410 *Protistology* **32**, 389-398.
- 411 **Estensoro, I, Álvarez-Pellitero, P and Sijta-Bobadilla, A** (2013) Antigenic characterization of
412 *Enteromyxum leei* (Myxozoa: Myxosporea). *Diseases of Aquatic Organisms* **106**, 149-162.
- 413 **Feist, SW, Longshaw, M, Canning, EU and Okamura, B** (2001) Induction of proliferative
414 kidney disease (PKD) in rainbow trout *Oncorhynchus mykiss* via the bryozoan *Fredericella*
415 *sultana* infected with *Tetracapsula bryosalmonae*. *Diseases of Aquatic Organisms* **45**, 61–68.
- 416 **Grabner, DS and El-Matbouli, M** (2010) Experimental transmission of malacosporean parasites
417 from bryozoans to common carp (*Cyprinus carpio*) and minnow (*Phoxinus phoxinus*).
418 *Parasitology* **137**, 629–639.
- 419 **Grabner, DS and El-Matbouli, M** (2008). Transmission of *Tetracapsuloides bryosalmonae*
420 (Myxozoa: Malacosporea) to *Fredericella sultana* (Bryozoa: Phylactolaemata) by various fish
421 species. *Diseases of Aquatic Organisms* **79**, 133–139.
- 422 **Hartikainen, H, Gruhl, A and Okamura, B** (2014) Diversification and repeated morphological
423 transitions in endoparasitic cnidarians (Myxozoa: Malacosporea). *Molecular Phylogenetics and*
424 *Evolution* **76**, 261–269.
- 425 **Hartikainen, H and Okamura, B** (2015) Ecology and evolution of malacosporean-bryozoan
426 interactions. In Okamura B, Gruhl A and Bartholomew JL (ed.) *Myxozoan evolution, ecology*
427 *and development*, Switzerland: Springer International Publishing, pp. 201–213.

- 428 **Hedrick, RP, Baxa DV, de Kinkelin, P and Okamura, B** (2004) Malacosporean-like spores in
429 urine of rainbow trout react with antibody and DNA probes to *Tetracapsuloides bryosalmonae*.
430 *Parasitology Research* **92**, 81–88.
- 431 **Hedrick, RP, MacConnell, E and Kinkelin, P** (1993) Proliferative kidney disease of salmonid
432 fish. *Annual Review of Fish Diseases* **3**, 227-290.
- 433 **Holland, JW, Okamura, B, Hartikainen, H and Secombes, CJ** (2011) A novel minicollagen
434 gene links cnidarians and myxozoans. *Proceedings of the Royal Society B* **278**, 546-553.
- 435 **Kaltner, H, Stipl, M, Knaus, M and El-Matbouli, M** (2007) Characterization of glycans in the
436 developmental stages of *Myxobolus cerebralis* (Myxozoa), the causative agent of whirling
437 disease. *Journal of Fish Diseases* **30**, 637–647.
- 438 **Kent, ML, Andree, KB, Bartholomew, JL, El-Matbouli, M, Desser, SS, Devlin, RH, Feist,**
439 **SW, Hedrick, RP, Hoffmann, RW, Khattra, J, Hallett, SL, Lester, RJ, Longshaw, M,**
440 **Palenzeula, O, Siddall, ME and Xiao, C** (2001) Recent advances in our knowledge of the
441 Myxozoa. *The Journal of Eukaryotic Microbiology* **48**, 395–413.
- 442 **Kent, ML and Hedrick, RP** (1986) Development of the PKX myxosporean in rainbow trout
443 *Salmo gairdneri*. *Diseases of Aquatic Organisms* **1**, 169-182.
- 444 **Kent, ML and Hedrick, RP** (1985) PKX, the causative agent of Proliferative Kidney Disease
445 (PKD) in Pacific salmonid fishes and its affinities with the Myxozoa. *Journal of Parasitology*
446 **32**, 254–260.
- 447 **Liu, Y, Zhou, Z, Miao, W, Zhang, Y, Cao, Y, et al** (2011) A chitinase from *Aeromonas veronii*
448 CD3 with the potential to control myxozoan disease. *PLoS ONE* **6**, e29091.
- 449 **Lom, J and Dyková, I** (2006) Myxozoan genera: definition and notes on taxonomy, life-cycle
450 terminology and pathogenic species. *Folia Parasitologica* **53**, 1–36.

- 451 **Lom, J and Dyková, I** (1992) Chapter 7 Myxosporidia (Phylum Myxozoa). In Lom, J and Dyková,
452 I (ed.) *Protozoan parasites of fishes*, Amsterdam: Developments in Aquaculture and Fisheries
453 Sciences, pp. 159-235.
- 454 **Lom, J, Feist, SW, Dyková, I and Kepr, T** (1989) Brain myxoboliasis of bullhead, *Cottus gobio*
455 L., due to *Myxobolus jiroveci* sp. nov.: light and electron microscope observations. *Journal of*
456 *Fish Diseases* **12**, 15-27.
- 457 **Lukeš, J, Volf, P and Lom, J** (1993) Detection of chitin in spores of *Myxobolus muelleri* and *M.*
458 *subepithelialis* (Myxosporidia, Myxozoa). *Parasitology Research* **79**, 439-440.
- 459 **Marcus, E** (1941) Sobre Bryozoa do Brasil. Boletim Faculdade de Filosofia, Ciências e Letras,
460 Universidade de São Paulo, *Serie Zoologia* **10**, 3–207.
- 461 **Medlin, L, Elwood, HJ, Stickel, S and Sogin, ML** (1988) The characterization of enzymatically
462 amplified eukaryotic 16S-like rRNA-coding regions. *Gene* **71**, 491–499.
- 463 **Moreira, GSA, Adriano, EA, Silva, MRM, Ceccarelli, PS, and Maia, AAM** (2014) The
464 morphological and molecular characterization of *Hennequya rotunda* n. sp., a parasite of the gill
465 arch and fins of *Salminus brasiliensis* from the Mogi Guaçu River, Brazil. *Parasitology*
466 *Research* **113**, 1703-1711.
- 467 **Morris, D and Freeman, Mark** (2009). Hyperparasitism has wide-ranging implications for
468 studies on the invertebrate phase of myxosporean (Myxozoa) life cycles. *International Journal*
469 *for Parasitology* **40**, 357-69.
- 470 **Morris, DJ and Adams, A** (2006) Transmission of *Tetracapsuloides bryosalmonae* (Myxozoa:
471 Malacosporidia), the causative organism of salmonid proliferative kidney disease, to the
472 freshwater bryozoan *Fredericella sultana*. *Parasitology* **133**, 701–709.

- 473 **Morris, DJ, Adams, A and Richards, RH** (2000) Observations on the electron-dense bodies of
474 the PKX parasite, agent of proliferative kidney disease in salmonids. *Diseases of Aquatic*
475 *Organisms* **39**, 201-209.
- 476 **Morris, DJ, Adams, A and Richards, RH** (1997) Studies of the PKX parasite in rainbow trout
477 via immunohistochemistry and immunogold electron microscopy. *Journal of Aquatic Animal*
478 *Health* **9**, 265–273.
- 479 **Morsy, K, Semmler, M, Al-Olayan, E and Mehlhorn, H** (2016) *Henneya collaris* sp. nov.,
480 (Myxosporidia), parasite of the Greenband Parrotfish *Scarus collana* Rüppell, 1835
481 (Actinopterygii, Scaridae) from the Red Sea, Egypt. A light and electron microscopic study.
482 *Parasitology Research* **115**, 2253-2261.
- 483 **Muñoz, P, Sitjà-Bobadilla, A and Álvarez-Pellitero, P** (2000) Ultrastructural localisation of
484 carbohydrates in four myxosporean parasites. *Parasite* **7**, 185–191.
- 485 **Muñoz, P, Palenzuela, O, Alvarez-Pellitero, P and Sitjà-Bobadilla, A** (1999) Comparative
486 studies on carbohydrates of several myxosporean parasites of fish using lectin histochemical
487 methods. *Folia Parasitologica* **46**, 241–247.
- 488 **Okamura, B, Hartigan, A and Naldoni, J** (2018) Extensive uncharted biodiversity: the parasite
489 dimension. *Integrative and Comparative Biology* 1-14. doi: 10.1093/icb/icy039
- 490 **Okamura, B, Gruhl, A and Bartholomew, JL** (2015a) An introduction to myxozoan evolution,
491 ecology and development. In Okamura B, Gruhl A and Bartholomew JL (ed.) *Myxozoan*
492 *evolution, ecology and development*, Switzerland: Springer International Publishing, pp. 1-20.
- 493 **Okamura, B, Gruhl, A and Reft, AJ** (2015b) Cnidarian origins of the Myxozoa. In Okamura B,
494 Gruhl A and Bartholomew JL (ed.) *Myxozoan evolution, ecology and development*, Switzerland:
495 Springer International Publishing, pp. 45-68.

- 496 **Patra, S, Bartošová-Sojková, P, Pecková, H, Fiala, I, Eszterbauer, E, and Holzer, AS** (2018)
497 Biodiversity and host-parasite cophylogeny of *Sphaerospora* (sensu stricto) (Cnidaria:
498 Myxozoa). *Parasites & Vectors* **11**, 347.
- 499 **Patra, S, Hartigan, A, Morris, DJ, Kodádková, A and Holzer, AS** (2016) Description and
500 experimental transmission of *Tetracapsuloides vermiformis* n. sp. (Cnidaria: Myxozoa) and
501 guidelines for describing malacosporan species including reinstatement of *Buddenbrockia*
502 *bryozoides* n. comb. (syn. *Tetracapsula bryozoides*). *Parasitology* **144**, 497-511.
- 503 **Redondo, MJ, Cortadellas, N, Palenzuela, O and Alvarez-Pellitero, P** (2008) Detection of
504 carbohydrate terminals in the enteric parasite *Enteromyxum scophthalmi* (Myxozoa) and
505 possible interactions with its fish host *Psetta maxima*, *Parasitology Research* **102**, 1257-1267.
- 506 **Rey, L** (2008) Parasitologia. 4^a edição. Editora Guanabara Koogan.
- 507 **Shpirer, E, Diamant, A, Cartwright, P and Huchon, D** (2018) A genome wide survey reveals
508 multiple nematocyst-specific genes in Myxozoa. *BMC Evolutionary Biology* **18**, 138.
- 509 **Shpirer, E, Chang, ES, Diamant, A, Rubinstein, N, Cartwright, P and Huchon, D** (2014)
510 Diversity and evolution of myxozoan minicollagens and nematogalectins. *BMC Evolutionary*
511 *Biology* **14**, 205.
- 512 **Shul'man, SS** (1990). Myxosporidia of the USSR. Russian Translations Series 75. A. A.
513 Balkema/Rotterdam.
- 514 **Supamattaya K, Fischer-Scherl T, Hoffmann RW and Boonyaratpalin S** (1993) Light and
515 electron microscope observations on presporogonic and sporogonic stages of *Sphaerospora*
516 *epinepheli* (Myxosporidia) in grouper (*Epinephelus malabaricus*). *The Journal of Eukaryotic*
517 *Microbiology* **40**, 71-80. doi:10.1111/j.1550-7408.1993.tb04885.x

518 **Tamura, K, Peterson, D, Peterson, N, Stecher, G, Nei, M and Kumar, S** (2011) MEGA5:
519 Molecular Evolutionary Genetics Analysis using maximum likelihood, evolutionary distance,
520 and maximum parsimony methods. *Molecular Biology and Evolution* **28**, 2731–2739.

521 **Whipps, CM and Kent, ML** (2006) Phylogeography of the cosmopolitan marine parasite *Kudoa*
522 *thyrsites* (Myxozoa: Myxosporea). *Journal of Eukaryotic Microbiology* **53**, 364–373.

523

524 **Table legends**

525 **Table 1.** Fish sampled from the UK and Switzerland that were infected with malacosporeans.
526 Details on collection sites, fish hosts, malacosporean species; prevalence of infection, SSU rDNA
527 sequence lengths; sequences that provided the closest match in GenBank, and **bryozoans (*) and**
528 **fishes** previously identified to be infected by the respective parasites. Archived samples of brown
529 and rainbow trout known to be infected by *Tetracapsuloides bryosalmonae* (**) or infected by *T.*
530 *bryosalmonae* through transmission studies (***). bp = base pairs.

531

532 **Table 2.** Fish sampled from the UK, Switzerland, the USA and Brazil in which infections were not
533 detected. Data on collection sites, fish species and the number of fish sampled (**No. sampled**).

534

535 **Table 3.** The detection of malacosporean infections in fish kidney material from the United
536 Kingdom (UK) and Switzerland (CF) according to PCR, sequencing and ultrastructural analysis.
537 Included are fish species and locality data, the malacosporean species inferred by sequencing, the
538 number (No.) of fish kidneys detected by PCR, the number of kidneys examined by ultrastructure,
539 and the number of kidneys in which spores were detected by ultrastructure. (*) degraded material.

540

541 **Table 4.** Dissimilarity matrix for small subunit ribosomal DNA (SSU-rDNA) of malacosporean
542 species found in this study and their closest matches in Genbank. The upper triangle shows
543 nucleotide differences in relation to the number of bases compared. The lower triangle shows %
544 pairwise distances. SSU rDNA = 18 Small Subunit Ribosomal DNA; UK: United Kingdom, CF:
545 Switzerland. (BS) = Bartošová-Sojková et al. 2014; (H) = Hartikainen et al. 2014.

546

547 **Figure legends**

548 **Fig. 1.** Photomicrography of kidney tubules (t) of brown trout collected in the River Stour, Kent,
549 UK, in semi-thin sections stained by toluidine blue (A-B). Transmission electron microscopy (C-
550 H) showing the development of *Tetracapsuloides bryosalmonae* spores in the lumen of the kidney
551 tubules (t) of brown trout (C-D) collected in the River Stour, Kent, UK, and **myxozoan**
552 **development (either malacosporean or sphaerosporid)** in white fish (E-H), collected in the Lake
553 Lucerne, Switzerland. (A) Note the presence of two early developmental stages (arrows)
554 developing attached to the kidney tubule wall. Scale bar = 10µm. (B-C) Advanced stage of spore
555 development (empty arrows) showing polar capsule (thin black arrows). Scale bar = 10µm. (D)
556 Mature **spore (s)** showing polar capsule (empty arrow) with polar filaments (white arrow). Scale
557 bar = 2µm. (E) **Primary cell (empty arrow) developing in the lumen of the kidney tubule. Note the**
558 **presence of sporoplasmosomes (box). Scale bar = 1µm.** (F) **High magnification of E showing the**
559 **sporoplasmosomes with a lucent area. Scale bar = 200 nm.** (G) Pseudoplasmodium (empty arrow)
560 developing attached to the kidney tubule wall. Scale bar = 10µm. (H) High magnification of G
561 showing pseudoplasmodium connected to the kidney tubule wall via pseudopodia (thin black
562 arrow). Note the two secondary cell nuclei (white arrows). Scale bar = 2µm.

563

564 **Fig. 2.** Photomicrography of kidney tubules (t) of roach collected in the River Stour, Kent, UK, in
565 semi-thin sections stained by toluidine blue (A), and by transmission electron microscopy (B-D)
566 showing spore development of *Tetracapsuloides* sp. 4. (A) Advanced stage of spore development
567 (empty arrow) with polar capsules (white arrow). Note a stage connected to the kidney tubule wall
568 (thin black arrow). Scale bar = 20 μ m. (B) Two pseudoplasmodia (p) showing secondary cell nuclei
569 (empty arrows) and a polar capsule (white arrow). Note pseudopodia anchoring the parasite to the
570 kidney tubule wall via pseudopodia (thin black arrows). Scale bar = 2 μ m. (C) Primary cell (p) with
571 scattered sporoplasmosomes (box). Scale bar = 2 μ m. (D) High magnification of box in figure C
572 showing details of sporoplasmosomes, each with a lucent area. Scale bar = 500nm.

573

574 **Fig. 3.** Photomicrography of kidney tubules (t) of stone loach collected in the River collected in
575 the River Stour, Kent, UK, in semi-thin sections stained by toluidine blue (A-B), and by
576 transmission electron microscopy (C-E) showing spore development of *Buddenbrockia* sp. 2. (A)
577 Note the different stages of development of spores (arrows) as well as earlier developmental stages.
578 Scale bar = 10 μ m. (B) Two spores in advanced developmental stages (empty arrows) with polar
579 capsules (black arrows). Scale bar = 10 μ m. (C) Note two young developmental stages of spores
580 (empty arrows), with a polar capsule in development (large white arrows). Scale bar = 2 μ m. (D)
581 Pseudoplasmodium (p) in the lumen of the kidney tubule. Scale bar = 5 μ m. (E) High magnification
582 of D showing pseudoplasmodium connected to the kidney tubule wall via pseudopodia (thin black
583 arrow). Note the secondary cell nucleus (white arrow). Scale bar = 1 μ m.

584

585 **Fig. 4.** Transmission electron microscopy showing the development of *Sphaerospora* spp. spores
586 in the lumen of kidney tubules of white fish (A-C) collected in the Lake Lucerne, Switzerland and
587 dace (D-F) collected in the River Stour, Kent, UK. (A) Sporogonic stages (empty arrows)
588 developing in the lumen of the kidney tubule of white fish. Scale bar = 10 μ m. (B) Early sporogonic

589 stage with electron dense material in **valvogenic cells** (black arrow) and a mature spore (empty
590 arrow) in the lumen of the kidney tubule. Scale bar = 5 μ m. (C) High magnification of C showing
591 the mature spore with electron dense material forming the hard valves (black arrow) and two polar
592 capsules (white arrows). Scale bar = 2 μ m. (D) Sporogonic stages (empty arrows) developing in the
593 lumen of the kidney tubule of dace. Scale bar = 5 μ m. (E) High magnification of D showing a
594 sporogonic stage with electron dense material in **the valvogenic cells** (black arrow), the nucleus of
595 the valvogenic cell (nvc), other nuclei (n) and a polar capsule (white arrow). Scale bar = 2 μ m. (F)
596 A sporogonic stage showing electron dense material **in the valvogenic cells** (black arrow) and the
597 nucleus of the valvogenic cell (nvc). Scale bar = 1 μ m.

For Peer Review

Table 1.

Country	Water body (latitude; longitude)	Fish species (and Family)	Malacosporean species	Prevalence	SSU rDNA sequence length	SSU rDNA sequence match	Bryozoan* and fishes associated with infection (and author)
United Kingdom	River Stour, Kent (51°13'49" N; 0°58'10" E)	Brown Trout, <i>Salmo trutta</i> (Salmonidae)	<i>Tetracapsuloides bryosalmonae</i>	77.7% (7/9)	1673 bp	KF731712	<i>Oncorhynchus mykiss</i> , <i>Salmo trutta</i> (Bartošová-Sojková <i>et al.</i> , 2014)
		Dace, <i>Leuciscus leuciscus</i> (Cyprinidae)	<i>Buddenbrockia plumatellae</i>	33.3% (4/12)	1691 bp	KF731682	<i>Abramis brama</i> , <i>Alburnus alburnus</i> , <i>Aspius aspius</i> , <i>Blicca bjoerkna</i> , <i>Chondrostoma nasus</i> , <i>Leuciscus idus</i> , <i>Leuciscus leuciscus</i> , <i>Perca fluviatilis</i> , <i>Rutilus rutilus</i> , <i>Scardinius erythrophthalmus</i> , <i>Squalius cephalus</i> (Bartošová-Sojková <i>et al.</i> , 2014)
		Roach, <i>Rutilus rutilus</i> (Cyprinidae)	<i>Tetracapsuloides</i> sp. 4	6.3% (1/16)	1579 bp	KF731727	<i>Alburnus alburnus</i> , <i>Rutilus rutilus</i> (Bartošová-Sojková <i>et al.</i> , 2014)
		Stone Loach, <i>Barbatula barbatula</i> (Nemacheilidae)	<i>Buddenbrockia</i> sp. 2	100% (20/20)	1693 bp	KJ150268	* <i>Fredericella sultana</i> (Hartikainen <i>et al.</i> , 2014)
		Gudgeon, <i>Gobio gobio</i> (Cyprinidae)	<i>Tetracapsuloides</i> sp. 5	42.9% (9/21)	1580 bp	KF731729	<i>Gobio gobio</i> (Bartošová-Sojková <i>et al.</i> , 2014)
	Blickling Lake, Norfolk (52°48'45.60" N; 1°13'52.90" E)	Roach, <i>Rutilus rutilus</i> (Cyprinidae)	<i>Buddenbrockia plumatellae</i>	53.8% (7/13)	1704 bp	KF731682	<i>Abramis brama</i> , <i>Alburnus alburnus</i> , <i>Aspius aspius</i> , <i>Blicca bjoerkna</i> , <i>Chondrostoma nasus</i> , <i>Leuciscus idus</i> , <i>Leuciscus leuciscus</i> , <i>Perca fluviatilis</i> , <i>Rutilus rutilus</i> , <i>Scardinius erythrophthalmus</i> , <i>Squalius cephalus</i> (Bartošová-Sojková <i>et al.</i> , 2014)
Switzerland	Lake Lucerne (47°04'02.72" N; 8°25'53.63" E)	White Fish, <i>Coregonus albula</i> (Salmonidae)	<i>Tetracapsuloides bryosalmonae</i>	85.7% (6/7)	1683 bp	KF731712	<i>Oncorhynchus mykiss</i> , <i>Salmo trutta</i> (Bartošová-Sojková <i>et al.</i> , 2014)
		Roach, <i>Rutilus rutilus</i> (Cyprinidae)	<i>Buddenbrockia plumatellae</i>	5.5% (1/18)	1692 bp	KF731682	<i>Abramis brama</i> , <i>Alburnus alburnus</i> , <i>Aspius aspius</i> , <i>Blicca bjoerkna</i> , <i>Chondrostoma nasus</i> , <i>Leuciscus idus</i> , <i>Leuciscus leuciscus</i> , <i>Perca fluviatilis</i> , <i>Rutilus rutilus</i> , <i>Scardinius erythrophthalmus</i> ,

							<i>Squalius cephalus</i> (Bartošová-Sojková <i>et al.</i> , 2014)
River Brubach (47°27'48" N; 9°07'15" E)	** Brown Trout, <i>Salmo trutta</i> (Salmonidae)	<i>Tetracapsuloides bryosalmonae</i>	88.9% (16/18)	1683 bp	KF731712		<i>Oncorhynchus mykiss</i> , <i>Salmo trutta</i> (Bartošová-Sojková <i>et al.</i> , 2014)
River Furtbach (47°27'05" N; 8°23'28" E)	***Rainbow Trout, <i>Oncorhynchus mykiss</i> (Salmonidae)	<i>Tetracapsuloides bryosalmonae</i>	100% (12/12)	1683 bp	KF731712		<i>Oncorhynchus mykiss</i> , <i>Salmo trutta</i> (Bartošová-Sojková <i>et al.</i> , 2014)

For Peer Review

Table 2

Country	Site	Fish species (and Family)	N. sampled
United Kingdom	River Stour	Pike, <i>Esox lucius</i> (Esocidae)	12
		Perch, <i>Perca fluviatilis</i> (Percidae)	14
		Minnow, <i>Phoxinus phoxinus</i> (Cyprinidae)	12
		Chub, <i>Squalius cephalus</i> (Cyprinidae)	1
	Felbrigg Lake	Pike, <i>Esox lucius</i> (Esocidae)	4
Switzerland	Lake Lucerne	Perch, <i>Perca fluviatilis</i> (Percidae)	6
		Arctic charr, <i>Salvelinus alpinus</i> (Salmonidae)	6
USA	Oklahoma	Paddle fish, <i>Polyodon spathula</i> (Polyodontidae)	10
Brazil	Fish farm, Campos do Jordão	Rainbow trout, <i>Oncorhynchus mykiss</i> (Salmonidae)	10
	Fish Farm, Santa Cruz da Conceição	Tambaqui, <i>Colossoma macropomum</i> (Serrasalminidae)	10
		Pacu, <i>Piaractus mesopotamicus</i> (Serrasalminidae)	10
	Lake Santa Fé, Pirassununga	Carp, <i>Cyprinus carpio</i> (Cyprinidae)	10
	Amazon River, Santarém	Tambaqui, <i>Colossoma macropomum</i> (Serrasalminidae)	3
		Tucunaré, <i>Cichla monoculus</i> (Cichlidae)	1
Caparari, <i>Pseudoplatystoma punctifer</i> (Pimelodidae)		1	

Table 3.

Fish species and water body	Malacosporean species	No. kidneys positive by PCR	No. kidneys examined by ultrastructure	No. kidneys with spores
Brown Trout, <i>Salmo trutta</i> River Stour, Kent (UK)	<i>Tetracapsuloides bryosalmonae</i>	7	5	1
*Dace, <i>Leuciscus leuciscus</i> River Stour, Kent (UK)	<i>Buddenbrockia plumatellae</i>	4	4	0
Roach, <i>Rutilus rutilus</i> River Stour, Kent (UK)	<i>Tetracapsuloides</i> sp. 4	1	1	1
Stone Loach, <i>Barbatula barbatula</i> River Stour, Kent (UK)	<i>Buddenbrockia</i> sp. 2	20	5	1
Gudgeon, <i>Gobio gobio</i> River Stour, Kent (UK)	<i>Tetracapsuloides</i> sp. 5	9	5	0
*Roach, <i>Rutilus rutilus</i> Blickling Lake, Norfolk (UK)	<i>Buddenbrockia plumatellae</i>	7	2	0
White Fish, <i>Coregonus albula</i> Lake Lucerne (CF)	<i>Tetracapsuloides bryosalmonae</i>	6	5	1
Roach, <i>Rutilus rutilus</i> Lake Lucerne (CF)	<i>Buddenbrockia plumatellae</i>	1	1	0
*Brown Trout, <i>Salmo trutta</i> River Brubach (CF)	<i>Tetracapsuloides bryosalmonae</i>	16	3	0
*Rainbow Trout, <i>Oncorhynchus mykiss</i> River Furtbach (CF)	<i>Tetracapsuloides bryosalmonae</i>	12	2	0
Total	-	83	33	4

Table 4.

Species	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1- <i>T. bryosalmonae</i> KF731712.1 (BS)	-	5/1673	4/1683	4/1683	2/1683	39/1726	39/1579	22/1724	23/1580	247/1726	235/1691	235/1692	244/1695	217/1601	223/1693
2- <i>T. bryosalmonae</i> (this study - host: brown trout - UK)	0.3	-	1/1673	1/1673	2/1673	42/1673	39/1579	25/1673	23/1580	238/1673	236/1668	237/1668	235/1649	219/1601	222/1673
3- <i>T. bryosalmonae</i> (this study - host: brown trout - CF)	0.2	0.1	-	0/1683	2/1683	41/1683	40/1579	24/1683	24/1580	241/1683	238/1683	239/1683	237/1679	218/1601	212/1683
4- <i>T. bryosalmonae</i> (this study - host: rainbow trout - CF)	0.2	0.1	0.0	-	2/1683	41/1683	40/1579	24/1683	24/1580	240/1683	237/1683	238/1683	236/1679	218/1601	222/1683
5- <i>T. bryosalmonae</i> (this study - host: white fish - CF)	0.1	0.1	0.1	0.1	-	41/1683	40/1579	24/1683	24/1580	240/1683	237/1672	238/1672	236/1657	218/1601	222/1671
6- <i>Tetracapsuloides</i> sp.4 KF731727 (BS)	2.3	2.5	2.4	2.4	2.4	-	1/1579	41/1724	41/1580	243/1725	231/1691	231/1692	240/1695	216/1601	222/1693
7- <i>Tetracapsuloides</i> sp.4 (this study - host: roach - UK)	2.5	2.5	2.5	2.5	2.5	0.1	-	41/1579	40/1577	225/1579	226/1579	226/1579	226/1579	217/1577	218/1579
8- <i>Tetracapsuloides</i> sp.5 KF731729 (BS)	1.3	1.5	1.4	1.4	1.4	2.4	2.6	-	1/1580	239/1724	227/1691	2227/1692	236/1695	218/1601	224/1693
9- <i>Tetracapsuloides</i> sp.5 (this study - host: gudgeon - UK)	1.5	1.5	1.5	1.5	1.5	2.6	2.5	0.1	-	222/1580	223/1580	223/1580	223/1580	219/1580	220/1580
10- <i>B. plumatellae</i> KF731682 (BS)	14.5	14.5	14.6	14.5	14.5	14.3	14.5	14.1	14.3	-	1/1691	1/1692	2/1695	67/1601	70/1693
11- <i>B. plumatellae</i> (this study - host: dace - UK)	14.3	14.4	14.5	14.4	14.4	14.1	14.6	13.8	14.4	0.1	-	1/1691	1/1666	68/1601	69/1691
12- <i>B. plumatellae</i> (this study - host: roach - CF)	14.3	14.4	14.5	14.5	14.5	14.1	14.6	13.8	14.4	0.1	0.1	-	2/1666	68/1601	70/1692
13- <i>B. plumatellae</i> (this study - host: roach - UK)	14.2	14.5	14.5	14.5	14.5	14.6	14.6	14.3	14.4	0.1	0.1	0.1	-	69/1601	70/1693
14- <i>Buddenbrockia</i> sp. 2 KJ150268 (H)	14.0	14.1	14.0	14.0	14.0	13.9	14.0	14.0	14.1	4.2	4.3	4.3	4.3	-	5/1601
15- <i>Buddenbrockia</i> sp. 2. (this study - host: loach - UK)	13.5	13.5	13.5	13.5	13.5	13.5	14.0	13.6	14.1	4.2	4.1	4.2	4.2	0.3	-




Fig. 1. Photomicrography of kidney tubules (t) of brown trout collected in the River Stour, Kent, UK, in semi-thin sections stained by toluidine blue (A-B). Transmission electron microscopy (C-H) showing the development of *Tetracapsuloides bryosalmonae* spores in the lumen of the kidney tubules (t) of brown trout (C-D) collected in the River Stour, Kent, UK, and myxozoan development (either malacosporean or sphaerosporid) in white fish (E-H), collected in the Lake Lucerne, Switzerland.




Fig. 2. Photomicrography of kidney tubules (t) of roach collected in the River Stour, Kent, UK, in semi-thin sections stained by toluidine blue (A), and by transmission electron microscopy (B-D) showing spore development of *Tetracapsuloides* sp. 4.




Fig. 3. Photomicrography of kidney tubules (t) of stone loach collected in the River collected in the River Stour, Kent, UK, in semi-thin sections stained by toluidine blue (A-B), and by transmission electron microscopy (C-E) showing spore development of *Buddenbrockia* sp. 2.




Fig. 4. Transmission electron microscopy showing the development of *Sphaerospora* spp. spores in the lumen of kidney tubules of white fish (A-C) collected in the Lake Lucerne, Switzerland and dace (D-F) collected in the River Stour, Kent, UK.