



Jefferys, J. G. R., Ashby-Lumsden, A., & Lovick, T. A. (2020). Cardiac effects of repeated focal seizures in rats induced by intrahippocampal tetanus toxin: bradyarrhythmias, tachycardias and prolonged interictal QT interval. *Epilepsia*. <https://doi.org/10.1111/epi.16479>

Peer reviewed version

Link to published version (if available):
[10.1111/epi.16479](https://doi.org/10.1111/epi.16479)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Wiley at <https://onlinelibrary.wiley.com/doi/abs/10.1111/epi.16479?af=R>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>



Cardiac effects of repeated focal seizures in rats induced by intrahippocampal tetanus toxin: bradyarrhythmias, tachycardias and prolonged interictal QT interval

Journal:	<i>Epilepsia</i>
Manuscript ID	EPI-01028-2019.R2
Manuscript Type:	Full length original research paper
Date Submitted by the Author:	n/a
Complete List of Authors:	Jefferys, John; University of Oxford, Pharmacology; Purdue University, Weldon School of Biomedical Engineering; University of Birmingham, School of Clinical and Experimental Medicine; Charles University Second Faculty of Medicine, Physiology Ashby-Lumsden, Alexander; University of Oxford, Pharmacology; University of Birmingham, School of Clinical and Experimental Medicine Lovick, Thelma; University of Bristol School of Physiology and Pharmacology, Physiology, Pharmacology, and Neuroscience; Purdue University, Weldon School of Biomedical Engineering; University of Birmingham, School of Clinical and Experimental Medicine
Key Words:	Cardiac dysfunction, QT correction, QT prolongation, cardiac arrhythmia, autonomic

1
2
3 **Cardiac effects of repeated focal seizures in rats induced by intrahippocampal tetanus**
4 **toxin: bradyarrhythmias, tachycardias and prolonged interictal QT interval**
5

6 John G. R. Jefferys^{1,2,3,4,6}, Alexander Ashby-Lumsden^{1,2} and Thelma A. Lovick^{3,5},
7

8
9 ¹ Department of Pharmacology, Oxford University, Oxford, UK

10 ² School of Clinical & Experimental Medicine, The University of Birmingham, Birmingham,
11 UK

12 ³ School of Biomedical Engineering, Purdue University, West Lafayette, Indiana

13 ⁴ Department of Physiology, 2nd Medical School, Motol, Charles University, Prague, Czech
14 Republic

15 ⁵ School of Physiology and Pharmacology, The University of Bristol, Bristol, UK

16 ⁶ Corresponding author
17
18
19
20
21
22

23 Contact Information:

24
25 John G. R. Jefferys

26 Department of Pharmacology

27 Oxford University, Oxford, OX1 3QT, UK

28 +44 (1865)556963

29 john.jefferys@pharm.ox.ac.uk
30
31

32 Key Words: Epilepsy, cardiac dysfunction, QT correction, QT prolongation,
33

34 Number of text pages: 14

35 Number of words: 4460

36 Number of references: 43

37 Number of figures: 6

38 Number of tables: 0
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Summary

Objective: To determine electrical changes in the heart in a chronic, non-status model of epilepsy.

Methods: Electrocorticogram (ECoG) and electrocardiogram (ECG) of 9 animals (5 made epileptic by intrahippocampal injection of tetanus neurotoxin (TeNT) and 4 control), are monitored continuously by radiotelemetry for up to 7 weeks.

Results: Epileptic animals develop a median of 168 seizures, with postictal tachycardias reaching a mean of 487 beats/min and lasting a mean of 661 seconds. Ictal changes in heartrate include tachycardia and in the case of convulsive seizures, bradyarrhythmias resembling Mobitz type 1 second degree atrioventricular block; notably the P-R interval increased before block. Postictally the amplitude of T wave increases. Interictally, QT dependence on RR is modest and conventional QT corrections prove ineffective. Interictal QT intervals, measured at a heartrate of 400 bpm, increased from 65 ms to 75 ms, an increase dependent on seizure incidence over the preceding 10-14 days.

Significance: Repeated seizures induce a sustained tachycardia and increase in QT interval of the ECG and evoke arrhythmias including periods of atrioventricular block during Racine Type 4 and 5 seizures. These changes in cardiac function may predispose to development in fatal arrhythmias and sudden death in humans with epilepsy.

Key Point Box:

- Repeated seizures are associated with increases in interictal QT interval, depending on seizure incidence over 10-14 days.
- Conventional QT correction methods prove counterproductive in our study.
- QT at heartrate 400 bpm (QT₄₀₀) are estimated either by selecting periods of ECG at that heartrate, or interpolating from the modest linear QT-RR relationship we find in our unanaesthetized rats.
- Seizures are associated with ictal and prolonged postictal tachycardia.
- Convulsive seizures are associated with arrhythmias including bradyarrhythmias, resembling Mobitz type 1 second degree AV block in humans.

Introduction

Cardiac, along with respiratory problems are the most likely causes of Sudden Unexpected Death in Epilepsy (SUDEP).¹ Repeated seizures have been associated with cardiac abnormalities, including prolongation of the QT interval of the ECG, in humans²⁻⁵ and in experimental rodents.⁶⁻⁹ QT interval reflects the duration of the cardiac ventricular action potential, and is important in cardiac pathophysiology, particularly when prolonged.

Temporal lobe epilepsy features prominently in SUDEP series in adults.^{1; 10} The main risk factor is the number of generalized tonic-clonic seizures;¹¹ arrhythmias during seizures also associate with risk of SUDEP.^{10; 12} The pathophysiological changes underlying this effect remain unresolved, yet an understanding is fundamental to the development of therapeutic interventions to minimize the risk of SUDEP.

The most commonly-used animal models of chronic epilepsy rely on systemic injection of epileptogenic agents.¹³ Direct actions on peripheral or central (e.g. brainstem) sites^{7; 14-16} may induce cardiac and respiratory changes independently of seizures, complicating interpretation of results.¹⁷ These models reveal prolongation of QTc (QT corrected to a standardized heartrate) following chronic epilepsy.^{7;8} The hippocampal kindling model avoids the potential problem of subcortical or systemic effects of the epileptogenic agent, but does not induce spontaneous recurrent seizures; repeatedly evoked seizures in kindling do result in prolonged QTc.⁹ Here, we use the tetanus neurotoxin (TeNT) model,^{18; 19} in which toxin is microinjected directly into the hippocampus. This has the advantage of restricting the toxin's direct effects (on VAMP) to the vicinity of the injection site in the hippocampus,¹⁸ inducing repeated spontaneous self-limiting seizures without inducing status epilepticus (SE). Thus, effects seen in peripheral organs must be due to central seizure-driven events.

The purpose of this study is to determine cardiac changes that occur during and between seizures, and their relationship with repeated seizures. Clinical measurements usually correct

1
2
3 for dependence on heartrate, but this process raises problems which we discuss below and
4 assess in our experimental evidence.^{20; 21} Following a single injection of TeNT into the
5 hippocampus in rats, we record electrocorticograms (ECoG), electrocardiograms (ECG) and
6 seizure-related behaviors in freely-moving animals continuously for periods of weeks.
7
8
9
10
11
12
13
14

15 **Materials and Methods**

16 **Ethics**

17
18
19
20
21 Experimental procedures complied with the UK Animals (Scientific Procedures) Act 1986,
22 and were approved by ethics committees of the Universities of Birmingham and Oxford.
23
24
25

26 **Animal housing**

27
28 Rats are housed two per cage (one experimental animal, one “buddy”) with free access to rat
29 chow and water. Lights turn on automatically from 07.00h -19.00h. Only authorized staff
30 have access to the room and are instructed to minimize disturbance to the rats. Data are
31 collected continuously via radiotelemetry and video recordings for analysis off-line.
32
33
34
35
36

37 **Electrophysiology and induction of focus**

38
39
40 *Surgery.* Under isoflurane anesthesia, male Wistar rats (220-320 g; Charles River) are
41 implanted with Telemetry Research TR50BB radiotelemeters (Millar Instruments, now Kaha
42 Sciences) into the abdomen following the manufacturer’s protocol. During aseptic surgery
43 under isoflurane anaesthesia: wires are tunneled to the chest and head to make electrodes for
44 ECG and ECoG, respectively; ECG leads are sutured onto chest musculature near the right
45 shoulder and left caudal rib cage for lead 2 ECG; bur holes are drilled over the dorsal
46 hippocampus and superficial neocortex 3.5 mm caudal and ± 3.0 mm lateral to bregma and
47 stainless steel ECoG electrodes cemented in place; a further bur hole is drilled over ventral
48 hippocampus 4.3 mm caudal and 4.4 mm lateral to bregma, and a guide cannula cemented in
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 place.²² Incisions are repaired with absorbable sutures. After recording 6-14 days of baseline
4 data, and after rats resumed normal growth (required as evidence of recovery from surgery),
5 they are briefly anesthetized to inject 2.5 ng TeNT (1 μ l in phosphate buffered saline with 2%
6 bovine serum albumin; produced by List Biological Inc, supplied by Quadrantech Diagnostics
7 Ltd) into the ventral hippocampus 7.5 mm below the cortical surface²² over 5 min.

8 Recordings resume following recovery from anesthesia. Previously we showed that injection
9 of TeNT into the hippocampus degrades its molecular target (VAMP) focally in the injected
10 hippocampus,¹⁸ showing that direct effects of TeNT are restricted to near the injection site.

11 *Electrophysiology.* Analogue outputs from the Telemetry Research system are recorded in
12 contiguous 6-hour epochs for up to 9 weeks using a Power 1401 acquisition system and
13 Spike2 software (Cambridge Electronic Design, Cambridge, UK (CED)), with time-locked
14 video recording in three epileptic rats. Seizure detection is assisted by measuring power of
15 the hippocampal recording in the 10-300 Hz band, which increases >4-fold during seizures.
16 The Pan Tompkins method²³ identifies the peaks of R waves to measure instantaneous
17 heartrate (reciprocal of R-R interval).

18 To relate cardiac pathophysiology to Racine score in the 3 rats with simultaneous video
19 recording, we use the Random Sequence Generator at www.random.org to randomly select 3
20 seizures each Racine stage 2-5 from all 43 seizures in one rat and from first, middle and last
21 thirds of seizures in two rats with 164 and 392 respectively. Measurements include ictal and
22 postictal tachycardia, and depth and duration of any abrupt bradyarrhythmia.

23 Finding R wave peaks also allows for averaging ECG waveforms centered on the peak of the
24 R wave. To measure ECG, we average 5-second epochs with steady heartrate, following a
25 period of 60 seconds with as little change in heart rate as possible to minimize QT adaptation
26 and hysteresis.²⁴ Adaptation of QT may occur in our data, but systematic study is impossible
27 given our lack of control over rat behavior and hence heart rate. One rat has step changes in
28

1
2
3 heart rate both before and 18 days after TeNT injection (330 to 421 and 360 to 416 bpm),
4 which allow repeated measures of QT at the new higher heartrate. No progressive changes
5 occur over 30 seconds either before or 18 days after induction of epilepsy. While the data are
6 very limited, they suggest that adaptation is not a major factor under our conditions.
7
8
9

10
11
12
13 QT interval is measured, using averaged ECGs, from the onset of Q to the end of T identified
14 as the intersect between the isopotential line and a tangential line fitted to its descending
15 slope between 80% and 20% amplitude. This method is implemented in a semi-automated
16 script for Spike2 provided by CED (Figure 1A). QT corrections aim to make the resultant
17 QTc independent of RR and heartrate. The Bazett formula, used clinically, normalizes to a
18 heartbeat duration of 1 second which is inappropriate for rodents. Application of the Bazett
19 clinical correction to our data (Figure 1B) reveals that it exacerbates the impact of the RR
20 interval on QTc and increases QTc to unphysiologically high values, which may explain the
21 high values of QTc found in some previous reports using Bazett's correction on rat models.
22 A rat-specific version²⁰ normalizes to a standard heartrate of 400 bpm (RR=150 ms), at least
23 in rats under anaesthesia. The unanaesthetized rats in our study have QT with limited,
24 generally inverse, dependence on RR. This effect is greater for very low heartrates (high RR).
25 The result is that the Kmecova-Klimas correction²⁰ overcorrects QT (Figure 1C).
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Superimposed ECGs confirm that QT is shorter for longer RR (Figure 1D).
The first QT measurements in the epilepsy syndrome examine the time course of changes
during interictal periods, avoiding corrections for heartrate by selecting periods when
heartrate is 400 beats/minute (5-s averages included 33 beats) to produce a measure "QT₄₀₀".
Measurements are restricted to the lights-on period 0700-1900 hours to avoid the risk of a
circadian variation in QT.^{25; 26} Controls comprised of two vehicle-injected rats and two
injected with subthreshold doses of TeNT that failed to induce seizure activity. QT₄₀₀ was
measured between 1100 and 1800 on 6 different days in each rat. There was no significant

1
2
3 difference in QT₄₀₀ between these two groups (ANOVA, $F_{1,22}=2.27$, $P=0.113$), so they were
4
5 pooled as a single set of seizure-free controls.
6
7

8 Although it is not possible, for safety reasons, to blind the experimenter to the toxin or
9
10 vehicle at the time of surgery, files are renamed using numbers generated by the Random
11
12 Sequence Generator at www.random.com before the relationship between QT and RR on
13
14 epileptic rats is measured blind in second series of measurements. One-hour epochs are
15
16 extracted from interictal periods 18 days after injection, and at least 30 min before or after the
17
18 closest seizure, from both the light and dark parts of the circadian cycle. Internal controls are
19
20 from similar epochs the day before injection in the same rats. QT, RR and mean RR over the
21
22 previous 60 seconds are measured blind to the nature of the rat, day and time; graphs of QT
23
24 against RR are made (Sigmaplot version 14) and QT₄₀₀ estimated by interpolation. Statistical
25
26 analyses use SPSS version 23 (IBM); values are presented as mean±SD unless otherwise
27
28 stated.
29
30
31
32
33
34
35
36

37 **Results**

38 **Cardiac electrophysiology**

39
40 Spontaneous seizures start 3-9 days after injection of 2.5 ng TeNT, and the animals are
41
42 recorded for several weeks. During seizures rats exhibit behavioral arrest, vibrissal twitching
43
44 and/or purposeless chewing, which can progress to head nodding and to generalized seizures
45
46 including forelimb clonus, rearing, and rearing and falling seizures (“Racine stage 5”).
47
48
49

50 Seizures recur, often in clusters of 10-40 seizures over 2-5 days separated by one or more
51
52 seizure-free days (Figure 2A). Seizure frequency usually decreases markedly 2-3 weeks after
53
54 induction, but seizures are still seen in some rats at up to 7 weeks, before remission occurred.
55
56
57

58 Total number of seizures experienced by each of the 5 rats ranges from 43 to 382. Individual
59
60

1
2
3 seizures last 83 ± 37 seconds (mean \pm SD; range 19 to 317 seconds; Figure 2A,B), often
4
5 followed by postictal suppression of cortical activity, similar to postictal generalized EEG
6
7 suppression (PGES)²⁷ (Figure 2C, upper trace).
8
9

10 **Heartrate changes during and after seizures**

11
12
13 Resting heart rate (the slowest heart rate measured during 24-hour periods) increases
14
15 progressively during the epileptic syndrome, from 313.8 ± 38.7 to 352.8 ± 43.0 bpm by 3 weeks
16
17 after TeNT injection (paired *t*-test, $p < 0.014$). By 4 weeks, heart rate decreases to 293.6 ± 32.9
18
19 bpm, not significantly different from the starting value.
20
21

22
23 Pronounced changes in heart rate also occur during and following seizures. The most
24
25 consistent is tachycardia, both ictal and postictal. Ictal tachycardia starts within 2-3 seconds
26
27 of electrographic seizure onset and can be interrupted by bradyarrhythmias. In 19 of the 20
28
29 Racine stage 5 seizures in the random sample of videoed rats (see Methods), the seizure starts
30
31 with behavioural immobility lasting 19.5 ± 5.7 seconds associated with an increase in heart
32
33 rate of 38 ± 21 bpm. In the example shown in Figure 2C, this increase terminates with the
34
35 onset of rearing (bradycardia at red arrow in Figure 2C) before falling and rearing repeatedly
36
37 (tonic-clonic convulsions) from 20 to 45 seconds after seizure onset (greyed section on plot
38
39 of heartrate corresponds to bradyarrhythmia, discussed below). The seizure in Figure 2C
40
41 stops at 65 seconds. At the start of the postictal period, the rat experiences tachycardia while
42
43 immobile in a prone position for 1 min until the dashed arrow on Figure 2C, when it starts
44
45 repeated jaw movements. At the solid arrow, about 260 seconds after seizure onset, it
46
47 becomes active and explores its cage for 5 min 42 seconds. This example shows that
48
49 tachycardia occurs independently of postictal active behaviors (exploration and grooming are
50
51 common), when postictal motor activity occurs it prolongs postictal tachycardia, reflecting
52
53 physiological modulation of heart rate. Postictal tachycardia with behavioural inactivity
54
55 occurs in 17 of the random sample of 20 convulsive Racine stage 5 seizures from 3 rats,
56
57
58
59
60

1
2
3 where the postictal rat was visible in the video (see Methods), peaking at 424 ± 46 bpm and
4
5 lasting 75 ± 55 seconds.
6
7

8 In a sample of three rats (including 2 without video), postictal heartrate for all 322 seizures
9
10 increases from 314 ± 39 bpm to reach a mean of 487 ± 34 bpm, ranging from 409 to 578 bpm
11
12 (paired *t*-test $p < 0.001$). It persists for 661 ± 484 seconds (range 28 to 3948 seconds). Only two
13
14 of a series of 127 seizures in three rats with continuous video recording lack postictal
15
16 tachycardia. In one of these seizures, the rat fights with its cage-mate before the seizure and
17
18 its heartrate is already at a high average of 505 bpm over the 5 minutes preceding the seizure,
19
20 which may mask postictal tachycardia. In the other case, heartrate does not change from the
21
22 pre-ictal value of 430 bpm; this seizure has brief and modest motor symptoms (mainly slow
23
24 rearing).
25
26
27
28

29 More detailed statistical analysis of the relationship between postictal tachycardia and Racine
30
31 stage is performed on a random sample of 84 seizures from the 3 rats with video recording
32
33 (see Methods). The peak heart rate is significantly related to Racine stage of the
34
35 corresponding seizure (Figure 2D; ANOVA $F_{3,80} = 5.0$; $P = 0.003$; Dunnett post hoc tests:
36
37 Racine stage 2 differs from stage 5 $P < 0.002$, and from stage 4 $P = 0.008$). For each Racine
38
39 stage maximal postictal heart rate is significantly greater than preictal (paired *t*-tests; $P < 0.001$
40
41 in each case). The increase in postictal heart rate from preictal does not depend on Racine
42
43 stage (ANOVA $F_{3,80} = 0.64$; $P = 0.59$; Supplemental Figure 2A), most likely due to variation in
44
45 preictal rat activity and heart rate.
46
47
48
49

50 Blinded measurements from a random sample of 23 Racine 5 seizures reveal that the T wave
51
52 amplitude expressed as a fraction of the QRS wave is significantly larger during postictal
53
54 tachycardia following tonic-clonic seizures (ratios: preictal, 0.34, early ictal 0.36, after tonic-
55
56 clonic seizures 0.57; ANOVA $F_{2,21} = 6.1$, $P = 0.008$; Dunnett post hoc test against preictal, ictal
57
58 $P = 0.91$, after tonic-clonic seizures $p < 0.01$); QT is unaffected ($P = 0.59$)(Figure 2E,F).
59
60

Bradyarrhythmias

Bradyarrhythmias occur during tonic-clonic components of motor seizures. (Figure 3A). They are absent in non-convulsive seizures (Figure 3B, note different heart rate scale from 3A). In each of the 3 rats in which we have video recording time-locked to the electrophysiological data, we randomly select seizures from each Racine stage 2-5, to assess ictal tachycardia and bradyarrhythmia (see Methods). Ictal tachycardia increases significantly with Racine stage (Figure 3C left; ANOVA $F_{3,80}=5.07$, $P=0.003$; Dunnett post hoc test, Racine stage 2 differs from stages 4 and 5 $P<0.001$). Peak ictal heart rate is significantly greater than preictal for each Racine stage (paired t -tests, $P<0.005$). The dependence of peak ictal heart rate on Racine stage could be related to the corresponding motor behaviours (Racine 2 head nodding, 3 to 5 progressively adding: forelimb clonus, rearing, and rearing and falling); it could also result from the neural circuits recruited during different degrees of seizure generalization. The increase of peak ictal from preictal heart rate was not significantly related to Racine stage (Figure 3C second panel from left; $F_{3,80}=1.6$, $P=0.2$). None of the 20 non-convulsive Racine 2 seizures exhibit bradyarrhythmias, nor do all but one of the 16 Racine 3 seizures (Figure 3B); the exception is an anomalous seizure where the rat makes 4 vertical jumps over a 10-second period starting 18 seconds after the end of a bradyarrhythmia which lacks clear motor signs (Supplemental Figure S1). Fifteen of 27 Racine 4 seizures and 18 of the 21 Racine 5 seizures are associated with substantial drops in heartrate from 367.6 ± 45.9 to 150 ± 111.7 bpm (related-samples Wilcoxon Signed Rank test, $p<0.001$), lasting 17.0 ± 5.3 seconds (Figure 3C, centre and right). Both bradyarrhythmia duration and heart rate decrease from preceding ictal tachycardia depend on Racine stages (Figure 3C centre and right; independent samples Median Test $P<0.001$ in both cases); these measures differ between Racine 5 seizures and each of Racine 2, 3 and 4 (Mann-Witney U test significance shown on boxplots in Figure

1
2
3 3C). The minimum heartrate reached during bradyarrhythmias is 118.7 ± 47.6 bpm, and is
4 independent of Racine scale and rat (data not shown; ANOVA $P=0.72$ and 0.83 respectively).
5
6

7
8 Close inspection of the ECG record during the bradyarrhythmias reveals that the heart misses
9 one beat in two, or two in three, consistent with development of Mobitz type-1 second degree
10 atrio-ventricular (AV) block in humans (Figure 4A-C).²⁸ Note the stable instantaneous
11 frequency of the ictal heartbeat (lower graphs Figure 4A,B), before bradyarrhythmia causes
12 dispersion with lower values settling on half or one third the preceding frequency. Expansion
13 of the dotted box in Figure 4B reveals periods of missed beats often start with an isolated P
14 wave (arrows, Figure 4C). The PR interval increases progressively shortly before the onset of
15 arrhythmias (Figure 4D). The illustrated case ends at the onset of bradyarrhythmia shown in
16 the central panel. Data from a total of 10 Racine stage 5 seizures from the 3 rats with video
17 show significantly increased PR intervals just before bradyarrhythmia onsets (paired *t*-test,
18 $P < 0.001$; Figure 4D, right). In 3/17 seizures periods of asystole up to 3s duration occur
19 (Figure 4B).
20
21
22
23
24
25
26
27
28
29
30
31
32
33

34 Interictal ECG waveform – QT segment

35
36
37
38
39 QT interval is an important measure of cardiac performance; changes are described in both
40 clinical and experimental epilepsies.^{2; 3; 6} Conventions exist for correcting QT for its
41 dependence on heartrate,^{20; 25; 26; 29} but have known limitations. During the present study we
42 find no correction method that satisfactorily succeeds in removing dependence on the R-R
43 interval (“RR”, inverse heartrate) in freely-moving rats (Figure 1), and therefore either
44 restrict measurements to periods when heart rate is stable at 400 bpm (QT_{400} ; Figure 5); or
45 interpolate QT_{400} from measurements at a range of heart rates (Figure 6).
46
47
48
49
50
51
52
53
54

55
56 First we examine the time course of any changes in interictal QT over the course of the
57 epilepsy syndrome. We avoid the risk of complications from circadian variation, by limiting
58
59
60

1
2
3 measurements to the lights-on period (0700-1900 hours, as close to midday as possible).

4
5 QT_{400} difference from the initial measurement removes inter-animal variation (Figure 5B).

6
7 The consistent finding in 5 epileptic rats is that interictal QT_{400} increases progressively during
8 the seizure syndrome, from a pre-injection mean of 63.9 ± 7.7 ms to 71.1 ± 7.3 ms 14 days after
9 injection ($p < 0.001$, paired t -test), and to 71.0 ± 9.9 ms 21 days after injection ($p = 0.022$, paired
10 t -test); means at 7 and 28 days do not differ from pre-injection values, but some individual
11 rats do, reflecting variations in the timing of the seizure syndrome (Figure 5B: symbols with
12 black outlines indicate $> 1.96 * SD$ estimated from 5 pre-injection measurements for each rat).

13
14 QT_{400} relaxes back to baseline 7 days after the last seizure, or as the seizure incidence
15 declines. QT_{400} does not change significantly over 14 days in 4 control rats (71.0 ± 6.6 ms v.
16 69.1 ± 8.1 ms; $p = 0.27$, paired t -test). Typical averaged ECG waveforms (selected as nearest
17 case to their respective group means; Figure 5C) show the prolongation of QT_{400} at 14 days in
18 an epileptic rat. The increase in QT_{400} correlates significantly with the number of seizures
19 over the preceding 3-14 days, but not with the cumulative total of seizures (Supplemental
20 Table 1). The strongest correlation is for a 10-day seizure history (Figure 5D; Pearson
21 correlation coefficient $r = 0.58$, $P = 0.008$ for all seizures and $r = 0.54$, $P = 0.01$ for convulsive
22 Racine 4/5). This suggests QT prolongation depends on seizure incidence over the previous
23 ~10 days.

24
25 To determine whether the epilepsy-related change in QT depends on heart rate we plot
26 interictal QT against RR measured blind during the light and dark periods of the day, before
27 and 18 days after injection of TeNT (see Methods; Figure 5, Supplemental Figure S2).

28
29 Vectors to each point indicate mean RR during the preceding minute to indicate risk of QT
30 adaptation. Interpolation of the QT-RR plots confirm that QT_{400} increases following repeated
31 seizures, from 62.5 ± 8.7 ms to 68.4 ± 2.9 ms 18 days after injection (paired t -test $P < .004$). QT
32 is longer during the light part of the day, 69.6 ± 7.2 ms vs 60.8 ± 9.8 ms in the dark (paired t -

1
2
3 test $P < 0.001$). These plots also reveal that the relationship between QT and RR does not
4
5 change qualitatively after the repeated seizures of the chronic epilepsy model; the increase in
6
7 QT with repeated seizures occurs across the range of heart rates measured (Figure 6;
8
9 Supplemental Figure S2). Note that since the range of heart rates available for measurement
10
11 is determined by the spontaneous activity of individual rats, complete datasets across a wide
12
13 range of heart rates are not possible. Very low levels of activity during the light period before
14
15 induction, for example, mean that moderate and high heart rates are rarely achieved.
16
17
18
19

20 **Discussion**

21 **Application of QT corrections for rat**

22
23 In humans the QT interval is accepted to be influenced by heart rate, requiring correction
24
25 methods to standard heart rate.²⁹ In the present study in rats, which shows tachycardia and
26
27 increased QT interval develop with progression of the seizure syndrome, we find that the
28
29 Bazett correction, which is widely applied to human ECG, is unsuitable for the high heart
30
31 rates of rats. A refinement of Bazett's formula, the Kmecova-Klimas correction, aims to
32
33 overcome this limitation.²⁰ Surprisingly, this correction exacerbates the relationship between
34
35 QT and RR in our data, mainly because the relationship between the raw measurements was
36
37 weak and, if anything, inverse, in marked contrast to the earlier work.²⁰ The earlier study
38
39 differs from ours in using anaesthetized rats, and controlling heart rate using drugs,
40
41 contrasting with the unanaesthetized, freely-moving rats with endogenously-controlled heart
42
43 rate in the present work. A study in unanaesthetized mice, also reports the relationship
44
45 between QT and RR to be relatively flat.³⁰ Another study in rats anaesthetized with
46
47 barbiturate, shows a modified measure of QT (to the peak of the T wave, "QaT") also is
48
49 independent of heart rate.³¹ Neither tribromoethanol²⁰ nor barbiturate³¹ are commonly used as
50
51 anaesthetics in studies of cardiovascular function in small rodents. The impact of anaesthesia
52
53 on the relationship between QT and RR needs further investigation, including for more
54
55
56
57
58
59
60

1
2
3 commonly used anaesthetics. We conclude that at present, most available correction methods
4 may be inappropriate in unanaesthetised rats. In the light of these findings we use the raw
5 data for the analysis of QT either measuring at, or interpolating to, 400 bpm.
6
7

8 **Changes in cardiac function during the seizure syndrome**

9
10 We use intrahippocampal TeNT to induce a seizure profile characterized by repeated self-
11 limiting seizures over a period of several weeks without SE. We are confident that
12 cardiovascular effects are due to locally induced seizure activity rather than direct actions on
13 remote cardiovascular control areas, because the loss of VAMP (the direct effect of TeNT) is
14 restricted to the injected hippocampus.¹⁸ The spontaneous self-limiting seizures of the TeNT
15 model also avoid potential complications from long-term consequences of SE or systemic
16 effects of the epileptogenic agents used in other models. The rat TeNT model resembles
17 temporal lobe epilepsy in humans, in that is associated with development of similar cardiac
18 abnormalities such as baseline (interictal) QT interval prolongation,^{3; 7; 32} increased heart
19 rate,⁴ ictal tachycardia,^{2; 33} and arrhythmias during generalized tonic-clonic seizures.^{10; 12}
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34

35
36 None of the rats die unexpectedly in the present study, so we do not have evidence for a
37 model of SUDEP, but it does provide a useful means to study the manner in which repeated
38 self-limiting seizures can impact on cardiac function. In acute rat models that induce SE,
39 animals do die, but the brain is challenged excessively.^{17; 34} In the present study only a
40 proportion of seizures were convulsive (Racine stages 4-5) and, apart from experiencing
41 seizures, the animals were in good health. In human epilepsy some individuals are at much
42 higher risk of SUDEP than others,³⁵ which may be linked to their inability to overcome the
43 challenge of the intense autonomic disturbances associated with exposure to repeated
44 seizures.³⁶
45
46
47
48
49
50
51
52
53
54

55
56
57 Over the course of the epilepsy syndrome evoked by TeNT in rats, the interictal heart rate
58 increases progressively and QT interval of the ECG becomes prolonged. The increase
59
60

1
2
3 depends on seizure incidence over the preceding 10 days (3-14 day periods all show
4
5 significant correlations with QT). As seizures remit or their incidence wanes, interictal heart
6
7 rate and QT recover over a period of a week; this suggests that seizure clusters may present
8
9 additional risk of repolarization abnormality. Previous reports of QT prolongation following
10
11 two weeks of convulsive seizures induced in a kindling model show increased susceptibility
12
13 to arrhythmias.⁹ A pharmacological investigation into the origin of seizure-related cardiac
14
15 changes is outside the scope of our study. However, others using a systemic kainic acid-
16
17 induced epilepsy model, which also induces increased QT, show that it could be prevented by
18
19 beta adrenergic blockade indicating a sympathetically-mediated effect.¹⁶ These authors⁹ also
20
21 show QT prolongation increases susceptibility to arrhythmias in rats following two weeks of
22
23 kindling-induced Racine 4 and 5 convulsive seizures. The sustained interictal tachycardia that
24
25 develops in the present study is consistent with prolonged sympathoactivation and, with the
26
27 prolongation of interictal QT, suggests that exposure to multiple short, intense seizures
28
29 (seizure clusters) could lead to cardiac dysfunction.
30
31
32
33
34
35

36 A striking finding in the present study is that convulsive (Racine stage 5) seizures, and to a
37
38 lesser extent stage 4 seizures, are associated with arrhythmias that include the development of
39
40 ictal bradyarrhythmia consistent with development of Mobitz type 1 second degree AV heart
41
42 block.
43
44

45 Several reports suggest this effect is parasympathetically mediated. In urethane-anaesthetized
46
47 rats, vagal stimulation induces frequency-dependent cardiac slowing and AV block.³⁷ At the
48
49 highest frequencies employed (50Hz), the heart may stop completely.³⁷ Increases in vagal
50
51 nerve activity during acute SE induced by systemic administration of KA, can also lead to
52
53 bradyarrhythmias and death due to AV block.³⁴ The onset of arrhythmias induced by vagal
54
55 activation is preceded by lengthening of the PR interval consistent with slowed conduction
56
57 though the AV node,³⁸ as it is in our animals. Thus it seems likely that the bradyarrhythmia
58
59
60

1
2
3 and periods of asystole seen during the periodic convulsive seizure activity in the present
4
5 study are a consequence of intense vagal activity.
6
7

8 The combination of sympathetic and parasympathetic over-activation of the heart is
9 particularly dangerous. The sympathetically-mediated increase in QT may increase
10 susceptibility of arrhythmia whilst at the same time vagal activation can evoke AV block.
11
12

13 This pattern of response: ictal tachycardia with periods of bradyarrhythmia and even asystole,
14
15 has been reported in a minority of seizures in individuals judged to be at high risk of
16
17 SUDEP.¹² Whilst this may not pose a grave risk to life in patients with healthy hearts, it could
18
19 be especially dangerous in those in which cardiac function is already compromised.
20
21
22

23
24 In addition to its effect on the heart, vagal activation during increased sympathetic drive has
25
26 been shown to predispose to periods of apnoea.³⁷ Although we did not measure respiration in
27
28 the chronic model in this study, it is interesting that acute SE induced by systemic or
29
30 intrahippocampal injection of kainic acid, also induces intense sympathoactivation
31
32 punctuated by apnoeas, which were often lethal in these models.^{17; 34; 39}
33
34
35
36

37 The mechanism via which focal seizure activity initiated in the hippocampus produces life-
38 threatening disturbances in cardiac and respiratory function is intriguing. Hippocampal
39 seizure activity is known to propagate to the amygdala,⁴⁰ a key forebrain region for
40
41 controlling cardiorespiratory function through its descending connection to key medullary
42
43 cardiovascular control centres.^{41; 42} Studies in patients at high risk for SUDEP show higher
44
45 than normal resting functional interconnectivity between the bilateral amygdalae and bilateral
46
47 hippocampi,^{42; 43} which may pose a risk of exaggerated and abnormal descending influences
48
49 on both cardiac and respiratory control centres. If the seizure syndrome continues,
50
51 pathophysiological changes in cardiac function may develop, which would represent a
52
53 significant risk for development of heart failure or fatal arrhythmias, especially in individuals
54
55 with pre-existing heart disease.
56
57
58
59
60

Acknowledgements

This work was supported by a grant from the James Lewis Foundation through Epilepsy Research UK (JGRJ, TAL), and by Dr Ivan Osorio of Kansas University. We confirm that we have read the Journal's position on issues involved in ethical publication and affirm that this report is consistent with those guidelines.

Conflicts of Interest

Authors declare that they have no conflicts of interest.

References

1. Ryvlin P, Nashef L, Lhatoo SD, et al. Incidence and mechanisms of cardiorespiratory arrests in epilepsy monitoring units (MORTEMUS): a retrospective study. *Lancet Neurol* 2013;12:966-977.
2. Surges R, Scott CA, Walker MC. Enhanced QT shortening and persistent tachycardia after generalized seizures. *Neurology* 2010;74:421-426.
3. Brotherstone R, Blackhall B, McLellan A. Lengthening of corrected QT during epileptic seizures. *Epilepsia* 2010;51:221-232.
4. Lamberts RJ, Blom MT, Novy J, et al. Increased prevalence of ECG markers for sudden cardiac arrest in refractory epilepsy. *J Neurol Neurosurg Psychiatry* 2015;86:309-313.
5. Neufeld G, Lazar JM, Chari G, et al. Cardiac repolarization indices in epilepsy patients. *Cardiology* 2009;114:255-260.
6. Stewart M. An explanation for sudden death in epilepsy (SUDEP). *J Physiol Sci* 2018;68:307-320.
7. Powell KL, Jones NC, Kennard JT, et al. HCN channelopathy and cardiac electrophysiologic dysfunction in genetic and acquired rat epilepsy models. *Epilepsia* 2014;55:609-620.
8. Brewster AL, Marzec K, Hairston A, et al. Early cardiac electrographic and molecular remodeling in a model of status epilepticus and acquired epilepsy. *Epilepsia* 2016;57:1907-1915.

- 1
2
3 9. Bealer SL, Little JG. Seizures following hippocampal kindling induce QT interval
4
5 prolongation and increased susceptibility to arrhythmias in rats. *Epilepsy Res*
6
7 2013;105:216-219.
- 8
9 10. Surges R, Adjei P, Kallis C, et al. Pathologic cardiac repolarization in pharmacoresistant
10
11 epilepsy and its potential role in sudden unexpected death in epilepsy: A case-control
12
13 study. *Epilepsia* 2010;51:233-242.
- 14
15 11. Manolis TA, Manolis AA, Melita H, et al. Sudden unexpected death in epilepsy: The
16
17 neuro-cardio-respiratory connection. *Seizure* 2019;64:65-73.
- 18
19 12. Rugg-Gunn FJ, Simister RJ, Squirrell M, et al. Cardiac arrhythmias in focal epilepsy: A
20
21 prospective long-term study. *Lancet* 2004;364:2212-2219.
- 22
23 13. Lidster K, Jefferys JG, Blümcke I, et al. Opportunities for improving animal welfare in
24
25 rodent models of epilepsy and seizures. *J Neurosci Methods* 2016;260:2-25.
- 26
27 14. Auzmendi J, Buchholz B, Salguero J, et al. Pilocarpine-Induced Status Epilepticus Is
28
29 Associated with P-Glycoprotein Induction in Cardiomyocytes, Electrocardiographic
30
31 Changes, and Sudden Death. *Pharmaceuticals (Basel)* 2018;11.
- 32
33 15. Lai YC, Li N, Lawrence W, et al. Myocardial remodeling and susceptibility to ventricular
34
35 tachycardia in a model of chronic epilepsy. *Epilepsia Open* 2018;3:213-223.
- 36
37 16. Little JG, Bealer SL. beta adrenergic blockade prevents cardiac dysfunction following
38
39 status epilepticus in rats. *Epilepsy Res* 2012;99:233-239.
- 40
41 17. Jefferys JGR, Arafat MA, Irazoqui PP, et al. Brainstem activity, apnea, and death during
42
43 seizures induced by intrahippocampal kainic acid in anaesthetized rats. *Epilepsia*
44
45 2019.
- 46
47 18. Ferecsko AS, Jiruska P, Foss L, et al. Structural and functional substrates of tetanus toxin
48
49 in an animal model of temporal lobe epilepsy. *Brain Struct Funct* 2015;220:1013-
50
51 1029.
- 52
53 19. Walker MC, Jefferys JGR, Wykes RC. Tetanus Toxin. In Pitkanen A, Buckmaster PS,
54
55 Galanopoulou AS, et al. (Eds) *Models of Seizures and Epilepsy*, 2nd Edition;
56
57 2017:589-598.
- 58
59 20. Kmecova J, Klimas J. Heart rate correction of the QT duration in rats. *Eur J Pharmacol*
60
2010;641:187-192.
21. Botelho AFM, Joviano-Santos JV, Santos-Miranda A, et al. Non-invasive ECG recording
and QT interval correction assessment in anesthetized rats and mice. *Pesqui Vet Bras*
2019;39:409-415.

- 1
2
3 22. Paxinos G, Watson C. The Rat Brain in Stereotaxic Coordinates. Academic Press:
4 London; 1998.
- 5
6 23. Pan J, Tompkins WJ. A Real-Time QRS Detection Algorithm. *IEEE Trans Biomed Eng*
7 1985;32:230-236.
- 8
9 24. Malik M, Garnett C, Hnatkova K, et al. Importance of QT/RR hysteresis correction in
10 studies of drug-induced QTc interval changes. *J Pharmacokinet Pharmacodyn*
11 2018;45:491-503.
- 12
13 25. Molnar J, Zhang F, Weiss J, et al. Diurnal pattern of QTc interval: How long is
14 prolonged?: Possible relation to circadian triggers of cardiovascular events. *J Am Coll*
15 *Cardiol* 1996;27:76-83.
- 16
17 26. Smetana P, Batchvarov V, Hnatkova K, et al. Circadian Rhythm of the Corrected QT
18 Interval: Impact of Different Heart Rate Correction Models. *Pacing Clin*
19 *Electrophysiol* 2003;26:383-386.
- 20
21 27. Seyal M, Bateman LM, Li CS. Impact of periictal interventions on respiratory
22 dysfunction, postictal EEG suppression, and postictal immobility. *Epilepsia*
23 2013;54:377-382.
- 24
25 28. Langendorf R, Pick A. Atrioventricular block, type II (Mobitz)--its nature and clinical
26 significance. *Circulation* 1968;38:819-821.
- 27
28 29. Funck-Brentano C, Jaillon P. Rate-corrected QT interval: techniques and limitations. *Am*
29 *J Cardiol* 1993;72:17b-22b.
- 30
31 30. Roussel J, Champeroux P, Roy J, et al. The Complex QT/RR Relationship in Mice.
32 *Scientific reports* 2016;6:25388.
- 33
34 31. Hayes E, Pugsley MK, Penz WP, et al. Relationship between QaT and RR intervals in
35 rats, guinea pigs, rabbits, and primates. *J Pharmacol Toxicol Methods* 1994;32:201-
36 207.
- 37
38 32. Ravindran K, Powell KL, Todaro M, et al. The pathophysiology of cardiac dysfunction in
39 epilepsy. *Epilepsy Res* 2016;127:19-29.
- 40
41 33. Scattolini M, Scorza CA, Cavalheiro EA, et al. Tachycardia and SUDEP: reassuring news
42 about beta blockers. *Epilepsy Behav* 2013;27:510-512.
- 43
44 34. Sakamoto K, Saito T, Orman R, et al. Autonomic consequences of kainic acid-induced
45 limbic cortical seizures in rats: peripheral autonomic nerve activity, acute
46 cardiovascular changes, and death. *Epilepsia* 2008;49:982-996.
- 47
48 35. Barot N, Nei M. Autonomic aspects of sudden unexpected death in epilepsy (SUDEP).
49 *Clin Auton Res* 2019;29:151-160.
- 50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 36. Page T, Rugg-Gunn FJ. Bitemporal seizure spread and its effect on autonomic
4 dysfunction. *Epilepsy Behav* 2018;84:166-172.
5
6 37. Hotta H, Koizumi K, Stewart M. Cardiac sympathetic nerve activity during kainic acid-
7 induced limbic cortical seizures in rats. *Epilepsia* 2009;50:923-927.
8
9 38. Schiereck P, Sanna N, Mosterd WL. AV blocking due to asynchronous vagal stimulation
10 in rats. *Am J Physiol Heart Circ Physiol* 2000;278:H67-73.
11
12 39. Nakase K, Kollmar R, Lazar J, et al. Laryngospasm, central and obstructive apnea during
13 seizures: Defining pathophysiology for sudden death in a rat model. *Epilepsy Res*
14 2016;128:126-139.
15
16 40. Toyoda I, Bower MR, Leyva F, et al. Early activation of ventral hippocampus and
17 subiculum during spontaneous seizures in a rat model of temporal lobe epilepsy. *J*
18 *Neurosci* 2013;33:11100-11115.
19
20 41. Schwaber JS, Kapp BS, Higgins GA, et al. Amygdaloid and basal forebrain direct
21 connections with the nucleus of the solitary tract and the dorsal motor nucleus. *J*
22 *Neurosci* 1982;2:1424-1438.
23
24 42. Cassell MD, Gray TS. The amygdala directly innervates adrenergic (C1) neurons in the
25 ventrolateral medulla in the rat. *Neurosci Lett* 1989;97:163-168.
26
27 43. Allen LA, Harper RM, Kumar R, et al. Dysfunctional Brain Networking among
28 Autonomic Regulatory Structures in Temporal Lobe Epilepsy Patients at High Risk of
29 Sudden Unexpected Death in Epilepsy. *Front Neurol* 2017;8:544.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Supplemental Figure Legends

Supplemental Figure S1.

Anomalous heart rate changes during a Racine 3 seizure. Rat lacks any evidence of clonic or tonic seizures through the period of the Mobitz-type arrhythmia, here dropping 3 in 4 cycles of ECG, with heart rate dropping from 400 to 100 bpm. Later in seizure the rat makes 4 jumps from immobility, which trigger further missed beats.

Supplemental Table S1

Dependence of interictal QT400 on seizure history. Pearson correlation coefficients and corresponding significance values between interictal QT₄₀₀ and the number of seizures during the preceding number of days shown in the row “History”. Seizure counts are divided into Racine 4/5 seizures, or those with ictal bradyarrhythmias where video recording is absent, and all seizures as shown in row “Type”. P<0.01 shown in bold red font; P<0.05 shown in brown font.

Supplemental Figure S2.

QT-RR plots for 4 epileptic rats (separated by thick dashed lines). Each plot presents data for 1-hour epochs during interictal periods 18 days after injection (“epi”), or a corresponding period before injection of TeNT. Epileptic epochs start at least 30 minutes after previous seizure and end at least 30 minutes before the next. “Dark” and “light” indicate the lighting conditions in the monitoring room. Linear regression for the dark pre-injection condition is reproduced in grey in the remaining 3 panels for each rat. Reference lines mark RR = 150 ms and QT for the dark pre-injection case.




Figure 1. QT corrections in rats. A. Averaged ECG waveform illustrates the method for estimating the QT interval. Red dashed line shows isopotential, blue lines indicate T wave amplitude and 80% and 20% heights used for fitting line (light green) to the descending phase of T. B. QT corrections exacerbate the impact of heartbeat duration (RR) on QT. Raw QT is plotted in red. The Bazett correction $QT_c = QT / (RR)^{1/2}$ (green, on extended y axis) is widely used in clinical practice, but is ill-suited to the faster rat heart rate. Kmevcova & Klimas adapt Bazett's correction for the rat (blue): $QT_c = QT / (RR/150)^{1/2}$. All variables in milliseconds. Both corrections increase the dependence of QT on heartrate. C. Raw and Kmevcova & Klimas correction in more detail. D. Averaged ECGs with short (red) and long (black) RR; data points shown in C by stars. Raw QT is little affected by RR until it slows below 160 ms, where it is prolonged. Broken lines show linear fit to T-wave intersecting the isopotential line at points marked by black and red arrows. Voltage scales were adjusted to match R wave amplitudes.

74x192mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 2. Seizures and ECGs during chronic epilepsy induced by intrahippocampal tetanus neurotoxin. A. Seizure profile for one rat, shows seizure duration (y-axis), Racine stage (circles, see key), and time after injection (x-axis). B. Expansion of seizure marked red in (A), 21 days after injection; triangles indicate start and end of seizure. C. Same seizure as (B) on slower timescale, shows electrographic activity (upper trace) and corresponding heart rate (lower trace). Heart rate increase at seizure onset coincides with raising head, red arrow indicates sinus bradycardia related to loss of muscle tone. Arrhythmia during convulsive stage is in grey lines (see also Figures 3, 4). Rat remained prone and immobile after seizure (~50 seconds), until the dashed blue arrow when it started making repeated jaw movements. At the solid blue arrow, it explores its cage. D. Boxplot shows postictal tachycardia increases with Racine stage (median is horizontal bar, interquartile range is grey box and 10-90% are vertical bars). E. Seizure ECoG recording aligned with heart rate and ratio of T to R waves measured from ECG averaged over 2 seconds. T to QRS ratio increases after bradyarrhythmia. In this case T waves remain elevated for approximately 200 seconds after seizure onset. E, insert. Averages of 5-second epochs of ECG aligned by R wave peaks, from before (blue) and after (red) bradyarrhythmia. F. Boxplot of T/QRS ratio for 3 randomly selected Racine 5 seizures from each of the 3 rats with video recording. Values normalized to the preictal T/QRS ratio, and scaled to the grand mean of the preictal ratios. Significant differences indicated by horizontal links: ** $P < 0.008$, *** $P < 0.002$.

165x185mm (300 x 300 DPI)

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60




Figure 3. Ictal tachycardia and bradyarrhythmia during seizures induced by intrahippocampal TeNT. A.

ECoG, ECG and heartrate during a Racine stage 5 convulsive seizure. Note prominent bradyarrhythmia, superimposed on ictal tachycardia, between 31 and 47 seconds on the timescale. B. Similar recording during a Racine stage 3 non-convulsive seizure. A modest bradyarrhythmia superimposed on the ictal tachycardia but heart remains in sinus rhythm. C. For a random sample of seizures classified on the Racine scale (x-axis, 2-5), quantification of, from left: peak ictal tachycardia, increase of postictal tachycardia from preictal heart rate, heart rate decrease during bradyarrhythmia from preceding ictal tachycardia peak, and duration of bradyarrhythmia. Box and whisker plots show median (thick horizontal bar, which appears on the x-axes for Racine scores 1-4), interquartile range (grey box) and 10-90% (whiskers or vertical bars). Outliers are marked by circles or stars, but are included in medians. Significant differences indicated by horizontal links: peak tachycardia by ANOVA with Dunnett post hoc test using Racine 2 as control; bradyarrhythmia heart rate drop and duration by Mann-Witney U Tests comparing Racine 5 with each of 2, 3, 4.

74x149mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 4. Ictal bradyarrhythmias. A. Detail of ECG and heartrate during period of Racine 5 seizure marked by thick line on time axis of Fig 3A. ECG shows deviation from its normal highly regular pattern. Instantaneous frequency (reciprocal of previous R-R interval) reveals intermittent loss of 1 beat in 2 (alternating 400 and 200 bpm) between 14 and 16 seconds on the timescale, and then loss of 2 beats in 3 (149 bpm) until 24 seconds. B. ECG from another rat experiencing a Racine stage 5 convulsive seizure reveals intermittent loss of single beats (alternating 391 and 185 bpm) before an asystole starts at 40 seconds. C. Section of ECG marked by dotted box in B expanded to show waveform. Note P waves at start of section of missed beats (black arrows). D. Increase in P-R interval preceding missed beats: left graph shows an example leading to the onset of the arrhythmia shown in the centre (similar to that shown in C). Dashed grey horizontal lines show 95% confidence intervals, and solid black line shows mean, for P-R during the preceding interictal period. Boxplot shows P-R intervals during ictal sinus rhythm (ictal) and during the beat preceding bradyarrhythmia (pre-A), for 10 Racine stage 5 seizures from 3 rats.

153x196mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 5. Changes in QT duration in epileptic and control rats. A. The QT segment of the ECG, measured from averages centred on R wave peaks during 5-second epochs when heartrate was 400 bpm, shows prolongation during the weeks following TeNT injection (red symbols and dashed red lines represent individual animals; continuous thick orange line, group mean). One rat (inverted triangles) stops seizing after 3 weeks. Local regulations prevented our collecting data for more than a few days after the seizures ended (dashed lines). The remaining 4 continue seizing for more than 4 weeks. Control rats (blue symbols and lines) and non-epileptic rats injected with TeNT (pale blue symbols and lines) failed to show an increase over 2 weeks (controls and non-epileptic rats did not differ significantly and are pooled to a single group mean shown as a black line). B. Same data as A, subtracting initial values to reduce inter-rat variability. Symbols for values more than 1.96 SD from the pre-injection mean QT400 have black outlines. C. Averages of ECGs at 2 weeks for the rat most closely matching the median QT400 for all epileptic rats (red), aligned by R wave with control rat (blue); arrows show duration of QT. D. Change in QT400 plotted against total seizures during 10 days preceding each measurement. Individual rats have their own symbol shapes and colours. Overall these measurements are correlated significantly (see Results and Supplemental Table 1).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

71x210mm (300 x 300 DPI)