



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

## Molecular and genetic control of plant thermomorphogenesis

**Citation for published version:**

Quint, M, Delker, C, Franklin, KA, Wigge, PA, Halliday, KJ & Van Zanten, M 2016, 'Molecular and genetic control of plant thermomorphogenesis', *Nature Plants*, vol. 2, 15190.  
<https://doi.org/10.1038/nplants.2015.190>

**Digital Object Identifier (DOI):**

[10.1038/nplants.2015.190](https://doi.org/10.1038/nplants.2015.190)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Peer reviewed version

**Published In:**

Nature Plants

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



## Molecular and genetic control of plant thermomorphogenesis

Marcel Quint\*<sup>1,2§</sup>, Carolin Delker\*<sup>2</sup>, Keara A. Franklin<sup>3</sup>, Philip A. Wigge<sup>4</sup>, Karen J. Halliday<sup>5</sup>, Martijn van Zanten<sup>6§</sup>

<sup>1</sup> Institute of Agricultural and Nutritional Sciences, Martin Luther University Halle-Wittenberg, Betty-Heimann Str. 5, 06120 Halle (Saale), Germany

<sup>2</sup> Department of Molecular Signal Processing, Leibniz Institute of Plant Biochemistry, Weinberg 3, 06120 Halle (Saale), Germany

<sup>3</sup> School of Biological Sciences, University of Bristol, Bristol BS8 1UG, United Kingdom

<sup>4</sup> The Sainsbury Laboratory, University of Cambridge, Cambridge CB2 1LR, United Kingdom

<sup>5</sup> Synthetic and Systems Biology (SynthSys), University of Edinburgh, CH Waddington Building, Mayfield Road, Edinburgh EH9 3JD, United Kingdom

<sup>6</sup> Molecular Plant Physiology, Institute of Environmental Biology, Utrecht University, Padualaan 8, 3584CH Utrecht, The Netherlands

\* Authors who contributed equally

§ Corresponding authors; Marcel Quint: marcel.quint@landw.uni-halle.de; Martijn van Zanten: m.vanzanten@uu.nl

Temperature is a major factor governing the distribution and seasonal behaviour of plants. Being sessile, plants are highly responsive to small differences in temperature and adjust their growth and development accordingly. The suite of morphological and architectural changes induced by high ambient temperature (up to ~29°C) is collectively called thermomorphogenesis. Understanding the molecular genetic circuitries underlying thermomorphogenesis is particularly relevant in the context of climate change, as this knowledge will be key to breed for thermo-tolerant crop varieties in a rational fashion. Until quite recently the fundamental mechanisms of temperature perception and signalling remained unknown. Our understanding of temperature signalling is now progressing, mainly by exploiting the model plant *Arabidopsis thaliana*. The transcription factor PHYTOCHROME INTERACTING FACTOR 4 (PIF4) has emerged as a major player to regulate phytohormone levels and their activity. To control thermomorphogenesis, multiple regulatory circuits are in place to modulate PIF4 levels, activity, and its downstream mechanisms. Thermomorphogenesis is integrally governed by various light signalling pathways, the circadian clock, epigenetic mechanisms and chromatin-level regulation. In this review we summarize recent progress in the field and discuss how the emerging knowledge in *A. thaliana* may be transferred to relevant crop systems.

2014 was the warmest year since systematic temperature measurements began in 1880<sup>1</sup>. In fact, the ten warmest years on record all occurred after 1998. The 5<sup>th</sup> report of the United Nations Intergovernmental Panel on Climate Change<sup>2</sup> projects an increase of 0.8-4.8°C in global mean surface temperature within the 21<sup>st</sup> century. Such figures are alarming as it is expected that this will strongly affect plant distribution and survival and therefore threaten

biodiversity<sup>3–11</sup>. Some studies already indicate that plant species unable to adjust flowering time in response to temperature are disappearing from certain environments<sup>5</sup> and species tend to shift to higher altitudes and latitudes<sup>12</sup>.

Likewise, crop productivity will probably greatly suffer from global warming, while food production is required to increase drastically to sustain a growing and more demanding world population<sup>9,13–15</sup>. A meta-analysis summarizing more than 1700 studies on the effects of climate change and adaptations on crop yields revealed consensus that in the second half of this century climate warming will likely have a negative effect on yields of important staple crops<sup>13</sup>.

Breeding for crop-level adaptations to cope with high temperatures could potentially reverse this negative trend<sup>9,13–15</sup>. In several plant species mechanisms evolved to adapt growth and morphology to stimulate mitigation of warmth through enhanced evaporative cooling, increased convection and direct avoidance of heat flux from the sun<sup>16–20</sup>. If understood, the underlying molecular processes of these so-called thermomorphogenesis responses could be attractive breeding targets for improving crops to withstand climate warming.

Although abundant literature is available on how plants tolerate extreme heat stress (reviewed in<sup>9,21</sup>), we are only beginning to understand the molecular mechanisms underlying thermomorphogenesis in response to moderately increased temperatures. A key breakthrough was the identification of the bHLH transcription factor PHYTOCHROME INTERACTING FACTOR 4 (PIF4) as a central regulator of ambient temperature signalling in *Arabidopsis thaliana*<sup>22</sup>. Recent findings implicated important roles for light signalling pathways, the circadian clock<sup>23–28</sup>, auxin<sup>22,29–31</sup> and other phytohormones<sup>31–34</sup> in PIF4-mediated temperature-induced growth. Furthermore, epigenetic mechanisms appear at the nexus of induction<sup>35</sup> and attenuation<sup>36</sup> of growth acclimation in response to high ambient temperatures.

In this review, we discuss and integrate recent findings on the molecular networks driving thermomorphogenic adaptations. We will furthermore highlight missing links and suggest how the knowledge on *A. thaliana* could be transferred to relevant crop systems. In addition to thermomorphogenesis, adaptation to high ambient temperature also involves physiological processes such as photosynthetic acclimation, respiration and changes in carbon balance. For discussions of these topics as well as on phenological changes including premature flowering, we refer the reader to reviews elsewhere<sup>20,37–39</sup>.

### **Thermomorphogenesis; growth and developmental processes affected by high temperature**

To the best of our knowledge, the term thermomorphogenesis was coined by Erwin and colleagues<sup>16</sup>, in analogy to photomorphogenesis (light-mediated growth), to describe the effects of temperature on plant morphology. In the context of this review, it is defined as the suite of morphological changes that together likely contribute to adaptive growth acclimation to otherwise detrimental high ambient temperature conditions.

Elongation of the hypocotyl is one of the earliest thermomorphogenic effects seen in seedlings across *A. thaliana* accessions in response to high ambient temperature<sup>22–36,40–50</sup> (Fig. 1a, Table 1). It has been suggested that hypocotyl elongation moves the sensitive meristematic and photosynthetically active tissues away from heat-absorbing soil and may promote cooling by allowing better access to moving air<sup>31</sup>.

Rosette leaves and cotyledons exhibit marked petiole elongation upon sensing of high ambient temperatures<sup>17–20,22,23,28,30,35,36,41,45,50</sup> and move upward; a process called hyponastic growth<sup>18–20,22,36,45,51–54</sup> (Fig. 1a,b, Table 1). It was

argued that hyponasty reduces direct heat flux from the sun and, again, allows a cooling breeze to reach the leaves<sup>17–20</sup>. Together with petiole elongation, hyponasty results in an open rosette structure. High ambient temperature-grown plants exhibiting these phenotypes showed greater transpiration rates and had cooler leaves than their cool-grown counterparts, when both groups were subjected to high temperature conditions<sup>17</sup>. These data suggest that thermomorphogenic adaptations may contribute to high temperature mitigation by enhancing leaf evaporative cooling<sup>17,18</sup>. This idea was supported by mathematical models, which predicted that a combination of petiole elongation and hyponastic growth may operate in concert to sufficiently separate leaves from both the soil and each other to assure optimal transpiration and leaf cooling under well-watered conditions<sup>17,18</sup>. In addition, high temperature-grown plants have fewer stomata and develop smaller and thinner leaves<sup>17,28,45,53,54</sup> (Table 1). These phenotypes may further facilitate cooling by reducing boundary layer thickness, which stimulates heat dissipation by evaporation and convection<sup>17–20</sup>.

### **PIF4 is a hub in ambient temperature signalling**

Changes in plant morphology initiated by high ambient temperature and by vegetation shade are very similar<sup>55</sup>, indicating the possibility of shared signalling elements. This idea led to the identification of the bHLH transcription factor PIF4 as a key regulator of thermomorphogenic phenotypes including hyponasty, hypocotyl and petiole elongation<sup>22,29,30,32,56,57</sup>. As discussed below, PIF4, and to a lesser extent PIF5, performs its pivotal function in high temperature signalling by orchestrating transcriptional changes which subsequently trigger primarily phytohormone-induced elongation responses.

Quickly after shifting plants to high ambient temperature, a notable increase in *PIF4* transcript has been observed, triggering thermomorphogenesis<sup>22,30,32</sup>. However, thermomorphogenesis needs to be precisely timed and restrained to, for example, balance elongation growth versus biomass production<sup>58</sup>. A complex circuitry of PIF4 regulation is therefore at play that includes gene expression, epigenetic regulation, protein stability, protein sequestration, promoter access and promoter competition (Fig. 2). This tight control of PIF4 activity and other coordinating factors is indispensable for the integration of various environmental signals into plant morphogenesis and growth control.

#### **Transcriptional regulation of PIF4**

Expression of *PIF4* itself is rhythmic and tightly regulated by the circadian clock (Fig. 2a)<sup>59–62</sup>. The clock regulates the rhythmic expression of *PIF4* and *PIF5* through repression by the so called evening complex (EC), consisting of the proteins EARLY FLOWERING 3 (ELF3), ELF4 and LUX ARRHYTHMO (LUX)<sup>59,62</sup>. Expression of core clock genes shows temperature-induced alterations in transcription profiles in extended dark periods<sup>25</sup>. However, in diurnal conditions, clock gene expression is largely robust over a wide range of ambient temperatures. This temperature compensation seems to be primarily maintained via the clock components LATE ELONGATED HYPOCOTYL (LHY) and GIGANTEA (GI)<sup>63</sup>. It is possible that clock and temperature information are transmitted to PIF4 directly via ELF3, since the ability of ELF3 to bind target genes is attenuated at 27°C<sup>26</sup>. Interestingly, two recent studies indicated that genetic variation in *ELF3* explains a large part of natural variation in temperature-induced *PIF4* expression and elongation growth among *A. thaliana* accessions<sup>26,28</sup>. When the EC peaks in the early night, *PIF4* expression is suppressed<sup>64,65</sup>. Reduction of EC during the progression of the night then leads to a rise in *PIF4* levels. However, post-dawn decrease of *PIF4* levels suggests the involvement of other transcriptional repressors. As an additional level of regulatory control, ELF3 can also directly bind to PIF4 protein<sup>66</sup>.

In the light, *PIF4* restriction likely involves a similar repression mechanism facilitated at least partially by the bZIP transcription factor LONG HYPOCOTYL 5 (HY5<sup>67-69</sup>; Fig. 2a). *hy5* mutants grown at standard growth temperatures (20°C) show increased *PIF4* expression at mid-day and a transiently increased expression in response to elevated temperature<sup>41</sup>. Genome-wide ChIP analyses have identified *PIF4* promoters as HY5 targets<sup>70</sup> and a temperature-insensitive quadruple *pif* mutant suppressed temperature-hypersensitivity of *hy5* mutants<sup>41</sup>. Interestingly, HY5 protein is less abundant at higher temperatures<sup>69</sup>, which presumably dampens HY5 control of *PIF4* in warm conditions. Thus, temperature-dependent transcriptional release of *PIF4* by reducing HY5 levels, likely via the DETIOLATED 1 (DET1) - CONSTITUTIVE PHOTOMORPHOGENIC 1 (COP1) regulatory cascade<sup>41</sup>, may represent a mechanism to control *PIF4* transcript levels in a light- and temperature-dependent manner.

#### Post-translational regulation of *PIF4* protein levels

In addition to control at the transcriptional level, *PIF4* is also subjected to post-translational control. *PIF4* interacts with several proteins, which can affect its activity or stability. The name-giving interaction with phytochrome B (phyB) in the light, for example, results in phosphorylation and subsequent ubiquitination followed by proteasomal degradation of *PIFs*<sup>71</sup> (Fig. 2b). The kinase BRASSINOSTEROID-INSENSITIVE 2 (BIN2) has also been shown to phosphorylate *PIF4* preferentially in the light, restricting the daytime impact of *PIF4* by depleting protein levels<sup>72</sup>. However, as high temperature triggers accumulation of phosphorylated *PIF4* in red and blue light, light-mediated phosphorylation does not necessarily result in degradation of the protein<sup>58</sup>. Possibly, differential phosphorylation patterns by independent kinases may occur in response to distinct stimuli, resulting in different fates of the protein.

Recently, interaction of *PIFs* with DET1, a repressor of photomorphogenesis, has been shown to stabilize *PIFs* and counteract their degradation<sup>73,74</sup> (Fig. 2b). Whether or not this process directly contributes to the regulation of *PIF* activity in response to elevated temperatures remains to be elucidated. However, *det1* mutants are impaired in temperature-induced hypocotyl elongation<sup>41</sup>, which could very well indicate a dual role of DET1 in temperature-dependent *PIF* regulation via direct interaction/stabilization, and also DET1-COP1-mediated HY5 degradation.

Interaction with other proteins can also sequester free *PIF4* protein, preventing its DNA-binding and downstream transcriptional regulation<sup>48,58,75</sup>. Among these, LONG HYPOCOTYL IN FAR-RED 1 (HFR1), which accumulates in a CRYPTOCHROME 1 (CRY1)-dependent manner, acts as a negative regulator in temperature responses under monochromatic blue light<sup>58</sup>. This process may also contribute to *PIF4* regulation in blue light-rich white light conditions (Fig. 2b).

In addition, *PIF4* access to target promoters seems to be under tight control as well. Here, competition for mutual regulatory DNA-binding sites can occur among *PIF4* and HY5, which differentially affects the transcriptional activity of target genes<sup>69</sup>. As increasing temperatures result in decreased HY5 and increased *PIF4* protein levels<sup>22,32,69</sup>, the alteration in protein ratios can quantitatively affect target gene expression levels.

#### **Thermomorphogenesis depends on *PIF4*-mediated regulation of phytohormone levels and activity**

Phytohormone biosynthesis and signalling genes represent prominent *PIF4* targets<sup>32</sup>, thereby connecting *PIF4* activity

with the long-known essential role of phytohormones in thermomorphogenesis<sup>31</sup> (Figure 2C).

Auxin and auxin signalling are required and sufficient for PIF4-mediated high temperature-induced hypocotyl elongation and other thermomorphogenic responses<sup>29–32</sup>. At high ambient temperatures, free IAA levels in aerial tissues are increased<sup>29–31</sup>. This is likely caused by temperature-mediated binding of PIF4 to promoters, and subsequent activation of auxin biosynthesis genes like *YUCCA 8 (YUC8)*, cytochrome P450 *CYP79B*, and *TRYPTOPHAN AMINOTRANSFERASE OF ARABIDOPSIS 1 (TAA1)*<sup>29,30</sup> (Fig. 2c). In support of this, IAA levels do not increase at high ambient temperatures in *pif4* mutants<sup>29–31</sup>.

Increased intracellular auxin levels initiate gene expression changes via the TRANSPORT INHIBITOR 1/AUXIN SIGNALING F-BOX proteins (TIR1/AFBs) signalling pathway<sup>77</sup>. Auxin binding by a co-receptor complex formed by TIR1/AFBs and members of the AUXIN/INDOLE-3-ACETIC ACID (AUX/IAA) protein family results in the subsequent degradation of AUX/IAAs and the initiation of transcriptional auxin responses<sup>78</sup>. Accordingly, mutants defective in one or more of the partially redundant TIR1/AFBs show reduced temperature-induced hypocotyl elongation<sup>31,41</sup>.

Among the temperature-inducible auxin response genes are the *SMALL AUXIN UP RNA 19-24 (SAUR19-24)* and *SAUR61-68* subfamilies<sup>29,32</sup>. Several members of this gene family have been shown to regulate elongation growth, likely by increasing H<sup>+</sup>-ATPase activity at the plasma membrane<sup>79–81</sup>. Accordingly, the overexpression of stabilized GFP-SAUR19 rescues the thermomorphogenic hypocotyl elongation defect of the *pif4* mutant<sup>29</sup>. Besides SAURs, EXPANSIN cell wall loosening enzymes directly affect cell elongation and interestingly, temperature-induced expression of an *EXPANSIN* gene was positively correlated with heat tolerance in the grass *Agrostis scabra*<sup>82</sup>. Furthermore, *EXPANSIN* expression in response to light and GA has been shown to depend on PIF4<sup>75</sup>, which makes it likely that temperature control of *EXPANSIN* also requires PIF4<sup>44</sup>.

In addition to auxin, brassinosteroids (BR) and gibberellins (GA) play crucial roles in high temperature-induced hypocotyl elongation<sup>22,31–34,48,83,84</sup> (Fig. 2c). The transcription factor *BRASSINAZOLE RESISTANT 1 (BZR1)*, for instance, is involved in the regulation of temperature-induced hypocotyl elongation in a PIF-dependent manner and directly interacts with PIF4<sup>33</sup>. Furthermore, the *det2-1* BR biosynthesis mutant displays defects in thermomorphogenic responses<sup>31</sup>, and pharmacological inhibition of BR signalling inhibits temperature-induced growth<sup>32</sup>. Consistent with the currently understood molecular mechanism for synergistic interaction of auxin and BRs, a highly active BR pathway might sensitize seedlings for the temperature-induced increase in auxin levels<sup>32</sup>. This might be mediated via the regulation of transcription factor activity. PIF4 and BZR1 directly interact with AUXIN RESPONSE FACTOR 6 (ARF6) and enhance its binding to promoters. Accordingly, the BZR1/ARF6/PIF4 (BAP) module synergistically regulates many shared target genes that may ultimately trigger elongation growth<sup>34,84</sup> (Fig. 2c). However, it remains unclear whether ARF6 has a role in thermomorphogenesis and also the exact role of BR requires further investigation.

Gibberellin (GA) presence leads to degradation of growth-repressive DELLA proteins that inhibit PIF action in light signalling<sup>75,85</sup>. Moreover, Stavang and colleagues<sup>32</sup> demonstrated a rapid up-regulation of the major GA biosynthesis genes *AtGA20ox1* and *AtGA3ox1* in *A. thaliana* seedlings subjected to elevated temperatures, whereas the prominent catabolism gene *AtGA2ox1* was down-regulated. Consistent with these observations, detailed mutant analyses showed that both GA biosynthesis and signalling are required for the promotion of thermomorphogenesis<sup>32</sup>. This suggests that the GA pathway is more active at high ambient temperatures, putatively as a result of increased GA levels and release of DELLA-dependent PIF4 sequestering. However, in contrast to auxin, the GA pathway appears not sufficient to induce thermomorphogenesis, since quintuple *della* mutant seedlings still show a partial hypocotyl

elongation response<sup>22,32</sup>. Interestingly, GA-mediated cell elongation requires BRs, auxin, BZR1 and PIF4, and it was shown that DELLA growth repressors directly interact with BZR1 and ARF6<sup>34,83</sup>. GA presence releases DELLA-mediated repression of BZR1 and ARF6 to allow BAP-module function and subsequent induction of hypocotyl elongation<sup>34,83</sup> (Fig. 2c). Hence, GA seems permissive, rather than regulatory, by modulation of PIF4 activity.

### **Multiple signalling pathways converge at PIF4 to balance auxin-mediated thermomorphogenesis**

Tight regulation of PIF4 and its downstream auxin biosynthesis and signalling targets is required to assure that cooling capacity is achieved, while physiological imbalance and exaggerated elongation growth is prevented. Therefore, several signal transduction pathways converge on PIF4 in addition to the (post-)transcriptional regulatory mechanisms discussed above.

One such pathway involves feedback regulation by *AUX/IAA* auxin signalling genes (Fig. 2c). Various *AUX/IAAs* (e.g. *IAA4* and *IAA29*) are induced under high ambient temperatures in a PIF4-dependent manner<sup>22,31</sup>. Auxin-mediated degradation of *AUX/IAAs* and subsequent release of ARF transcription factors is essential for thermomorphogenesis. Yet, the TIR1/AFB-independent direct and rapid induction of the genes encoding *AUX/IAA* transcriptional repressors by PIF4 also provides the possibility of a fast and timely attenuation of the auxin stimulus when auxin levels decrease. Consistent with this idea, gain-of-function mutations in several *AUX/IAAs* (e.g. *SHY2/IAA3* and *IAA19/MSG2*) can suppress PIF4-mediated hypocotyl elongation at high temperatures<sup>30,47</sup>.

A recent study described the involvement of epigenetic silencing of the auxin biosynthesis gene *YUC8* to attenuate thermomorphogenesis<sup>36</sup>. Mutants in the RNA-binding protein FLOWERING TIME CONTROL PROTEIN A (FCA) exhibited increased PIF4 binding to the *YUC8* promoter (Fig. 2c). Accordingly, *fca* mutants displayed increased auxin levels and exhibited enhanced hypocotyl and petiole elongation as well as hyponasty under both control and elevated temperatures<sup>36</sup>. Furthermore, enhanced levels of the activating epigenetic histone mark H3K4me2 on chromatin of the *YUC8* promoter were observed at high temperatures, which was further stimulated in the *fca* mutant background<sup>36</sup>. Taken together, the results suggest that PIF4 binds to the *YUC8* promoter and stimulates auxin biosynthesis driving thermomorphogenesis shortly after high temperature sensing, followed by PIF4-mediated recruitment of FCA. This leads to removal of activating H3K4me2 marks and subsequent dissociation of PIF4 from the *YUC8* locus, resulting in attenuation of thermomorphogenesis<sup>36</sup>.

Additional regulation of PIF4 may be conferred via HLH factors (Fig. 2c). The non DNA-binding HLH factor PHYTOCHROME RAPIDLY REGULATED 1 (PAR1) attenuates high temperature-mediated elongation responses through direct inactivation of PIF4<sup>48</sup>, resulting in decreased high temperature-induced hypocotyl elongation<sup>48</sup>. Furthermore, the BAP module stimulates the expression of another non-DNA-binding HLH factor *PACLOBUTRAZOL RESISTANCE 1 (PRE1)*<sup>34,83</sup>. PRE1 acts as a positive regulator of thermomorphogenesis as part of a module of three HLH/bHLH factors, together with ILI1 BINDING BHLH1 (IBH1) and HOMOLOG OF BEE2 INTERACTING WITH IBH1 (HBI1)<sup>34,44,83</sup>. Sequestration of IBH1 by PRE1 facilitates the binding of HBI1 to the promoters of *EXPANSIN* genes<sup>44</sup>, promoting cell wall loosening and hypocotyl elongation (Fig. 2c). Consistent with this model, high temperature-induced hypocotyl elongation is severely reduced in transgenic lines displaying reduced PRE1/HBI1 or enhanced IBH1 levels<sup>34,44,83</sup>.

### **Modelling-based integration of light, circadian and temperature signals in the control of thermomorphogenesis**

The studies outlined above illustrate that PIF4 associates with a number of proteins, that collectively integrate multiple environmental and endogenous stimuli to control thermomorphogenesis. While we already have detailed knowledge of some molecular events, we are still some way from understanding how the network operates at a whole system level. When striving to do this, lab-to-lab variation in experimental regimes, and limited access to quantitative data, can provide additional obstacles. Thus, linking new and published data to gain a comprehensive understanding of thermo-regulation is not a trivial process. Despite these constraints, mathematical modelling has emerged as a valuable approach to learn how complex biological systems work. Modelling provides a formal means to consolidate knowledge, challenge our current understanding and derive new and experimentally testable hypotheses. Recently, a combination of modelling and experimental approaches was successfully applied to address the complex regulatory circuitry underlying morphogenesis by connecting the circadian clock, light and temperature to identify new regulators and interconnections and to explain regulatory switches in response to multiple conflicting stimuli<sup>27,43,86</sup>.

Initial groundwork in this area was laid by Rausenberger and colleagues<sup>87</sup>, who constructed the first kinetic model for light signalling. This model captured key aspects of phyB photochemistry including photoreceptor protein dynamics to hypocotyl length<sup>87,88</sup>. The model also highlighted the combined network features that were required to deliver fluence rate dependency of phyB. A more recent study extended the Rausenberger<sup>87</sup> model to incorporate PIF control of hypocotyl elongation<sup>43</sup>. This revised model provided a framework to understand how changes in the light and temperature environment alter signalling through the phyB-PIF circuit. The study revealed that temperature has a strong impact on how light regulates hypocotyl elongation by showing that fluence rate-dependent hypocotyl elongation is attenuated at 22°C compared to 17°C. Furthermore, at 27°C increasing fluence rates do not inhibit, but instead, promotes, elongation above a low irradiance threshold. This infers that temperature can completely switch the mode of light action, possibly by increased photoconversion between active Pfr and inactive Pr forms at higher fluence rates, resulting in less efficient phyB signalling. This scenario predicts that phyB would be less effective at degrading PIF proteins at increased fluence rates at 27°C. However, this is not the case, as a strong fluence rate-dependent depletion of PIF4 (and PIF3) protein levels was observed at both 22°C and 27°C<sup>43</sup>. Model analysis provided an alternative hypothesis; that fluence rate-dependent factors are required to modulate PIF activity. At moderate temperatures these factors suppress PIF action, but at higher temperatures they activate PIFs. This hypothesis was partially validated, as HY5 was shown to be a strong PIF suppressor at cooler temperatures, particularly as fluence rates increase<sup>43,69</sup>. Nevertheless, the molecular or biochemical entity that mediates light activation of PIFs at higher temperatures has yet to be determined.

Although such steady-state hypocotyl models provide useful formats to conduct network structure-function analyses, rhythmicity of PIF-mediated hypocotyl elongation requires integration of the circadian clock and natural photoperiods<sup>59,62,60,61</sup>. A study by Seaton and colleagues<sup>27</sup> constructed the first external coincidence model for hypocotyl growth. This was accomplished by integrating the evening complex (EC) and light regulation of PIF4, PIF5 and their direct targets, *ARABIDOPSIS THALIANA HOMEBOX 2 (ATHB2)* and *IAA29*<sup>59,76,61</sup>. This model configuration matched observed photoperiod responses of *ATHB2* and *IAA29* in wild type and simulated clock mutants. As temperature modulates *PIF4* expression through the EC<sup>25,26</sup>, the authors<sup>27</sup> tested whether this response could be captured by the model. By introducing temperature modulation of EC affinity for the *PIF4* promoter, the model was able to match the temperature-induced early rise of *PIF4* expression, and the associated changes in *ATHB2* and *IAA29*, substantiating the proposed mode of thermal *PIF4* regulation through the EC.

Based on the described examples, it is evident that combining modelling and experimental approaches has proved to



be important in deciphering biological complexity. The highlighted studies<sup>27,43,68</sup> provide conceptual frameworks to understand how the mode of PIF4 control by light is switched by temperature; and the temperature-dependent nocturnal rise in PIF4 transcription in a diurnal cycle. The latter study<sup>27</sup> also provides a systems level understanding of how temperature and photoperiodic signals integrate to control growth.

### **Chromatin level regulation at the nexus of thermomorphogenesis**

Temperature influences virtually every biological process and a key feature of investigating the impact of temperature on any given organism is that passive, thermodynamic effects of temperature on biomolecules needs to be separated from active thermal perception and signalling<sup>89</sup>. Among the processes that are tightly controlled by temperature are gene transcription and mRNA degradation. Sidaway-Lee and colleagues noted that both transcription and mRNA decay rates passively increased in response to higher ambient temperatures in *A. thaliana*<sup>90</sup>. In an effort to dissect active and passive thermal regulation, they found that active temperature-directed changes in mRNA abundances could be assigned to temperature-mediated regulation of transcription, rather than mRNA decay<sup>90</sup>. The authors next determined which epigenetic modifications were related to temperature-mediated transcriptional regulation and found that H3K27me3 was associated with genes exhibiting both high and low temperature-dependent transcriptional regulation. This epigenetic mark was depleted from genes showing passive temperature-mediated regulation only<sup>90</sup>. Global changes in several other epigenetic marks, including H3K4me3, H3K9Ac and DNA methylation, were however not inferred in active thermo-regulation of gene expression<sup>90</sup>, but contribution of these marks on specific thermomorphogenesis-regulating genes cannot be excluded. The prominent role for epigenetic modifications in thermomorphogenesis control was recently supported by the above-described example of FCA-mediated H3K4me2 removal from the *YUC8* promoter, which likely restricts PIF4 binding and thereby attenuates thermomorphogenesis<sup>36</sup>.

In addition to epigenetic modifications, chromatin remodelling has a prominent role in thermomorphogenesis. *ACTIN RELATED PROTEIN 6 (ARP6)* controls H2A.Z-nucleosome incorporation into chromatin<sup>91</sup> and plants carrying mutations in *ARP6* display several aspects consistent with a constitutive thermomorphogenic response such as longer hypocotyls and petioles and a transcriptome profile typical for high ambient temperatures, even at lower growth temperatures<sup>35</sup> (Fig. 2c). This implies a role for H2A.Z-containing nucleosomes in thermal regulation of transcription. H2A.Z-nucleosomes are highly enriched at the beginning of genes at the +1 position, adjacent to the transcription start site. For some genes, such as *HEAT SHOCK PROTEIN 70 (HSP70)*, it has been shown that the occupancy of the +1 H2A.Z-nucleosome is rate-limiting for expression. Consequently, *HSP70* was more highly expressed in the *arp6* background compared to wild type at low ambient temperatures. Based on these observations, it was hypothesized that the observed high temperature-induced H2A.Z eviction may provide thermal information to the cell by allowing better accessibility for transcriptional regulators that ultimately orchestrate thermomorphogenesis<sup>35</sup>. H2A.Z eviction therefore appears to enable temperature-dependent expression at least for some - and possibly many - genes. A key question is whether H2A.Z-nucleosome eviction is a direct response to temperature (suggesting it is thermosensory) or whether it is mediated indirectly, for example via a temperature-responsive chromatin remodelling factor. Notably, however, *arp6* mutants still show an increase in hypocotyl elongation at warmer temperatures, suggesting that H2A.Z-nucleosomes themselves do not transmit all temperature information.

### **Future challenges, knowledge transfer and conclusions**

Numerous open questions about temperature signalling and response networks remain to be resolved before comprehensive understanding of how thermomorphogenesis regulation is achieved. Likely, many relevant thermomorphogenesis regulators remain to be identified and their signalling hierarchies need to be investigated to understand how multiple conflicting signals are integrated in coordinated plant growth and development. Importantly, the thermomorphogenesis mechanisms described here are probably operating across a broad range of non-damaging temperatures, beyond the somewhat rigid temperature range of ~20 to ~29°C normally used in thermomorphogenesis research in *A. thaliana* (Table 1). To fully understand plant acclimation to warmer temperatures, a broader temperature range needs to be taken into account. Above all, however, the exact mechanisms by which small changes in ambient temperature are sensed remain enigmatic. H2A.Z eviction and subsequent changes in chromatin suggest a possible temperature sensing mechanism, but this needs to be confirmed. The data are consistent with a model whereby H2A.Z-nucleosomes at the transcriptional start site<sup>35</sup> and/or the gene body<sup>90</sup> may be rate-limiting for the expression of other key genes in the thermomorphogenesis pathway, such as *PIF4* or *PIF4* targets. Alternatively, the enhanced elongation phenotype of *arp6* may arise from a parallel pathway.

Our currently rather limited understanding of ambient temperature perception is in contrast to many other signal transduction pathways. This may be due in part to the involvement of numerous processes, prohibiting the elucidation of a 'temperature receptor'. Among these, temperature effects on transcriptional rates, protein-protein interaction, protein turn-over, changes in subcellular localization and changes in rates of metabolism might intricately contribute to altered physiological read-outs of known and unknown signalling processes. The recent identification of natural *CRYPTOCHROME 2* alleles and their role in thermomorphogenesis<sup>50</sup> emphasizes that the identification of additional, yet unknown rate-limiting and crucial signalling hubs within this network of sensors and response elements constitutes a major challenge, as does experimental design and interpretation. In this respect, the role of metabolism in thermomorphogenesis deserves more attention. Carbon starvation occurs in plants shifted to high ambient temperatures and this correlates with thermomorphogenesis phenotypes<sup>54</sup>. Moreover, PIFs including *PIF4*, are required for sucrose-induced hypocotyl elongation and *PIF5* has been shown to be stabilized by sucrose<sup>92,93</sup>. Sugars induce auxin biosynthesis by stimulating auxin biosynthesis genes<sup>94</sup>, an effect that might potentially be counteracted or enhanced by PIFs depending on specific growth conditions. Such data underscore that temperature, light, sugars, PIFs and auxin are part of a complex, not yet well understood circuitry integrating environmental and metabolic cues into a coordinated growth response. Genetic analysis can be used to provide novel insight into the complex molecular networks underlying thermomorphogenesis, but major advances will require the combination of wet lab genetic, physiological and biochemical approaches together with *in silico* modelling of dynamic structural plant phenotypes and the underlying genetic circuitries.

One important aspect that needs particular consideration is the interaction of thermomorphogenesis with other environmental stresses. The relationship with drought deserves more attention, since thermomorphogenesis facilitates cooling by enhanced transpiration, which is only favourable under well-irrigated conditions<sup>17</sup>. Water is already growth-limiting in many parts of the world<sup>95</sup> and high temperatures and drought often occur simultaneously, suggesting that thermomorphogenic acclimation is not beneficial, and can be even detrimental in these conditions. Accordingly, when combined, high temperatures and drought result in a more severe inhibition of growth in plants than observed if only one individual stress is experienced<sup>53</sup>. Both stresses have impact on growth via partly separate and partly parallel mechanisms that become additive when experienced together. Therefore, it is important to assess

the contribution of thermomorphogenesis-regulatory networks on plant acclimation to other stresses and their combinations.

Climate change already has caused large-scale changes in distribution and behaviour of wild species, and unseasonably hot weather led to global disruptions in crop productivity, for example in 2003 and 2012. Further temperature increases during this century are forecast to exacerbate these problems<sup>3-9,13-15</sup>.

Crop-level adaptations have the potential to reverse projected detrimental effects of climate change on agricultural yield<sup>13-15</sup>. Such adaptations could include the use of alternate varieties or even species, planting times, irrigation and fertilization regimes. Of all possibilities, cultivar adaptations are predicted to have the greatest positive impact on yields under the projected climate change<sup>13</sup>. If understood, one promising and socially accepted way to improve thermomorphogenic acclimation would be allele-mining combined with marker-assisted breeding approaches. In this respect, the general conservation of thermomorphogenesis responses in crop species is certainly promising (Fig. 3). However, in a study on genetic variability in developmental rates in 18 species, including the 14 most cultivated crops world-wide, it was found that temperature dose-response curves of developmental processes are strikingly similar between cultivars/lines even if these originated from very different climates<sup>96</sup>. It is therefore likely that current crop-breeding approaches will need to be complemented with more directed genetic engineering approaches that enable genes from a wider range of backgrounds, as well as potentially synthetically designed genes with optimized temperature response properties, to be introduced into key crops. A considerable advance making this approach feasible is the advent of CRISPR/Cas9 technology enabling genome-wide targeting of genetic alterations. Additionally, it may be necessary to combine multiple genes or entire pathways to obtain desired crop protection, something which may not be feasible with conventional breeding approaches alone.

Potential targets for mining of favourable natural alleles could include the receptor-like kinase *ERECTA*, which was recently shown to play a critical role in high temperature stress tolerance<sup>97</sup>. *ERECTA* likely acts by protecting against temperature-induced cellular damage, since overexpression of *ERECTA* conferred high temperature tolerance to *A. thaliana*, tomato and rice in greenhouse and field conditions, without compromising growth and yield. Also, major thermomorphogenesis regulators such as PIF4 and elements of the EC are good candidates. Allelic variation in ELF3, ELF4, LUX and other clock components, for example, has contributed to the domestication of several crop species in terms of flowering time adaptation<sup>98</sup>. Based on the experimental work in *A. thaliana*, allelic variation of EC components can significantly impact on thermomorphogenesis under controlled environmental conditions<sup>26,28</sup>. It remains to be investigated whether these alleles also cause differential temperature responses under natural environmental conditions and if similar differences can be observed in different crop species. On the bright side, the observation that H2A.Z-nucleosome-mediated temperature responses in the monocot model species *Brachypodium distachyon*<sup>99</sup> are similar to those observed in the dicot *A. thaliana*, suggests that at least some of the major molecular circuitries underlying thermomorphogenesis are functionally conserved.

Meeting future challenges to plant productivity imposed by globally increasing temperatures will require basic research in model plant species as well as applied approaches in crops. Integration of these ends of the spectrum will require directed efforts of the academic plant research community and private companies. Further development of thermomorphogenesis as a research area could ultimately provide efficient and timely leads for the initiation of appropriate breeding efforts to generate much required thermo-tolerant crops.

### **Acknowledgements**

The authors thank Julia Bellstädt for graphical support and Jana Trenner for the photographs in Fig. 3. This work was supported by a grant from the Deutsche Forschungsgemeinschaft to MQ (Qu 141/3-1) and NWO VENI grant 863.11.008 to MvZ.

## References

1. American Meteorological Society. State of the climate in 2014. *Bull. Am. Meteorol. Soc.* **96**, (2015).
2. IPCC. Climate change 2013: The physical science basis. Fifth assessment report. UNEP/WMO.
3. Fitter, A. H. & Fitter, R. S. R. Rapid Changes in Flowering Time in British Plants. *Science* **296**, 1689–1691 (2002).
4. Thuiller, W., Lavorel, S., Araújo, M. B., Sykes, M. T. & Prentice, I. C. Climate change threats to plant diversity in Europe. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 8245–8250 (2005).
5. Willis, C. G., Ruhfel, B., Primack, R. B., Miller-Rushing, A. J. & Davis, C. C. Phylogenetic patterns of species loss in Thoreau’s woods are driven by climate change. *Proc. Natl. Acad. Sci.* **105**, 17029–17033 (2008).
6. Nicotra, A. B. *et al.* Plant phenotypic plasticity in a changing climate. *Trends Plant Sci.* **15**, 684–692 (2010).
7. Bellard, C., Bertelsmeier, C., Leadley, P., Thuiller, W. & Courchamp, F. Impacts of climate change on the future of biodiversity. *Ecol. Lett.* **15**, 365–377 (2012).
8. Peñuelas, J. *et al.* Evidence of current impact of climate change on life: a walk from genes to the biosphere. *Glob. Change Biol.* **19**, 2303–2338 (2013).
9. Bitá, C. & Gerats, T. Plant tolerance to high temperature in a changing environment: scientific fundamentals and production of heat stress tolerant crops. *Front. Plant Sci.* **4**, 273 (2013).
10. Ovaskainen, O. *et al.* Community-level phenological response to climate change. *Proc. Natl. Acad. Sci.* **110**, 13434–13439 (2013).
11. CaraDonna, P. J., Iler, A. M. & Inouye, D. W. Shifts in flowering phenology reshape a subalpine plant community. *Proc. Natl. Acad. Sci.* **111**, 4916–4921 (2014).
12. Pauli, H. *et al.* Recent Plant Diversity Changes on Europe’s Mountain Summits. *Science* **336**, 353–355 (2012).
13. Challinor, A. J. *et al.* A meta-analysis of crop yield under climate change and adaptation. *Nat. Clim Change* **4**, 287–291 (2014).
14. Battisti, D. S. & Naylor, R. L. Historical Warnings of Future Food Insecurity with Unprecedented Seasonal Heat. *Science* **323**, 240–244 (2009).
15. Lobell, D. B. & Gourdjí, S. M. The Influence of Climate Change on Global Crop Productivity. *Plant Physiol.* **160**, 1686–1697 (2012).
16. Erwin, J. E., Heins, R. D. & Karlsson, M. G. Thermomorphogenesis in *Lilium longiflorum*. *Am. J. Bot.* **76**, 47–52 (1989).
17. Crawford, A. J., McLachlan, D. H., Hetherington, A. M. & Franklin, K. A. High temperature exposure increases plant cooling capacity. *Curr. Biol.* **22**, R396–R397 (2012).
18. Bridge, L. J., Franklin, K. A. & Homer, M. E. Impact of plant shoot architecture on leaf cooling: a coupled heat and mass transfer model. *J. R. Soc. Interface* **10**, (2013).
19. van Zanten, M., Pons, T. L., Janssen, J. A. M., Voesenek, L. A. C. J. & Peeters, A. J. M. On the Relevance and Control of Leaf Angle. *Crit. Rev. Plant Sci.* **29**, 300–316 (2010).
20. van Zanten, M., Bours, R., Pons, T. L. & Proveniers, M. C. G. in *Temperature and Plant Development* 49–78 (John Wiley & Sons, Inc, 2014).
21. Kotak, S. *et al.* Complexity of the heat stress response in plants. *Physiol. Metab. Ed. Clint Chapple Malcolm M Campbell* **10**, 310–316 (2007).
22. Koini, M. A. *et al.* High Temperature-Mediated Adaptations in Plant Architecture Require the bHLH Transcription Factor PIF4. *Curr. Biol.* **19**, 408–413 (2009).

23. Nomoto, Y. *et al.* A Circadian Clock- and PIF4-Mediated Double Coincidence Mechanism is Implicated in the Thermosensitive Photoperiodic Control of Plant Architectures in *Arabidopsis thaliana*. *Plant Cell Physiol.* **53**, 1965–1973 (2012).
24. Yamashino, T. *et al.* Verification at the protein level of the PIF4-mediated external coincidence model for the temperature-adaptive photoperiodic control of plant growth in *Arabidopsis thaliana*. *Plant Signal. Behav.* **8**, e23390 (2013).
25. Mizuno, T. *et al.* Ambient Temperature Signal Feeds into the Circadian Clock Transcriptional Circuitry Through the EC Night-Time Repressor in *Arabidopsis thaliana*. *Plant Cell Physiol.* **55**, 958–976 (2014).
26. Box, M. S. *et al.* ELF3 Controls Thermoresponsive Growth in Arabidopsis. *Curr. Biol.* **25**, 194–199 (2015).
27. Seaton, D. D. *et al.* Linked circadian outputs control elongation growth and flowering in response to photoperiod and temperature. *Mol. Syst. Biol.* **11**, (2015).
28. Raschke, A. *et al.* Natural Variants of ELF3 Affect Thermomorphogenesis by Transcriptionally Modulating PIF4-Dependent Auxin Response Genes. *BMC Plant Biol* **15**, 197 (2015).
29. Franklin, K. A. *et al.* PHYTOCHROME-INTERACTING FACTOR 4 (PIF4) regulates auxin biosynthesis at high temperature. *Proc. Natl. Acad. Sci.* **108**, 20231–20235 (2011).
30. Sun, J., Qi, L., Li, Y., Chu, J. & Li, C. PIF4-Mediated Activation of YUCCA8 Expression Integrates Temperature into the Auxin Pathway in Regulating Arabidopsis Hypocotyl Growth. *PLoS Genet* **8**, e1002594 (2012).
31. Gray, W. M., Östin, A., Sandberg, G., Romano, C. P. & Estelle, M. High temperature promotes auxin-mediated hypocotyl elongation in Arabidopsis. *Proc. Natl. Acad. Sci.* **95**, 7197–7202 (1998).
32. Stavang, J. A. *et al.* Hormonal regulation of temperature-induced growth in Arabidopsis. *Plant J.* **60**, 589–601 (2009).
33. Oh, E., Zhu, J.-Y. & Wang, Z.-Y. Interaction between BZR1 and PIF4 integrates brassinosteroid and environmental responses. *Nat Cell Biol* **14**, 802–809 (2012).
34. Oh, E. *et al.* Cell elongation is regulated through a central circuit of interacting transcription factors in the Arabidopsis hypocotyl. *eLife* **3**, e03031 (2014).
35. Kumar, S. V. & Wigge, P. A. H2A.Z-Containing Nucleosomes Mediate the Thermosensory Response in Arabidopsis. *Cell* **140**, 136–147 (2010).
36. Lee, H.-J. *et al.* FCA mediates thermal adaptation of stem growth by attenuating auxin action in Arabidopsis. *Nat Commun* **5**, (2014).
37. Yamori, W., Hikosaka, K. & Way, D. Temperature response of photosynthesis in C3, C4, and CAM plants: temperature acclimation and temperature adaptation. *Photosynth. Res.* **119**, 101–117 (2014).
38. Capovilla, G., Schmid, M. & Posé, D. Control of flowering by ambient temperature. *J. Exp. Bot.* **66**, 59–69 (2014).
39. Verhage, L., Angenent, G. C. & Immink, R. G. H. Research on floral timing by ambient temperature comes into blossom. *Trends Plant Sci.* **19**, 583–591 (2014).
40. Orbovic, V. & Poff, K. L. Growth Distribution during Phototropism of *Arabidopsis thaliana* Seedlings. *Plant Physiol.* **103**, 157–163 (1993).
41. Delker, C. *et al.* The DET1-COP1-HY5 Pathway Constitutes a Multipurpose Signaling Module Regulating Plant Photomorphogenesis and Thermomorphogenesis. *Cell Rep.* **9**, 1983–1989 (2014).
42. Delker, C. *et al.* Natural Variation of Transcriptional Auxin Response Networks in *Arabidopsis thaliana*. *Plant Cell* **22**, 2184–2200 (2010).
43. Johansson, H. *et al.* Arabidopsis cell expansion is controlled by a photothermal switch. *Nat Commun* **5**, 4848 (2014).

44. Bai, M.-Y., Fan, M., Oh, E. & Wang, Z.-Y. A Triple Helix-Loop-Helix/Basic Helix-Loop-Helix Cascade Controls Cell Elongation Downstream of Multiple Hormonal and Environmental Signaling Pathways in Arabidopsis. *Plant Cell* **24**, 4917–4929 (2012).
45. Ibañez, C. *et al.* Developmental plasticity of Arabidopsis thaliana accessions across an ambient temperature range. *bioRxiv* (2015). doi:10.1101/017285
46. Miyazaki, Y. ZEITLUPE positively regulates hypocotyl elongation at warm temperature under light in Arabidopsis thaliana. *Plant Signal. Behav.* **10**, e998540 (2015).
47. Maharjan, P. & Choe, S. High Temperature Stimulates DWARF4 (DWF4) Expression to Increase Hypocotyl Elongation in Arabidopsis. *J. Plant Biol.* **54**, 425–429 (2011).
48. Hao, Y., Oh, E., Choi, G., Liang, Z. & Wang, Z.-Y. Interactions between HLH and bHLH Factors Modulate Light-Regulated Plant Development. *Mol. Plant* **5**, 688–697 (2012).
49. Zhu, W. *et al.* Natural Variation Identifies ICARUS1, a Universal Gene Required for Cell Proliferation and Growth at High Temperatures in Arabidopsis thaliana. *PLoS Genet* **11**, e1005085 (2015).
50. Sanchez-Bermejo, E. *et al.* Genetic Architecture of Natural Variation in Thermal Responses of Arabidopsis. *Plant Physiol.* **169**, 647–659 (2015).
51. Millenaar, F. F. *et al.* Ethylene-Induced Differential Growth of Petioles in Arabidopsis. Analyzing Natural Variation, Response Kinetics, and Regulation. *Plant Physiol.* **137**, 998–1008 (2005).
52. van Zanten, M., Voeseek, L. A. C. J., Peeters, A. J. M. & Millenaar, F. F. Hormone- and Light-Mediated Regulation of Heat-Induced Differential Petiole Growth in Arabidopsis. *Plant Physiol.* **151**, 1446–1458 (2009).
53. Vile, D. *et al.* Arabidopsis growth under prolonged high temperature and water deficit: independent or interactive effects? *Plant Cell Environ.* **35**, 702–718 (2012).
54. Vasseur, F., Pantin, F. & Vile, D. Changes in light intensity reveal a major role for carbon balance in Arabidopsis responses to high temperature. *Plant Cell Environ.* **34**, 1563–1576 (2011).
55. Franklin, K. A. Shade avoidance. *New Phytol.* **179**, 930–944 (2008).
56. Proveniers, M. C. G. & van Zanten, M. High temperature acclimation through PIF4 signaling. *Trends Plant Sci.* **18**, 59–64 (2013).
57. de Wit, M., Lorrain, S. & Fankhauser, C. Auxin-mediated plant architectural changes in response to shade and high temperature. *Physiol. Plant.* **151**, 13–24 (2014).
58. Foreman, J. *et al.* Light receptor action is critical for maintaining plant biomass at warm ambient temperatures. *Plant J.* **65**, 441–452 (2011).
59. Nozue, K. *et al.* Rhythmic growth explained by coincidence between internal and external cues. *Nature* **448**, 358–361 (2007).
60. Niwa, Y., Yamashino, T. & Mizuno, T. The Circadian Clock Regulates the Photoperiodic Response of Hypocotyl Elongation through a Coincidence Mechanism in Arabidopsis thaliana. *Plant Cell Physiol.* **50**, 838–854 (2009).
61. Kunihiro, A. *et al.* PHYTOCHROME-INTERACTING FACTOR 4 and 5 (PIF4 and PIF5) Activate the Homeobox ATHB2 and Auxin-Inducible IAA29 Genes in the Coincidence Mechanism Underlying Photoperiodic Control of Plant Growth of Arabidopsis thaliana. *Plant Cell Physiol.* **52**, 1315–1329 (2011).
62. Nusinow, D. A. *et al.* The ELF4-ELF3-LUX complex links the circadian clock to diurnal control of hypocotyl growth. *Nature* **475**, 398–402 (2011).
63. Gould, P. D. *et al.* The Molecular Basis of Temperature Compensation in the Arabidopsis Circadian Clock. *Plant Cell* **18**, 1177–1187 (2006).
64. Covington, M. F. *et al.* ELF3 Modulates Resetting of the Circadian Clock in Arabidopsis. *Plant Cell* **13**,

- 1305–1316 (2001).
65. Liu, X. L., Covington, M. F., Fankhauser, C., Chory, J. & Wagner, D. R. ELF3 Encodes a Circadian Clock–Regulated Nuclear Protein That Functions in an Arabidopsis PHYB Signal Transduction Pathway. *Plant Cell* **13**, 1293–1304 (2001).
  66. Nieto, C., López-Salmerón, V., Davière, J.-M. & Prat, S. ELF3-PIF4 Interaction Regulates Plant Growth Independently of the Evening Complex. *Curr. Biol.* **25**, 187–193 (2015).
  67. Koornneef, M., Rolff, E. & Spruit, C. J. P. Genetic Control of Light-inhibited Hypocotyl Elongation in *Arabidopsis thaliana* (L.) Heynh. *Z. Für Pflanzenphysiol.* **100**, 147–160 (1980).
  68. Oyama, T., Shimura, Y. & Okada, K. The Arabidopsis HY5 gene encodes a bZIP protein that regulates stimulus-induced development of root and hypocotyl. *Genes Dev.* **11**, 2983–2995 (1997).
  69. Toledo-Ortiz, G. *et al.* The HY5-PIF Regulatory Module Coordinates Light and Temperature Control of Photosynthetic Gene Transcription. *PLoS Genet* **10**, e1004416 (2014).
  70. Lee, J. *et al.* Analysis of Transcription Factor HY5 Genomic Binding Sites Revealed Its Hierarchical Role in Light Regulation of Development. *Plant Cell* **19**, 731–749 (2007).
  71. Lorrain, S., Allen, T., Duek, P. D., Whitelam, G. C. & Fankhauser, C. Phytochrome-mediated inhibition of shade avoidance involves degradation of growth-promoting bHLH transcription factors. *Plant J.* **53**, 312–323 (2008).
  72. Bernardo-García, S. *et al.* BR-dependent phosphorylation modulates PIF4 transcriptional activity and shapes diurnal hypocotyl growth. *Genes Dev.* **28**, 1681–1694 (2014).
  73. Dong, J. *et al.* Arabidopsis DE-ETIOLATED1 Represses Photomorphogenesis by Positively Regulating Phytochrome-Interacting Factors in the Dark. *Plant Cell* **26**, 3630–3645 (2014).
  74. Shi, H. *et al.* Arabidopsis DET1 degrades HFR1 but stabilizes PIF1 to precisely regulate seed germination. *Proc. Natl. Acad. Sci.* **112**, 3817–3822 (2015).
  75. de Lucas, M. *et al.* A molecular framework for light and gibberellin control of cell elongation. *Nature* **451**, 480–484 (2008).
  76. Hornitschek, P. *et al.* Phytochrome interacting factors 4 and 5 control seedling growth in changing light conditions by directly controlling auxin signaling. *Plant J.* **71**, 699–711 (2012).
  77. Quint, M. & Gray, W. M. Auxin signaling. *Curr. Opin. Plant Biol.* **9**, 448–453 (2006).
  78. Delker, C., Raschke, A. & Quint, M. Auxin dynamics: the dazzling complexity of a small molecule’s message. *Planta* **227**, 929–941 (2008).
  79. Chae, K. *et al.* Arabidopsis SMALL AUXIN UP RNA63 promotes hypocotyl and stamen filament elongation. *Plant J.* **71**, 684–697 (2012).
  80. Spartz, A. K. *et al.* The SAUR19 subfamily of SMALL AUXIN UP RNA genes promote cell expansion. *Plant J.* **70**, 978–990 (2012).
  81. Spartz, A. K. *et al.* SAUR Inhibition of PP2C-D Phosphatases Activates Plasma Membrane H<sup>+</sup>-ATPases to Promote Cell Expansion in Arabidopsis. *Plant Cell* **26**, 2129–2142 (2014).
  82. Xu, J., Tian, J., Belanger, F. C. & Huang, B. Identification and characterization of an expansin gene AsEXP1 associated with heat tolerance in C3 Agrostis grass species. *J. Exp. Bot.* **58**, 3789–3796 (2007).
  83. Bai, M.-Y. *et al.* Brassinosteroid, gibberellin and phytochrome impinge on a common transcription module in Arabidopsis. *Nat Cell Biol* **14**, 810–817 (2012).
  84. Wang, W., Bai, M.-Y. & Wang, Z.-Y. The brassinosteroid signaling network — a paradigm of signal integration. *SI Cell Signal. Gene Regul.* **21**, 147–153 (2014).
  85. Feng, S. *et al.* Coordinated regulation of Arabidopsisthaliana development by light and gibberellins. *Nature* **451**, 475–479 (2008).
  86. Hersch, M. *et al.* Light intensity modulates the regulatory network of the shade avoidance response



- in Arabidopsis. *Proc. Natl. Acad. Sci.* **111**, 6515–6520 (2014).
87. Rausenberger, J. *et al.* An Integrative Model for Phytochrome B Mediated Photomorphogenesis: From Protein Dynamics to Physiology. *PLoS ONE* **5**, e10721 (2010).
  88. Chew, Y. H. *et al.* Mathematical Models Light Up Plant Signaling. *Plant Cell* **26**, 5–20 (2014).
  89. Arrhenius, Svante. *Quantitative laws in biological chemistry.* (London :G. Bell, 1915).
  90. Sidaway-Lee, K., Costa, M., Rand, D., Finkenstadt, B. & Penfield, S. Direct measurement of transcription rates reveals multiple mechanisms for configuration of the Arabidopsis ambient temperature response. *Genome Biol.* **15**, R45 (2014).
  91. Deal, R. B., Topp, C. N., McKinney, E. C. & Meagher, R. B. Repression of Flowering in Arabidopsis Requires Activation of FLOWERING LOCUS C Expression by the Histone Variant H2A.Z. *Plant Cell* **19**, 74–83 (2007).
  92. Stewart, J. L., Maloof, J. N. & Nemhauser, J. L. PIF Genes Mediate the Effect of Sucrose on Seedling Growth Dynamics. *PLoS ONE* **6**, e19894 (2011).
  93. Liu, Z. *et al.* Phytochrome interacting factors (PIFs) are essential regulators for sucrose-induced hypocotyl elongation in Arabidopsis. *J. Plant Physiol.* **168**, 1771–1779 (2011).
  94. Sairanen, I. *et al.* Soluble Carbohydrates Regulate Auxin Biosynthesis via PIF Proteins in Arabidopsis. *Plant Cell* **24**, 4907–4916 (2012).
  95. Mueller, N. D. *et al.* Closing yield gaps through nutrient and water management. *Nature* **490**, 254–257 (2012).
  96. Parent, B. & Tardieu, F. Temperature responses of developmental processes have not been affected by breeding in different ecological areas for 17 crop species. *New Phytol.* **194**, 760–774 (2012).
  97. Shen, H. *et al.* Overexpression of receptor-like kinase ERECTA improves thermotolerance in rice and tomato. *Nat Biotech* **33**, 996–1003 (2015).
  98. Nakamichi, N. Adaptation to the Local Environment by Modifications of the Photoperiod Response in Crops. *Plant Cell Physiol.* **56**, 594–604 (2014).
  99. Boden, S., Kavanova, M., Finnegan, E. & Wigge, P. Thermal stress effects on grain yield in *Brachypodium distachyon* occur via H2A.Z-nucleosomes. *Genome Biol.* **14**, R65 (2013).
  100. Hanzawa, T. *et al.* Cellular Auxin Homeostasis under High Temperature Is Regulated through a SORTING NEXIN1–Dependent Endosomal Trafficking Pathway. *Plant Cell* **25**, 3424–3433 (2013).

**Table 1: Thermomorphogenesis in *Arabidopsis thaliana*.** Typical phenotypes associated with thermomorphogenesis, the effect direction: increase ( $\wedge$ ), decrease ( $\vee$ ), or equal (=), the temperature treatment that was commenced in the experiments and the accessions used in the respective studies.

Trait	Effect	Range (°C)	Ref.	Arabidopsis accessions used
Hypocotyl elongation	$\wedge$	17 - 27	43	Col-0
	$\wedge$	16 - 24	45	Col-0, Bay-0, C24, CVi-0, Got-7, Rrs-7, Sha, Ws-2
	$\wedge$	20 - 28	28,29,33,41	Col-0, Ws-2, Ler, Rrs-7, Bay-0, Sha, Sf-2, Zu-0
	$\wedge$	20 - 29	31,32,42,44	Col-0, Ler, <sup>a</sup>
	$\wedge$	22 - 27	26,35	<sup>b</sup> , Col-0
	$\wedge$	22 - 28	22-25,46	Col-0
	$\wedge$	22 - 29	30,47,48	Col-0, Ws-2, Ler
	$\wedge / = / \vee$	23 - 27	49,50	Col-0 ( $\wedge$ ), Sij-4 (=), <sup>c</sup>
	$\wedge$	23 - 28	36	Col-0
	$\wedge$	Various <sup>d</sup>	40	Estland
Petiole elongation	$\wedge$	16 - 24	45	Col-0, Bay-0, C24, CVi-0, Sha, Ws-2
	$\wedge$	20 - 28	28,41	Col-0, Rrs-7, Bay-0, Sha
	$\wedge$	22 - 27	35	Col-0
	$\wedge$	22 - 28	17,22,23	Col-0
	$\wedge$	23 - 28	37	Col-0
	$\wedge$	22 - 29	30	Col-0
Hyponastic growth	$\wedge$	16 - 24	45	Col-0, Bay-0, C24, CVi-0, Got-7, Rrs-7, Sha, Ws-2
	$\wedge$	Various <sup>e</sup>	52	Col-0, Ws-2, Ler
	$\wedge$	22 - 28	22	Col-0
	$\wedge$	23 - 28	36	Col-0
	$\wedge$	20 - 30	51,53,54	<sup>f</sup> , Col-0, Ler, An-1, Cvi-0
Stomatal density	$\vee$	22 - 28	17	Col-0
	$\vee$	20 - 30	53,54	<sup>f</sup>
Leaf area	$\wedge$	20 - 28	28	Bay-0, Sha
	$\vee$	22 - 28	17	Col-0
Leaf thickness	$\vee$	22 - 28	17	Col-0
	$\vee / =$	20 - 30	53,54	<sup>f</sup> , Col-0, Ler, An-1, Cvi-0
Specific leaf area (cm <sup>2</sup> g <sup>-1</sup> )	$\wedge / =$	20 - 30	53,54	<sup>f</sup> , Col-0, Ler, An-1, Cvi-0
Blade length/total leaf length	$\vee / =$	20 - 30	53,54	<sup>f</sup> , Col-0, Ler, An-1, Cvi-0
Root elongation	$\wedge$	16 - 24	45,	Col-0, Bay-0, C24, Cvi-0, Got-7, Rrs-7, Sha, Ws-2
	$\wedge$	23 - 29	100	Col-0

#### Footnotes

- <sup>a</sup> Delker et al., 2010<sup>42</sup> used 20 accessions, that all elongated
- <sup>b</sup> Box et al., 2015<sup>26</sup> used 19 accessions, that all elongated
- <sup>c</sup> Sanchez-Bermejo<sup>50</sup> used 139 accessions with the majority of accessions displaying elongation
- <sup>d</sup> Orbovic & Poff (2007)<sup>40</sup> shifted plants between various temperatures
- <sup>e</sup> Van Zanten et al. 2009<sup>52</sup> used a range between 20 and 42°C
- <sup>f</sup> Effects derived from Vile et al. (2012)<sup>53</sup> are based on averages of 10 accessions, each accessions showed the same trend

## Figure legends

**Figure 1: Typical thermomorphogenesis phenotypes of *Arabidopsis thaliana* plants.** Artist impression of thermomorphogenic phenotypes at the **(a)** young seedling stage and **(b)** vegetative stage. Note the occurrence of temperature-induced hypocotyl and petiole elongation and hyponasty in both seedlings and vegetative plants, resulting in an open rosette structure favouring leaf cooling capacity.

**Figure 2: Simplified model of the central role of PIF4 in the molecular genetic circuitries underlying thermomorphogenesis (center).** **(a)** In darkness, transcriptional regulation of *PIF4* involves gating via the evening complex (EC) of the circadian clock. In the light, transcriptional repression via HY5 is relieved by the COP1-SPA E3 ubiquitin ligase and the COP10-DDB1-DET1 (CDD) complex. **(b)** PIF4 post-translational regulation contributing to temperature signalling involves phosphorylation by yet unidentified kinase activity and sequestering of free PIF4. Whether or not other PIF4-interactors/modifiers known from the light signalling context contribute to temperature signalling, remains to be established. **(c)** PIF4 mediates transcriptional regulation of its target genes via binding to G-box promoter elements. This regulation is counteracted by HY5, which competes for the same binding sites. In addition, FCA can attenuate PIF4-G-box binding by removing H3K4Me2 chromatin marks. Further chromatin modifications such as eviction of H2A.Z-containing nucleosomes have been shown to contribute to thermomorphogenesis. However, whether this process directly affects PIF4-target genes remains to be established. Elongation growth is subsequently triggered via PIF4-mediated induction of the auxin biosynthesis and auxin signalling pathway resulting in SAUR-mediated elongation growth and by a cascade involving PAR1, PRE1, IBH1 and HBI1, ultimately resulting in the induction of *EXPANSIN* genes. Both downstream pathways involve feedback regulations and, at least partially, the transcription factors BZR1 and ARF6 (BAP module) are involved. Other phytohormones are involved in thermomorphogenesis with brassinosteroids (BR) and gibberellic acid (GA) having an essential or permissive signalling function, respectively, involving the DELLA repressor proteins. (A-C) Mechanisms with demonstrated relevance in temperature signalling are depicted by solid black lines while connections known from other biological processes which may potentially contribute to temperature signalling are shown as dashed grey lines.

**Figure 3: Thermomorphogenesis in crop species.** Compared to the situation in the model plant *Arabidopsis thaliana* (Figure 1a), temperature-induced hypocotyl elongation seems widely conserved among crop species. Shown here are cabbage (*Brassica oleracea*) and tomato (*Solanum lycopersicum*). Both have been grown for 7 days at 20°C vs. 7 days at 28°C under long day conditions with 90  $\mu\text{mol m}^{-2} \text{s}^{-1}$  white light.