



# THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### **A chimeric virus created by DNA shuffling of the capsid genes of different subtypes of porcine circovirus type 2 (PCV2) in the backbone of the non-pathogenic PCV1 induces protective immunity against the predominant PCV2b and the emerging PCV2d in pigs**

#### **Citation for published version:**

Matzinger, SR, Opriessnig, T, Xiao, C-T, Catanzaro, N, Beach, NM, Slade, DE, Nitzel, GP & Meng, X-J 2016, 'A chimeric virus created by DNA shuffling of the capsid genes of different subtypes of porcine circovirus type 2 (PCV2) in the backbone of the non-pathogenic PCV1 induces protective immunity against the predominant PCV2b and the emerging PCV2d in pigs', *Virology*, vol. 498, pp. 82-93.  
<https://doi.org/10.1016/j.virol.2016.08.011>

#### **Digital Object Identifier (DOI):**

[10.1016/j.virol.2016.08.011](https://doi.org/10.1016/j.virol.2016.08.011)

#### **Link:**

[Link to publication record in Edinburgh Research Explorer](#)

#### **Document Version:**

Peer reviewed version

#### **Published In:**

Virology

#### **General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

#### **Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1        **A chimeric virus created by DNA shuffling of the capsid genes of**  
2        **different subtypes of porcine circovirus type 2 (PCV2) in the backbone**  
3        **of the non-pathogenic PCV1 induces protective immunity against the**  
4        **predominant PCV2b and the emerging PCV2d in pigs**

6        Shannon R. Matzinger <sup>a</sup>, Tanja Opriessnig <sup>b</sup>, Chao-Ting Xiao <sup>c</sup>, Nicholas Cantanzaro <sup>a</sup>,  
7        Nathan M. Beach <sup>a</sup>, David E. Slade <sup>d</sup>, Gregory P. Nitzel <sup>d</sup>, and Xiang-Jin Meng <sup>a\*</sup>

9        <sup>a</sup>Department of Biomedical Sciences and Pathobiology, College of Veterinary Medicine,  
10        Virginia Polytechnic Institute and State University, Blacksburg, VA; <sup>b</sup>The Roslin  
11        Institute and The Royal (Dick) School of Veterinary Studies, University of Edinburgh,  
12        Midlothian, EH25 9RG, Scotland, UK; <sup>c</sup>Department of Veterinary Diagnostic and  
13        Production Animal Medicine, Iowa State University, Ames, Iowa, USA; <sup>d</sup>Zoetis Inc,  
14        Kalamazoo, MI

16        **Running title:** Shuffled PCV2 protects against emerging strains

17        \*Author for correspondence

18        X.J. Meng, M.D., Ph.D., University Distinguished Professor  
19        College of Veterinary Medicine, Virginia Tech  
20        1981 Kraft Drive, Blacksburg, VA 24061-0913

21        email: xjmeng@vt.edu

22        **Word counts**

23        Text: 5,721; Abstract: 150; Figures: 8; Tables: 2; Supplemental table: 1

1  
2  
3  
4  
5  
6  
7  
8  
9 24  
10  
11 25 **Abstract**  
12  
13 26 Porcine circovirus type 2 (PCV2) is the primary causative agent of porcine circovirus-  
14  
15 27 | associated disease (PCVAD). Available commercial vaccines all target ~~the~~ PCV2a  
16  
17 28 | subtype, although the circulating predominant subtype worldwide is PCV2b, and the  
18  
19 29 | emerging PCV2d subtype is also increasingly associated with PCVAD. Here we  
20  
21 30 | molecularly bred genetically-divergent strains representing PCV2a, PCV2b, PCV2c,  
22  
23 31 | PCV2d, and “divergent PCV2aPCV2e” subtypes by DNA-shuffling of the capsid genes  
24  
25 32 | to produce a chimeric virus representing PCV2 global genetic diversity. When placed  
26  
27 33 | in the PCV2a backbone, one chimeric virus (PCV2-3c114) induced higher neutralizing  
28  
29 34 | antibody titers against different PCV2 subtypes. Subsequently, a candidate vaccine  
30  
31 35 | (PCV1-3c114) was produced by cloning the shuffled 3c114 capsid into the backbone of  
32  
33 36 | the non-pathogenic PCV1. A vaccine efficacy study revealed that chimeric virus PCV1-  
34  
35 37 | 3c114 induces protective immunity against challenge with PCV2b or PCV2d in pigs. The  
36  
37 38 | chimeric PCV1-3c114 virus is a strong candidate for a novel vaccine in pigs infected with  
38  
39 39 | variable PCV2 strains.  
40  
41  
42 40  
43  
44 41 **Keywords:** Porcine circovirus type 2 (PCV2); porcine circovirus-associated disease  
45  
46 42 (PCVAD); DNA shuffling; capsid; vaccine  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9 43 **Introduction**


10 44 Porcine circovirus (PCV) is a small, non-enveloped, single-stranded DNA virus  
11  
12 45 which belongs to the family *Circoviridae* (1). PCV type 1 (PCV1) was originally  
13  
14 46 identified as a cell culture contaminant of the porcine kidney cell line PK-15 in the  
15  
16 47 1970's, and was later found to be non-pathogenic in pigs (2, 3). In 1997, a pathogenic  
17  
18 48 variant designated as PCV type 2 (PCV2) was identified in wasting piglets shortly after  
19  
20 49 weaning (2, 4-9). As more cases were identified worldwide, PCV2 was determined to be  
21  
22 50 the primary causative agent of porcine circovirus-associated disease (PCVAD), which  
23  
24 51 includes a broad spectrum of clinical symptoms such as wasting, reproductive failure,  
25  
26 52 respiratory signs and enteritis, and PCV2 may also have a role in the porcine dermatitis  
27  
28 53 and nephropathy syndrome (10).  
29  
30

31 54 PCV2 is one of the most economically devastating viral pathogens to affect the  
32  
33 55 global pig industry to date, and vaccination has been an effective strategy to reduce the  
34  
35 56 economic losses associated with PCV2 infection (11). Currently, all commercially  
36  
37 57 available inactivated or subunit vaccines ~~target the~~consist of a single PCV2a ~~subtype~~  
38  
39 58 capsid antigen (11-14). However, since 2005, a new subtype, PCV2b, has taken over as  
40  
41 59 the most prevalent PCV2 strain associated with PCVAD cases in the U.S. and other  
42  
43 60 countries (15-17). In addition, newly emerging PCV2d strains (previously referred to as  
44  
45 61 "mutant PCV2b"), have been identified in an increasing number of cases in vaccinated  
46  
47 62 herds worldwide, leading to the speculation by some that the emerging PCV2d strains are  
48  
49 63 able to overcome vaccine protection (18-20). A recent study showed that animals  
50  
51 64 vaccinated with recombinant PCV2a capsid protein had lower viral loads and generated  
52  
53 65 higher neutralizing antibodies against a PCV2d-1 strain than vaccination with either a  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

66 PCV2b or homologous PCV2d-1 recombinant capsid protein, suggesting that PCV2  
67 capsid immunogenicity varies (21). However this could not fully explain how PCV2d  
68 infections are emerging in PCV2a vaccinated herds.

69           Until recently, only three PCV2 subtypes were recognized, including PCV2a,  
70 PCV2b, and PCV2c, the last of which was identified in Denmark during the 2000's, is  
71 recognized but not very prevalent (22, 23). While the majority of the PCVAD cases in  
72 the United States are now associated with PCV2b, the emerging PCV2d subtype has been  
73 slowly increasing in the U.S since its initial discovery in 2012 and is now more prevalent  
74 than PCV2a (24). Although the exact reason for the emergence of PCV2d remains  
75 unclear, it can be commonly found in vaccinated herds, leading to the speculation of  
76 either reduced protection against this emerging PCV2d or vaccination failure of  
77 individual animals (18). While the introduction of PCV2a based vaccine strategies has  
78 resulted in a drastic decline in PCV2 prevalence (25), the increased genetic diversity of  
79 PCV2 strains is concerning, and is suggestive of selective pressure promoting genetic  
80 diversity. In fact, a recent report has demonstrated the increasing genetic diversity  
81 amongst the PCV2d subtype, as the majority of isolates identified from 1999-2011 can be  
82 classified under the subclade "PCV2d-1," and the majority of isolates identified recently,  
83 from 2006-2014, diverge from the PCV2d-1 subclade and are now designated "PCV2d-  
84 2" (24). In addition, *in vitro* evidence suggests distinct antigenic differences among  
85 PCV2 subtypes, which may help explain the emergence of new strains (26, 27).  
86 Therefore, in order to address the concern of emerging PCV2d as well as the predominant  
87 PCV2b now circulating in global swine herds, as well as the possibility for the generation  
88 of increasingly divergent PCV2 strains that cannot be controlled by vaccination with a



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

89 | PCV2a antigen alone, future vaccine strategies should focus on broadening the protection  
90 | of a single vaccine by targeting emerging strains such as PCV2d and the predominant  
91 | PCV2b subtype.

92 | DNA shuffling has been shown to be a powerful tool to introduce genetic  
93 | diversity into the virus of interest (28, 29). In fact, recently our group has successfully  
94 | shuffled the structural genes of porcine reproductive and respiratory syndrome virus  
95 | (PRRSV) and developed chimeric virus vaccine candidates with broadly protective  
96 | properties against heterologous PRRSV strains (30-33). Therefore, in the present study  
97 | we aimed to molecularly breed by DNA shuffling the capsid genes of 5 genetically  
98 | diverse PCV2 subtypes including PCV2a, PCV2b, PCV2c, PCV2d and a capsid sequence  
99 | representing a recently identified divergent PCV2a virus previously referred to as  
100 | “PCV2e.” ~~“PCV2e”~~ The “PCV2e” genotype was originally identified by phylogenetic  
101 | analysis of the capsid sequence (34), but was later determined be included in the  
102 | divergent PCV2a genotype based on full sequence phylogenetic analysis (35)). While the  
103 | “PCV2e” strains identified are not divergent enough from PCV2a strains to be referred to  
104 | as their own genotype, this strain was included in this study to increase genetic diversity  
105 | of the PCV2 capsids utilized for DNA shuffling, and will be referred to as “divergent  
106 | PCV2a” in this paper to separate it from the classic PCV2a strain used in this study. In ~~in~~  
107 | order to create a chimeric virus that can induce broad cross-protection against different  
108 | PCV2 subtypes especially the emerging PCV2d and the currently predominant circulating  
109 | PCV2b.

110 | We were able to successfully generate four viable chimeric viruses with shuffled  
111 | capsid gene sequences in the backbone of PCV2a. An *in vivo* pilot study was first

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

112 conducted in pigs to assess the infectivity and cross-neutralizing activities of these 4  
113 chimeric viruses. The chimeric virus (3c114) exhibiting the highest level of cross-  
114 neutralizing activity against different PCV2 subtypes were subsequently selected for a  
115 challenge and efficacy study in pigs against the currently predominant circulating PCV2b  
116 strain as well as the emerging PCV2d strain. We demonstrated that the capsid-shuffled  
117 chimeric virus 3c114 induces protective immunity in conventional pigs against challenges  
118 with both PCV2b and PCV2d.

119

120 **Materials and Methods**

121 *Cells:* A subclone of the PK-15 cell line that is free of PCV1 contamination was  
122 produced previously by end-point dilution of PK-15 cells (ATCC CCL-33) (36). This  
123 subclone PK-15 cell line was cultured in Minimal Essential Medium (MEM)  
124 supplemented with 10% Fetal Bovine Serum (FBS) and antibiotics and was used in the  
125 serum virus neutralization assay and to propagate all virus stocks for this study.

126

127 *DNA shuffling of the capsid genes from 5 different PCV2 subtypes:* The capsid gene  
128 sequences representing each of the 5 genetically-diversified PCV2 subtypes were selected  
129 for DNA shuffling, including PCV2a (strain 40895, GenBank accession number  
130 AF264042), PCV2b (strain NC16845, accession number GU799576), PCV2c (accession  
131 number EU148503), PCV2d-1 (accession number AY181947), and “[PCV2edivergent](#)  
132 [PCV2a](#)” (accession number EF524533). The PCV2a and PCV2b strains were isolated  
133 from U.S. pigs and described previously (12, 37), while the PCV2c, PCV2d-1, and

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

134 | “divergent PCV2aPCV2e” capsid genes were synthesized by GenScript (Piscataway,  
135 | NJ).

136 |       Traditional DNA shuffling was used to shuffle the 5 different PCV2 capsid genes  
137 | essentially as previously described for PRRSV (31), with slight modifications. Briefly,  
138 | the capsid gene DNAs from each of the five PCV2 strains were mixed in equimolar  
139 | amounts with a total of 5 µg DNA and diluted in 50 µl of 50 mM Tris-HCl (pH 7.4) and  
140 | 10 mM MgCl<sub>2</sub>. The mixture was incubated at 15°C for 3 min with 0.15 U of DNase I  
141 | (Sigma). DNA fragments ranging from 50 to 150 bp in size were purified from 2%  
142 | agarose gels, and subsequently added to the Pfu PCR mixture consisting of 1X Pfu  
143 | buffer, 0.2 mM each deoxynucleoside triphosphate (dNTP), and 0.06 U Pfu polymerase.  
144 | A PCR program without using primers (95°C for 4 min; 40 cycles of 95°C for 30s, 60°C  
145 | for 30s, 57°C for 30s, 54°C for 30s, 51°C for 30s, 48°C for 30s, 45°C for 30s, 42°C for  
146 | 30s, and 72°C for 2 min; and finally, 72°C for 7 min) was performed to reassemble the  
147 | digested DNA fragments. Subsequently, specific primers flanking the shuffled PCV2  
148 | capsid region, UniRep-F and 2aORF2-R (**Table S1**), were used to amplify the shuffled  
149 | PCV2 capsid using Pfu Ultra II Hotstart PCR Master Mix (Agilent Technologies) per the  
150 | manufacturer’s instructions (95°C for 4 min, 10 cycles of 95°C for 30s, 50°C for 30s,  
151 | 72°C for 30s, 25 cycles of 95°C for 30s, 54°C for 30s, 72°C for 30s, and finally 72°C for  
152 | 7 min).

153 |  
154 | ***Construction of infectious DNA clones of chimeric PCV2a and PCV1 viruses with***  
155 | ***shuffled PCV2 capsid genes:*** The shuffled capsid gene product libraries were cloned  
156 | into the blunt end cloning vector, pCR-Blunt II, using the Zero Blunt® TOPO® PCR



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

157 Cloning kit (Life Technologies, Carlsbad), per manufacturer’s instructions. Selected  
158 clones were sequenced and analyzed for DNA shuffling efficiency, and well-shuffled  
159 capsid genes containing regions from all 5 PCV2 subtypes were amplified and  
160 subsequently cloned into the infectious DNA clone backbone of the PCV2a strain 40895  
161 by fusion PCR, essentially as previously described (38). Briefly, the shuffled PCV2  
162 capsids were amplified using primers UniRep-F and 2aORF2-R (**Table S1**). The PCV2a  
163 infectious DNA clone backbone sequence was amplified in two fragments that flank the  
164 PCV2 capsid region using primers SacII-uni-F and UniRep-R, and primers 2aORF2F and  
165 SacII-uni-R, for PCV2a fragments 1 and 2, respectively (**Table S1**). All three PCR  
166 reactions were performed using ACCUZYME MIX™ (Bioline) at 95°C 10 min, 35  
167 cycles of 95°C for 30s, 54°C for 30s, and 68°C for 1.5 min. The first fusion PCR was  
168 performed with the PCV2 fragment 1 and the shuffled PCV2 capsid sequence using the  
169 external primers SacII-uni-F and 2aORF2-R. Subsequently, a second fusion PCR reaction  
170 was performed with the product of the first fusion PCR reaction and the PCV2a fragment  
171 2, using the external primers SacII-uni-F and SacII-uni-R (**Table S1**). All fusion PCR  
172 reactions were performed using ACCUZYME MIX™ at 95°C 10 min, 35 cycles of 95°C  
173 for 30s, 60°C for 30s, and 68°C for 4 min. The full-length chimeric PCV2a containing  
174 each individual shuffled PCV2 capsid was amplified, and cloned into the pCR-Blunt II  
175 TOPO plasmid using the Zero Blunt® cloning kit to produce infectious DNA clones of  
176 chimeric PCV2a with shuffled capsid genes.

177         The shuffled PCV2 capsid 3c114 was cloned into the infectious DNA clone  
178 backbone of the non-pathogenic PCV1 to create the vaccine candidate PCV1-3c114 by a  
179 similar fusion PCR protocol. Briefly, the shuffled PCV2 capsid 3c114 was amplified

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

180 using primers PCV1-BB-F and PCV1-DS-ORF2-R (**Table S1**). The infectious DNA  
181 clone PCV1 backbone sequence was amplified from the PBSK+ plasmid containing  
182 PCV1 in two fragments that flank the PCV1 capsid region using primers M13F (-20) and  
183 PCV-BB-R, and primers PCV-DS-ORF2-F and M13R, for PCV1 fragments 1 and 2,  
184 respectively (**Table S1**). All three PCR reactions were performed using Platinum® PCR  
185 Supermix (Thermo Scientific) at 94°C 3 min, 35 cycles of 94°C for 30s, 55°C for 30s,  
186 and 68°C for 1 min. Fusion PCR was performed first with the PCV1 fragment 1 and the  
187 shuffled PCV2 capsid 3c114 fragment using the external primers M13F and PCV1-DS-  
188 ORF2-R (**Table S1**). A second fusion PCR reaction was performed with the product of  
189 the first fusion PCR reaction and PCV1 fragment 2, using the external primers M13F and  
190 M13R. The full-length chimeric PCV1 virus containing the shuffled capsid 3c114 was  
191 cloned into pCR-Blunt II TOPO using the Zero Blunt® cloning kit to produce the  
192 infectious DNA clone of vaccine candidate chimeric PCV1 virus 3c114.

193

194 ***Preparation of virus stocks:*** The infectious virus stocks of PCV2b strain NC16845, U.S.  
195 PCV2d-2 strain JX535296, and each of the PCV2a capsid-shuffled chimeric viruses were  
196 produced by transfecting PK-15 cells with concatemered viral genomes from the  
197 respective infectious DNA clones. Briefly, the respective PCV2 genomes were excised  
198 from pCR-Blunt II TOPO by SacII digestion, concatemered, and transfected into PK-15  
199 cells to determine the viability and infectivity by immunofluorescence assay (IFA) as  
200 previously described (36, 37, 39). The virus stocks for the chimeric PCV1-2a and  
201 chimeric PCV1 containing shuffled 3c114 capsid (PCV1-3c114) were prepared similarly

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

202 as described above except that the viral genome was excised from the pCR-Blunt II  
203 TOPO vector by digestion with KpnI prior to concatemerization.

204

205 ***Determination of the infectivity and cross-neutralizing activities of the PCV2 capsid-***

206 ***shuffled viruses:*** To initially identify viable PCV2 capsid-shuffled viruses with improved  
207 cross-neutralizing activities against different PCV2 subtypes, we first conducted a pilot  
208 pig infection study with a limited number of animals (n=3). A total of 18, 4-week-old,  
209 cross-breed conventional pigs were purchased from a commercial farm that is known to  
210 be free of PRRSV and M. hyo without active PCV2 circulation as determined by  
211 regular PCV2 PCR on selected batches of pigs. Sows have low amounts of antibodies  
212 against PCV2 or are seronegative and we selected litters from negative sows without  
213 cross-fostering. The piglets were randomly assigned to six groups of 3 pigs each, and  
214 each group of pigs was housed separately. Prior to inoculation, each pig was weighed,  
215 bled, and confirmed to be negative for PCV2 by PCR and serology. Five groups were  
216 inoculated intramuscularly each with 5 ml ( $10^{3.66}$  TCID<sub>50</sub>/mL) of either chimeric virus  
217 PCV1-2a or one of the four PCV2 capsid-shuffled viruses (PCV2-3c113, PCV2-3c114,  
218 PCV2-3c14-2, or PCV2-3c112-2). One group was mock-inoculated similarly with 5 mL of  
219 PBS buffer (**Table 1**). Blood was collected weekly, and animals were monitored for  
220 seroconversion to PCV2 capsid antibodies by ELISA and evidence of PCV2 infection by  
221 qPCR. Animals were necropsied at 56 days post-infection (dpi). The weekly serum  
222 samples were used to perform serum virus neutralization test against strains representing  
223 different PCV2 subtypes (data not shown for 0-49 dpi). The animal study was approved  
224 by Virginia Tech IACUC.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

225

226 ***Serum virus neutralization assay:*** Serum samples collected from infected pigs were  
227 tested for neutralizing antibody titers against the wild-type PCV2a, PCV2b, PCV2d-1,  
228 and PCV2d-2 strains by IFA. Briefly, the serum samples were serially diluted 1:2 in PBS  
229 and mixed with 150 TCID<sub>50</sub> of PCV2a, PCV2b, PCV2d-1, or PCV2d-2 virus stocks,  
230 respectively, at an equal volume ratio and incubated for 1 hr at 37°C. The serum-virus  
231 mixture was then added to PK-15 cells in a 96 well plate in duplicate. After 72 hrs  
232 incubation at 37°C, an IFA was performed using pig sera against PCV2a diluted 1:1000,  
233 as the primary antibody and FITC-conjugated goat anti-pig IgG (KPL) diluted 1:50 as the  
234 secondary antibody. The 50% serum neutralizing antibody titers were determined as the  
235 highest dilution at which there was 50% or greater reduction in virus titer compared with  
236 the average of the serum from PBS control pig group at that dilution.

237

238 ***Vaccination efficacy and challenge study in conventional pigs:*** The virus containing  
239 shuffled capsid 3cl14 in the backbone of PCV2a induced significantly higher neutralizing  
240 antibody responses against different PCV2 strains. Therefore, the shuffled capsid  
241 sequence 3cl14 was subsequently cloned into the infectious DNA clone backbone of non-  
242 pathogenic PCV1 to produce a PCV1-3cl14 shuffled capsid chimeric virus as the vaccine  
243 candidate. Subsequently, a pig challenge study was conducted to evaluate the efficacy of  
244 the candidate PCV1-3cl14 chimeric virus vaccine against infection with currently  
245 predominant circulating PCV2b as well as the emerging PCV2d-2. This experiment was  
246 a subset of a larger study. However, wild type exposure prevented completion and  
247 analysis of other groups.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

248 Briefly, a total of 32, 3-week-old, cross-breed conventional pigs were purchased  
249 from a commercial farm that is known to be free of PRRSV and *M. hyopneumoniae*, and  
250 is negative for PCV2. The animal study was approved by Iowa State University IACUC  
251 as well as by Virginia Tech IACUC. The piglets were randomly assigned to 4 groups of  
252 8 pigs each. Prior to inoculation, each pig was weighed, bled, and confirmed to be  
253 negative for PCV2. Groups 1 and 2 pigs were each vaccinated intramuscularly (IM) in  
254 the neck region with 5 ml of the candidate PCV1-3c114 chimeric virus vaccine ( $10^{3.7}$   
255 TCID<sub>50</sub>/mL per pig). Groups 3 and 4 pigs were each mock-vaccinated IM with 5 ml PBS  
256 buffer (**Table 2**). All animals were monitored daily for clinical signs including wasting,  
257 respiratory distress, and behavioral changes such as lethargy and inappetence. Blood  
258 samples were collected prior to inoculation, and weekly thereafter from each pig through  
259 42 days post-vaccination (dpv).

260 At 42 dpv, groups 1 (vaccinated) and 3 (mock-vaccinated) pigs were each  
261 challenged with  $10^{4.8}$  TCID<sub>50</sub> (2.5 ml intranasally and 2.5ml IM) of the PCV2b NC16845  
262 virus strain, and groups 2 (vaccinated) and 4 (unvaccinated) were each similarly  
263 challenged with  $10^{4.8}$  TCID<sub>50</sub> of the PCV2d-2 JX535296 virus strain. Blood samples  
264 were collected weekly through 20 days post-challenge (dpc) (or 62 dpv), at which time  
265 all pigs were weighed and necropsied. A panel of serum and tissue samples was collected  
266 for quantification of viral DNA loads and for histological examination of PCV2-  
267 associated lesions.

269 **Gross pathology and histopathology evaluation:** Necropsies were performed at 20 dpc  
270 on all pigs in a treatment status blinded fashion. Estimates of macroscopic lung lesions

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

271 (ranging from 0 to 100% of the lung affected) and lymph node size (ranging from 0  
272 [normal] to 3 [four times the normal size]) were obtained for each pig (40, 41). Sections  
273 of lung, lymph nodes (superficial inguinal, mediastinal, tracheobronchial, and  
274 mesenteric), tonsil, heart, thymus, kidney, spleen, and liver were collected during  
275 necropsy and processed routinely for histological examination and PCV2  
276 immunohistochemistry (IHC) (Iowa State University Veterinary Diagnostic Lab). Also,  
277 samples of tracheobronchial lymph node (TBLN) were collected from each pig for DNA  
278 extraction and quantification of PCV2 viral genomes by real-time quantitative PCR.  
279 Microscopic lesions in the lymphoid tissues, lungs, heart, liver, kidney, ileum, and colon  
280 were scored in a treatment status blinded manner, as described previously (40).  
281 Specifically, lymph nodes, spleen, and tonsil were evaluated for presence and degree of  
282 lymphoid depletion and histiocytic replacement.

283

#### 284 *Quantitative PCR to quantify viral DNA loads in serum and tissues*

285 For both animal experiments we used a previously published protocol to extract DNA  
286 from serum and lymph node samples and a previously published qPCR SYBR green  
287 assay to quantify viral loads in these samples (37, 42). For the pilot infection study  
288 (**Table 1**) and for the challenge experiment (**Table 2**), PCV2 specific primers were used  
289 to amplify a conserved region spanning the origin of replication and a portion of the  
290 replicase gene, as previously reported (37), using primers PCV2-83F and PCV2-83R  
291 (**Table S1**). For the detection of the PCV1-3cl.14 vaccine strain in the challenge study  
292 (**Table 2**), primers PCV1-qRepF and PCV1-qRepR primers (**Table S1**) were used to  
293 amplify only the PCV1 backbone based vaccine virus DNA.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

294

295 **Serology:** A PCV2-specific ELISA using PCV2a capsid antigen (Iowa State University  
296 Veterinary Diagnostic Lab) was used to detect anti-PCV2 ORF2 IgG in each serum  
297 sample as previously described (43).

298

299 **Sequence confirmation of virus recovered from infected pigs:** DNA extracts from  
300 serum samples collected at 20 dpc from selected pigs in each group were tested by PCR  
301 for PCV2 capsid sequences, and the amplified PCR products were sequenced to verify  
302 that the virus recovered from the infected pigs was the same virus inoculated into the  
303 animals. PCR primers Unirep-F and 2aORF-2 were used to amplify the PCV2 capsid  
304 gene in these samples using the same PCR program as described above for cloning  
305 (**Table S1**). Additionally, DNA extracts of TBLN tissues from selected pigs in each  
306 group were also tested to confirm that the virus detected by PCR from infected pigs was  
307 the same virus that was inoculated into the animals. PCV2b was amplified and  
308 sequenced using primers specific for PCV2b as previously described (37). The PCV2d-2  
309 vDNA was amplified and sequenced using the same forward primer as for PCV2b and a  
310 PCV2d-specific reverse primer NB-56-m2b (**Table S1**).

311

312 **Statistical Analysis:** Statistical analysis was performed using Prism v6.0 (Graphpad, La  
313 Jolla CA). A one-tailed t-test was used to analyze statistical significance between two  
314 groups, while a one-way ANOVA and then t-tests corrected for multiple comparisons  
315 were used to determine significance between three or more groups.

316

## Results

*Generation of infectious chimeric viruses containing the shuffled capsid from 5 genetically distinct PCV2 strains:* Traditional DNA shuffling was used to molecularly breed the capsid genes from five genetically distinct PCV2 strains representing different subtypes PCV2a PCV2b, PCV2c, and PCV2d-1, as well as “~~divergent PCV2a~~PCV2e” (~~divergent PCV2a~~)(**Fig. 1**). Although the general consensus is that previously classified “~~divergent PCV2a~~PCV2e” virus isolates do not diverge enough from identified PCV2a strains to be considered their own subtype (35), a “~~divergent PCV2a~~PCV2e” capsid sequence was chosen to help increase the genetic diversity of the resulting shuffled capsid. The capsid gene sequences from these 5 strains were shuffled using DNase I digestion and reassembled by PCR without primers. A PCR product of the expected size was then generated after a second round of PCR with specific primers spanning the capsid gene. The shuffled capsid gene library was then cloned into the infectious clone backbone of PCV2a (strain 40985) to screen for viable viruses. Of the more than 50 clones with “well-shuffled” capsids (containing regions from all 5 parental PCV2 strains), only 4 of them successfully rescued infectious virus when transfected into PK-15 cells (data not shown).

The four viruses with shuffled capsids contain a range of combinations of the genetic signatures of PCV2 genomes from all 5 parental strains (**Fig. 1**). The majority of the unique amino acid signatures introduced into the shuffled capsids originated from PCV2c, which is not surprising since PCV2c is the most genetically distinct of the 5 parental strains, based on a phylogenetic analysis (**Fig. 2**). Therefore, we demonstrated



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

339 here that traditional DNA shuffling successfully generated viable infectious chimeric  
340 viruses with shuffled capsid genes from 5 different PCV2 subtypes.

341

342 *PCV2-3c114 with shuffled capsid genes induces cross-neutralizing antibodies against*

343 *different PCV2 subtypes:* To determine the viability and screen for the best virus with

344 shuffled capsids for subsequent challenge and efficacy study, we experimentally infected

345 conventional pigs with each of the four viruses (PCV2-3c113, PCV2-3c114, PCV2-

346 3c14\_2, and PCV2-3c112) as well as with the chimeric PCV1-2a virus (12). Serum

347 samples were collected prior to infection and weekly thereafter, and all animals were

348 monitored for seroconversion to PCV2a capsid by an ELISA (**Table 1**). All animals

349 experimentally inoculated with PCV1-2a or with PCV2-3c114 seroconverted to PCV2

350 antibodies by 49 days post-inoculation (dpi), however only 2 out of 3 animals in the

351 PCV2-3c112\_2 and 1 of 3 pigs inoculated with either virus PCV2-3c14 or PCV2-3c14\_2

352 were seropositive at 49 dpi (**Table 1**).

353 Serum samples collected from 56 dpi were tested by a serum virus neutralization

354 assay in PK15 cells for cross-neutralizing antibodies against wild-type PCV2a, PCV2b, a

355 PCV2d-1, and PCV2d-2 virus strains (**Fig. 3**). The neutralization assay was not

356 performed against the parental PCV2c and divergent PCV2aPCV2e strains because

357 PCV2c viruses have not associated with PCV2-induced disease and attempts to grow the

358 divergent PCV2aPCV2e wild type virus in PK-15 cells was unsuccessful in our hands

359 (data not shown). Infections of pigs with 3 PCV2 viruses with shuffled capsid genes

360 (PCV2-3c113, PCV2-3c14\_2, and PCV2-3c112) did not induce higher levels of


361 neutralizing antibody when compared to the chimeric PCV1-2a virus which is the basis

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

362 for the current Foster<sup>TM</sup> PCV commercial vaccine. However, infection of pigs with the  
363 chimeric virus PCV2-3c114 with shuffled capsid genes from different PCV2 subtypes  
364 induced significantly higher neutralizing antibody titers against PCV2a and PCV2d-2  
365 when compared to PCV1-2a (p<0.05) (**Fig. 3**). In addition, although not statistically  
366 significant, the chimeric virus PCV2-3c114 also induced higher levels of neutralizing  
367 antibody than the PCV1-2a against both PCV2b and PCV2d-2. Taken together, this pilot  
368 animal study suggests that the viruses with shuffled capsid genes are viable and  
369 infectious in pigs, and that one shuffled capsid virus PCV2-3c114 induces significantly  
370 higher levels of neutralizing antibodies against genetically distinct PCV2 strains when  
371 compared to the other chimeric viruses as well as to the PCV1-2a vaccine virus.  
372 Therefore, the virus PCV2-3c114 was selected for the subsequent challenge and efficacy  
373 study in pigs to evaluate its potential use as a novel vaccine.

374

375 *The chimeric virus PCV1-3c114 induces protective immunity in conventional pigs*  
376 *against challenge with PCV2b and PCV2d-2.* PCV2a is the genomic backbone for the  
377 virus PCV2-3c114. Therefore, in order to produce a novel vaccine candidate, we  
378 subsequently transferred the shuffled capsid gene from the virus PCV2-3c114, identified  
379 in the initial cross-neutralization study, to the genomic backbone of the non-pathogenic  
380 PCV1 to produce a new chimeric virus PCV1-3c114 with a shuffled capsid. To assess  
381 whether the chimeric virus PCV1-3c114 vaccine candidate protects against challenge with  
382 different PCV2 subtypes, two groups of pigs (n=8) were each vaccinated with the PCV1-  
383 3c114 chimeric virus, and another two groups of pigs (n=8) were mock-vaccinated with  
384 PBS as controls (**Table 2**). Blood samples were taken weekly and animals were



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

385 monitored for seroconversion to PCV2 capsid antibody. At 42 days post-vaccination, one  
386 group of vaccinated and one group of mock-vaccinated animals were challenged with the  
387 predominant field strain PCV2b currently circulating in swine herds worldwide.  
388 Similarly, one vaccinated group and one mock-vaccinated group of pigs were challenged  
389 with the emerging PCV2d-2 virus. Blood samples were taken weekly after challenge and  
390 all animals were necropsied at 20 dpc.

391 As expected, pigs in the two vaccinated groups started to seroconvert to PCV2  
392 capsid antibody by 42 dpv, whereas mock-vaccinated groups did not seroconvert until 7-  
393 14 dpc with PCV2b or PCV2d-2 (or 49 or 56 dpv, **Table 2, Fig. 4**). A qPCR assay  
394 targeting the PCV1 replicase gene (ORF1) was used to test for PCV1-3c114 viral DNA  
395 from weekly sera, but PCV1-3c114 viral DNA was undetectable and below the detection  
396 limit of the assay in any group after vaccination (data not shown). This is consistent with  
397 previous reports of the attenuated chimeric PCV1-2 virus infections in pigs (12, 37).

398 Only 2 out of 8 animals vaccinated and subsequently challenged with PCV2b had  
399 detectable viremia, and only at 14 dpc, compared to 4 and 7 out of 8 PCV2b challenge  
400 control animals at 14 and 20 dpc, respectively (**Table 2**). This difference was statistically  
401 significant, as the vaccinated and PCV2b challenged group had significantly lower levels  
402 of viral DNA loads in sera at 20 dpc, compared to mock-vaccinated and PCV2b  
403 challenged animals ( $p<0.01$ ) (**Fig. 5**). For animals vaccinated and subsequently  
404 challenged with PCV2d-2, 1/8 at 14 dpc and 2/8 at 20 dpc had detectable viremia, while  
405 7/8 PCV2d-2 challenged control animals were positive for serum viral DNA at 14 dpc  
406 and 20 dpc (**Table 2**). Also, the vaccinated and PCV2d-2 challenged group had serum  
407 viral DNA loads that were significantly reduced at 14 and 20 dpc ( $p<0.001$ ,  $p<0.05$ ,

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

408 respectively), as compared to PCV2d-2 challenge only controls (**Fig. 5**). All vaccinated  
409 and subsequently challenged groups had significantly lower levels of PCV2 viremia at  
410 the peak of virus replication compared to control groups. In addition, all vaccinated and  
411 subsequently challenged groups had significantly lower levels of detectable PCV2 DNA  
412 in lymph nodes compared to mock-vaccinated and challenged groups (PCV2b =  $p < 0.001$ ,  
413 PCV2d-2 =  $p < 0.0001$ , **Fig. 6**). These results indicated that vaccination with PCV1-3c114  
414 chimeric virus significantly reduces the level of virus replication in pigs when challenged  
415 with the predominant PCV2b subtype or with an emerging PCV2d-2 strain.

416 In addition to reducing viral DNA loads in sera and lymphoid tissues, vaccinated  
417 animals also had a decreased PCVAD lesion score compared to unvaccinated animals  
418 (**Fig. 7**). Vaccinated pigs that were subsequently challenged with PCV2b had  
419 significantly reduced pathological lesion scores for all measures of PCVAD, which  
420 includes lymphoid depletion and histiocytic replacement in lymph nodes, spleen, and  
421 tonsil tissues, as compared to unvaccinated but PCV2b challenged controls (**Fig. 7**).  
422 Similarly, pigs vaccinated and subsequently challenged with PCV2d-2 had significantly  
423 lower pathological lesion scores for lymph node measures, as well as tonsil lymphoid  
424 depletion (**Fig. 8A, 8B, 8E**) as compared to unvaccinated but PCV2d-2 challenged  
425 controls. Consistent with the results for serum and lymph node viral DNA detection, both  
426 vaccinated and subsequently challenged groups had significantly lower viral antigen  
427 scores in lymph node, spleen, and tonsil, compared to challenge only controls (**Fig. 8**).  
428 Overall, these results suggest that vaccination with PCV1-3c114 chimeric virus vaccine  
429 candidate protects against two genetically distinct and relevant PCV2 strains, the

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

430 predominant PCV2b subtype currently circulating in pig farms worldwide and the  
431 emerging PCV2d-2 strain.

432

433

### Discussion

434 PCVAD is arguably one of the most economically-important diseases affecting  
435 the global swine industry. Characterized by progressive wasting, hallmark histological  
436 lesions of lymphoid depletion with histiocytic infiltration, and the presence of PCV2  
437 antigen or DNA in the lesions, PCVAD is caused by PCV2 infection, although co-  
438 infection with other pathogens are usually necessary for the development of the full-  
439 spectrum of clinical PCVAD (44-46). Several commercial vaccines against PCV2 are  
440 currently available, all of which are based on the PCV2a subtype (11), which prior to  
441 2005 was the main subtype (15-17). However, now PCV2b has surpassed PCV2a as the  
442 most prevalent strain associated with PCVAD losses in the swine industry (15-17). ~~In~~  
443 ~~addition, recently, speculation of vaccination failures has been reported, and though no~~  
444 ~~direct evidence has been found as of yet, these~~ ~~And~~ ~~though all current vaccines have~~  
445 ~~been proven effective at preventing clinical signs and global economic loss due to~~  
446 ~~PCVAD, events have been associated with~~ the emergence of the PCV2d (or mutant  
447 PCV2b) subtype (18, 19, 24), as well as the replacement of PCV2a with PCV2b as the  
448 predominant circulating subtype, cannot be ignored. ~~-(18, 19, 24).~~ Therefore, it is logical  
449 to develop the next generation of vaccines especially against the emerging PCV2 strains.

450 The objectives of this study were to molecularly breed the capsid genes from  
451 different PCV2 subtypes by DNA shuffling, and to develop a candidate chimeric virus  
452 vaccine based on the non-pathogenic PCV1 backbone and shuffled capsid genes of

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

453 divergent PCV2 subtypes. Traditional DNA shuffling approach was undertaken in this  
454 study, in which 5 genetically distinct capsid sequences from each of the 4 known PCV2  
455 subtypes, as well as from the “[divergent PCV2aPCV2e](#)” type (47), which is now  
456 generally considered as a divergent PCV2a strain (35), were used for the DNA shuffling.  
457 Of the more than 50 shuffled PCV2 capsids that were cloned and sequenced, infectious  
458 chimeric viruses were rescued in PK15 cells only in 4 of them, suggesting that the small  
459 PCV2 genome cannot support a large number of forced random reassortment within the  
460 capsid gene.

461           The four viable viruses with shuffled PCV2 capsids generated by traditional DNA  
462 shuffling contained antigenic epitopes from all 5 genetically divergent PCV2 strains,  
463 although most of the variability in the shuffled capsids could be found in the PCV2c  
464 parental strain. This was not unexpected, as the PCV2c subtype is the most divergent  
465 strain from the rest of the PCV2 subtypes identified thus far, based on a phylogenetic  
466 analysis (24, 48). Alignment of the 5 selected parental strains revealed that the PCV2c  
467 does, in fact, contain the most genetically distinct amino acid variations, though some of  
468 these amino acids overlap with the parental PCV2d strain, including the addition of a  
469 terminal lysine residue. The presence of amino acid residues unique to PCV2c and  
470 PCV2d strains suggests that, although the PCV2c subtype has not associated with any  
471 clinical disease, this subtype could possibly have contributed to the evolutionary  
472 emergence of the current PCV2d subtype. In fact, the PCV2c subtype was recently  
473 isolated from feral pigs in Brazil for first time since it was originally described in  
474 Denmark in the early 90s. The feral pig populations were also infected with the other  
475 three PCV2 subtypes, suggesting the possibility of recombination (23). Therefore, these

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

476 findings support the inclusion of PCV2c for DNA shuffling in the current study in order  
477 to increase the breadth of protection of the resulting candidate vaccine against currently  
478 emerging and future possible emerging PCV2 strains.

479 In order to determine the *in vivo* infectivity of the shuffled viruses and to screen  
480 for the best chimera for subsequent challenge and efficacy study, conventional pigs were  
481 experimentally inoculated in a pilot study with each of the 4 viruses with shuffled capsids  
482 in the PCV2a backbone as well as with the chimeric PCV1-2a vaccine virus (12). The  
483 results showed that virus PCV2-3c1.14 induced higher levels of neutralizing antibody  
484 titers when compared to the chimeric PCV1-2a virus, as well as the other 3 shuffled  
485 capsid viruses. The chimeric virus PCV2-3c1.14 also induced significantly higher  
486 neutralizing antibody titers against PCV2a and PCV2d-2 strains. The fact that the PCV2-  
487 3c11.14 shuffled capsid virus induced higher neutralizing antibody titers against PCV2a  
488 compared to a homologous vaccination with the PCV1-2a chimeric vaccine strain was  
489 unexpected. However others have demonstrated this phenomenon with PCV2 viruses  
490 before. Although they demonstrate opposing results, there are many differences in the  
491 experimental design, which could explain these ~~diserepeneies~~discrepancies (21, 49). In  
492 addition, the PCV2-3c114 virus strain grew to the lowest titer of 10<sup>3.33</sup> TCID<sub>50</sub>/mL  
493 compared to the other PCV2-shuffled capsid strains and the PCV1-2a vaccine strain *in*  
494 *vitro* on multiple occasions (data not shown), suggesting that the increase in total and  
495 breadth of neutralizing antibody titers compared to the other strains tested was not simply  
496 due to increased replication efficiency. -Taken together, these results demonstrate that  
497 more research is needed to understand the complicated nature of PCV2 capsid  
498 immunogenicity. Comparison of the amino acid sequences of the shuffled capsid 3c114

Formatted: Font: Italic





1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

522 since PCV1-3c114 protects against PCV2d-2 infection in the challenge and efficacy  
523 experiment. While it is possible that the properties of 3c114 capsid sequence discussed  
524 above are important for production of cross-protective neutralizing antibodies in pigs,  
525 additional research is warranted to determine the important amino acid residues that may  
526 play a critical role in conferring cross-neutralizing activities against different PCV2  
527 subtypes.

528         Based on induction of significantly higher cross-neutralizing antibody titers,  
529 compared to the other shuffled capsid candidates, the shuffled 3c114 capsid sequence was  
530 subsequently selected to produce a chimeric virus PCV1-3c114 vaccine candidate. The  
531 protective efficacy of the PCV1-3c114 chimeric virus as a potential vaccine was evaluated  
532 by challenging vaccinated pigs with PCV2b or PCV2d, respectively. PCV2b is the  
533 predominant subtype currently infecting pigs worldwide, whereas the PCV2d is an  
534 emerging subtype (24). We previously have demonstrated the attenuation of chimeric  
535 PCV1-2a and PCV1-2b viruses in the genomic backbone of the non-pathogenic PCV1 *in*  
536 *vivo* (12, 37, 39). Consistent with these previous reports, there was no detectable PCV1-  
537 3c114 viremia in vaccinated pigs throughout the duration of the study, and no detectable  
538 clinical disease prior to challenge with either PCV2b or PCV2d (data not shown), even  
539 though the vaccinated pigs are infected as evidenced by seroconversion to PCV2 capsid  
540 antibody. It is also possible that the standard PCV2a capsid-based PCV2 ORF2 ELISA  
541 assay is less sensitive for detection of the PCV1-shuffle capsid induced antibodies,  
542 possibly leading an underrepresentation of the antibody titers in the PCV1-3c114  
543 vaccinated groups, however further research is needed to determine if this is the case.  
544 Whether the serology data is indeed blunted due to the limitations of the assay, the

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

545 reduction in challenge virus levels shows a significant effect of vaccination with the  
546 PCV1-3c114 vaccine candidate on PCV2b and PCV2d challenge strains.

547 Vaccination with the chimeric virus PCV1-3c114 vaccine candidate resulted in  
548 significantly reduced PCV2b or PCV2d viral DNA loads at the peak of viremia as well as  
549 reduced viral DNA loads in lymphoid tissues at termination of the study. Furthermore,  
550 the lymphoid lesions were also significantly reduced in vaccinated groups subsequently  
551 challenged with PCV2b compared to mock-vaccinated and challenged controls. Though  
552 the vaccinated animals showed no statistically significant reduction in spleen lymphoid  
553 depletion and spleen and tonsil histiocytic replacement when challenged with PCV2d,  
554 they did have significant reduction for the rest of the PCVAD-associated scores, as well  
555 as reduced viral DNA loads in serum and lymph node tissues, indicating that the PCV1-  
556 3c114 chimeric virus vaccine candidate induced protection against both PCV2b and  
557 PCV2d challenge in conventional pigs.

558

559 **Conclusion**

560 To our knowledge, this is the first report of construction of viable chimeric PCV2  
561 vaccine candidate by shuffling the capsid gene of 5 divergent PCV2 strains belonging to  
562 different subtypes. Importantly, vaccination of pigs with a chimeric virus PCV1-3c114  
563 with shuffled capsid genes induced protective immunity against challenge with the  
564 predominant PCV2b subtype and the emerging PCV2d subtype. Therefore, this chimeric  
565 virus is a potential candidate for further development into the next generation of vaccine  
566 against PCV2.

567

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

568

### **Conflict of Interest Declaration**


569 X.J. Meng is the lead inventor of the chimeric PCV1-2a upon which the current  
570 commercial vaccines Fosterera™ PCV and Fosterera™ PCV MH are based. Greg Nitzel and  
571 David Slade are both employees of Zoetis Inc.

572

573

### **Acknowledgement**

574 The authors thank Iowa State University for animal care and Olusegun Awe, Brett Burke  
575 and Danielle Evenson for assistance with the animal study. We thank Dr. Patrick G.  
576 Halbur for assistance with the necropsy. This work was supported by a grant from  
577 Zoetis Inc, Kalamazoo, MI.



## References

1. **Todd D BM, Biagini P, Hino S, Mankertz A, Mishiro S, Niel C, Okamoto H, Raidal S, Ritchie BW, Teo GC.** 2005. *Circoviridae*. Press EA, San Diego.
2. **Allan GM, McNeilly F, Kennedy S, Daft B, Clarke EG, Ellis JA, Haines DM, Meehan BM, Adair BM.** 1998. Isolation of porcine circovirus-like viruses from pigs with a wasting disease in the USA and Europe. *J Vet Diagn Invest* **10**:3-10.
3. **Tischer I, Miels W, Wolff D, Vagt M, Griem W.** 1986. Studies on epidemiology and pathogenicity of porcine circovirus. *Arch Virol* **91**:271-276.
4. **Meehan BM, McNeilly F, Todd D, Kennedy S, Jewhurst VA, Ellis JA, Hassard LE, Clark EG, Haines DM, Allan GM.** 1998. Characterization of novel circovirus DNAs associated with wasting syndromes in pigs. *J Gen Virol* **79 ( Pt 9)**:2171-2179.
5. **Ellis J, Hassard L, Clark E, Harding J, Allan G, Willson P, Strokappe J, Martin K, McNeilly F, Meehan B, Todd D, Haines D.** 1998. Isolation of circovirus from lesions of pigs with postweaning multisystemic wasting syndrome. *Can Vet J* **39**:44-51.
6. **Choi C, Chae C, Clark EG.** 2000. Porcine postweaning multisystemic wasting syndrome in Korean pig: detection of porcine circovirus 2 infection by immunohistochemistry and polymerase chain reaction. *J Vet Diagn Invest* **12**:151-153.
7. **Edwards S, Sands JJ.** 1994. Evidence of circovirus infection in British pigs. *Vet Rec* **134**:680-681.
8. **Larochelle R, Morin M, Antaya M, Magar R.** 1999. Identification and incidence of porcine circovirus in routine field cases in Quebec as determined by PCR. *Vet Rec* **145**:140-142.
9. **Onuki A, Abe K, Togashi K, Kawashima K, Taneichi A, Tsunemitsu H.** 1999. Detection of porcine circovirus from lesions of a pig with wasting disease in Japan. *J Vet Med Sci* **61**:1119-1123.
10. **Opriessnig T, Meng XJ, Halbur PG.** 2007. Porcine circovirus type 2 associated disease: update on current terminology, clinical manifestations, pathogenesis, diagnosis, and intervention strategies. *J Vet Diagn Invest* **19**:591-615.
11. **Beach NM, Meng XJ.** 2012. Efficacy and future prospects of commercially available and experimental vaccines against porcine circovirus type 2 (PCV2). *Virus Res* **164**:33-42.
12. **Fenaux M, Opriessnig T, Halbur PG, Elvinger F, Meng XJ.** 2004. A chimeric porcine circovirus (PCV) with the immunogenic capsid gene of the pathogenic PCV type 2 (PCV2) cloned into the genomic backbone of the nonpathogenic PCV1 induces protective immunity against PCV2 infection in pigs. *J Virol* **78**:6297-6303.
13. **Opriessnig T, Madson DM, Prickett JR, Kuhar D, Lunney JK, Elsener J, Halbur PG.** 2008. Effect of porcine circovirus type 2 (PCV2) vaccination on

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

621 porcine reproductive and respiratory syndrome virus (PRRSV) and PCV2  
622 coinfection. *Vet Microbiol* **131**:103-114.

623 14. **Chae C.** 2012. Commercial porcine circovirus type 2 vaccines: efficacy and  
624 clinical application. *Vet J* **194**:151-157.

625 15. **Cheung AK, Lager KM, Kohutyuk OI, Vincent AL, Henry SC, Baker RB,**  
626 **Rowland RR, Dunham AG.** 2007. Detection of two porcine circovirus type 2  
627 genotypic groups in United States swine herds. *Arch Virol* **152**:1035-1044.

628 16. **Gagnon CA, Tremblay D, Tijssen P, Venne MH, Houde A, Elahi SM.** 2007.  
629 The emergence of porcine circovirus 2b genotype (PCV-2b) in swine in  
630 Canada. *Can Vet J* **48**:811-819.

631 17. **Firth C, Charleston MA, Duffy S, Shapiro B, Holmes EC.** 2009. Insights into  
632 the evolutionary history of an emerging livestock pathogen: porcine  
633 circovirus 2. *J Virol* **83**:12813-12821.

634 18. **Opriessnig T, Xiao CT, Gerber PF, Halbur PG.** 2013. Emergence of a novel  
635 mutant PCV2b variant associated with clinical PCVAD in two vaccinated pig  
636 farms in the U.S. concurrently infected with PPV2. *Vet Microbiol* **163**:177-  
637 183.

638 19. **Seo HW, Park C, Kang I, Choi K, Jeong J, Park SJ, Chae C.** 2014. Genetic and  
639 antigenic characterization of a newly emerging porcine circovirus type 2b  
640 mutant first isolated in cases of vaccine failure in Korea. *Arch Virol*  
641 **159**:3107-3111.

642 20. **Eddicks M, Fux R, Szikora F, Eddicks L, Majzoub-Altweck M, Hermanns**  
643 **W, Sutter G, Palzer A, Banholzer E, Ritzmann M.** 2015. Detection of a new  
644 cluster of porcine circovirus type 2b strains in domestic pigs in Germany. *Vet*  
645 *Microbiol* **176**:337-343.

646 21. **Huang L, Wang Y, Wei Y, Chen D, Liu D, Du W, Xia D, Wu H, Feng L, Liu C.**  
647 2016. Capsid proteins from PCV2a genotype confer greater protection  
648 against a PCV2b strain than those from PCV2b genotype in pigs: evidence for  
649 PCV2b strains becoming more predominant than PCV2a strains from 2000 to  
650 2010s. *Appl Microbiol Biotechnol* doi:10.1007/s00253-016-7459-y.

651 22. **Allan G, McNeilly F, Meehan B, McNair I, Ellis J, Krakowka S, Fossum C,**  
652 **Watrang E, Wallgren P, Adair B.** 2003. Reproduction of postweaning  
653 multisystemic wasting syndrome in pigs experimentally inoculated with a  
654 Swedish porcine circovirus 2 isolate. *J Vet Diagn Invest* **15**:553-560.

655 23. **Franzo G, Cortey M, de Castro AM, Piovezan U, Szabo MP, Drigo M,**  
656 **Segales J, Richtzenhain LJ.** 2015. Genetic characterisation of Porcine  
657 circovirus type 2 (PCV2) strains from feral pigs in the Brazilian Pantanal: An  
658 opportunity to reconstruct the history of PCV2 evolution. *Vet Microbiol*  
659 **178**:158-162.

660 24. **Xiao CT, Halbur PG, Opriessnig T.** 2015. Global molecular genetic analysis  
661 of porcine circovirus type 2 (PCV2) sequences confirms the presence of four  
662 main PCV2 genotypes and reveals a rapid increase of PCV2d. *J Gen Virol*  
663 **96**:1830-1841.

664 25. **Dvorak CM, Yang Y, Haley C, Sharma N, Murtaugh MP.** 2016. National  
665 reduction in porcine circovirus type 2 prevalence following introduction of  
666 vaccination. *Vet Microbiol* **189**:86-90.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

667 26. **Saha D, Huang L, Bussalleu E, Lefebvre DJ, Fort M, Van Doorselaere J, Nauwynck HJ.** 2012. Antigenic subtyping and epitopes' competition analysis of porcine circovirus type 2 using monoclonal antibodies. *Vet Microbiol* **157**:13-22.

668  
669  
670  
671 27. **Saha D, Lefebvre DJ, Ooms K, Huang L, Delputte PL, Van Doorselaere J, Nauwynck HJ.** 2012. Single amino acid mutations in the capsid switch the neutralization phenotype of porcine circovirus 2. *J Gen Virol* **93**:1548-1555.

672  
673  
674 28. **Stemmer WP.** 1994. DNA shuffling by random fragmentation and reassembly: in vitro recombination for molecular evolution. *Proc Natl Acad Sci U S A* **91**:10747-10751.

675  
676  
677 29. **Stemmer WP.** 1994. Rapid evolution of a protein in vitro by DNA shuffling. *Nature* **370**:389-391.

678  
679 30. **Zhou L, Ni YY, Pineyro P, Sanford BJ, Cossaboom CM, Dryman BA, Huang YW, Cao DJ, Meng XJ.** 2012. DNA shuffling of the GP3 genes of porcine reproductive and respiratory syndrome virus (PRRSV) produces a chimeric virus with an improved cross-neutralizing ability against a heterologous PRRSV strain. *Virology* **434**:96-109.

680  
681  
682  
683  
684 31. **Ni YY, Opriessnig T, Zhou L, Cao D, Huang YW, Halbur PG, Meng XJ.** 2013. Attenuation of porcine reproductive and respiratory syndrome virus by molecular breeding of virus envelope genes from genetically divergent strains. *J Virol* **87**:304-313.

685  
686  
687  
688 32. **Tian D, Ni YY, Zhou L, Opriessnig T, Cao D, Pineyro P, Yugo DM, Overend C, Cao Q, Lynn Heffron C, Halbur PG, Pearce DS, Calvert JG, Meng XJ.** 2015. Chimeric porcine reproductive and respiratory syndrome virus containing shuffled multiple envelope genes confers cross-protection in pigs. *Virology* **485**:402-413.

689  
690  
691  
692  
693 33. **Zhou L, Ni YY, Pineyro P, Cossaboom CM, Subramaniam S, Sanford BJ, Dryman BA, Huang YW, Meng XJ.** 2013. Broadening the heterologous cross-neutralizing antibody inducing ability of porcine reproductive and respiratory syndrome virus by breeding the GP4 or M genes. *PLoS One* **8**:e66645.

694  
695  
696  
697  
698 34. **Wang F, Guo X, Ge X, Wang Z, Chen Y, Cha Z, Yang H.** 2009. Genetic variation analysis of Chinese strains of porcine circovirus type 2. *Virus Res* **145**:151-156.

699  
700  
701 35. **Cortey M, Olvera A, Grau-Roma L, Segales J.** 2011. Further comments on porcine circovirus type 2 (PCV2) genotype definition and nomenclature. *Vet Microbiol* **149**:522-523.

702  
703  
704 36. **Fenaux M, Halbur PG, Haqshenas G, Royer R, Thomas P, Nawagitgul P, Gill M, Toth TE, Meng XJ.** 2002. Cloned genomic DNA of type 2 porcine circovirus is infectious when injected directly into the liver and lymph nodes of pigs: characterization of clinical disease, virus distribution, and pathologic lesions. *J Virol* **76**:541-551.

705  
706  
707  
708  
709 37. **Beach NM, Ramamoorthy S, Opriessnig T, Wu SQ, Meng XJ.** 2010. Novel chimeric porcine circovirus (PCV) with the capsid gene of the emerging PCV2b subtype cloned in the genomic backbone of the non-pathogenic PCV1

710  
711

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

712 is attenuated in vivo and induces protective and cross-protective immunity  
713 against PCV2b and PCV2a subtypes in pigs. *Vaccine* **29**:221-232.

714 38. **Opriessnig T, Xiao CT, Gerber PF, Halbur PG, Matzinger SR, Meng XJ.**  
715 2014. Mutant USA strain of porcine circovirus type 2 (mPCV2) exhibits  
716 similar virulence to the classical PCV2a and PCV2b strains in caesarean-  
717 derived, colostrum-deprived pigs. *J Gen Virol* **95**:2495-2503.

718 39. **Fenaux M, Opriessnig T, Halbur PG, Meng XJ.** 2003. Immunogenicity and  
719 pathogenicity of chimeric infectious DNA clones of pathogenic porcine  
720 circovirus type 2 (PCV2) and nonpathogenic PCV1 in weanling pigs. *J Virol*  
721 **77**:11232-11243.

722 40. **Opriessnig T, Thacker EL, Yu S, Fenaux M, Meng XJ, Halbur PG.** 2004.  
723 Experimental reproduction of postweaning multisystemic wasting syndrome  
724 in pigs by dual infection with *Mycoplasma hyopneumoniae* and porcine  
725 circovirus type 2. *Vet Pathol* **41**:624-640.

726 41. **Halbur PG, Paul PS, Frey ML, Landgraf J, Eernisse K, Meng XJ, Lum MA,**  
727 **Andrews JJ, Rathje JA.** 1995. Comparison of the pathogenicity of two US  
728 porcine reproductive and respiratory syndrome virus isolates with that of  
729 the Lelystad virus. *Vet Pathol* **32**:648-660.

730 42. **Beach NM, Smith SM, Ramamoorthy S, Meng XJ.** 2011. Chimeric porcine  
731 circoviruses (PCV) containing amino acid epitope tags in the C terminus of  
732 the capsid gene are infectious and elicit both anti-epitope tag antibodies and  
733 anti-PCV type 2 neutralizing antibodies in pigs. *J Virol* **85**:4591-4595.

734 43. **Nawagitgul P, Harms PA, Morozov I, Thacker BJ, Sorden SD,**  
735 **Lekcharoensuk C, Paul PS.** 2002. Modified indirect porcine circovirus (PCV)  
736 type 2-based and recombinant capsid protein (ORF2)-based enzyme-linked  
737 immunosorbent assays for detection of antibodies to PCV. *Clin Diagn Lab*  
738 *Immunol* **9**:33-40.

739 44. **Segales J, Allan GM, Domingo M.** 2005. Porcine circovirus diseases. *Anim*  
740 *Health Res Rev* **6**:119-142.

741 45. **Albina E, Truong C, Hutet E, Blanchard P, Cariolet R, L'Hospitalier R,**  
742 **Mahe D, Allee C, Morvan H, Amenna N, Le Dimna M, Madec F, Jestin A.**  
743 2001. An experimental model for post-weaning multisystemic wasting  
744 syndrome (PMWS) in growing piglets. *J Comp Pathol* **125**:292-303.

745 46. **Magar R, Larochelle R, Thibault S, Lamontagne L.** 2000. Experimental  
746 transmission of porcine circovirus type 2 (PCV2) in weaned pigs: a sequential  
747 study. *J Comp Pathol* **123**:258-269.

748 47. **Jantafong T, Boonsoongnarn A, Poolperm P, Urairong K, Lekcharoensuk**  
749 **C, Lekcharoensuk P.** 2011. Genetic characterization of porcine circovirus  
750 type 2 in piglets from PMWS-affected and -negative farms in Thailand. *Virol J*  
751 **8**:88.

752 48. **Franzo G, Cortey M, Olvera A, Novosel D, De Castro AM, Biagini P,**  
753 **Segales J, Drigo M.** 2015. Revisiting the taxonomical classification of Porcine  
754 Circovirus type 2 (PCV2): still a real challenge. *Virol J* **12**:131.

755 49. **Opriessnig T, O'Neill K, Gerber PF, de Castro AM, Gimenez-Lirola LG,**  
756 **Beach NM, Zhou L, Meng XJ, Wang C, Halbur PG.** 2013. A PCV2 vaccine  
757 based on genotype 2b is more effective than a 2a-based vaccine to protect

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

758 against PCV2b or combined PCV2a/2b viremia in pigs with concurrent PCV2,  
759 PRRSV and PPV infection. *Vaccine* **31**:487-494.


760 50. **Ge M, Yan A, Luo W, Hu YF, Li RC, Jiang DL, Yu XL.** 2013. Epitope screening  
761 of the PCV2 Cap protein by use of a random peptide-displayed library and  
762 polyclonal antibody. *Virus Res* **177**:103-107.

763 51. **Constans M, Ssemadaali M, Kolyvushko O, Ramamoorthy S.** 2015.  
764 Antigenic Determinants of Possible Vaccine Escape by Porcine Circovirus  
765 Subtype 2b Viruses. *Bioinform Biol Insights* **9**:1-12.

766 52. **Trible BR, Kerrigan M, Crossland N, Potter M, Faaberg K, Hesse R,**  
767 **Rowland RR.** 2011. Antibody recognition of porcine circovirus type 2 capsid  
768 protein epitopes after vaccination, infection, and disease. *Clin Vaccine*  
769 *Immunol* **18**:749-757.

770 53. **Trible BR, Ramirez A, Suddith A, Fuller A, Kerrigan M, Hesse R, Nietfeld**  
771 **J, Guo B, Thacker E, Rowland RR.** 2012. Antibody responses following  
772 vaccination versus infection in a porcine circovirus-type 2 (PCV2) disease  
773 model show distinct differences in virus neutralization and epitope  
774 recognition. *Vaccine* **30**:4079-4085.

775 54. **Opriessnig T, Gerber PF, Xiao CT, Halbur PG, Matzinger SR, Meng XJ.**  
776 2014. Commercial PCV2a-based vaccines are effective in protecting naturally  
777 PCV2b-infected finisher pigs against experimental challenge with a 2012  
778 mutant PCV2. *Vaccine* **32**:4342-4348.





## Figure Legends

**Fig. 1. Amino acid sequence alignment of the capsid proteins from the five parental PCV2 wild-type strains and the four candidate DNA-shuffled capsids evaluated in this study.** The first five sequences represent the parental strains [including PCV2a \(strain 40895, GenBank accession number AF264042\)](#), [PCV2b \(strain NC16845, accession number GU799576\)](#), [PCV2c \(accession number EU148503\)](#), [PCV2d \(accession number AY181947\)](#), and [“divergent PCV2a” \(accession number EF524533\)](#), while the bottom four sequences represent the DNA-shuffled PCV2 capsids. Amino acids that differ from the consensus are shown in black.

**Fig. 2. A phylogenetic tree of the capsid genes of selected PCV2 strains from different subtypes.** The phylogenetic tree was constructed using the neighbor-joining method with bootstraps in 1,000 replicates. The number above each major branch indicates the bootstrap value. The bold italicized sequence names represent the PCV2 sequences of the 5 parental strains used for DNA shuffling in the study.

**Fig. 3. Comparison of 50% neutralizing antibody titers against four PCV2 wild-type strains from sera of pigs experimentally inoculated with chimeric viruses PCV2-3c13, PCV2-3c14, PCV2-3c14\_2, and PCV2-3c12\_2, or PCV1-2a with shuffled capsid genes.** *In vitro* 50% neutralization assay of respective sera collected at 56 days post-infection against three parental PCV2 strains: (A) PCV2a, (B) PCV2d-1, (C) PCV2b, and (D) PCV2d-2 isolate. The NA titers were calculated as the highest 2-fold dilution ( $2^n$ ) of the serum sample that showed a 50% or greater reduction in the number

1  
2  
3  
4  
5  
6  
7  
8 of positive fluorescent foci, compared to the serum samples from the mock (PBS)  
9 inoculated control group in the same dilution. Asterisk (\*) sign indicates  $p < 0.05$  analyzed  
10 using one-way ANOVA.  
11  
12  
13  
14

15  
16 **Fig. 4. PCV2 capsid-specific antibody response in conventional pigs experimentally**  
17 **inoculated with the chimeric virus PCV1-3c114 vaccine candidate and challenged**  
18 **with the wild-type virus strains PCV2b or PCV2d-2.** The mean S/P ratio  $\pm$  SEM is  
19 plotted for each treatment group throughout the duration of the study. The virus challenge  
20 took place at 42 days post-vaccination (dpv). The dashed line at 0.2 S/P ratio denotes the  
21 lower end cutoff for a positive sample in this assay.  
22  
23  
24  
25  
26  
27  
28  
29  
30

31 **Fig. 5. Quantification of PCV2 viral DNA loads in sera from pigs vaccinated with**  
32 **the chimeric PCV1-3c114 virus and subsequently challenged with PCV2b or PCV2d-**  
33 **2 compared to challenge only controls.** Quantification of PCV2 ORF1 viral DNA loads  
34 in sera using qPCR in (A) PCV2b challenged and (B) PCV2d-2 challenged animals.  
35 Group means  $\pm$  SEM are plotted for each time point post-challenge. The limit of  
36 detection for the assay was  $10^{4.2}$  copies/mL serum of ORF1 DNA determined by a  
37 standard curve for  $10^1 - 10^{10}$  copies of the wild-type PCV2b genome. (\*) Indicates  
38 statistical significance between groups (Student's t-test, corrected for multiple tests).  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

50 **Fig. 6: Quantification of PCV2 viral DNA loads in lymph nodes from pigs**  
51 **vaccinated with the chimeric PCV1-3c114 virus and challenged with PCV2b or**  
52 **PCV2d-2 compared to challenge only controls.** Quantification of PCV2 ORF1 viral  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65


1  
2  
3  
4  
5  
6  
7  
8 DNA loads in lymph nodes using qPCR in (A) PCV2b challenged (B) and PCV2d-2  
9 challenged animals. Group means  $\pm$  SEM are plotted for each time point post-challenge.  
10  
11 The limit of detection for the assay was  $10^{7.1}$  copies/mg tissue of ORF1 viral DNA, as  
12 determined by a standard curve for  $10^1 - 10^{10}$  copies of the wild-type PCV2b genome.  
13  
14 (\*) Indicates statistical significance between groups at that time point (Student's t-test,  
15 corrected for multiple tests).  
16  
17  
18  
19  
20  
21  
22

23 **Fig. 7. Comparison of lymphoid tissues in pigs vaccinated with the chimeric PCV1-**  
24 **3cl.14 virus and subsequently challenged with PCV2b or PCV2d-2 with those of**  
25 **challenge only controls.** Lymphoid depletion and histiocytic replacement for (A, B)  
26 lymph nodes, (C, D) spleen, and (E, F) tonsils at necropsy were compared for vaccinated  
27 and challenged animals (■) with those of challenge only controls (○). Individual animal  
28 scores are represented by individual symbols and group means  $\pm$  SEM are displayed.  
29 Asterisk (\*) sign indicates statistically significant differences between groups (student's  
30 t-test).  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 **Fig. 8. Quantification of PCV2 viral antigen in lymphoid tissues by PCV2**  
42 **immunohistochemistry (IHC). The tissues were obtained from pigs vaccinated with**  
43 **the chimeric PCV1-3cl.14 virus and subsequently challenged with PCV2b or**  
44 **PCV2d-2 compared to challenge only controls.** PCV2 viral antigen scores determined  
45 for (A) lymph nodes, (B) spleen, and (C) tonsils at necropsy were compared for  
46 vaccinated and challenged animals (■) with those of challenge only controls (○).  
47 Individual animal scores are represented by individual symbols and group means  $\pm$  SEM  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

are displayed. Asterisk (\*) sign indicates statistically significant differences between groups (student's t-test).



Figures 1-8

[Click here to download Figure: Figures 1-8\\_V2.pptx](#)

Figure 1




Figure 2




Figure 3







Figure 4







Figure 5

**A**

PCV2b



**B**

PCV2d-2




Figure 6





Figure 7





Figure 8



**Table 1****Table 1.** Seroconversion to PCV2-specific antibodies in pigs experimentally infected with chimeric PCV2 viruses containing shuffled capsids or with the PCV1-2a vaccine virus

Group	Inocula	No. of pigs positive for PCV2 antibodies/total on DPV <sup>a</sup> :							
		0	7	14	21	28	35	42	49
1	PCV2-3c113	0/3	0/3	0/3	0/3	0/3	1/3	2/3	1/3
2	PCV2-3c114	0/3	0/3	0/3	0/3	1/3	3/3	3/3	3/3
3	PCV2-3c14_2	0/3	0/3	0/3	0/3	0/3	1/3	1/3	1/3
4	PCV2-3c112_2	0/3	0/3	0/3	0/3	0/3	1/3	1/3	2/3
5	PCV1-2a	0/3	0/3	0/3	0/3	1/3	2/3	3/3	3/3
6	PBS	0/3	0/3	0/3	0/3	0/3	0/3	0/3	0/3

<sup>a</sup> PCV2 antibody was measured at different days post-inoculation (DPI) with an ELISA using the recombinant PCV2 capsid protein as the antigen. Animals were considered to have seroconverted when samples from two or more consecutive time points were seropositive. Seropositive time points are shown in grey.

**Table 2****Table 2.** Seroconversion to PCV2-specific antibodies by ELISA and detection of viremia by PCR in pigs vaccinated with PCV1-3cl14 virus and challenged with PCV2b or PCV2d-2

Group	Vaccine	Challenge Virus	No. of pigs positive for PCV2 antibodies/total on DPV <sup>a</sup> :							No. of pigs positive for PCV2 antibodies/total on DPC <sup>a,c</sup> :			No. of pigs positive for viremia/total on DPC <sup>b,c</sup> :		
			0	7	14	21	28	35	42	7	14	20	7	14	20
1	PCV1-3cl.14	PCV2b	0/8	0/8	0/8	0/8	1/8	3/8	3/8	7/8	8/8	8/8	0/8	2/8	0/8
2	PCV1-3cl.14	PCV2d-2	0/8	0/8	0/8	0/8	0/8	2/8	4/8	5/8	5/8	5/8	0/8	1/8	2/8
3	None (PBS)	PCV2b	0/8	0/8	0/8	0/8	0/8	0/8	0/8	2/8	5/8	8/8	0/8	4/8	7/8
4	None (PBS)	PCV2d-2	0/8	0/8	0/8	0/8	0/8	0/8	0/8	1/8	4/8	6/8	1/8	7/8	7/8

<sup>a</sup> PCV2 antibody was measured with an ELISA with the recombinant PCV2 capsid antigen. Animals were considered to have seroconverted when samples from two or more consecutive time points were seropositive

<sup>b</sup> Results represent detection by real-time PCR of wild-type PCV2 DNA

<sup>c</sup> At 42 days post-vaccination (DPV), the animals in all four groups were challenged with the wild-type PCV2 virus indicated above

**Table S1: Oligonucleotide primers used in this study**

<b><i>Primer ID</i></b>	<b><i>Primer Sequence (5'--&gt;3')</i></b>
uniRep_F	TTACTGAGTCTTTTTTATCACTTCGTAATGG
2aORF2_R	CTTTCGTTTTTCAGATATGACGTATCCAAGGAGGCG
uniRep_R	ACCCATTACGAAGTGATAAAAAAAGACTCAG
SacII_uni_R	AGCCCGCGGAAATTTCTGACAAACGTTAC
SacII_uni_F	TTTCCGCGGGCTGGCTGAACTTTTGAAAG
PCV1_DSORF2_F	CTTTTTTGTATCACATCGTAATGGTTTTTATT
PCV1_DSORF2_R	TTCTTTCACTTTTATAGGATGACGTATCCAAGGA
PCV1_BB_F	CCTCCTTGGATACGTCATCCTATAAAACTGAAAGAA
PCV1_BB_R	AAATAAAAACCATTACGATGTGATAACAAAAAAG
NB-56-m2b	GAGGTGTTTCGGCCCTCCTCA
PCV2-83F	AAAAGCAAATGGGCTGCTAA
PCV2-83R	TGGTAACCATCCCACCACTT
PCV1 qRepF	TGGAGAAGAAGTTGTTGT
PCV1 qRepR	TCTACAGTCAATGGATACC