



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Identification of candidate host cell factors required for actin-based motility of *Burkholderia pseudomallei*

Citation for published version:

Jitprasutwit, N, Zainal-abidin, N, Vander Broek, C, Thekkedath Kurian, D, Korbsrisate, S, Stevens, MP & Stevens, JM 2016, 'Identification of candidate host cell factors required for actin-based motility of *Burkholderia pseudomallei*', *Journal Of Proteome Research*, vol. 15, no. 12, pp. 4675-4685.
<https://doi.org/10.1021/acs.jproteome.6b00760>

Digital Object Identifier (DOI):

[10.1021/acs.jproteome.6b00760](https://doi.org/10.1021/acs.jproteome.6b00760)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Journal Of Proteome Research

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Identification of candidate host cell factors required for actin-based motility of *Burkholderia pseudomallei*

Niramol Jitprasutwit, Nurhamimah Zainal-Abidin, Charles Vander Broek, Dominic Kurian, Sunee Korbsrisate, Mark P Stevens, and Joanne M. Stevens

J. Proteome Res., **Just Accepted Manuscript** • DOI: 10.1021/acs.jproteome.6b00760 • Publication Date (Web): 14 Nov 2016

Downloaded from <http://pubs.acs.org> on November 19, 2016

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

1
2
3
4 **Identification of candidate host cell factors required for actin-based motility of**
5 ***Burkholderia pseudomallei***
6
7

8
9 Niramol Jitprasutwit^{1&2}, Nurhamimah Zainal-Abidin¹, Charles Vander Broek¹, Dominic
10 Kurian¹, Sunee Korbsrisate², Mark P. Stevens¹ and Joanne M. Stevens^{1*}
11
12

13
14
15
16
17
18 ¹ The Roslin Institute and Royal (Dick) School of Veterinary Studies, University of Edinburgh,
19 Easter Bush, Midlothian, EH25 9RG, United Kingdom.
20

21
22
23 ² Department of Immunology, Faculty of Medicine Siriraj Hospital, Mahidol University
24 Bangkok, Thailand.
25
26

27
28
29
30 * **Corresponding author.** The Roslin Institute and Royal (Dick) School of Veterinary Studies,
31 University of Edinburgh, Easter Bush, Midlothian, EH25 9RG, United Kingdom. Tel. +44 131
32 651 9100. Fax. +44 131 651 9105. E-mail jo.stevens@roslin.ed.ac.uk
33
34
35
36
37
38
39
40

41 **Keywords:** *Burkholderia pseudomallei*, melioidosis, intracellular bacterial pathogen, actin-
42 based motility, BimA, IQGAP1
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Intracellular actin-based motility of the melioidosis pathogen *Burkholderia pseudomallei* requires the bacterial factor BimA. Located at one pole of the bacterium, BimA recruits and polymerises cellular actin to promote bacterial motility within and between cells. Here we describe an affinity approach coupled with Mass Spectrometry to identify cellular proteins recruited to BimA-expressing bacteria under conditions that promote actin polymerisation. We identified a group of cellular proteins that are recruited to the *B. pseudomallei* surface in a BimA-dependent manner, a subset of which were independently validated with specific antisera including the ubiquitous scaffold protein Ras GTPase-activating-like protein (IQGAP1). IQGAP1 integrates several key cellular signalling pathways including those involved in actin dynamics and has been shown to be involved in the adhesion of attaching and effacing *Escherichia coli* to infected cells and invasion of host cells by *Salmonella enterica* serovar Typhimurium. Whilst a direct interaction between BimA and IQGAP1 could not be detected using either conventional pulldown or yeast two hybrid techniques, confocal microscopy revealed that IQGAP1 is recruited to *B. pseudomallei* actin tails in infected cells and siRNA-mediated knockdown highlighted a role for this protein in controlling the length and actin density of *B. pseudomallei* actin tails.

Introduction

Burkholderia pseudomallei causes melioidosis, a severe invasive infection of humans and animals endemic in Southeast Asia and Northern Australia. Human infections are acquired by inhalation, ingestion or via abraded skin and may produce a spectrum of disease ranging from rapidly fatal septicaemia and acute pneumonia, through chronic or localised abscess formation to subacute infections (reviewed in Wiersinga *et al.*¹). Relapse and latency are common in melioidosis and it is believed that this reflects the ability of *B. pseudomallei* to persist in an intracellular niche and its intrinsic resistance to many classes of antibiotic.

Facultative intracellular pathogens of several bacterial genera have evolved mechanisms to enter and exit eukaryotic cells by harnessing the power of actin polymerisation. Some species of *Burkholderia*, *Listeria*, *Mycobacterium*, *Rickettsia* and *Shigella* are propelled in the cytosol and into adjacent cells by polar nucleation of actin, a process termed actin-based motility. These bacteria use distinct mechanisms for actin-based motility, converging on activation of the cellular Arp2/3 complex by mimicry or recruitment of its regulators (reviewed in Stevens *et al.*²) or by mimicking Arp2/3-independent actin nucleators such as cellular formins or WH2-domain-containing proteins (reviewed in Qualmann and Kessels³). In the case of *Burkholderia* we discovered that this process requires BimA⁴, a putative Type V secreted protein that is conserved amongst *B. pseudomallei* isolates in the endemic area⁵. *B. pseudomallei* BimA influences intracellular survival, intercellular spread⁶ and virulence in mice⁷, however the mechanism by which it recruits and activates cellular factors to assemble actin is not fully understood. BimA homologues of divergent N-terminal amino acid sequence exist in the closely related species *B. mallei* and *B. thailandensis*⁸. These proteins vary in their composition and number of actin-binding WH2 domains, the presence of proline-rich motifs, PDAST repeats and Arp2/3-binding CA domains implying that different *Burkholderia* species have evolved distinct strategies of actin-based motility^{6, 8-9}. For example only *B. thailandensis* BimA sequesters Arp3, consistent with the presence of a unique CA domain^{9a}. Despite the variation in sequence both *B. mallei* and *B. thailandensis* BimA proteins can functionally substitute for the actin-based motility defect of a *B. pseudomallei* *bimA* mutant⁸. Similarly, BimA proteins from *B. pseudomallei* and *B. mallei* can restore actin-based motility to a *B. thailandensis*

1
2
3 *bimA* mutant^{9b, 10}. In *B. mallei* and *B. pseudomallei*, *bimA* gene expression is co-regulated
4 with the virulence-associated Type VI Secretion System cluster through a two-component
5 sensor-regulator VirAG which senses host cytosolic glutathione levels¹¹.
6
7

8 In common with other bacterial pathogens capable of actin-based motility, the Arp2/3
9 complex can be detected in *B. pseudomallei* actin-rich tails¹². However, we have previously
10 shown that *B. pseudomallei* BimA does not require the Arp2/3 complex to promote actin
11 polymerisation *in vitro*^{4, 9a}, a finding confirmed recently by Benanti *et al.*^{9b} and supported by
12 observations by Lu and colleagues in which Arp2/3 depletion failed to inhibit *B.*
13 *pseudomallei* actin-based motility¹⁰. Taken together with the findings that N(neural)-WASP
14 and the vasodilator-stimulated phosphoprotein (VASP) are not required for actin-based
15 motility of *B. pseudomallei*¹², and that actin-based motility is insensitive to overexpression
16 of the WA fragment of Scar1 (which inhibits actin-based motility of *L. monocytogenes*, *S.*
17 *flexneri* & *R. conorii*)¹², it may be inferred that *B. pseudomallei* employs a distinct
18 mechanism for intracellular motility to other pathogens. Indeed it has recently been
19 suggested that BimA mimics cellular Ena/VASP proteins promoting the polymerisation and
20 elongation of actin tails^{9b}.
21
22
23
24
25
26
27
28
29
30

31 In a similar manner described for the characterisation of host cell proteins in the actin
32 tails of *Listeria monocytogenes*¹³, we have utilised an affinity purification approach to
33 identify the cellular proteins recruited to *B. pseudomallei* under conditions that promote
34 actin assembly *in vitro*. This relied on overexpression of VirAG to drive surface expression of
35 BimA in *B. pseudomallei* and pulldown of host proteins from a murine splenic extract. Using
36 this approach we have identified a number of proteins common to both the actin-rich tails
37 of *L. monocytogenes* and *B. pseudomallei*, together with a number of proteins uniquely
38 associated with the actin tails of *B. pseudomallei*. Association of a subset of these proteins
39 with BimA-expressing bacteria was validated by immunoblotting. Confocal microscopy
40 confirmed the recruitment of the cellular cytoskeletal scaffold protein Ras GTPase-
41 activating-like protein (IQGAP1) to the actin tails of *B. pseudomallei* in infected cells. Using
42 siRNA-mediated knockdown we also demonstrate that IQGAP1 plays a role in regulating the
43 actin density and tail length of *B. pseudomallei* actin tails in infected cells.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Materials and methods

Bacterial strains

Burkholderia pseudomallei strain 10276 (NCTC 10276) was isolated from a British seaman suffering from the chronic form of melioidosis after likely exposure in Bangladesh¹⁴. Strain 10276 and our *bimA* insertion mutant⁴ were transformed with pBHR2-*virAG*^{11b} (a kind gift from Paul Brett and Mary Burtnick, University of South Alabama, US) by electroporation¹⁵ to give strains 10276 pBHR2-*virAG* and 10276 *bimA*::pDM4 pBHR2-*virAG*. Bacteria were cultured in Luria Bertani broth supplemented with appropriate antibiotics at 37°C with shaking for ~16 hours.

To generate a *bimA* deletion mutant, approximately 400 base pairs immediately upstream of the *bimA* gene (annotated as BPSS1492 in the reference *B. pseudomallei* K96243 genome) was amplified using primers P1 (ATATATCTCGAGACCCGACACGCCGTGGACAGAA) and P2 (ATATATGGGCCCATATCGATTGGCAGTGCCGT) using GC advantage polymerase (Takara Biotech Europe, Saint-Germain-en-Laye, France) following the manufacturer's instructions. Genomic DNA from strain NCTC 10276 was used as template. The PCR product was digested with *XhoI* and *ApaI* and ligated into similarly digested pDM4, an oriR6K *sacB* positive-selection suicide replicon¹⁶. The resulting plasmid was designated pDM4-*bimA1*. Next, approximately 540 base pairs downstream of the *bimA* gene was amplified from 10276 genomic DNA by PCR using primers P3 (ATATATGGGCCCTAAGCACCCGCAAACCCCCCGGGCATC) and P4 (ATATATAGATCTCAGGCTGCAGAACGCAGGCT). The PCR product was restricted with *ApaI* and *BglII* prior to ligation with similarly digested pDM4-*bimA1*. The resulting plasmid was designated pDM4- Δ *bimA*.

Construction of the *B. pseudomallei* mutant was performed according to the method described by Logue *et al.*¹⁷. Briefly, pDM4- Δ *bimA* was introduced into *B. pseudomallei* strain NCTC 10276 by conjugation from *E. coli* S17.1 λ pir. Colonies were selected on LB agar plates containing 50 μ g/ml kanamycin and 50 μ g/ml chloramphenicol. Colonies were screened by PCR with P1 and P4 primers to detect integration of the plasmid. A *B. pseudomallei* 10276 pDM4- Δ *bimA* merodiploid was next cultured in the absence of selection for the integrated plasmid then plated on LA plates lacking NaCl and containing 15% (w/v) sucrose to positively

1
2
3 select for a second recombination event involving excision of pDM4 and the *bimA* allele.
4 Resulting colonies were then screened by PCR using flanking primers BimA screen 1
5 (GATGTCGCCGACGAAAGCAG) and BimA screen 2 (AGTGGGCGCGATTCTCGCGGCT) for the
6 presence of a truncated *bimA* gene (of around 1Kb) instead of the full length gene (of
7 around 2.5Kb). The resulting *bimA* deletion mutant was designated 10276 Δ *bimA*.
8
9
10
11

12 13 14 **Preparation of murine splenic lysates**

15 Splensens were harvested from BALB/c or VM mice at post mortem. Splensens were rinsed
16 in ice-cold polymerisation buffer (10mM Tris pH 7.5, 50mM KCl, 2mM MgCl₂). 1ml of ice-
17 cold polymerisation buffer supplemented with protease and phosphatase inhibitors (1mM
18 phenylmethanesulfonyl fluoride (PMSF), 2mM Na₂VO₃, 2mM NaF, 2mM Na pyrophosphate,
19 1μg/ml aprotinin, 10μg/ml leupeptin, 1μg/ml pepstatin A) was added per spleen and the
20 tissues homogenised. Supernatants were clarified by ultracentrifugation at 100,000x g for 2
21 hours at 4°C. The clarified supernatants were carefully pipetted and stored at -70°C
22 between assays. The protein concentration of the murine splenic lysate was determined to
23 be ~7mg/ml using a bicinchoninic acid protein assay kit (Thermo Fisher Scientific,
24 Cramlington, UK).
25
26
27
28
29
30
31
32
33
34

35 **Affinity purification of *Burkholderia*-associated proteins**

36 Interacting proteins were isolated essentially as described by David *et al.*^{13a} with some
37 modifications. Approximately 1 x 10⁹ bacteria were pelleted and suspended in 1.5ml murine
38 splenic lysate supplemented with 5mM ATP and 30mM creatine phosphate. Bacteria were
39 incubated for 1 hour at 37°C with gentle agitation. Bacteria were washed gently with 1ml
40 ice-cold 10mM PIPES pH7.25 containing 40mM KCl, 5mM ATP and 5mM MgCl₂. Interacting
41 proteins were eluted from the bacterial surface using 0.5ml 10mM PIPES pH7.25 containing
42 1M KCl, 5mM ATP and 5mM MgCl₂. After repeating the elution, the eluate was filter-
43 sterilised by passing through a 0.2μm membrane filter before concentrating the proteins
44 using 30 μl StrataClean resin (Agilent Technologies UK Ltd., Stockport, UK). Resin were
45 suspended in Laemmli sample treatment buffer containing 2% (v/v) β-mercaptoethanol.
46 After incubation at 95°C for 10 minutes, samples were separated by SDS-PAGE using a 4-
47 15% gradient gel. After electrophoresis bands were stained using a SilverSnap kit following
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 the manufacturer's instructions (Thermo Fisher Scientific, Cramlington, UK). Bands were
4 excised and subjected to LC-MS/MS.
5
6
7

8 **LC-MS/MS**

9
10 Gel slices were destained with 15mM potassium ferricyanide/50mM sodium
11 thiosulphate for 5 minutes, reduced with 10mM DTT/100mM ammonium bicarbonate for 30
12 minutes and alkylated with 55mM iodoacetamide/100mM ammonium bicarbonate for 20
13 minutes. Gel slices were washed with 100mM ammonium bicarbonate and dehydrated with
14 100% (v/v) acetonitrile. Proteins were digested with 6 ng/ml trypsin/50mM ammonium
15 bicarbonate for 5 hours at 37°C and peptides were extracted in 1% (v/v) formic acid/2%
16 (v/v) acetonitrile followed by 50% (v/v) acetonitrile. LC-MS/MS analysis of peptides were
17 performed on a nanoAcquity UPLC system coupled to Q-ToF Premier Mass Spectrometer
18 (Waters Corporation, Milford, Massachusetts, USA). Tryptic peptides were desalted and
19 concentrated on a C₁₈ TRAP column (180µm x 20mm, 5µm Symmetry, Waters), for 3
20 minutes at 10µl/ minute and resolved on a 1.7µm BEH 130 C₁₈ column (100µm x 100mm,
21 Waters Corporation) using a Waters Corporation nanoAcquity UPLC. Peptides were eluted at
22 400 nl/ minute with a linear gradient of 0-50% (v/v) acetonitrile/ 0.1% (v/v) formic acid over
23 30 minutes, followed by 85% (v/v) acetonitrile/ 0.1% (v/v) formic acid for 7 minutes. Eluted
24 peptides were analysed on a Q-ToF Mass Spectrometer in 'data directed' acquisition mode,
25 where an MS survey scan was used to automatically select double and multiple charged
26 peptides for further MS/MS fragmentation. From each survey scan up to three of the most
27 intense peptides were selected for fragmentation. MS/MS collision energy was dependent
28 on precursor ion mass and charge state. A reference spectrum was collected every 30
29 seconds from the Glu-fibrinopeptide B (785.8426 m/z), introduced via a reference sprayer.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45 The raw mass spectral data was processed with ProteinLynx Global Server 2.3 (Waters
46 Corporation) to generate peaklist files. The mass accuracy of the spectra was further
47 corrected by using the reference spectra from Glu-fibrinopeptide B. The resulting peaklist
48 files were searched against a locally installed protein sequence database (5855 entries) of
49 the *B. pseudomallei* reference strain K96243 and mouse International Protein Index (IPI)
50 database¹⁸ using Mascot ver. 2.3 (Matrix Science, London, UK). Precursor and fragment ion
51 mass tolerance were set to +/- 100 ppm and +/- 0.1 Da respectively. Trypsin specificity was
52
53
54
55
56
57
58
59
60

1
2
3 used for allowing up to one missed cleavage. Carbamidomethylation of cysteines and
4 oxidation of methionines were selected as fixed and variable modifications respectively. The
5 interpretation and presentation of MS/MS data was performed according to published
6 guidelines¹⁹.
7
8
9

10 11 12 **Immunoblotting of bacterial and affinity-purified host proteins**

13
14 Protein samples were denatured by heating to 95°C for 5 minutes in Laemmli buffer
15 containing a final concentration of 1% (v/v) β -mercaptoethanol. Proteins were resolved by
16 SDS-PAGE and transferred to PVDF or nitrocellulose membranes using a semi-dry transfer
17 system (Bio-Rad, Hemel Hempstead, UK). Membranes were then blocked and incubated
18 sequentially with primary antibodies at a concentration of 0.5-1 μ g/ml followed by species-
19 specific antibodies conjugated with horseradish peroxidase (for standard ECL detection, GE
20 Healthcare, Chalfont St. James, UK) or fluorescently labelled secondary antibodies (α -rabbit
21 IgG DyLight⁸⁰⁰, Cell Signalling Technology, Leiden, The Netherlands), followed by detection
22 using a Biosciences Odyssey infrared imaging system (LI-COR Biosciences, Cambridge, UK).
23 Primary antibodies used in immunoblotting were: mouse anti-BimA monoclonals AF8, FB5
24 and FG11⁴; rabbit anti-BopE²⁰; mouse anti-*B. pseudomallei* capsule (a kind gift from Dstl,
25 Porton Down, UK²¹); goat anti-actin (Source Bioscience, Nottingham, UK); goat anti-IQGAP1
26 (Abcam, Cambridge, UK); rabbit anti-IQGAP1 (Insight Biotechnology, Wembley, UK); rabbit
27 anti-vinculin (Source Bioscience, Nottingham, UK); rabbit anti-HSP90 (Source Bioscience,
28 Nottingham, UK) and rabbit anti-L-Plastin (Source Bioscience, Nottingham, UK).
29
30
31
32
33
34
35
36
37
38
39
40
41

42 **Cell infection, siRNA knockdown and fluorescence microscopy**

43
44 HeLa cells were cultured on sterile glass coverslips and infected at a multiplicity of
45 infection (MOI) of \sim 100 essentially as described in Sitthidet *et al.*⁶. At the indicated time
46 points post-infection coverslips were rinsed with phosphate-buffered saline (PBS) and
47 incubated for a minimum of 16 hours in PBS containing 4% (w/v) paraformaldehyde before
48 staining for confocal microscopy. Coverslips were treated with 0.5% (v/v) Triton-X100 in PBS
49 for 15 minutes to permeabilise infected cells, blocked for 30 minutes in 0.5% (w/v) BSA in
50 PBS and then sequentially incubated in 0.5 μ g/ml primary and fluorescently labelled
51 secondary antibodies or cell stains as indicated in figure legends. Primary antibodies used in
52
53
54
55
56
57
58
59
60

1
2
3 fluorescence microscopy: mouse anti-BimA AF8, FB5 and FG11⁴; mouse anti-*B. pseudomallei*
4 lipopolysaccharide (LPS) (Camlab, Cambridge, UK); rabbit anti-IQGAP1 (Insight
5 Biotechnology, Wembley, UK). Actin filaments were stained with Phalloidin Alexa Fluor⁴⁸⁸
6 (Life Technologies Ltd., Paisley, UK). Images were captured using a Leica Microsystems
7 (Milton Keynes, UK) LSM710 confocal scanning microscope and Zen 2011 software (Carl
8 Zeiss Ltd., Cambridge, UK).

9
10 To reduce expression of IQGAP1 in HeLa cells, approximately 30,000 cells plated per
11 well of a 24-well tissue culture plate were transfected with 6pmol Silencer Select IQGAP1
12 siRNA duplex (Life Technologies Ltd., Paisley, UK) using Lipofectamine RNAiMax (Life
13 Technologies Ltd., Paisley, UK) essentially as described by the manufacturer's instructions.
14 Cells were lysed at 72 hours post-transfection to detect IQGAP1 knockdown by
15 immunoblotting. Whilst optimising conditions for IQGAP1 knockdown cells transfected with
16 MISSION negative control siRNA (Sigma-Aldrich, Dorset, UK) were used for reference. For
17 infection experiments, IQGAP1 siRNA-transfected cells were incubated for 56 hours prior to
18 infection with *B. pseudomallei* strain 10276. Cells were infected at an MOI of ~100 for a
19 further 16 hours before fixation and staining for confocal microscopy analysis.

20
21 Images representing maximal projection z-stacks were captured using identical laser
22 settings from 3 independent experiments and analysed using Image J software
23 (<http://imagej.nih.gov/ij/>). The maximal calliper function was used to calculate the length of
24 tails and the corrected total cell fluorescence (CTCF) formula to determine the intensity of
25 actin staining throughout each tail. Data from a total of 100 tails from each condition were
26 collated and analysed using a student t-test in GraphPad Prism.

27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 **Yeast 2 Hybrid**

45 A fragment of the *bimA* gene encoding amino acids 54 to 455 of the BimA protein
46 (capable of binding and polymerising actin *in vitro*) was cloned into pGBKT7 (Clontech, Saint-
47 Germain-en-Laye, France) to express an in-frame fusion protein with the GAL4 DNA-binding
48 domain. The *bimA* gene was amplified by PCR from 10276 genomic DNA using primers F-
49 BimA (GCGCGCCATATGAATCCCCCGAACCGCCGGGC) and R-BimA
50 (GCGCGCGAATTCTTAGCGCGGGTGTCTGGTG). The product was purified, digested with *NdeI*
51 and *EcoRI*, and ligated into similarly digested vector to give pGBKT7-*bimA*. This plasmid was

1
2
3 introduced into *Saccharomyces cerevisiae* strain AH109 using a standard lithium acetate
4 mediated transformation protocol outlined in the Matchmaker GAL4 Yeast Two Hybrid
5 System 3 handbook (Clontech, Saint-Germain-en-Laye, France) with selection on SD media
6 lacking tryptophan (Clontech, Saint-Germain-en-Laye, France), to give strain AH109 pGBKT7-
7 *bimA*. The *iqgap1* gene was amplified by PCR from HeLa cell cDNA (AMS Biotechnology Ltd,
8 Abingdon, UK) using primers F-IQGAP1 (TTTTCATATGATGTCCGCCGACAGACGAG) and R-
9 IQGAP1 (TTTTCTCGAGTTACTTCCCGTAGAACTTTTTGTTGA). Similarly the *actb* gene was
10 amplified by PCR from HeLa cell cDNA using primers F-actin
11 (TTTTCATATGATGGATGATGATATCGCCG) and R-actin
12 (TTTTCTCGAGCTAGAAGCATTGCGGTGG). PCR products were purified, digested with *NdeI*
13 and *XhoI*, and ligated into similarly digested pGADT7 (Clontech, Saint-Germain-en-Laye,
14 France) to create pGADT7-*iqgap1* and pGADT7-*actin*, encoding in-frame fusion proteins with
15 the GAL4 activation domain. All plasmids were verified by sequencing prior to
16 transformation into AH109 strains and selection on SD media lacking leucine. The pGADT7-
17 *iqgap1* and pGADT7-*actb* plasmids were separately introduced into AH109 pGBKT7-*bimA* by
18 lithium acetate with selection on SD media lacking leucine and tryptophan (DDO: Double
19 Drop Out media, Clontech, Saint-Germain-en-Laye, France) to give strains AH109 pGBKT7-
20 *bimA* pGADT7-*iqgap1* and AH109 pGBKT7-*bimA* pGADT7-*actin*.

21
22
23
24
25
26
27
28
29
30
31
32
33
34
35 To identify protein: protein interactions, single yeast colonies from DDO agar plates
36 were suspended in 20µl water and 5µl dotted onto SD agar plates lacking leucine,
37 tryptophan, adenine and histidine (QDO: Quadruple Drop Out) supplemented with 20µg/ml
38 X-α-galactosidase. Plates were incubated at 30°C for 24 hours before observing the
39 appearance of the colonies. The strains AH109 pGBKT7-*bimA* pGADT7-*iqgap1* and AH109
40 pGBKT7-*bimA* pGADT7-*actin* were tested alongside yeast strains transformed with positive
41 and negative control vectors (pGADT7-T antigen pGBKT7-p53 and pGADT7-T antigen
42 pGADT7-Lamin C respectively) supplied with the Matchmaker GAL4 Y2H System 3 (Clontech,
43 Saint-Germain-en-Laye, France).

Results

Engineering a *Burkholderia pseudomallei* strain that constitutively expresses BimA *in vitro*

BimA (BPSS1492), a predicted Type V autotransporter protein of *Burkholderia pseudomallei*, is required for actin-based motility of the bacterium in the cytosol of infected cells^{4, 6}. We have found that BimA expression is below the limit of detection in bacteria cultured in common laboratory media (Figure 1a) but detectable from 6 hours post-infection by immunoblotting infected cell lysates with antibodies specific to BimA⁴ (Figure 1b). The presence of intracellular bacteria in all samples was confirmed by immunoblotting with antibody reactive to the *B. pseudomallei* capsule²¹. In the closely related species *B. mallei*, *bimA* gene transcription and BimA protein expression is positively regulated by a two-component system, VirAG, encoded adjacent to the *bimA-E* genes on chromosome 2^{11b}. In order to induce BimA expression in LB-cultured bacteria we therefore introduced the pBHR2-*virAG* plasmid constitutively expressing the *B. mallei virAG* genes^{11b} into strain NCTC 10276¹⁴ or our *bimA* insertion mutant giving strains 10276 pBHR2-*virAG* and 10276 *bimA*::pDM4 pBHR2-*virAG*. The VirA and VirG proteins of *B. mallei* and *B. pseudomallei* share 99.4% and 99.6% identity respectively and introduction of the pBHR2-*virAG* plasmid into these strains resulted in expression of BimA protein in wild-type bacteria cultured in laboratory medium (Figure 1a). Protein loading was confirmed using an antibody against the *B. pseudomallei* Bsa Type III Secretion System effector protein BopE²⁰. Localisation of BimA on the surface of the 10276 pBHR2-*virAG* strain was confirmed by confocal microscopy of intact bacteria (data not shown).

Isolation of candidate host cell proteins involved in actin-based motility of *B. pseudomallei*

Having established an *in vitro* system in which *B. pseudomallei* bacteria were expressing BimA, we set out to identify host cell proteins that associated with 10276 pBHR2-*virAG* (BimA-expressing bacteria) and common bacterial proteins released from the bacterial surface following elution from 10276 pBHR2-*virAG* and 10276 *bimA*::pDM4 pBHR2-*virAG*. Bacteria were incubated with murine splenic lysates under conditions that promote BimA: actin binding and actin polymerisation⁴ and interacting proteins eluted and concentrated for SDS-PAGE. Following silver staining gel slices were excised and subjected to in-gel trypsin

1
2
3 digestion, extraction and LC-MS/MS analysis. A representative silver stained gel is shown in
4 Figure 2. Spectra were then separately searched against the mouse International Protein
5 Index (IPI) database¹⁸ and an 'in-house' *B. pseudomallei* K96243 protein database to obtain
6 protein identifications. Data were collected for 3 independent experiments with both
7 bacterial and host cell proteins identified in the eluted proteins.
8
9

10
11 We sought to identify host cell proteins that were solely associated with, or identified at
12 a higher frequency with, the BimA-expressing bacteria 10276 pBHR2-*virAG* compared to the
13 *bimA* mutant strain 10276 *bimA*::pDM4 pBHR2-*virAG*. Spectra were screened against the
14 mouse IPI database¹⁸ and protein identifications assigned when the proteins were identified
15 in at least two of the three biological replicates. From this analysis a set of 30 host proteins
16 were isolated (Table 1). In contrast to a previous study on the proteome of *L.*
17 *monocytogenes* actin tails^{13b}, there was a notable lack of microtubule components,
18 intermediate filaments or myosins. The most abundant and commonly isolated proteins
19 recruited in a *VirAG*-dependent manner were those of the actin family, consistent with the
20 known actin-binding activity of BimA⁴. Several of the proteins are associated with areas of
21 dynamic actin assembly such as membrane ruffles and cell junctions, or acting as scaffolds
22 to integrate signalling complexes. A selection of host cell proteins recruited by BimA-
23 expressing bacteria contained FERM domains, which are known to localise at the interface
24 between the cytoskeleton and the plasma membrane. The scaffold proteins Talin, 14-3-
25 3 protein zeta/delta and IQGAP1 were identified, as were components of the Arp2/3
26 complex (specifically subunits 1b, 2 and 4), previously described as being in the actin tails of
27 *B. pseudomallei*¹². In addition to actin, we identified several F-actin binding proteins
28 implicated in stabilising F-actin networks through bundling and cross-linking activities (α -
29 actinins 1 and 4, Filamin A, Spectrin/ fodrin, Vinculin, Tropomyosin, Plastin-2/ L-Plastin,
30 Transgelin), several F-actin capping proteins (Spectrin/ fodrin, ERM proteins, Macrophage
31 capping protein, CapZ) and an actin-depolymerising factor (ADF) in the form of cofilin. We
32 also identified the proteins WD repeat-containing protein 1 (AIP1) and serine/ threonine
33 phosphatase 2A, known regulators of cofilin activity.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

52 Several *B. pseudomallei* proteins were consistently identified in eluates from both wild-
53 type and *bimA* mutant bacteria (Table 2), including the Type VI secreted protein Hcp
54 (BPSS1498), a number of chaperone proteins and proteases. A number of the *B.*
55
56
57
58
59
60

1
2
3 *pseudomallei* proteins identified in this study have previously been identified in the *B.*
4 *pseudomallei* outer membrane proteome²², the total secretome²³ or as surface proteins by
5 surface labelling approaches²⁴, including the molecular chaperones GroEL (BPSL2697 and
6 BPSS0477), GroES (BPSL2698), DnaK (BPSL2827) and the transcription/ translation factors
7 Tuf (BPSL3215) (labelled OM in Table 2). Several have also been shown to be recognised by
8 the sera of convalescent melioidosis patients (labelled I in Table 2)²⁴⁻²⁵. The remainder of
9 proteins ranged in function from metabolic enzymes involved in glycolysis, amino acid
10 metabolism and the TCA cycle to RNA polymerase subunits. These may represent
11 particularly abundant proteins of the bacteria released upon cell lysis or proteins contained
12 within outer membrane vesicles (OMVs). OMVs have recently been reported to have
13 protective efficacy against pulmonary melioidosis and *B. pseudomallei*-induced sepsis²⁶,
14 however a proteomic analysis of the contents of such OMVs has yet to be described.
15
16
17
18
19
20
21
22
23
24
25

26 **Validation of the dataset**

27
28 The dataset described in Table 1 was validated by immunoblotting of independently-
29 generated samples for a subset of host cell proteins. An equal volume of eluted protein
30 generated from identical treatment of similar numbers of bacteria was probed in each
31 instance. Proteins that were identified in all 3 experiments, with high MASCOT scores, and
32 were novel when compared to proteins involved in *Listeria* or *Shigella* actin-based motility
33 were confirmed (HSP90 and IQGAP1) (Figure 3). We also chose to validate one of the
34 proteins identified in only 2 experiments with a moderate MASCOT score (vinculin) or
35 identified in all 3 experiments with a low MASCOT score (L-Plastin) (Figure 3). As a positive
36 control the recruitment of actin to BimA-expressing bacteria was also verified (Figure 3). All
37 chosen targets were detected in murine splenic lysate and samples derived from bacteria
38 expressing BimA *in vitro* (10276 pBHR2-*virAG*), and were absent from those samples lacking
39 BimA (10276 *bimA*::pDM4 pBHR2-*virAG*).
40
41
42
43
44
45
46
47
48
49
50

51 **IQGAP1 is present in actin tails formed by intracellular *B. pseudomallei***

52 The Ras GTPase-activating-like protein IQGAP1 is a ubiquitously expressed scaffold
53 protein that integrates multiple host signalling pathways including actin cytoskeleton
54 dynamics, cell cycle and cell adhesion. To date this protein has not been shown to be
55
56
57
58
59
60

1
2
3 involved in the actin-based motility of intracellular bacteria, although it has been shown to
4 be involved in entry of *Salmonella* into host cells and for actin pedestal formation of
5 attaching and effacing *E. coli* pathotypes to the apical surface of cells (reviewed by Kim *et*
6 *al.*²⁷). Our data implied that in contrast to *Listeria* and other pathogens that display actin-
7 based motility, IQGAP1 may be a component of *B. pseudomallei* actin tails. Indeed this was
8 supported by immunocytochemistry and confocal microscopy which revealed the presence
9 of IQGAP1 throughout the tails of *B. pseudomallei* 10276 at 16 hours post-infection of HeLa
10 cells (Figure 4(a)-(c)). In agreement with a recent report by Benanti *et al.*^{9b}, we could also
11 elucidate the architecture of the *B. pseudomallei* actin tail by fluorescent actin staining
12 coupled with confocal microscopy, albeit at low resolution (Figure 4(d)). The tails consisted
13 of loosely bundled F-actin filaments reminiscent of those described for certain *Rickettsia*
14 species that rely on the formin-like Sca2 autotransporter for actin polymerisation²⁸.
15
16
17
18
19
20
21
22
23
24
25

26 **Yeast 2 Hybrid analysis detects a direct interaction between BimA and actin but not** 27 **IQGAP1** 28

29
30 Whilst IQGAP1 could readily be detected in the actin tails of all bacteria displaying actin-
31 based motility in infected cells, we sought to determine if this is through direct interaction
32 with the BimA protein using a Yeast 2 Hybrid approach. The Yeast 2 Hybrid assay is a widely
33 utilised and sensitive molecular genetic approach to detect even weak or transient protein:
34 protein interactions *in vivo*. It relies on the modular nature of transcriptional trans-
35 activators where the DNA-binding and activation domains can be physically separated and
36 expressed as fusions with proteins of interest. If the proteins interact they bring the DNA-
37 binding and activation domains into close proximity thereby resulting in transcription of
38 certain reporter genes. In our Yeast 2 Hybrid system the reporter genes for histidine and
39 adenine biosynthesis (which allow growth on media lacking these two amino acids) and α -
40 galactosidase (which produces blue colonies on media containing X- α -Gal substrate) are
41 utilised. We have expressed amino acid residues 54 to 455 of the BimA protein as a DNA-
42 binding protein fusion protein in the reporter yeast strain AH109 pGBKT7-BimA. When β -
43 actin is also expressed in this strain as an activation domain fusion protein (AH109 pGBKT7-
44 BimA pGADT7-actin), the yeast generates blue colonies on media containing X- α -Gal
45 substrate but lacking leucine, tryptophan, adenine and histidine (QDO) (Figure 5), indicating
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 an interaction between these two proteins as we have previously described⁴. Conversely,
4 introduction of a vector expressing IQGAP1 as an activation domain fusion protein into the
5 BimA yeast strain (AH109 pGBKT7-BimA pGADT7-IQGAP1) resulted in growth of the yeast
6 strain on QDO media, however the colonies remained white even after extended incubation
7 times (Figure 5). Growth on selective media in the absence of detectable α -galactosidase
8 activity indicates that these two proteins do not directly interact with each other. Growth
9 under these circumstances can be attributed to a low level of intrinsic *trans*-activation in
10 yeast strains harbouring the pGBKT7-BimA plasmid as shown in Figure 5.

11
12 Consistent with the Yeast 2 Hybrid data presented in Figure 5, we were unable to detect
13 a direct interaction between BimA and IQGAP1 by conventional pulldown assay using
14 affinity-purified glutathione S-transferase (GST)-BimA₅₄₋₄₅₅ and maltose-binding protein
15 (MBP)-IQGAP1 fusion proteins expressed by *E. coli* under conditions previously used to
16 detect the direct interaction of actin with BimA⁴ (data not shown).

27 28 **Effect of IQGAP1 knockdown on actin density and length of BimA-induced actin tails**

29
30 Given the presence of IQGAP1 in the tails of *B. pseudomallei* in infected cells we next
31 utilised an IQGAP1 siRNA approach to determine the functional relevance of this host cell
32 protein in *B. pseudomallei* actin-based motility. Transfection of Silencer Select siRNA against
33 IQGAP1 into HeLa cells consistently resulted in efficient knockdown of around 70% as
34 assessed by IQGAP1 immunoblotting (Figure 6a and b). Following infection of siRNA
35 knockdown cells and control non-transfected cells, over 2000 bacteria were imaged from
36 each infected cell type across 3 biological replicates. There was no significant difference in
37 the proportion of tailing bacteria in either cell type under the assay conditions used (~42%).
38 It was evident that bacteria could still form actin tails in the knockdown cells however this
39 may be a result of the residual IQGAP1 expression in the cells, or compensation by the
40 related IQGAP proteins IQGAP2 or IQGAP3 which are also expressed in this cell type.

41
42 Using Image J software we next measured the length and actin density of tails as
43 described in the Methods section. 100 tails from control and siRNA transfected HeLa cells
44 across 3 biological replicates were analysed. The data showed a statistically significant
45 increase in overall tail length ($p=0.0033$) with concomitant decrease in actin density
46 ($p=0.0001$) in IQGAP1 knockdown cells compared with control cells (Figure 6c and d). In cells

1
2
3 with reduced levels of IQGAP1 tails were significantly longer with a mean length of 20.5µm
4 compared to a mean length of 14.8 µm in control cells (Figure 6c). Although longer in length
5 the actin density of tails generated in IQGAP1 knockdown cells was lower with a mean signal
6 of 1036.0 A.U. compared to a mean signal of 2522.4 A.U. measured for tails in control cells.
7
8
9

10 11 12 **Discussion**

13
14 *Burkholderia pseudomallei* is a facultative intracellular bacterium capable of escape
15 from the endocytic vacuole into the cytosol of host cells. In common with several other
16 bacteria, namely *Listeria monocytogenes*, *Shigella flexneri*, *Mycobacterium marinum* and
17 several *Rickettsia* spp., it is capable of harnessing cellular actin to promote its movement
18 within and between host cells by actin-based motility. It is clear that these bacteria have
19 evolved several independent mechanisms to utilise cellular actin and accessory proteins for
20 actin-based motility. For example *L. monocytogenes* expresses ActA, a protein which recruits
21 the host cell Arp2/3 complex to its surface where it directly interacts with actin and
22 stimulates its polymerisation²⁹. Conversely certain rickettsial species express Sca2, a protein
23 which acts through functional mimicry of eukaryotic formins²⁸. We have previously
24 characterised the *B. pseudomallei* encoded protein required for actin-based motility. Known
25 as BimA, this predicted Type V auto-transporter protein is expressed on a single pole of the
26 bacterium where it recruits cellular actin and stimulates its polymerisation to mediate
27 intracellular movement^{4, 6}. We and others have shown that this protein can stimulate actin
28 polymerisation *in vitro* in a manner that is independent of other host cell factors such as the
29 Arp2/3 complex^{4, 9}. Despite localisation of Arp2/3 components to the actin-rich tails of *B.*
30 *pseudomallei* in infected cells¹², tail formation does not require the activity of this cellular
31 actin nucleation complex since it can be observed in cells depleted of Arp2/3¹⁰. Whilst it is
32 known that actin-based motility of *B. pseudomallei* does not require Arp2/3, the potential
33 involvement of other cellular factors is unclear.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

49 Here we have used an *in vitro* affinity approach to identify cellular factors recruited to
50 BimA over-expressing bacteria under actin polymerising conditions. We report the
51 identification of 30 cellular proteins, including actin, actin-depolymerising factor and
52 capping protein, the minimal requirement for polymerisation in the presence of a bacterial
53 actin polymerisation factor *in vitro*³⁰. The identification of vinculin and α -actinin is
54
55
56
57
58
59
60

1
2
3 corroborated by the work of Breitbach *et al.* who demonstrated the presence of these
4 proteins in *B. pseudomallei* actin tails by immunofluorescence microscopy¹². Whilst α -
5 actinin is commonly detected in the tails of bacteria that display actin-based motility,
6 vinculin has been reported to be absent from *R. rickettsii* tails³¹.
7
8
9

10 In comparison to previous studies, it was notable that our dataset lacked any
11 microtubules, intermediate filaments or myosins that were shown to be present in *Listeria*
12 tails^{13b}, or VASP or profilin, which have been shown to be components of *R. rickettsia* actin
13 tails³¹. Two proteins that have not previously been associated with actin-based motility of
14 intracellular pathogens, HSP90 and IQGAP1, were identified in this analysis and the
15 presence of IQGAP1 detected in actin-rich tails formed by *B. pseudomallei* in HeLa cells.
16 IQGAP1 is a ubiquitously expressed scaffold protein that integrates numerous signalling
17 pathways and regulates multiple cellular processes. Involved in *Salmonella* Typhimurium
18 host cell invasion and *Chlamydomphila pneumoniae*-induced vascular smooth cell migration,
19 IQGAP1 has also been implicated in the adhesion of EPEC to the cell surface and viral
20 trafficking and replication of Mu-MuLV (reviewed in Kim *et al.*²⁷). IQGAP1 has been shown to
21 interact directly with the EPEC Ibe and Tir proteins and Salmonella T3SS effector protein
22 SseI, although the yeast 2 hybrid and pulldown results are not consistent with a direct
23 interaction between IQGAP1 and BimA. Indeed, the finding that the protein localises along
24 the length of the actin-rich tail in *B. pseudomallei* infected tails, together with reports that
25 IQGAP1 is an actin-binding protein, suggests it is recruited indirectly through its interaction
26 with actin filaments³². Furthermore we demonstrate that IQGAP1 plays a functional role in
27 organising the architecture of the actin tails formed by *B. pseudomallei*. Indeed siRNA
28 mediated knockdown resulted in a significant lengthening of the tails concomitant with a
29 reduction in the actin density. This warrants further investigation, as does the role of the
30 other validated and candidate host cell proteins in actin-based motility of this fascinating
31 bacterial pathogen.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

50 **Conclusion**

51 We have demonstrated a hitherto unknown role for the cellular protein IQGAP1 in
52 modulating the length and actin density of actin tails induced by the intracellular bacterial
53 pathogen *Burkholderia pseudomallei*.
54
55
56
57
58
59
60

Acknowledgements

This work was supported by the BBSRC (grant number BBSRC BB/E021212/1); and Institute Strategic Grant funding from the BBSRC (JS, MS, CVB, DK). Niramol Jitprasutwit was supported by a Siriraj Development Scholarship (Bangkok, Thailand). Nurhamimah Zainal-Abidin was supported by a scholarship awarded by Majlis Amanah Rakyat (MARA) (Council of Trust for the People, Malaysia). We thank Joann Prior and Andrew Scott at Dstl, Porton Down, UK for supplying the capsule specific antibody. We also thank Paul Brett and Mary Burtnick at the University of South Alabama, US for providing the pBHR2-*virAG* plasmid used in this study.

Statement on Dual Use Research of Concern

The work described in this manuscript was undertaken in the United Kingdom through funding obtained from UK sources and in strict accordance with UK Health and Safety Executive (HSE) guidance. Our work in relation to genetic modification has been extensively reviewed both by our own University of Edinburgh Biosafety Unit, the HSE and by the UK Counter Terrorism and Security Agency. The genetically modified *B. pseudomallei* strains described in this paper do not constitute a US Government Dual Use of Research Concern since these modifications do not enhance the pathogenicity of the micro-organism, do not interfere with immunity or immunization strategies, do not confer resistance to useful prophylactic or therapeutic agents and do not alter the transmissibility, host range or tropism of the pathogen.

Figure legends

Figure 1: *Burkholderia pseudomallei* BimA expression is not constitutive and can be induced by expression of VirAG *in trans*. (a) Western blot analysis of BimA expression in whole cell lysates of bacteria cultured in LB overnight at 37°C. Bacterial lysates were probed for BimA and BopE (as a protein loading control) using specific antibodies, and show that BimA is not expressed by wild-type *B. pseudomallei* under standard culture conditions. Expression *in trans* of the two component regulator VirAG results in BimA expression. Note that approximately 10-fold more 10276 and *bimA*::pDM4 bacteria (than their *pvirAG*-containing counterparts) were loaded onto the protein gel for analysis. (b) Western blot analysis of RAW cells infected with wild-type *B. pseudomallei* and a *bimA* mutant at an MOI of 100 at 3, 6 and 8 hours post-infection. Lysates were probed for BimA and *Burkholderia* capsule (as a protein loading control) using specific antibodies and indicate detection of BimA in lysates at the later time points post-infection.

Figure 2: Host cell and bacterial proteins are eluted from the surface of *B. pseudomallei* strains incubated with murine splenic lysate. A representative silver stained SDS-polyacrylamide gel showing the profile of proteins eluted from 10276 pBHR2-*virAG* and 10276 *bimA*::pDM4 pBHR2-*virAG* bacteria incubated with murine splenic lysates. The abundant presence of actin in the 10276 pBHR2-*virAG* sample is indicated by the arrow. Protein bands from samples from 3 independent experiments were processed for LC-MS/MS analysis.

Figure 3: Independent validation confirms the presence of a subset of host cell proteins in eluates from *B. pseudomallei* strains incubated with murine splenic lysate. Independent protein eluates from 10276 pBHR2-*virAG* and 10276 *bimA*::pDM4 pBHR2-*virAG* bacteria incubated with murine splenic lysates were prepared and immunoblotted to confirm the specific presence of IQGAP1, vinculin, HSP90 and L-plastin in the 10276 pBHR2-*virAG* samples. 5µg and 10µg of total splenic lysate were loaded as positive controls for the antibodies.

1
2
3 **Figure 4: IQGAP1 is recruited to *B. pseudomallei* actin tails in infected HeLa cells.** HeLa cells
4 were infected with *B. pseudomallei* at an MOI of 100 and fixed at 16 hours post-infection for
5 visualisation of actin tails by immunohistochemistry and confocal microscopy.
6 Representative images of bacteria (blue) and actin (green) are shown in panel (a), and
7 bacteria (blue) and IQGAP1 (red) in panel (b). The white arrows indicate areas of co-
8 localisation of actin and IQGAP1 in bacterial tails. Panel (c) represents a merged image of (a)
9 and (b). (d) illustrates the architecture of the actin tails, where the scale bar is 5µm.
10
11
12
13
14
15
16

17 **Figure 5: BimA and IQGAP1 do not interact in a yeast two-hybrid assay.** To identify protein:
18 protein interactions, single yeast colonies from SD agar plates lacking leucine and
19 tryptophan (DDO: Double Drop Out) were suspended in 20µl water and 5µl dotted onto SD
20 agar plates lacking leucine, tryptophan, adenine and histidine (QDO: Quadruple Drop Out)
21 supplemented with 20µg/ml X-α-galactosidase. Plates were incubated at 30°C for 24 hours
22 before observing the appearance of the colonies. The strains AH109 pGBKT7-*bimA* pGADT7-
23 *iqgap1* (BimA X IQGAP1) and AH109 pGBKT7-*bimA* pGADT7-*actin* (BimA X actin) were tested
24 alongside yeast strains transformed with positive and negative control vectors (pGADT7-T
25 antigen pGBKT7-p53 and pGADT7-T antigen pGADT7-Lamin C, respectively). The blue
26 colouration of the yeast indicates a direct interaction between the two proteins, as
27 demonstrated by the positive control and BimA X actin strains. White colouration indicates a
28 lack of interaction between the two proteins, as demonstrated by the negative control and
29 BimA X IQGAP1 strains.
30
31
32
33
34
35
36
37
38
39
40
41

42 **Figure 6: siRNA-mediated knockdown of IQGAP1 in HeLa cells affects *B. pseudomallei* tail**
43 **morphology.** (a) Western blot analysis of HeLa cell lysates from three independent
44 experiments. IQGAP1 and actin (as a protein loading control) were detected in lysates
45 prepared from cells treated only with transfection reagent (control) and cells transfected
46 with 6pmol siIQGAP1 for 72 hours (IQGAP1 siRNA). (b) Quantitation of IQGAP1 protein
47 levels in control and IQGAP1 siRNA cells indicates that IQGAP1 expression is reduced by
48 ~70% in siRNA-treated cells. Control and IQGAP1 siRNA cells were infected with *B.*
49 *pseudomallei* at an MOI of 100 and fixed at 16 hours post-infection for visualisation of actin
50 tails by immunohistochemistry and confocal microscopy. Images were collected from three
51
52
53
54
55
56
57
58
59
60

1
2
3 independent experiments and tail length in μm (c) and actin density in A.U. of phalloidin
4 intensity (d) assessed using ImageJ software.
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Host cell proteins isolated from BimA-expressing *Burkholderia pseudomallei*.

| Gene ID | Gene Name | Description | Protein Accession No. | Protein Score | # Unique Peptides | Notes |
|--------------------|---------------------------|--|-----------------------|---------------|-------------------|-----------------------|
| IPI00131138 | <i>flna</i> | Filamin A | Q8BTM8 | 104 | 32 | Lm |
| IPI00753793 | <i>spna2</i> | Spectrin alpha chain/ fodrin alpha chain | P16546 | 176 | 50 | Lm |
| IPI00319830 | <i>spnb2</i> | Spectrin beta chain/ fodrin beta chain | Q62261 | 79 | 8 | Lm |
| IPI00465786 | <i>tln1</i> | Talin 1 | P26039 | 91 | 8 | Lm |
| IPI00467447 | <i>iqgap1</i> | IQ motif containing GTPase activating protein 1, IQGAP1 | Q9JKF1 | 107 | 49 | |
| IPI00405227 | <i>vcl</i> | Vinculin | Q64727 | 67 | 12 | Lm |
| IPI00118899 | <i>actn4</i> | Alpha-actinin-4 | P57780 | 109 | 31 | Lm |
| IPI00380436 | <i>actn1</i> | Alpha-actinin-1 | Q7TPR4 | 116 | 38 | Lm |
| IPI00229080 | <i>hsp90ab1</i> | Heat shock protein 90kDa alpha | Q71LX8 | 79 | 17 | |
| IPI00169731 | <i>fermt3</i> | Fermitin family homolog 3/ kindlin 3 | Q8K1B8 | 55 | 1 | |
| IPI00118892 | <i>lcp1</i> | Lymphocyte cytosolic protein 1/ plastin -2/ L-plastin | Q61233 | 58 | 3 | Lm (T-plastin) |
| IPI00338302 | <i>pgm2</i> | Phosphoglucomutase-2 | Q7TSV4 | 43 | 2 | |
| * | <i>ezr/msn/rdx</i> | Ezrin/ Radixin/ Moesin** | ** | 61 | 4 | Lm (Ezrin) |
| IPI00314748 | <i>wdr1</i> | WD repeat-containing protein 1/ AIP1 | O88342 | 61 | 2 | Lm |
| IPI00110827 | <i>acta1</i> | Alpha actin | P68134 | 98 | 34 | Lm |
| IPI00125143 | <i>arpc1b</i> | Actin-related protein 2/3 complex subunit 1b | Q91Z25 | 58 | 2 | |
| IPI00473320 | <i>actb</i> | Beta actin-like protein | Q3U804 | 140 | 117 | Lm |
| IPI00874482 | <i>actg1</i> | Gamma actin | P63260 | 140 | 121 | Lm |
| IPI00110850 | <i>actb</i> | Beta actin | P60710 | 103 | 34 | Lm |
| IPI00277930 | <i>capg</i> | Macrophage-capping protein | Q99LB4 | 96 | 6 | |
| IPI00830701 | <i>tpm1</i> | Tropomyosin 1 | P58771 | 102 | 11 | Lm |
| IPI00111556 | <i>ppp2cb</i> | Serine/ threonine phosphatase 2A catalytic subunit | P62715 | 62 | 1 | |
| IPI00111265 | <i>capza2</i> | F-actin-capping protein subunit alpha-2 | P47754 | 77 | 1 | Lm |
| IPI00874728 | <i>tpm2</i> | Tropomyosin 2 | P58774 | 102 | 13 | Lm |

| | | | | | | | |
|----|---|----------------------|--|---------------|-----------|----------|-----------|
| 1 | | | | | | | |
| 2 | | | | | | | |
| 3 | | | | | | | |
| 4 | | | | | | | |
| 5 | IPI00775844 | <i>arpc2</i> | Actin-related protein 2/3 complex subunit 2 | Q9CVB6 | 71 | 3 | Lm |
| 6 | IPI00230044 | <i>tpm3</i> | Tropomyosin 3 | Q58E70 | 88 | 7 | Lm |
| 7 | IPI00116498 | <i>ywhaz</i> | 14-3-3 protein zeta/delta | P63101 | 71 | 6 | Lm |
| 8 | | | | | | | |
| 9 | IPI00125778 | <i>tagln2</i> | Transgelin 2 | Q9WVA4 | 69 | 2 | Lm |
| 10 | IPI00138691 | <i>arpc4</i> | Actin-related protein 2/3 complex subunit 4 | P59999 | 51 | 5 | Lm |
| 11 | IPI00890117 | <i>cfl1</i> | Cofilin-1 | P18760 | 100 | 3 | Lm |
| 12 | | | | | | | |
| 13 | | | | | | | |
| 14 | | | | | | | |
| 15 | *Peptides matched ezrin (IPI00330862, P26040), moesin (IPI00110588, P26041) and radixin (IPI00308324, P26043) | | | | | | |
| 16 | | | | | | | |
| 17 | | | | | | | |
| 18 | | | | | | | |
| 19 | | | | | | | |
| 20 | | | | | | | |
| 21 | | | | | | | |
| 22 | | | | | | | |
| 23 | | | | | | | |
| 24 | | | | | | | |
| 25 | | | | | | | |
| 26 | | | | | | | |
| 27 | | | | | | | |
| 28 | | | | | | | |
| 29 | | | | | | | |
| 30 | | | | | | | |
| 31 | | | | | | | |
| 32 | | | | | | | |
| 33 | | | | | | | |
| 34 | | | | | | | |
| 35 | | | | | | | |
| 36 | | | | | | | |
| 37 | | | | | | | |
| 38 | | | | | | | |
| 39 | | | | | | | |
| 40 | | | | | | | |
| 41 | | | | | | | |
| 42 | | | | | | | |
| 43 | | | | | | | |
| 44 | | | | | | | |
| 45 | | | | | | | |
| 46 | | | | | | | |
| 47 | | | | | | | |
| 48 | | | | | | | |
| 49 | | | | | | | |

1
2
3 List of murine proteins isolated either solely or at a higher frequency from eluates from
4 BimA-expressing bacteria 10276 pBHR2-*virAG* compared to the *bimA* mutant strain 10276
5 *bimA::pDM4* pBHR2-*virAG*. Proteins were assigned when peptides were isolated in at least 2
6
7 experiments, with a MASCOT score of at least 30, at least one rank 1 peptide, where the
8 predicted mass of the protein matched the observed mass and where the reported cellular
9 localisation was listed as cytoplasm or plasma membrane. Proteins highlighted in grey and
10 shown in bold were isolated in all 3 experiments. Proteins are listed in order of molecular
11 weight (highest to lowest) with 'Lm' in the Notes column denoting the identification of this
12 protein in similar proteomic studies of *Listeria monocytogenes* tails.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 2: *Burkholderia pseudomallei* proteins identified in the samples.

| Gene ID | Gene Name | Description | Protein Accession No. | Protein Score | # Unique Peptides | Notes |
|----------|--------------|--|-----------------------|---------------|-------------------|-------|
| bpsl0779 | <i>sucC</i> | Succinyl-CoA synthetase subunit beta | YP_107404.1 | 88 | 2 | |
| bpsl1013 | <i>ppc</i> | Phosphoenolpyruvate carboxylase | YP_107641.1 | 67 | 4 | |
| bpsl1087 | <i>htpG</i> | Heat shock protein 90 | YP_107709.1 | 103 | 4 | |
| bpsl1402 | <i>tig</i> | Trigger factor | YP_108024.1 | 69 | 6 | |
| bpsl1405 | <i>lon</i> | ATP-dependent protease | YP_108027.1 | 96 | 6 | |
| bpsl2192 | <i>aceB</i> | Malate synthase | YP_108787.1 | 80 | 4 | |
| bpsl2270 | <i>eno</i> | Phosphopyruvate hydratase | YP_108866.1 | 128 | 3 | |
| bpsl2305 | | Oligopeptidase A | YP_108901.1 | 117 | 2 | |
| bpsl2697 | <i>groEL</i> | Chaperonin GroEL | YP_109293.1 | 157 | 38 | OM, I |
| bpsl2698 | <i>groES</i> | Co-chaperonin GroES | YP_109294.1 | 169 | 8 | I |
| bpsl2827 | <i>dnaK</i> | Molecular chaperone DnaK | YP_109422.1 | 107 | 6 | OM, I |
| bpsl2953 | <i>tktA</i> | Transketolase | YP_109547.1 | 101 | 8 | |
| bpsl3004 | <i>rpmA</i> | 50S ribosomal protein L27 | YP_109599.1 | 109 | 2 | |
| bpsl3196 | <i>rpsE</i> | 30S ribosomal protein S5 | YP_109790.1 | 78 | 5 | |
| bpsl3215 | <i>tuf</i> | Elongation factor Tu | YP_109809.1 | 104 | 9 | |
| bpsl3220 | <i>rpoC</i> | DNA-directed RNA polymerase subunit beta' | YP_109814.1 | 102 | 23 | |
| bpsl3221 | <i>rpoB</i> | DNA-directed RNA polymerase subunit beta | YP_109815.1 | 117 | 22 | |
| bpss0421 | <i>rfbH</i> | Lipopolysaccharide biosynthesis protein (O-antigen-related) | YP_110445.1 | 45 | 1 | I |
| bpss0879 | | Porin protein | YP_110888.1 | 83 | 2 | OM, I |
| bpss0913 | | Methionine gamma-lyase | YP_110922.1 | 104 | 4 | |
| bpss1356 | | Hypothetical protein | YP_111366.1 | 133 | 22 | OM |
| bpss1498 | <i>hcp</i> | Hypothetical protein (Type VI secretion system secreted protein Hcp) | YP_111505.1 | 112 | 14 | |
| bpss1715 | <i>glTA</i> | Type II citrate synthase | YP_111721.1 | 81 | 7 | |
| bpss1722 | <i>mdh</i> | Malate dehydrogenase | YP_111728.1 | 107 | 5 | I |
| bpss1726 | <i>acnA</i> | Aconitate hydratase | YP_111732.1 | 118 | 20 | |

1
2
3 List of *Burkholderia pseudomallei* proteins isolated from eluates from both 10276 pBHR2-
4 *virAG* and 10276 *bimA*::pDM4 pBHR2-*virAG* bacteria. Proteins were assigned when peptides
5 were isolated from both samples in at least 2 experiments, with a MASCOT score of at least
6 30, at least one rank 1 peptide and where the predicted mass of the protein matched the
7 observed mass. Proteins highlighted in grey and shown in bold were isolated in all 3
8 experiments. In the Notes column, 'OM' indicates that the protein has previously been
9 described as surface associated or a component of the *B. pseudomallei* outer membrane
10 proteome^{22, 24}. 'I' indicates the protein is recognised by recovering human melioidosis
11 patient sera²⁴⁻²⁵. Proteins are listed in order of *B. pseudomallei* gene number with
12 chromosome 1 encoded proteins (BPSLxxxx) listed first.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1. Wiersinga, W. J.; van der Poll, T.; White, N. J.; Day, N. P.; Peacock, S. J., Melioidosis: insights into the pathogenicity of *Burkholderia pseudomallei*. *Nature reviews. Microbiology* **2006**, *4* (4), 272-82.
2. Stevens, J. M.; Galyov, E. E.; Stevens, M. P., Actin-dependent movement of bacterial pathogens. *Nature reviews. Microbiology* **2006**, *4* (2), 91-101.
3. Qualmann, B.; Kessels, M. M., New players in actin polymerization--WH2-domain-containing actin nucleators. *Trends in cell biology* **2009**, *19* (6), 276-85.
4. Stevens, M. P.; Stevens, J. M.; Jeng, R. L.; Taylor, L. A.; Wood, M. W.; Hawes, P.; Monaghan, P.; Welch, M. D.; Galyov, E. E., Identification of a bacterial factor required for actin-based motility of *Burkholderia pseudomallei*. *Molecular microbiology* **2005**, *56* (1), 40-53.
5. Sitthidet, C.; Stevens, J. M.; Chantratita, N.; Currie, B. J.; Peacock, S. J.; Korbsrisate, S.; Stevens, M. P., Prevalence and sequence diversity of a factor required for actin-based motility in natural populations of *Burkholderia* species. *Journal of clinical microbiology* **2008**, *46* (7), 2418-22.
6. Sitthidet, C.; Korbsrisate, S.; Layton, A. N.; Field, T. R.; Stevens, M. P.; Stevens, J. M., Identification of motifs of *Burkholderia pseudomallei* BimA required for intracellular motility, actin binding, and actin polymerization. *Journal of bacteriology* **2011**, *193* (8), 1901-10.
7. Lazar Adler, N. R.; Stevens, M. P.; Dean, R. E.; Saint, R. J.; Pankhania, D.; Prior, J. L.; Atkins, T. P.; Kessler, B.; Nithichanon, A.; Lertmemongkolchai, G.; Galyov, E. E., Systematic Mutagenesis of Genes Encoding Predicted Autotransported Proteins of *Burkholderia pseudomallei* Identifies Factors Mediating Virulence in Mice, Net Intracellular Replication and a Novel Protein Conferring Serum Resistance. *PLoS one* **2015**, *10* (4), e0121271.
8. Stevens, J. M.; Ulrich, R. L.; Taylor, L. A.; Wood, M. W.; Deshazer, D.; Stevens, M. P.; Galyov, E. E., Actin-binding proteins from *Burkholderia mallei* and *Burkholderia thailandensis* can functionally compensate for the actin-based motility defect of a *Burkholderia pseudomallei* bimA mutant. *Journal of bacteriology* **2005**, *187* (22), 7857-62.
9. (a) Sitthidet, C.; Stevens, J. M.; Field, T. R.; Layton, A. N.; Korbsrisate, S.; Stevens, M. P., Actin-based motility of *Burkholderia thailandensis* requires a central acidic domain of BimA that recruits and activates the cellular Arp2/3 complex. *Journal of bacteriology* **2010**, *192* (19), 5249-52; (b) Benanti, E. L.; Nguyen, C. M.; Welch, M. D., Virulent *Burkholderia* species mimic host actin polymerases to drive actin-based motility. *Cell* **2015**, *161* (2), 348-60.
10. Lu, Q.; Xu, Y.; Yao, Q.; Niu, M.; Shao, F., A polar-localized iron-binding protein determines the polar targeting of *Burkholderia* BimA autotransporter and actin tail formation. *Cellular microbiology* **2015**, *17* (3), 408-24.
11. (a) Wong, J.; Chen, Y.; Gan, Y. H., Host Cytosolic Glutathione Sensing by a Membrane Histidine Kinase Activates the Type VI Secretion System in an Intracellular Bacterium. *Cell host & microbe* **2015**, *18* (1), 38-48; (b) Schell, M. A.; Ulrich, R. L.; Ribot, W. J.; Brueggemann, E. E.; Hines, H. B.; Chen, D.; Lipscomb, L.; Kim, H. S.; Mrazek, J.; Nierman, W. C.; Deshazer, D., Type VI secretion is a major virulence determinant in *Burkholderia mallei*. *Molecular microbiology* **2007**, *64* (6), 1466-85.

- 1
2
3 12. Breitbach, K.; Rottner, K.; Klocke, S.; Rohde, M.; Jenzora, A.; Wehland, J.; Steinmetz, I., Actin-based motility of *Burkholderia pseudomallei* involves the Arp 2/3 complex, but not N-WASP and Ena/VASP proteins. *Cellular microbiology* **2003**, *5* (6), 385-93.
- 7 13. (a) David, V.; Gouin, E.; Troys, M. V.; Grogan, A.; Segal, A. W.; Ampe, C.; Cossart, P., Identification of cofilin, coronin, Rac and capZ in actin tails using a *Listeria* affinity approach. *Journal of cell science* **1998**, *111* (Pt 19), 2877-84; (b) Van Troys, M.; Lambrechts, A.; David, V.; Demol, H.; Puype, M.; Pizarro-Cerda, J.; Gevaert, K.; Cossart, P.; Vandekerckhove, J., The actin propulsive machinery: the proteome of *Listeria monocytogenes* tails. *Biochemical and biophysical research communications* **2008**, *375* (2), 194-9.
- 14 14. Maegraith, B. G.; Leithead, C. S., Melioidosis: A Case-Report. *Lancet* **1964**, *1* (7338), 862-3.
- 17 15. Choi, K. H.; Kumar, A.; Schweizer, H. P., A 10-min method for preparation of highly electrocompetent *Pseudomonas aeruginosa* cells: application for DNA fragment transfer between chromosomes and plasmid transformation. *Journal of microbiological methods* **2006**, *64* (3), 391-7.
- 21 16. Milton, D. L.; O'Toole, R.; Horstedt, P.; Wolf-Watz, H., Flagellin A is essential for the virulence of *Vibrio anguillarum*. *Journal of bacteriology* **1996**, *178* (5), 1310-9.
- 25 17. Logue, C. A.; Peak, I. R.; Beacham, I. R., Facile construction of unmarked deletion mutants in *Burkholderia pseudomallei* using *sacB* counter-selection in sucrose-resistant and sucrose-sensitive isolates. *Journal of microbiological methods* **2009**, *76* (3), 320-3.
- 29 18. Kersey, P. J.; Duarte, J.; Williams, A.; Karavidopoulou, Y.; Birney, E.; Apweiler, R., The International Protein Index: an integrated database for proteomics experiments. *Proteomics* **2004**, *4* (7), 1985-8.
- 33 19. Taylor, G. K.; Goodlett, D. R., Rules governing protein identification by mass spectrometry. *Rapid communications in mass spectrometry : RCM* **2005**, *19* (23), 3420.
- 37 20. Stevens, M. P.; Friebel, A.; Taylor, L. A.; Wood, M. W.; Brown, P. J.; Hardt, W. D.; Galyov, E. E., A *Burkholderia pseudomallei* type III secreted protein, BopE, facilitates bacterial invasion of epithelial cells and exhibits guanine nucleotide exchange factor activity. *Journal of bacteriology* **2003**, *185* (16), 4992-6.
- 41 21. Cuccui, J.; Milne, T. S.; Harmer, N.; George, A. J.; Harding, S. V.; Dean, R. E.; Scott, A. E.; Sarkar-Tyson, M.; Wren, B. W.; Titball, R. W.; Prior, J. L., Characterization of the *Burkholderia pseudomallei* K96243 capsular polysaccharide I coding region. *Infection and immunity* **2012**, *80* (3), 1209-21.
- 47 22. Schell, M. A.; Zhao, P.; Wells, L., Outer membrane proteome of *Burkholderia pseudomallei* and *Burkholderia mallei* from diverse growth conditions. *Journal of proteome research* **2011**, *10* (5), 2417-24.
- 51 23. Vander Broek, C. W.; Chalmers, K. J.; Stevens, M. P.; Stevens, J. M., Quantitative Proteomic Analysis of *Burkholderia pseudomallei* Bsa Type III Secretion System Effectors Using Hypersecreting Mutants. *Molecular & cellular proteomics : MCP* **2015**, *14* (4), 905-16.

- 1
2
3 24. Harding, S. V.; Sarkar-Tyson, M.; Smither, S. J.; Atkins, T. P.; Oyston, P. C.; Brown, K. A.; Liu,
4 Y.; Wait, R.; Titball, R. W., The identification of surface proteins of *Burkholderia pseudomallei*.
5 *Vaccine* **2007**, *25* (14), 2664-72.
6
- 7 25. (a) Su, Y. C.; Wan, K. L.; Mohamed, R.; Nathan, S., A genome level survey of *Burkholderia*
8 *pseudomallei* immunome expressed during human infection. *Microbes and infection / Institut*
9 *Pasteur* **2008**, *10* (12-13), 1335-45; (b) Felgner, P. L.; Kayala, M. A.; Vigil, A.; Burk, C.; Nakajima-
10 Sasaki, R.; Pablo, J.; Molina, D. M.; Hirst, S.; Chew, J. S.; Wang, D.; Tan, G.; Duffield, M.; Yang, R.;
11 Neel, J.; Chantratita, N.; Bancroft, G.; Lertmemongkolchai, G.; Davies, D. H.; Baldi, P.; Peacock, S.;
12 Titball, R. W., A *Burkholderia pseudomallei* protein microarray reveals serodiagnostic and cross-
13 reactive antigens. *Proceedings of the National Academy of Sciences of the United States of America*
14 **2009**, *106* (32), 13499-504.
15
16
- 17 26. (a) Nieves, W.; Asakrah, S.; Qazi, O.; Brown, K. A.; Kurtz, J.; Aucoin, D. P.; McLachlan, J. B.;
18 Roy, C. J.; Morici, L. A., A naturally derived outer-membrane vesicle vaccine protects against lethal
19 pulmonary *Burkholderia pseudomallei* infection. *Vaccine* **2011**, *29* (46), 8381-9; (b) Nieves, W.;
20 Petersen, H.; Judy, B. M.; Blumentritt, C. A.; Russell-Lodrigue, K.; Roy, C. J.; Torres, A. G.; Morici, L. A.,
21 A *Burkholderia pseudomallei* outer membrane vesicle vaccine provides protection against lethal
22 sepsis. *Clinical and vaccine immunology : CVI* **2014**, *21* (5), 747-54.
23
24
- 25 27. Kim, H.; White, C. D.; Sacks, D. B., IQGAP1 in microbial pathogenesis: Targeting the actin
26 cytoskeleton. *FEBS letters* **2011**, *585* (5), 723-9.
27
- 28 28. Haglund, C. M.; Choe, J. E.; Skau, C. T.; Kovar, D. R.; Welch, M. D., *Rickettsia Sca2* is a
29 bacterial formin-like mediator of actin-based motility. *Nature cell biology* **2010**, *12* (11), 1057-63.
30
- 31 29. (a) Welch, M. D.; Iwamatsu, A.; Mitchison, T. J., Actin polymerization is induced by Arp2/3
32 protein complex at the surface of *Listeria monocytogenes*. *Nature* **1997**, *385* (6613), 265-9; (b)
33 Welch, M. D.; Rosenblatt, J.; Skoble, J.; Portnoy, D. A.; Mitchison, T. J., Interaction of human Arp2/3
34 complex and the *Listeria monocytogenes* ActA protein in actin filament nucleation. *Science* **1998**, *281*
35 (5373), 105-8.
36
37
- 38 30. Loisel, T. P.; Boujemaa, R.; Pantaloni, D.; Carlier, M. F., Reconstitution of actin-based motility
39 of *Listeria* and *Shigella* using pure proteins. *Nature* **1999**, *401* (6753), 613-6.
40
- 41 31. Van Kirk, L. S.; Hayes, S. F.; Heinzen, R. A., Ultrastructure of *Rickettsia rickettsii* actin tails and
42 localization of cytoskeletal proteins. *Infection and immunity* **2000**, *68* (8), 4706-13.
43
44
- 45 32. (a) Mateer, S. C.; Morris, L. E.; Cromer, D. A.; Bensenor, L. B.; Bloom, G. S., Actin filament
46 binding by a monomeric IQGAP1 fragment with a single calponin homology domain. *Cell motility and*
47 *the cytoskeleton* **2004**, *58* (4), 231-41; (b) Erickson, J. W.; Cerione, R. A.; Hart, M. J., Identification of
48 an actin cytoskeletal complex that includes IQGAP and the Cdc42 GTPase. *The Journal of biological*
49 *chemistry* **1997**, *272* (39), 24443-7.
50
51
52
53
54
55
56
57
58
59
60

Synopsis description and image

The BimA protein of the melioidosis pathogen *Burkholderia pseudomallei* recruits and polymerises cellular actin to promote motility within the host cell cytosol. The mechanism by which BimA subverts the cellular cytoskeletal machinery is understudied. We have used an affinity approach coupled with Mass Spectrometry to identify cellular proteins involved in tail formation, demonstrating that IQGAP1 participates in determining both the length and actin density of these structures.

For TOC only




Figure 1




Figure 1

254x190mm (96 x 96 DPI)

Figure 2




Figure 2

254x190mm (96 x 96 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 3




Figure 3

254x190mm (96 x 96 DPI)

Figure 4




Figure 4

254x190mm (96 x 96 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 5




Figure 5

254x190mm (96 x 96 DPI)

Figure 6




Figure 6

254x190mm (96 x 96 DPI)