



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Anosognosia for chronic dysphasia

Citation for published version:

Beschin, N, Reverberi, C & Della Sala, S 2018, 'Anosognosia for chronic dysphasia', *Cortex*.
<https://doi.org/10.1016/j.cortex.2018.08.023>

Digital Object Identifier (DOI):

[10.1016/j.cortex.2018.08.023](https://doi.org/10.1016/j.cortex.2018.08.023)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Cortex

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Clinical Postcard

ANOSOGNOSIA FOR CHRONIC DYSPHAGIA

Nicoletta Beschin ^a, Cristina Reverberi ^b and Sergio Della Sala ^{c*}

a Neuropsychological Service, Rehabilitation Unit, ASST Valle Olona, Somma Lombardo Hospital,
Italy

b SLT, HR, Azienda USL-IRCCS di Reggio Emilia, Italy

c Human Cognitive Neuroscience, Psychology, University of Edinburgh, Edinburgh, UK

* Corresponding author.

Human Cognitive Neuroscience, University of Edinburgh, 7 George Square, Edinburgh EH8 9JZ,
UK. E-mail address: sergio@ed.ac.uk (S. Della Sala).

Aldo (not real name), a man with 17 years of formal education, was 60-years old at the time of our assessment. He was affected by a probable Progressive Supranuclear Palsy which started four years before our examination. He was treated only with antidepressants. His neuropsychological profile was characterised by executive impairments (see Table 1), which reflected in some everyday behaviour (apathy, withdrawal, social inappropriateness and fatuity). Aldo also presented with balance difficulties, clear dysarthria and severe dysphagia, characterised by altered oral and pharyngeal phases, resulting in food sticking in his throat, choking, hawking and coughing.

--- Table 1 about here---

Aldo's dysphagia was accompanied by tongue protrusion and profuse sialorrhea, causing considerable embarrassment to his wife, but apparently not to him. Curiously, he apologised when his drooling was dripping on other people or on the testing desk solely when he was perceiving negative feedback from other people. Even drinking caused substantial anterior leakage and drooling, with nasal penetration, predeglutitive aspiration and oral residues. However, on the Movement Disorder Society - Unified Parkinson's Disease Rating Scale (MDS-UPDRS – Goetz et al. 2008), section 2.2 questioning about drooling, he self-rated himself zero, i.e. no drooling. He also rated himself normal (i.e., score zero) on sections 2.3 and 2.4 of the same scale, inquiring respectively about chewing/swallowing and eating tasks, indicating a complete denial of his clear dysphagia.

On the Functional Oral Intake Scale (Crary, Carnaby Mann & Groher, 2005) measuring severity of dysphagia, he was scored 4/7, indicating that he could be fed an oral diet of a single consistency. He was referred to a speech and language therapist who instructed him (and his wife) to adopt appropriate dietary and swallowing strategies and postural adjustments, to modify food texture and to increase the viscosity of liquids.

Aldo was perfectly aware of his motor and balance difficulties, as well as of his dysarthria, for which he required insistently to be treated and exercised thoroughly following the speech therapist's advice with his wife's assistance. On a self-rating questionnaire assessing dysarthria (Schindler & Gullí, 2002) he scored 68/140, whereas both his wife and his therapist scored his problems 46/140, indicating that he slightly overestimated his articulation and language problems.

On the contrary, he strongly denied his dysphagia often arguing with his wife who worried about his frequent choking, coughing, vomiting and breathing gasps. On the Swallowing Disturbance Questionnaire (Cohen & Manor, 2011) he scored 5/45, whereas his therapist scored his performance 28/45 and his wife 29/45, indicating a clear underestimation of his swallowing problems. For example, to the question “Do you have any difficulty when eating solid food?” he responded, “No problems at all, I am always hungry”; “Does food get stuck in your throat?”, “No, it never happens”. To the question “Do you sometime cough when drinking?” he replied, “No, never”. Unlike paresis or dysarthria, which are always present, responding to questionnaires about one’s own dysphagia requires the testee to rely on their memory of the symptom. However, Aldo was often confronted by his wife when he was coughing and choking whilst eating or drinking. He denied his dysphagia even when facing it. For example, his wife reported a recent argument they had, as he was gasping for air and almost suffocating after having swigged out of a bottle of water, a habit he was advised to avoid on numerous occasions. Pressed by his wife, disquieted by his irrepressible coughing, he denied any problems maintaining that it was normal to cough sometimes.

The only item in the questionnaire that he acknowledged as a potential problem was having a hoarse voice after eating, which paradoxically is a symptom rarely noticed by dysphagics.

He refused any help to improve his swallowing difficulties with either food or liquids, and never implemented any strategy to avoid possible suffocation. Spontaneously, he often complained of his language deficits, but never of his swallowing problems. Aldo kept denying his swallowing problems even after he was alerted to them by doctors, nurses, speech and physio therapists as well as by his relatives, nor did he acknowledge them after he was shown the outcome of the relevant instrumental diagnostic procedures. His denial for his swallowing difficulties did not change after the training, and he refused to adopt compensatory techniques.

Two years later his conditions further deteriorated yet his anosognosia for his dysphagia remained, whereas he continued to be overtly apprehensive about his dysarthria for which he sought insistent advice.

Dysphagia is not rare in Parkinson-like diseases (Suttrup & Warnecke, 2016; Simons 2017) and can be denied by the patients (Bayés-Rusiñol et al., 2011). Dysphagia has also been described in PSP (Müller et al., 2001). However, in all the cases one can glean from the literature, patients were asked to evaluate their swallowing problems before being confronted with objective assessments or

simultaneously to them. Therefore, their denial has been interpreted as due to inaccurate perception of swallowing abilities (Ding & Logemann, 2008).

The case of Aldo shows that anosognosia for dysphagia can manifest also in the face of clear evidence of the problem and repeated advice from the caring professionals and relatives, to the point of hampering training and resulting in the refusal of helping strategies. Aldo's denial of his very severe dysphagia was not part of a general picture of unawareness of his own deficits, as testified by his preoccupation for his dysarthria and his insistence in being treated for it, even if his articulatory deficits were far less severe and limiting in his daily and social life than his swallowing impairments.

References

- Bayés-Rusiñol, À., Forjaz, M.J., de la Cruz Crespo, A.A.M., et al. (2011). Consciencia de disfagia en la enfermedad de Parkinson. *Rev Neurol*, 53 (11), 664-672.
- Cohen, J.T., & Manor, Y. (2011). Swallowing disturbance questionnaire for detecting dysphagia. *Laryngoscope*, 121(7), 1383-1387.
- Crary, M., Carnaby Mann, G.D., & Groher, M.E. (2005). Initial psychometric assessment of a functional oral intake scale for dysphagia in stroke patients. *Archives Physical Medicine Rehabilitation*, 86, 1516-1520.
- Ding, R., & Logemann, J. A. (2008). Patient self-perceptions of swallowing difficulties as compared to expert ratings of videofluorographic studies. *Folia Phoniat Logop*, 60(3), 142-150.
- Goetz, C.G., Tilley, B.C., Shaftman, S.R., Stebbins, G.T., Fahn, S. et al. for the Movement Disorder Society UPDRS Revision Task Force. (2008). Movement Disorder Society-Sponsored Revision of the Unified Parkinson's Disease Rating Scale (MDS-UPDRS): Scale Presentation and Clinimetric Testing Results. *Movement Disorders*, 23(15), 2129-2170.
- Müller J1, Wenning GK, Verny M, McKee A, Chaudhuri KR, Jellinger K, Poewe W, & Litvan I. (2001). Progression of dysarthria and dysphagia in postmortem-confirmed parkinsonian disorders. *Arch Neurol.*, 58(2):259-64.
- Schindler, A., & Gulli, M. (2002). L'autovalutazione della disartria. In: Schindler, O., Venero, I., Gambino, M., & Schindler, A. (Eds.), *Cartella Logopedica. Età adulta ed involutiva*. Torino: Omega Edizioni.
- Simon J.A. (2017). Swallowing Dysfunction in Parkinson's Disease. *Int Rev Neurobiol*. 134, 1207-1238.
- Suttrup, I., & Warnecke, T. (2016). Dysphagia in Parkinson's disease. *Dysphagia*, 31(1), 24–32.

Table 1. Aldo's neuropsychological assessment four years post-onset. (References in Supplementary Material).

Test	Score
<i>General Cognition</i>	
MMSE	29/30
Milan overall dementia assessment (MODA)	93.9/100
<i>Intelligence and Reasoning</i>	
Verbal Judgements	51/60
Coloured Progressive Matrices (Raven)	27/36
<i>Attention and Executive Functions</i>	
Visual Search	52/60
Trail Making Test B-A	169
Stroop Test errors	9.5*
time	45.5*
Frontal Assessment Battery (FAB)	13/18*
Cognitive Estimation	20/42*
Bizarre responses	7/21*
Card Sorting Test (Wisconsin)	88/128*
Phonological (Letter) Fluency	13*
Semantic (Category) Fluency	28
<i>Language</i>	
Sentences Generation	17/18
Token Test	33/36
Picture Naming	43/48
<i>Memory</i>	
Prose memory Immediate recall	7.9/8
Delayed recall	8/8
Pair Associate Learning	12.5/22.5
Digit Span	5
Spatial (Corsi) Span	5
Complex Figure Recall (Rey)	9/36
<i>Perception</i>	
Complex Figure Copy (Rey)	27/36*
<i>Praxis</i>	
Object use	14/14
Oro-facial praxis	19/20
Gesture Imitation Right arm	72/72
Left arm	71/72
<i>Mood</i>	
Zung Depression Scale	54/80*#

* = Performance below cut-off

= Mild depression

SUPPLEMENTARY MATERIALS

References for the neuropsychological tests administered (see Table 1).

- Appollonio, I., Leone, M., Isella, V., Piamarta, F., Consoli, T., Villa, M.L., Forapani, E., Russo, A., & Nichelli, P. (2005). The Frontal Assessment Battery (FAB): normative values in an Italian population sample. *Neurological Sciences*, 26(2), 108-116.
- Basso, A., Capitani, E., & Laiacona, M. (1987). Raven's Coloured Progressive Matrices: normative values on 305 adult normal controls. *Functional Neurology*, 2, 189-194.
- Brazzelli, M., Capitani E., Della Sala, S., Spinnler, H., & Zuffi, M. (1996). A neuropsychological instrument adding to the description of patients with suspected cortical dementia: the Milan overall dementia assessment. *Journal of Neurology, Neurosurgery and Psychiatry*, 57, 1510-1517.
- Cafarra, P., Vezzadini, G., Dieci F., Zonato, A., & Venneri, A. (2002). Rey-Osterrieth Complex Figure: normative values in an Italian population sample. *Neurological Sciences*, 22, 443-447.
- Cafarra, P., Vezzadini, G., Dieci, F., Zonato, A., & Venneri, A. (2002). Una versione abbreviata del test di Stroop: dati normativi nella popolazione italiana. *Nuova Rivista di Neurologia*, 12(4), 111-115.
- Carlesimo, G.A., Buccione, I., Fadda, L., Graceffa, A., Mauri, M., Lorusso, S., Bevilacqua, G., & Caltagirone, C. (2002). Standardizzazione di due test di memoria per uso clinico. Breve racconto e Figura di Rey. *Nuova Rivista di Neurologia*, 12(1), 1-13.
- Catricalá, E., Della Rosa, P.A., Ginex, V., Mussetti, Z., Plebani, V., & Cappa, S.F. (2013). An Italian battery for the assessment of semantic memory disorders. *Neurological Sciences*, 34(6), 985-993.
- Della Sala, S., MacPherson, S. E., Phillips, L. H., Sacco, L., & Spinnler, H. (2003). How many camels are there in Italy? Cognitive estimates standardised on the Italian population. *Neurological Sciences*, 24, 10-15.
- De Renzi, E., Pieczuro, A., & Vignolo, L.A. (1968). Ideational Apraxia. *Neuropsychologia*, 6, 41-52.
- De Renzi, E., Motti, F., & Nichelli P. (1980). Imitating gestures. A quantitative approach to ideomotor apraxia. *Archives of Neurology*, 37, 6-10.

- Giovagnoli, A.R., Del Pesce, M., Mascheroni, S., Simoncelli, M., Laiacona, M., & Capitani, E. (1996). Trail making test: normative values from 287 normal adult controls. *Italian Journal of Neurological Sciences*, 17, 305-309.
- Laiacona, M., Inzaghi, M.G., De Tanti, A., & Capitani, E. (2000). Wisconsin Card Sorting Test: a new global score, with Italian norms, and its relationship with the Weigl sorting test. *Neurological Sciences*, 21, 279-291.
- Measso, G., Cavarzeran, F., Zappalà, G., Lebowitz, B.D., Crook, T.H., Pirozzolo, F.J. & et al. (1993). The Mini-Mental State Examination: Normative study of Italian random sample. *Developmental Neuropsychology* 1993, 9, 77-85.
- Novelli, G., Papagno, C., Capitani, E., Laiacona, M., Cappa, S.F., & Vallar, G. (1986). Tre test clinici di memoria verbale a lungo termine. Taratura su soggetti normali. *Archivio di Psicologia, Neurologia e Psichiatria*, (4)47, 278-295.
- Novelli, G., Papagno, C., Capitani, E., Laiacona, M., Vallar, G., & Cappa, S.F. (1986). Tre test clinici di ricerca e produzione lessicale. Taratura su soggetti normali. *Archivio di Psicologia, Neurologia e Psichiatria*, (4)47, 477-506.
- Orsini, A., Grossi, D., Capitani, E., Laiacona, M., Papagno, C., & Vallar, G. (1987). Verbal and spatial immediate memory span: Normative data from 1355 adults and 1112 children. *The Italian Journal of Neurological Sciences*, 8, 539-548.
- Spinnler, H.R., & Tognoni G. (Eds) (1987). Standardizzazione e taratura italiana di test neuropsicologici. *The Italian Journal of Neurological Sciences* suppl. 8.
- Zung, W.K., Richards, C.B., & Short, M.J. (1965). Self-Rating Depression Scale in an Outpatient Clinic. *Archives Gen. Psychiatry*, 13(6), 508-515.

