



**Manchester
Metropolitan
University**

Redfern, J and Enright, MC (2020) Further understanding of *Pseudomonas aeruginosa*'s ability to horizontally acquire virulence: possible intervention strategies. *Expert Review of Anti-Infective Therapy*. ISSN 1478-7210

Downloaded from: <http://e-space.mmu.ac.uk/625668/>

Version: Accepted Version

Publisher: Taylor and Francis

DOI: <https://doi.org/10.1080/14787210.2020.1751610>

Please cite the published version

<https://e-space.mmu.ac.uk>

Article highlights

A review of the state of the art of the fast-moving field of *Pseudomonas aeruginosa* genomics

Insight into the species population structure that shows the genetic diversity underlying its evolutionary and environmental flexibility.

Examination of high-risk multidrug resistant clone evolution and penetration into healthcare systems.

Consideration of alternatives and possible adjuncts to antibiotic chemotherapy including environmental control of AMR isolates.

Review possible interventions with potential to control or reduce the ability for *P. aeruginosa* to gain virulence via HGT.

Expert Opinion

Analysis of the genome of *Pseudomonas aeruginosa* has greatly intensified in recent years following the publication of the type strain PAO1 genome sequence in 2000. Phenotypically diverse and a coloniser of a large variety of natural habitats, this species is able to infect both plants and humans using the same virulence armamentarium. The genomes of 4660 *P. aeruginosa* are in the public domain and these have started to reveal the reasons for the species evolutionary success. For a species with a relatively large genome of about 6.3 million nucleotides coding for an estimated 5600 genes the essential core genome appears surprisingly small – only 321 genes. The vast majority of *P. aeruginosa* genes are shared between a small number of isolates with approximately half found in only one isolate. This unusually high level of genetic diversity is driven by frequent horizontal gene transfer events between members of the species as well as extra-species associations that are evidenced by, for example the acquisition of antibiotic resistance genes on mobile genetic elements such as plasmids containing carbapenem resistance and even colistin resistance determinants. Perhaps thankfully in this species, although multiple drug resistance (MDR) is commonplace even in environmental isolates, extreme (XDR) and pan-drug resistances are much less common and largely restricted to a relatively small number of genetic backgrounds. These high-risk clones can be easily identified by genome sequencing and there is good potential for this to become routine so that rapid diagnostics can provide antibiotic sensitivity and even virulence potential data in a clinically useful timeframe that can minimise the untargeted, empirical use of that valuable resources - antibiotics. Genomic sequencing is now used to trace *P. aeruginosa* outbreaks retrospectively but technology is currently sufficiently advanced to allow real-time outbreak analyses in hospitals and other healthcare settings. Currently this is financially unviable in most settings but the near future promises declining costs for genomic sequencing and future innovations will reduce the human resource requirements of current epidemiological analyses.

Pseudomonas aeruginosa exists almost exclusively in biofilms which provide challenges for therapeutic intervention due to the low metabolic activity of some cells, the inability of some antibiotics to adequately penetrate the biofilm matrix and the strong attachment between biofilms and human tissue and indwelling device materials. Biofilms are also obviously also the place where intra- and inter-species gene exchange occurs allowing virulence genes and antibiotic resistance genes to move. With the current dearth in antibiotic drug discovery for gram-negative nosocomial pathogens in particular the future of antimicrobial chemotherapy will need to be supplemented by other

1
2
3 approaches to target pathogen biofilms in humans and also perhaps in environmental reservoirs of
4 antibiotic resistance. Controlling the diverse biofilms associated with *P. aeruginosa* will be challenging
5 due to the multispecies nature of many infective biofilms however targeting the species quorum-
6 sensing network could be a possible anti-virulence strategy. Future therapies based on depleting
7 biofilm directly using recombinant dispersins or phage-encoded polysaccharide depolymerases could
8 also provide additional tools to limit the spread of *P. aeruginosa*. The removal or reduction of *P.*
9 *aeruginosa* from water sources or wastewater processes through photocatalysis or other low-energy
10 disinfection processes could also offer a promising future approach to reducing the reservoir of AMR
11 *P. aeruginosa* **before** they impact on human health.
12
13
14
15
16
17

18 **1. Introduction**

19 *Pseudomonas aeruginosa* is ubiquitous in many natural environments such as soils and bodies of
20 water, in addition to being a major cause of infections in humans, other animals and a cause of soft
21 rot infections in plants. This is due to its ability to survive and even thrive under a wide range of
22 temperatures, on different nutrient sources and the species inherent and acquired virulence and
23 resistances to antibiotics, disinfectants and other antimicrobial compounds [1]. It has been classified
24 as a major species associated with multiple antimicrobial resistance of urgent public health concern
25 by the Infectious Diseases Society of America [2], Centers for Disease Control and Prevention and the
26 World Health Organization. It is a leading cause of severe, life-threatening nosocomial human
27 infections and the major pathogen associated with lung infections in patients with cystic fibrosis [3, 4]
28 due to the pathogens various and successful virulence mechanisms.
29
30
31
32
33
34
35

36 **1.1 Virulence**

37
38 Damage to the bacterial host, and hence bacterial pathogenicity, is due to the action of virulence
39 factors, a multifactorial process whereby bacterially-derived molecules enable entry, replication, and
40 persistence that may lead to tissue damage [5]. A recent review by Diggle and Whiteley [6] described
41 the major virulence systems of *P. aeruginosa* virulence and these are shown in Figure 2.
42
43
44
45

46 Understanding virulence in *P. aeruginosa* has been a key research field for decades, with many factors
47 identified as common across infection of both humans and plants, for example *toxA*, *plcS* and *gacA*,
48 as well as the phenazine pigment compound pyocyanin [7, 8]. A measure of the complexity and our
49 lack of knowledge of the gene repertoire of the species is that we cannot discern human pathogenic
50 genotypes from those infecting plants. This is illustrated in Figure 1 by the very high genetic similarity
51 between clinical and non-clinical (environmental) isolates and the plant isolate PA14. Quorum sensing
52 (QS), the ability for bacteria to produce and detect signalling molecules (autoinducers) is known to
53 play an important role in virulence (Figure 3), with mutations in QS genes or specific blocking/targeting
54 of QS molecules able to reduce virulence of *P. aeruginosa* [9, 10, 11], although notably, such QS
55
56
57
58
59
60

1
2
3 mutants are not always avirulent [10]. Siderophores are small molecular weight molecules secreted
4 by bacteria for iron-chelating, and are also responsible for regulation of virulence. Pyoverdine, a major
5 siderophore in *P. aeruginosa*, is itself a virulence factor and regulates at least two others factors
6 (endotoxin A and an endoprotease) [12, 13]. Virulence factors can also be activated by
7
8
9
10 mechanochemical sensory systems (e.g. type-IV pili) once the *P. aeruginosa* cell recognises a surface
11
12 (for example the chp system) [14] aiding in colonisation and biofilm formation. Virulence factors can
13 also be secreted by *P. aeruginosa* and packaged into outer-membrane vesicles and delivered to a
14
15 target host at relatively long distances [15].
16

17
18 Horizontal gene transfer (HGT), the process where genetic material is passed between different
19 bacterial cells, is the most significant driver in shaping the genome of *P. aeruginosa* [16]. HGT can
20 occur due to a number of intra- and even inter-species processes including conjugation, transduction
21 and transformation. Transduction, the process by which genetic information is obtained as a
22
23 consequence of viral (bacteriophage) infection has long been studied in *P. aeruginosa* [17], considered
24
25 a process naturally occurring in the environment, and has even been demonstrated with 'large' jumbo-
26 phage [18]. Similarly, conjugation, the process by which bacteria transfer genetic material between
27
28 cells, is also well studied in *P. aeruginosa* [19]. Recent literature has examined the effect of conjugation
29
30 on antibiotic resistance [20], and how biofilm phenotype may encourage this [21]. Natural
31
32 transformation in *P. aeruginosa*, meaning the uptake of genetic material from the environment, is less
33
34 studied/observed in the literature [22], with the ability to uptake and recombine naked DNA has not
35
36 been proven conclusively to occur in this species [23] although numerous circumstantial pieces of
37
38 evidence support this mechanism of HGT. Bacteria also have mechanisms to resist HGT. Restriction-
39
40 modification (R-M) systems are a highly diverse group on enzymes that can be considered 'bacterial
41
42 immunity' by defending bacterial cells from extra-cellular DNA. It is also hypothesised that R-Ms play
43
44 a broader role in the bacterial cell, with a recent study demonstrating *P. aeruginosa* R-M systems
45
46 potentially impacting virulence [24]. CRISPR-Cas is another cellular mechanism that can defend a
47
48 bacterial cell from DNA uptake and integration by protecting against bacteriophage transduction [25].
49
50 One study of CRISPR-Cas systems within 672 genomes found reduced genome size to be associated
51
52 with particular CRISPR-Cas types compared to those without such a system [26], however, the wider
53
54 impact on HGT in this species is still unclear [27, 28].
55
56

57 Virulence in *Pseudomonas aeruginosa* is largely shaped by high rates of horizontal gene transfer (HGT)
58 [29] that transfers regions of DNA between cells. Pathogenicity islands (PI) are horizontally acquired
59

1
2
3 genetic elements present in the accessory genome [16] that are frequently found in virulent clinical
4 isolates of *P. aeruginosa*. The first such, called PAPI-1 and PAPI-2, contain homologues of genes found
5 in other human and plant pathogens [30]. A recent study found 11 out of 21 identified GI in an ocular
6 multi-drug resistant isolate of *P. aeruginosa* contained virulence genes. A recent analysis of 1,311 *P.*
7
8
9
10 *aeruginosa* pangenomes Freschi, Vincent [29] concluded HGT has a non-negligible effect on the
11 evolution of the bacteria, and are potentially implicated in the emergence of virulence. It has also
12 been suggested that virulence traits in *P. aeruginosa* are selected for and maintained even when
13 interaction with a human host is absent, potentially driven by interaction with other eukaryotic
14 organisms in the wider environment, e.g. amoeba [22]. As greater numbers of genomes for *P.*
15 *aeruginosa* become available, pangenome/comparative genome studies will continue shedding light
16 on the genomic diversity of virulence.
17
18
19
20
21

22
23 However, there is little discussion in the literature on the control of HGT, and its impact on virulence
24 and the wider issues of *P. aeruginosa* in healthcare settings. This review therefore seeks to discuss
25 these considerations in the wider context of *P. aeruginosa* as a human pathogen with consideration
26 of relevant original research on *Pseudomonas aeruginosa* pathogenicity and population structure with
27 a strong emphasis on recent large genomic and pan-genomic studies on this very successful, diverse
28 and adaptive bacterial species. All literature searches were performed in PubMed and Web of Science
29 in the period May 2019 – March 2020.
30
31
32
33

34 35 36 37 **1.2 Lessons from Genomics?**

38 Since the first genome sequence of a *P. aeruginosa* isolate (PAO1) was published in 2000 [31] the
39 species has been the subject of many genome sequencing studies - at the time of writing 4660
40 genomes are currently in the public domain making it one of the most studied pathogen species. PAO1
41 is the most commonly used *P. aeruginosa* laboratory strain and the subject of ongoing international
42 efforts to ascribe functions to its 5570 open reading frames, in a genome of 6,264,403bp, through the
43 Pseudomonas Genome DB project (accessible at <http://www.pseudomonas.com>) [32]. *P. aeruginosa*
44 genomes are approximately twice the size of that of the average bacterial species with around 6.3
45 million base pairs [31, 33]. What is clearly apparent from genomic sequencing efforts to date is that
46
47 the species has a very large pan-genome, the full complement of genes found in the species, with a
48 very small number of core - genes found in every isolate and an extremely large accessory genome. A
49 2019 study [34] using genomics and analysis of gene knockouts from isolates grown under different
50 conditions calculated that the number of core essential genes to be 321, which comprises only 6.6%
51 of the *P. aeruginosa* genome. A study the previous year by Freschi and colleagues [29] estimated that
52
53
54
55
56
57
58
59
60

1
2
3 the core genome was only 1.2% of the pan-genome based upon genomes that they sequenced or were
4 publicly available in 2016. Both of these studies and other large genome sequencing projects describe
5 a species with a minimal essential genome (compared with other bacterial species such as *S.*
6 *pneumoniae* and *C. jejuni*, which has approximately 851 and 866 core genes out of an estimated 1990
7 and 1623 genes respectively [35]). According to the Freschi study, 50.1% of unique genes are found in
8 single isolates.
9
10
11
12

13
14
15 Early DNA sequencing studies suggested *P. aeruginosa* was divided into two main groups represented
16 by PAO1 [31] - isolated from a chronic wound in the 1950s and PA14 [36]- an isolate from a plant soft-
17 rot infection, however recent studies show a much more complex population structure comprising at
18 least five major clades according to one genomic analysis [37] although data from a 2015 study of
19 human infection and environmental isolates not included in this analysis suggests a more complex
20 picture [26] (Figure 1). The particularly large accessory genome of *P. aeruginosa* is obviously a major
21 contributor to the species versatility that allows it to exploit different environmental niches as a
22 ubiquitous organism in soil and water and also as a commensal that can on occasion cause infections
23 in plants and animals including humans. The *P. aeruginosa* pan-genome contains many genes that are
24 unique to a particular isolate and provides the species with an extremely large variety of possible
25 phenotypes [38].
26
27
28
29
30
31
32
33
34

35 **1.3 Antibiotic Resistance**

36
37 Although antibiotic resistance (ABR) is not considered to be a virulence factor, however,
38 understanding ABR is essential due to the increasingly important role both play in the clinical setting
39 [39]. In its 2013 publication “Antibiotic Resistant Threats in the United States, 2013”, CDC reported
40 51,000 cases of *P. aeruginosa* infections a year. 6700 (13.1%) of these were multidrug resistant (MDR),
41 that is, untreatable with three or more antibiotic classes commonly used to treat such infections.
42 These were found to have caused 440 deaths a year in the study period [40]. European CDC data for
43 2018 show a variable rate of MDR in Europe overall, with 32.1% of *P. aeruginosa* isolates resistant to
44 at least one antibiotic, and 19.2% isolates resistant to two or more antibiotics [41]. This report
45 suggests large inter-country variations for resistance across all antibiotics, with generally higher
46 resistance percentages reported in southern and eastern Europe compared to northern Europe and
47 resistance to carbapenems, the most commonly used antibiotics to treat nosocomial gram-negative
48 infections, remains common in many EU/EEA countries. Resistance to polymyxin antibiotics (colistin
49 and polymyxin B), often referred to as the antibiotic class of last resort for gram-negative infections
50 still appears to be rare in Europe, the United States, Canada and Latin America with 0.5 – 1.1% of
51
52
53
54
55
56
57
58
59
60

1
2
3 isolates being resistant [42] although resistance rates were higher in China with 22.2% resistance
4 among XDR (eXtremely Drug-Resistant, ie, resistant to four or more antibiotic classes) *P. aeruginosa*
5 from bloodstream infections, according to a 2016 report [42, 43].
6
7

8 **1.4 Population Structure**

9
10 Recombination is the major mechanism driving sequence divergence in this species and this enables
11 the frequent acquisition and dissemination of both virulence and antibiotic resistance genes. The very
12 large pan-genome and large number of genes found in only one isolate indicate a very fluid species
13 [26, 37]. Analysis of the core genome also shows a highly recombining population. Multilocus
14
15 sequence typing (MLST), first described in 1998 for the characterization of *Neisseria meningitidis* [44],
16 has been applied to the study of all common bacterial pathogens, as well as fungal species. As the
17 method relies on DNA sequencing of specific regions of (typically) seven core metabolic /
18 housekeeping gene loci and the online deposition of sequence- and meta- data, it can be used to study
19 pathogen epidemiology at the local, national and international level. It has also been an invaluable
20 tool for studying the evolution and global spread of antibiotic resistant and / or hyper-virulent lineages
21 of species such as *Staphylococcus aureus* (including methicillin-resistant [MRSA] clones) [45],
22 *Streptococcus pneumoniae* [46], *Escherichia coli* [47] and of course *Pseudomonas aeruginosa* [48].
23
24 MLST has also been used to examine and compare evolutionary parameters between species.
25
26 Comparison of the alleles for individual species allows an estimation to be made of the ratio of
27 recombination (R) to point mutation (M) per MLST allele or per nucleotide. Two studies using this
28 methodology estimated that MLST loci nucleotides were changed 8.4 - 9.5 times more often by
29 recombination compared to point mutation which is much lower than the naturally transformable *S.*
30 *pneumoniae* (50 times) and *N. meningitidis* (80 times), species that have epidemic population
31 structures but are demonstrably naturally transformable [49]. However these latter two species have
32 much smaller pan-genomes presumably due to their niche adaptation to almost exclusively colonise
33 and infect mammals and have much lower numbers of prophage, plasmids and other mobile genetic
34 elements compared to *P. aeruginosa*.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59

60
In an era in which rapid, inexpensive genome sequencing is now widely available traditional MLST is rarely used and most epidemiological studies nowadays extract core-genome (cg)MLST data from assembled or unassembled whole genome sequences [50, 51] for comparison with database sequences and isolate information held at, for example, the PubMLST website [50, 51, 52]. This allows the identification of multiple antibiotic resistant “high risk” clones (see below) and phylogenetic and phylogeographic comparisons of users isolates with the >3300 isolates present in the *P. aeruginosa* database.

1.5 High Risk Clones

In common with other pathogens with epidemic population structures, *P. aeruginosa* is notable for the presence of genotypes that are over-represented in samples from human disease and carriage.

These so-called “high risk” clones are indicative of an epidemic population structure where genotypes increase in frequency through rapid transmission in hospital and other environments probably due to antibiotic selection and enhanced virulence in certain vulnerable patient populations. Clonal spread of *P. aeruginosa* genotypes belonging to clonal complex (CCs) 111, 175, 235 and 395 are associated with the global dissemination of multidrug resistant genes including metallo-beta lactamases *bla*_{VIM}, IMP, SPM and NDM that confer carbapenem resistance [53, 54] (Figure 1). Carbapenemase genes are much more common in high-risk clones compared to other isolates and these may also have enhanced virulence – one study found isolates belonging to CC111 and 235 clones were highly virulent in comparison to ST175, a third, less significant clone in an invertebrate infection model [55].

1.6 Biofilm / Secretion Systems

In common with many bacterial pathogens *P. aeruginosa* cells exist within complex microbial communities known as biofilm; found at surface-liquid, surface-air and liquid-air interfaces. *P. aeruginosa* can thrive in biofilm due to the extracellular polymeric substances (EPS) it can produce, helping the bacteria resist desiccation, mechanical removal and the actions of antibiotics and disinfectants [56]. When *P. aeruginosa* forms biofilm in human tissue, cells are difficult to eradicate and represent infectious foci that can lead to severe systemic disease. Polymicrobial biofilms are considered a “melting pot” for horizontal gene transfer within and between bacterial species [57], recognized as important niches for MDR evolution. Isolates belonging to the two high-risk CC 111 and 235 described earlier have been shown to produce significantly more biofilm compared to those from non-high risk clones [58]. Numerous links between virulence and biofilm have been reported in the literature, with the presence of *P. aeruginosa* in polymicrobial increasing virulence of other bacterial species [59]. For example, it is suggested that the virulence factor phenazinethe, the enzyme polyphosphate kinase (ppk) and the HD-GYP domain proteins can modulate/regulate biofilm in addition to virulence [60, 61, 62]. Whilst there is still considerable work to be done to unpick the relationship between *P. aeruginosa*, virulence, HGT and biofilm production, it is clear that acknowledging this interplay will be an important aspect of future research.

1.7 Bacteriophage and HGT of virulence: friend or foe?

1
2
3 Bacteriophage (phage) are viruses that infect bacteria, and are considered of significant interest for
4 their antibiofilm mechanism. Once a phage enters the cell, one of two life-cycles can occur. The lytic
5 cycle results in the destruction (or lysis) of the infected cell, preceded by transcription of the phage
6 DNA in order to synthesise new phage. The resultant destruction of the cell releases the newly formed
7 phage into the environment to continue infection of the bacterial community. Therefore, there is
8 significant interest in the ability for phage to infect and destroy bacterial cells (and particularly those
9 in biofilm), with numerous review articles available describing the breadth of research in this area e.g.
10 [63, 64].
11
12

13
14
15
16
17
18 However, some phage follow the second life cycle, known as the lysogenic cycle. Here, following
19 infection, the phage DNA integrates with the bacterial DNA by a process known as transduction (a
20 form of HGT), whilst the bacterial cell continues its usual lifestyle and remains able to reproduce. This
21 DNA, known as a mobile genetic element (MGE), can also carry genes that may increase a bacterial
22 cells ability to survive, including virulence factors and antimicrobial resistance mechanisms. MGE,
23 from phage and other sources, are well documented as an important avenue of evolution in *P.*
24 *aeruginosa* e.g. [65, 66, 67]. One large genomic study suggests that the presence in a *P. aeruginosa*
25 genome of phage transduction - targeting CRISPR-Cas systems is associated with reduced antibiotic
26 resistance [26]. Therefore, understanding this process and targeting the mobile genetic elements of
27 disease to stop the spread of virulence factors by HGT is of significant interest [68].
28
29
30
31
32
33

34 35 36 37 38 39 **2. Control Strategies**

40 41 42 **2.1 Antibiofilm approaches**

43
44 As described above, *P. aeruginosa* biofilm provides an ideal scenario to share virulence genes through
45 HGT. Therefore, understanding how to remove or control biofilm, and therefore limit potential HGT
46 of virulence genes, is an important and active research field. Various molecules have been identified
47 as having both antibiofilm and anti-virulence effects. For example, the flavanol quercetin [69], ZnO
48 nanoparticles [70], garlic [71] and cinnamon oil [72] all have been shown to have both antibiofilm and
49 anti-virulence effects. The genetic links between antibiofilm and anti-virulence activity has also been
50 explored, for example, in a *typA* (a GTPase) mutant that has demonstrably lower virulence and
51 reduced biofilm [73].
52
53
54
55
56
57
58
59

1
2
3 A more targeted approach to biofilm control relies on the ability for bacterial cells to revert from
4 biofilm to planktonic (unattached and free-floating), causing dispersal of cells. However, this process
5 is not easy, due to the cementing of cells into complex 3D structures which form biofilm [74].
6 Dispersins are molecules that trigger cell dispersal. For example, the signalling molecules nitric oxide
7 (NO) [75], succinate, glutamate, glucose [76] and the dendrimer FD2 [77] have all been shown to
8 trigger enhanced dispersal of *P. aeruginosa* cells from biofilm. Innovations in electrochemical
9 engineering may also provide one avenue for researchers to find new dispersins. In 2018, Robb
10 demonstrated the use of pyrolytic graphite electrodes to assess potential dispersal-inducing
11 properties of 2-aminoimidazole (2-AI) [78]. Such analytical technologies may enable microbiologists
12 to better understand and measure the dispersal potential of novel antimicrobial and anti-biofilm
13 molecules.
14
15
16
17
18
19
20
21
22
23

24 Another targeted approach for biofilm control is the use of depolymerase, enzymes which are able to
25 breakdown polymeric substances. Depolymerase produced by *P. aeruginosa* has been a topic of
26 research for decades [79], with much focus on bacteriophage-encoded depolymerases [80, 81]. It is
27 believed that phage with depolymerase can degrade extracellular polymeric substances (EPS - the
28 'cement' of the biofilm), gaining easier access to host receptors. This ability to degrade the EPS also
29 means the phage are likely able to infect cells within a biofilm more efficiently. For example, PT-6, a
30 *P. aeruginosa*-infecting phage was able to reduce viscosity of four different alginate polymers by two-
31 thirds within 15 minutes [82]. A recent review found 160 putative depolymerases from 143 phage,
32 including sialidases, levanases, xylosidases, dextranases, hyaluronidases, peptidases as well as
33 pectate/pectin lyases [83]. These represent potential anti-virulence strategies for the reduction of
34 prevention of *P. aeruginosa* biofilm formation. Whilst there is a lack of literature describing the impact
35 of antibiofilm measures on HGT, given the potential for HGT in biofilm (as discussed earlier),
36 controlling biofilm should be considered a potential intervention strategy for the potential of *P.*
37 *aeruginosa* to acquire virulence genes (and other genetic material). This is particularly important in
38 environments where HGT of virulence genes could lead to particularly negative outcomes e.g. clinical
39 biofilm.
40
41
42
43
44
45
46
47
48
49
50
51

52 **2.2 Controlling Quorum Sensing**

53 Quorum sensing is well known to be important in *P. aeruginosa* biofilm development [84], which as
54 described previously, can be considered an important enabler for HGT. Therefore, strategies to control
55 or reduce QS, and therefore reduce biofilm formation, may provide an indirect strategy for reducing
56 HGT. *P. aeruginosa* has at least four quorum-sensing signalling networks, Las, Rhl, Pqs and Iqs [85].
57
58
59
60

1
2
3 The presence of QS molecules is associated with the production of extracellular DNA [86] and elevated
4 rates of horizontal gene transfer (HGT) [87]. A study by Rossmann, Racek [88] found that high
5 concentrations of the universal AI-2 compound in an already established biofilm promoted the release
6 of phage capable of infecting other bacteria, increasing their virulence. In *E. coli*, transfer of antibiotic
7 resistant plasmids can be enhanced by the presence of autoinducers, however, QS inhibitors were
8 able to inhibit the transfer [89]. Various QS inhibitors have been demonstrated to block *P. aeruginosa*
9 QS molecules. A synthetic derivative of a naturally occurring furanone has demonstrated antagonistic
10 activity against *P. aeruginosa* QS, increasing susceptibility of biofilm to antibiotics and increased
11 clearance in a mouse model [90]. The compound meta-bromo-thiolactone (mBTL), has also
12 demonstrated inhibition of virulence factor production in *P. aeruginosa*, in addition to inhibiting
13 biofilm formation by partially inhibiting LasR and RhIL [91]. Other compounds such as the essential oil
14 6-Gingerol, produced by fresh ginger, has similarly demonstrated QS inhibition [92]. Furthermore,
15 presence of QS inhibitors has been shown to suppress the CRISPR-Cas9 system in *P. aeruginosa*, making
16 the bacterial cell more prone to death resulting from phage infection [93]. These are only an example
17 of the wide range of research currently ongoing to understand the impact of QS in *P. aeruginosa* and
18 many other microorganisms, a new and fast paced field [94]. Whilst specific literature describing the
19 impact of QS control on HGT of virulence genes is lacking, it is clear that QS control has huge potential
20 to indirectly limit HGT of virulence and the inevitable increase of isolate virulence.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36

37 2.3 Molecular Diagnostics

38 Rapid genomic or metagenomic sequencing from patient samples should soon deliver a future where
39 key features such as species diversity antibiotic sensitivity and virulence gene complement can be
40 determined in a clinically relevant timescale to guide appropriate and specific intervention strategies
41 including patient isolation and specific antibiotic regimens. Rapid, real-time outbreak analysis and *in*
42 *silico* resistome characterisation will lead to more targeted and thus more effective antibiotic
43 administration that will also reduce the diversifying selection on microbial communities that is
44 imposed by inappropriate antibiotic administration. The potential for Nextgen genomic sequencing to
45 deliver epidemiologically useful information within a clinically useful timeframe was first
46 demonstrated during an outbreak of *Staphylococcus aureus* in a paediatric setting in Cambridge, UK
47 in 2012 [95]. The potential for *in silico* resistome analysis as a tool to determine the antibiotic
48 sensitivity of *P. aeruginosa* has been successfully demonstrated in two separate studies [53, 96] and
49 NGS is becoming a commonly used tool in retrospective analyses of *P. aeruginosa* outbreaks, for
50 example in a 2014 study tracing an outbreak due to an XDR ST111 clone in the United Kingdom [97].
51
52
53
54
55
56
57
58
59
60

1
2
3 Therefore, there do not seem to be any significant barriers (except cost) to the routine use of genome
4 sequencing in hospital laboratories for outbreak control and antibiotic prescribing purposes. With few
5 therapeutic options for treating some *P. aeruginosa* infections, rapid information on antibiotic
6 sensitivity of an individual isolate could be crucial and it could be used to guide and minimise
7 inappropriate use of new antibiotics as they are developed.
8
9

10
11
12
13 The use of new agents such as therapeutic antibodies and antimicrobial peptides, perhaps co-
14 administered with antibiotics, could also be effectively targeted for the control of “high-risk” *P.*
15 *aeruginosa* genotypes based on rapid genomic profiling in hospital settings.
16
17
18

19 20 **2.4 Controlling environmental conditions likely to encourage HGT - photocatalysis**

21
22 AMR is increasingly recognised as a problem outside of the clinic with many organisms acquiring and
23 transmitting resistance and virulence genes within and between environmental, agricultural and
24 animal (including human) reservoirs. Thus, control of key human pathogens such as *P. aeruginosa* in
25 the environment could play a major role in reducing HGT of resistance and virulence loci.
26
27
28

29
30 Wastewater is considered a hotspot for horizontal gene transfer [98]. Overuse and potential misuse
31 of antibiotics in farming, industry and medicine (both human and veterinary) as well as irresponsible
32 use of pollutants such as heavy metals and disinfectants have resulted in the inevitable discharge into
33 waste water [99]. These conditions cause cellular stress, which alongside high bacterial numbers (and
34 subsequent populations of bacteriophage) and biofilm formation [100] have led to a uniquely blurred
35 boundary between the human and natural world where HGT is readily facilitated. Isolates of *P.*
36 *aeruginosa*, is a common member of the wastewater community, have demonstrated an increase in
37 QS production linked with potential virulence and biofilm formation [101]. More generally, transfer of
38 antimicrobial resistance genes (ARG) in wastewater has become a dominant line of research [102],
39 with suggestion that bacteria gaining ARGs via phage may also have an increased capacity for virulence
40 [103]. Therefore, interventions that target conditions within aquatic environments that promote HGT
41 should be a priority.
42
43
44
45
46
47
48
49
50
51

52
53 Advanced treatment technologies such as photocatalysis have demonstrated efficiency at removing
54 contaminants from wastewater [102]. Over recent years, photocatalysis, a relatively simple, yet
55 efficient technique of mineralizing organic contaminants to carbon dioxide and water without the
56 production of secondary pollutants, has gained popularity as a safe, cheap and sustainable method
57 for water decontamination [104, 105]. Titanium dioxide (Titania, TiO₂) is the most studied
58
59
60

1
2
3 photocatalytic material for decontamination, depollution and disinfection processes [105, 106], killing
4
5 *P. aeruginosa* (and other bacteria) through the generation of radical oxygen species [107, 108, 109,
6
7 110], and can be favoured over other antimicrobial active substances due to its long-term
8
9 antimicrobial action [111]. In addition to killing bacteria, photocatalysis has also been shown to break
10
11 down viruses, and a wide variety of organic materials [112, 113, 114]. All of these properties target
12
13 and reduce the factors that can increase HGT (large number of bacterial cells, pollutants and biofilm).

14
15 Despite the obvious benefits of such an antimicrobial material, fundamental questions relating to the
16
17 effect of photocatalytic coatings on the microbiology of water have been raised in the literature.
18
19 Critically, some advanced treatment technologies do not provide a rapid bacterial kill, and will create
20
21 conditions likely to cause cellular stress and potentially leading to the cell SOS response, encouraging
22
23 horizontal gene transfer.

24 25 **3. Conclusion**

26
27 *Pseudomonas aeruginosa* occupies a huge variety of environmental niches being a ubiquitous in many
28
29 soils and watercourse habitats. These isolates are frequently indistinguishable from those colonising
30
31 and infecting humans. The species diversity and genome plasticity reflects this – the species has a very
32
33 small core genome of a few hundred genes representing 1% of the species genes discovered thus far.
34
35 Approximately half of the genes found in *P. aeruginosa* have only been seen in one isolate. A
36
37 significant proportion of this pan- genome especially that related to antibiotic resistance is mobile,
38
39 present on elements such as plasmids, transposons and insertion sequences however the
40
41 chromosome itself is also subject to high rates of recombinational change through HGT. The rapid
42
43 detection and characterization of globally disseminated “High Risk” genotypes (clones) that are
44
45 particularly associated with enhanced virulence and MDR, XDR and PDR resistance to antibiotics,
46
47 through molecular diagnostics should enable more targeted treatments based on inferred antibiotic
48
49 sensitivity from genome data or where no antibiotic is available, treatment with other agents in
50
51 development such as therapeutic monoclonal antibodies or antimicrobial peptides. Anti-virulence
52
53 molecules that interfere with inter-cell quorum sensing and signalling or approaches that destroy
54
55 mixed strain and species biofilms where gene exchange occurs at high frequency represent future
56
57 approaches for limiting the proliferation of *P. aeruginosa* genes for antibiotic resistance and virulence
58
59 that represent an increasing threat to human health.

60 61 **Funding statement**

The authors received no specific funding for this work.

References

1. Silby MW, Winstanley C, Godfrey SAC, et al. *Pseudomonas* genomes: diverse and adaptable. *FEMS Microbiol Rev.* 2011;35(4):652-680. doi: 10.1111/j.1574-6976.2011.00269.x.
2. Boucher HW, Talbot GH, Bradley JS, et al. Bad bugs, no drugs: no ESKAPE! An update from the Infectious Diseases Society of America. *Clin Infect Dis.* 2009 Jan 1;48(1):1-12. doi: 10.1086/595011. PubMed PMID: 19035777.
3. Davies JC. *Pseudomonas aeruginosa* in cystic fibrosis: pathogenesis and persistence. *Paediatr Respir Rev.* 2002 2002/06/01;3(2):128-134. doi: [https://doi.org/10.1016/S1526-0550\(02\)00003-3](https://doi.org/10.1016/S1526-0550(02)00003-3).
4. Sousa AM, Pereira MO. *Pseudomonas aeruginosa* Diversification during Infection Development in Cystic Fibrosis Lungs—A Review. *Pathogens.* 2014;3(3):680-703. PubMed PMID: doi:10.3390/pathogens3030680.
5. Cross AS. What is a virulence factor? *Crit Care.* 2008;12(6):196-196. doi: 10.1186/cc7127. PubMed PMID: 19090973; eng.
6. Diggle SP, Whiteley M. Microbe Profile: *Pseudomonas aeruginosa*: opportunistic pathogen and lab rat. *Microbiology.* 2019 Oct 10. doi: 10.1099/mic.0.000860. PubMed PMID: 31597590.
7. Rahme LG, Stevens EJ, Wolfort SF, et al. Common Virulence Factors for Bacterial Pathogenicity in Plants and Animals. *Science.* 1995;268(5219):1899-1902.
8. Mahajan-Miklos S, Tan M-W, Rahme LG, et al. Molecular Mechanisms of Bacterial Virulence Elucidated Using a *Pseudomonas aeruginosa* – *Caenorhabditis elegans* Pathogenesis Model. *Cell.* 1999;96(1):47-56. doi: 10.1016/S0092-8674(00)80958-7.
9. and MBM, Bassler BL. Quorum Sensing in Bacteria. *Annu Rev Microbiol.* 2001;55(1):165-199. doi: 10.1146/annurev.micro.55.1.165. PubMed PMID: 11544353.
10. Smith RS, Iglewski BH. *P. aeruginosa* quorum-sensing systems and virulence. *Curr Opin Microbiol.* 2003 2003/02/01;6(1):56-60. doi: [https://doi.org/10.1016/S1369-5274\(03\)00008-0](https://doi.org/10.1016/S1369-5274(03)00008-0).
11. Paczkowski JE, Mukherjee S, McCready AR, et al. Flavonoids suppress *Pseudomonas aeruginosa* virulence through allosteric inhibition of quorum-sensing receptors. *J Biol Chem.* 2017 January 24, 2017. doi: 10.1074/jbc.M116.770552.
12. Visca P, Imperi F, Lamont IL. Pyoverdine siderophores: from biogenesis to biosignificance. *Trends in Microbiology.* 2007;15(1):22-30. doi: 10.1016/j.tim.2006.11.004.
13. Lamont IL, Beare PA, Ochsner U, et al. Siderophore-mediated signaling regulates virulence factor production in *Pseudomonas aeruginosa*. *Proc Natl Acad Sci U S A.* 2002;99(10):7072-7077. doi: 10.1073/pnas.092016999. PubMed PMID: 11997446; eng.
14. Persat A, Inclan YF, Engel JN, et al. Type IV pili mechanochemically regulate virulence factors in *Pseudomonas aeruginosa*. *Proceedings of the National Academy of Sciences.* 2015;112(24):7563-7568. doi: 10.1073/pnas.1502025112.
15. Bomberger JM, MacEachran DP, Coutermarsh BA, et al. Long-Distance Delivery of Bacterial Virulence Factors by *Pseudomonas aeruginosa* Outer Membrane Vesicles. *PLoS Pathog.* 2009;5(4):e1000382. doi: 10.1371/journal.ppat.1000382.

16. Kung VL, Ozer EA, Hauser AR. The Accessory Genome of *Pseudomonas aeruginosa*. *Microbiology and Molecular Biology Reviews* : MMBR. 2010;74(4):621-641. doi: 10.1128/MMBR.00027-10. PubMed PMID: PMC3008168.
17. Morrison WD, Miller RV, Saylor GS. Frequency of F116-mediated transduction of *Pseudomonas aeruginosa* in a freshwater environment. *Appl Environ Microbiol*. 1978;36(5):724-730.
18. Monson R, Foulds I, Foweraker J, et al. The *Pseudomonas aeruginosa* generalized transducing phage ϕ PA3 is a new member of the ϕ KZ-like group of ‘jumbo’ phages, and infects model laboratory strains and clinical isolates from cystic fibrosis patients. *Microbiology*. 2011;157(3):859-867. doi: <https://doi.org/10.1099/mic.0.044701-0>.
19. Stanisich V, Holloway BW. Conjugation in *Pseudomonas aeruginosa*. *Genetics*. 1969;61(2):327-339. PubMed PMID: 4980017; eng.
20. van der Zee A, Kraak WB, Burggraaf A, et al. Spread of Carbapenem Resistance by Transposition and Conjugation Among *Pseudomonas aeruginosa* [Original Research]. *Front Microbiol*. 2018 2018-September-05;9(2057). doi: 10.3389/fmicb.2018.02057. English.
21. Tanner WD, Atkinson RM, Goel RK, et al. Horizontal transfer of the bla_{NDM-1} gene to *Pseudomonas aeruginosa* and *Acinetobacter baumannii* in biofilms. *FEMS Microbiol Lett*. 2017;364(8). doi: 10.1093/femsle/fnx048.
22. Qiu X, Kulasekara BR, Lory S. Role of Horizontal Gene Transfer in the Evolution of *Pseudomonas aeruginosa* Virulence. *Genome dynamics*. 2009;6:126-139. doi: 10.1159/000235767. PubMed PMID: 19696498; eng.
23. Kung VL, Ozer EA, Hauser AR. The accessory genome of *Pseudomonas aeruginosa*. *Microbiol Mol Biol Rev*. 2010 Dec;74(4):621-41. doi: 10.1128/MMBR.00027-10. PubMed PMID: 21119020; PubMed Central PMCID: PMC3008168.
24. Doberenz S, Eckweiler D, Reichert O, et al. Identification of a *Pseudomonas aeruginosa* PAO1 DNA Methyltransferase, Its Targets, and Physiological Roles. *mBio*. 2017;8(1):e02312-16. doi: 10.1128/mBio.02312-16.
25. Barrangou R, Fremaux C, Deveau H, et al. CRISPR provides acquired resistance against viruses in prokaryotes. *Science*. 2007 Mar 23;315(5819):1709-12. doi: 10.1126/science.1138140. PubMed PMID: 17379808; eng.
26. van Belkum A, Soriaga LB, LaFave MC, et al. Phylogenetic Distribution of CRISPR-Cas Systems in Antibiotic-Resistant *Pseudomonas aeruginosa*. *MBio*. 2015 Nov 24;6(6):e01796-15. doi: 10.1128/mBio.01796-15. PubMed PMID: 26604259; PubMed Central PMCID: PMC3008168.
27. Gophna U, Kristensen DM, Wolf YI, et al. No evidence of inhibition of horizontal gene transfer by CRISPR-Cas on evolutionary timescales. *ISME J*. 2015;9(9):2021-2027. doi: 10.1038/ismej.2015.20. PubMed PMID: 25710183; eng.
28. Du Toit A. CRISPR–Cas enhances HGT by transduction. *Nature Reviews Microbiology*. 2018 2018/04/01;16(4):186-186. doi: 10.1038/nrmicro.2018.28.
29. Freschi L, Vincent AT, Jeukens J, et al. The *Pseudomonas aeruginosa* Pan-Genome Provides New Insights on Its Population Structure, Horizontal Gene Transfer, and Pathogenicity. *Genome Biol Evol*. 2018;11(1):109-120. doi: 10.1093/gbe/evy259.
30. He J, Baldini RL, Déziel E, et al. The broad host range pathogen *Pseudomonas aeruginosa* strain PA14 carries two pathogenicity islands harboring plant and animal virulence genes. *Proc Natl Acad Sci U S A*. 2004;101(8):2530-2535. doi: 10.1073/pnas.0304622101.
31. Stover CK, Pham XQ, Erwin AL, et al. Complete genome sequence of *Pseudomonas aeruginosa* PAO1, an opportunistic pathogen. *Nature*. 2000 2000/08/01;406(6799):959-964. doi: 10.1038/35023079.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
1
15
16
17
18
19
20
2
22
23
24
25
26
27
28
2
30
31
32
33
34
35
36
3
38
39
40
41
42
43
4
45
46
47
48
49
50
51
5
53
54
55
56
57
58
59
60
32. Winsor GL, Griffiths EJ, Lo R, et al. Enhanced annotations and features for comparing thousands of *Pseudomonas* genomes in the *Pseudomonas* genome database. *Nucleic Acids Res.* 2016 Jan 4;44(D1):D646-53. doi: 10.1093/nar/gkv1227. PubMed PMID: 26578582; PubMed Central PMCID: PMC4702867.
 33. diCenzo GC, Finan TM. The Divided Bacterial Genome: Structure, Function, and Evolution. *Microbiol Mol Biol Rev.* 2017;81(3):e00019-17. doi: 10.1128/membr.00019-17.
 34. Poulsen BE, Yang R, Clatworthy AE, et al. Defining the core essential genome of *Pseudomonas aeruginosa*. *Proceedings of the National Academy of Sciences.* 2019;116(20):10072-10080. doi: 10.1073/pnas.1900570116.
 35. van Tonder AJ, Mistry S, Bray JE, et al. Defining the Estimated Core Genome of Bacterial Populations Using a Bayesian Decision Model. *PLoS Comput Biol.* 2014;10(8):e1003788. doi: 10.1371/journal.pcbi.1003788.
 36. He J, Baldini RL, Déziel E, et al. The broad host range pathogen *Pseudomonas aeruginosa* strain PA14 carries two pathogenicity islands harboring plant and animal virulence genes. *Proceedings of the National Academy of Sciences.* 2004;101(8):2530-2535.
 37. Freschi L, Vincent AT, Jeukens J, et al. The *Pseudomonas aeruginosa* Pan-Genome Provides New Insights on Its Population Structure, Horizontal Gene Transfer, and Pathogenicity. *Genome Biol Evol.* 2019 Jan 1;11(1):109-120. doi: 10.1093/gbe/evy259. PubMed PMID: 30496396; PubMed Central PMCID: PMC6328365.
 38. Wiedenbeck J, Cohan FM. Origins of bacterial diversity through horizontal genetic transfer and adaptation to new ecological niches. *FEMS Microbiol Rev.* 2011;35(5):957-976. doi: 10.1111/j.1574-6976.2011.00292.x.
 39. Beceiro A, Tomás M, Bou G. Antimicrobial resistance and virulence: a successful or deleterious association in the bacterial world? *Clin Microbiol Rev.* 2013;26(2):185-230. doi: 10.1128/CMR.00059-12. PubMed PMID: 23554414; eng.
 40. Anonymous. Antibiotic / Antimicrobial Resistance (AR / AMR) / Biggest Threats and Data.
 41. ECDC. Surveillance of antimicrobial resistance in Europe 2018. Antimicrobial resistance surveillance in Europe. Solna, Sweden: European Centre for Disease Prevention and Control,; 2019.
 42. Jeannot K, Bolard A, Plesiat P. Resistance to polymyxins in Gram-negative organisms. *Int J Antimicrob Agents.* 2017 May;49(5):526-535. doi: 10.1016/j.ijantimicag.2016.11.029. PubMed PMID: 28163137.
 43. Xu A, Zheng B, Xu YC, et al. National epidemiology of carbapenem-resistant and extensively drug-resistant Gram-negative bacteria isolated from blood samples in China in 2013. *Clin Microbiol Infect.* 2016 Mar;22 Suppl 1:S1-8. doi: 10.1016/j.cmi.2015.09.015. PubMed PMID: 26846351.
 44. Maiden MC, Bygraves JA, Feil E, et al. Multilocus sequence typing: a portable approach to the identification of clones within populations of pathogenic microorganisms. *Proc Natl Acad Sci U S A.* 1998 Mar 17;95(6):3140-5. PubMed PMID: 9501229; PubMed Central PMCID: PMC19708. eng.
 45. Enright MC, Day NP, Davies CE, et al. Multilocus sequence typing for characterization of methicillin-resistant and methicillin-susceptible clones of *Staphylococcus aureus*. *J Clin Microbiol.* 2000 Mar;38(3):1008-15. PubMed PMID: 10698988; PubMed Central PMCID: PMC86325.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
46. Enright MC, Spratt BG. A multilocus sequence typing scheme for *Streptococcus pneumoniae*: identification of clones associated with serious invasive disease. *Microbiology*. 1998 Nov;144 (Pt 11):3049-60. PubMed PMID: 9846740.
 47. Reid SD, Herbelin CJ, Bumbaugh AC, et al. Parallel evolution of virulence in pathogenic *Escherichia coli*. *Nature*. 2000 Jul 6;406(6791):64-7. doi: 10.1038/35017546. PubMed PMID: 10894541.
 48. Curran B, Jonas D, Grundmann H, et al. Development of a multilocus sequence typing scheme for the opportunistic pathogen *Pseudomonas aeruginosa*. *J Clin Microbiol*. 2004 Dec;42(12):5644-9. doi: 10.1128/JCM.42.12.5644-5649.2004. PubMed PMID: 15583294; PubMed Central PMCID: PMC535286.
 49. Feil EJ, Enright MC, Spratt BG. Estimating the relative contributions of mutation and recombination to clonal diversification: a comparison between *Neisseria meningitidis* and *Streptococcus pneumoniae*. *Res Microbiol*. 2000 Jul-Aug;151(6):465-9. PubMed PMID: 10961460.
 50. Dekker JP, Frank KM. Next-Generation Epidemiology: Using Real-Time Core Genome Multilocus Sequence Typing To Support Infection Control Policy. *J Clin Microbiol*. 2016 Dec;54(12):2850-2853. doi: 10.1128/JCM.01714-16. PubMed PMID: 27629902; PubMed Central PMCID: PMC5121370.
 51. Royer G, Fourreau F, Boulanger B, et al. Local outbreak of extended-spectrum beta-lactamase SHV2a-producing *Pseudomonas aeruginosa* reveals the emergence of a new specific sub-lineage of the international ST235 high-risk clone. *J Hosp Infect*. 2020 Jan;104(1):33-39. doi: 10.1016/j.jhin.2019.07.014. PubMed PMID: 31369808.
 52. Jolley KA, Bray JE, Maiden MCJ. Open-access bacterial population genomics: BIGSdb software, the PubMLST.org website and their applications. *Wellcome Open Res*. 2018;3:124. doi: 10.12688/wellcomeopenres.14826.1. PubMed PMID: 30345391; PubMed Central PMCID: PMC6192448.
 53. Kos VN, Deraspe M, McLaughlin RE, et al. The resistome of *Pseudomonas aeruginosa* in relationship to phenotypic susceptibility. *Antimicrob Agents Chemother*. 2015 Jan;59(1):427-36. doi: 10.1128/AAC.03954-14. PubMed PMID: 25367914; PubMed Central PMCID: PMC4291382.
 54. Oliver A, Mulet X, López-Causapé C, et al. The increasing threat of *Pseudomonas aeruginosa* high-risk clones. *Drug Resistance Updates*. 2015 2015/07/01;21-22(Supplement C):41-59. doi: <https://doi.org/10.1016/j.drug.2015.08.002>.
 55. Sánchez-Diener I, Zamorano L, López-Causapé C, et al. Interplay among Resistance Profiles, High-Risk Clones, and Virulence in the *Caenorhabditis elegans* *Pseudomonas aeruginosa* Infection Model. *Antimicrob Agents Chemother*. 2017 December 1, 2017;61(12). doi: 10.1128/aac.01586-17.
 56. Rasamiravaka T, Labtani Q, Duez P, et al. The Formation of Biofilms by *Pseudomonas aeruginosa*: A Review of the Natural and Synthetic Compounds Interfering with Control Mechanisms. *BioMed Research International*. 2015;2015:17. doi: 10.1155/2015/759348.
 57. Wendel AF, Ressina S, Kolbe-Busch S, et al. Species Diversity of Environmental GIM-1-Producing Bacteria Collected during a Long-Term Outbreak. *Appl Environ Microbiol*. 2016 June 15, 2016;82(12):3605-3610. doi: 10.1128/aem.00424-16.
 58. Mulet X, Cabot G, Ocampo-Sosa AA, et al. Biological Markers of *Pseudomonas aeruginosa* Epidemic High-Risk Clones. *Antimicrob Agents Chemother*. 2013 November 1, 2013;57(11):5527-5535. doi: 10.1128/aac.01481-13.
 59. Pastar I, Nusbaum AG, Gil J, et al. Interactions of methicillin resistant *Staphylococcus aureus* USA300 and *Pseudomonas aeruginosa* in polymicrobial

- wound infection. PLoS One. 2013;8(2):e56846-e56846. doi: 10.1371/journal.pone.0056846. PubMed PMID: 23451098; eng.
60. Das T, Kutty SK, Tavallaie R, et al. Phenazine virulence factor binding to extracellular DNA is important for *Pseudomonas aeruginosa* biofilm formation [Article]. Sci Rep. 2015 02/11/online;5:8398. doi: 10.1038/srep08398 <https://www.nature.com/articles/srep08398#supplementary-information>.
61. Ryan RP, Lucey J, O'Donovan K, et al. HD-GYP domain proteins regulate biofilm formation and virulence in *Pseudomonas aeruginosa*. Environmental Microbiology. 2009;11(5):1126-1136. doi: 10.1111/j.1462-2920.2008.01842.x.
62. Rashid MH, Rumbaugh K, Passador L, et al. Polyphosphate kinase is essential for biofilm development, quorum sensing, and virulence of *Pseudomonas aeruginosa*. Proceedings of the National Academy of Sciences. 2000;97(17):9636-9641. doi: 10.1073/pnas.170283397.
63. Pires DP, Vilas Boas D, Sillankorva S, et al. Phage Therapy: a Step Forward in the Treatment of *Pseudomonas aeruginosa* Infections. J Virol. 2015;89(15):7449-7456. doi: 10.1128/jvi.00385-15.
64. Krylov VN. Chapter Five - Bacteriophages of *Pseudomonas aeruginosa*: Long-Term Prospects for Use in Phage Therapy. In: Maramorosch K, Murphy FA, editors. Advances in Virus Research. Vol. 88: Academic Press; 2014. p. 227-278.
65. Lobato-Márquez D, Díaz-Orejas R, García-del Portillo F. Toxin-antitoxins and bacterial virulence. FEMS Microbiol Rev. 2016;40(5):592-609. doi: 10.1093/femsre/fuw022.
66. Hernández-Ramírez KC, Reyes-Gallegos RI, Chávez-Jacobo VM, et al. A plasmid-encoded mobile genetic element from *Pseudomonas aeruginosa* that confers heavy metal resistance and virulence. Plasmid. 2018 2018/06/01;98:15-21. doi: <https://doi.org/10.1016/j.plasmid.2018.07.003>.
67. Gaillard M, Pernet N, Vogne C, et al. Host and invader impact of transfer of the *clc* genomic island into *Pseudomonas aeruginosa* PAO1. Proceedings of the National Academy of Sciences. 2008;105(19):7058-7063. doi: 10.1073/pnas.0801269105.
68. Keen EC. Paradigms of pathogenesis: targeting the mobile genetic elements of disease. Frontiers in cellular and infection microbiology. 2012;2:161-161. doi: 10.3389/fcimb.2012.00161. PubMed PMID: 23248780; eng.
69. Ouyang J, Sun F, Feng W, et al. Quercetin is an effective inhibitor of quorum sensing, biofilm formation and virulence factors in *Pseudomonas aeruginosa*. Journal of Applied Microbiology. 2016;120(4):966-974. doi: 10.1111/jam.13073.
70. Lee J-H, Kim Y-G, Cho MH, et al. ZnO nanoparticles inhibit *Pseudomonas aeruginosa* biofilm formation and virulence factor production. Microbiol Res. 2014 2014/12/01;169(12):888-896. doi: <https://doi.org/10.1016/j.micres.2014.05.005>.
71. Harjai K, Kumar R, Singh S. Garlic blocks quorum sensing and attenuates the virulence of *Pseudomonas aeruginosa*. Pathogens and Disease. 2010;58(2):161-168. doi: 10.1111/j.1574-695X.2009.00614.x.
72. Kalia M, Yadav VK, Singh PK, et al. Effect of Cinnamon Oil on Quorum Sensing-Controlled Virulence Factors and Biofilm Formation in *Pseudomonas aeruginosa*. PLOS ONE. 2015;10(8):e0135495. doi: 10.1371/journal.pone.0135495.
73. Neidig A, Yeung ATY, Rosay T, et al. TypA is involved in virulence, antimicrobial resistance and biofilm formation in *Pseudomonas aeruginosa*. BMC Microbiology. 2013 2013/04/09;13(1):77. doi: 10.1186/1471-2180-13-77.

- 1
2
3 74. Kim S-K, Lee J-H. Biofilm dispersion in *Pseudomonas aeruginosa* [journal article].
4 Journal of Microbiology. 2016 February 01;54(2):71-85. doi: 10.1007/s12275-016-
5 5528-7.
6
7 75. Li Y, Heine S, Entian M, et al. NO-Induced Biofilm Dispersion in <span
8 class="named-content genus-species" id="named-content-1">*Pseudomonas*
9 *aeruginosa* Is Mediated by an MHYT Domain-Coupled Phosphodiesterase.
10 Journal of Bacteriology. 2013;195(16):3531-3542. doi: 10.1128/jb.01156-12.
11
12 76. Sauer K, Cullen MC, Rickard AH, et al. Characterization of Nutrient-Induced
13 Dispersion in *Pseudomonas aeruginosa* PAO1 Biofilm. Journal of
14 Bacteriology. 2004;186(21):7312-7326. doi: 10.1128/jb.186.21.7312-7326.2004.
15
16 77. Johansson EMV, Crusz SA, Kolomiets E, et al. Inhibition and Dispersion of
17 *Pseudomonas aeruginosa* Biofilms by Glycopeptide Dendrimers Targeting the
18 Fucose-Specific Lectin LecB. Chemistry & Biology. 2008 2008/12/22;15(12):1249-
19 1257. doi: <https://doi.org/10.1016/j.chembiol.2008.10.009>.
20
21 78. Robb AJ, Vinogradov S, Danell AS, et al. Electrochemical detection of small
22 molecule induced *Pseudomonas aeruginosa* biofilm dispersion. Electrochimica Acta.
23 2018 2018/04/01;268:276-282. doi: <https://doi.org/10.1016/j.electacta.2018.02.113>.
24
25 79. Dunne WM, Buckmire FL. Partial purification and characterization of a
26 polymannuronic acid depolymerase produced by a mucoid strain of *Pseudomonas*
27 *aeruginosa* isolated from a patient with cystic fibrosis. Appl Environ Microbiol.
28 1985;50(3):562-567.
29
30 80. Bartell PF, Orr TE, Lam GKH. Polysaccharide Depolymerase Associated with
31 Bacteriophage Infection. J Bacteriol. 1966;92(1):56-62.
32
33 81. Bartell PF, Lam GKH, Orr TE. Purification and Properties of Polysaccharide
34 Depolymerase Associated with Phage-infected *Pseudomonas aeruginosa*. J Biol
35 Chem. 1968 May 10, 1968;243(9):2077-2080.
36
37 82. Glonti T, Chanishvili N, Taylor PW. Bacteriophage-derived enzyme that
38 depolymerizes the alginate capsule associated with cystic fibrosis isolates of
39 *Pseudomonas aeruginosa*. J Appl Microbiol. 2010;108(2):695-702. doi:
40 10.1111/j.1365-2672.2009.04469.x.
41
42 83. Pires DP, Oliveira H, Melo LDR, et al. Bacteriophage-encoded depolymerases: their
43 diversity and biotechnological applications [journal article]. Applied Microbiology
44 and Biotechnology. 2016 March 01;100(5):2141-2151. doi: 10.1007/s00253-015-
45 7247-0.
46
47 84. De Kievit TR. Quorum sensing in *Pseudomonas aeruginosa* biofilms. Environ
48 Microbiol. 2009;11(2):279-288. doi: 10.1111/j.1462-2920.2008.01792.x.
49
50 85. Lee J, Zhang L. The hierarchy quorum sensing network in *Pseudomonas aeruginosa*.
51 Protein & cell. 2015;6(1):26-41.
52
53 86. Spoering AL, Gilmore MS. Quorum sensing and DNA release in bacterial biofilms.
54 Current Opinion in Microbiology. 2006 2006/04/01;9(2):133-137. doi:
55 <https://doi.org/10.1016/j.mib.2006.02.004>.
56
57 87. Antonova ES, Hammer BK. Quorum-sensing autoinducer molecules produced by
58 members of a multispecies biofilm promote horizontal gene transfer to *Vibrio*
59 *cholerae*. FEMS Microbiol Lett. 2011;322(1):68-76. doi: 10.1111/j.1574-
60 6968.2011.02328.x.
61
62 88. Rossmann FS, Racek T, Wobser D, et al. Phage-mediated Dispersal of Biofilm and
63 Distribution of Bacterial Virulence Genes Is Induced by Quorum Sensing. PLoS
64 Pathog. 2015;11(2):e1004653. doi: 10.1371/journal.ppat.1004653.


- 1
2
3
4
5
6
7
8
9
10
11
12
13
1
15
16
17
18
19
20
2
22
23
24
25
26
27
28
2
30
31
32
33
34
35
36
3
38
39
40
41
42
43
4
45
46
47
48
49
50
51
5
53
54
55
56
57
58
5
60
89. Zhang Y, Ma Q, Su B, et al. A study on the role that quorum sensing play in antibiotic-resistant plasmid conjugative transfer in *Escherichia coli*. *Ecotoxicology*. 2018;27(2):209-216.
90. Hentzer M, Wu H, Andersen JB, et al. Attenuation of *Pseudomonas aeruginosa* virulence by quorum sensing inhibitors. *The EMBO Journal*. 2003;22(15):3803-3815. doi: 10.1093/emboj/cdg366.
91. O'Loughlin CT, Miller LC, Siryaporn A, et al. A quorum-sensing inhibitor blocks *Pseudomonas aeruginosa* virulence and biofilm formation. *Proceedings of the National Academy of Sciences*. 2013;110(44):17981-17986.
92. Kim H-S, Lee S-H, Byun Y, et al. 6-Gingerol reduces *Pseudomonas aeruginosa* biofilm formation and virulence via quorum sensing inhibition. *Sci Rep*. 2015;5:8656-8656. doi: 10.1038/srep08656. PubMed PMID: 25728862; eng.
93. Høyland-Kroghsbo NM, Paczkowski J, Mukherjee S, et al. Quorum sensing controls the *Pseudomonas aeruginosa* CRISPR-Cas adaptive immune system. *Proceedings of the National Academy of Sciences*. 2017;114(1):131-135. doi: 10.1073/pnas.1617415113.
94. Whiteley M, Diggle SP, Greenberg EP. Progress in and promise of bacterial quorum sensing research. *Nature*. 2017;551(7680):313-320. doi: 10.1038/nature24624. PubMed PMID: 29144467; eng.
95. Koser CU, Holden MT, Ellington MJ, et al. Rapid whole-genome sequencing for investigation of a neonatal MRSA outbreak. *N Engl J Med*. 2012 Jun 14;366(24):2267-75. doi: 10.1056/NEJMoa1109910. PubMed PMID: 22693998; PubMed Central PMCID: PMC3715836.
96. Jaillard M, van Belkum A, Cady KC, et al. Correlation between phenotypic antibiotic susceptibility and the resistome in *Pseudomonas aeruginosa*. *Int J Antimicrob Agents*. 2017 Aug;50(2):210-218. doi: 10.1016/j.ijantimicag.2017.02.026. PubMed PMID: 28554735.
97. Witney AA, Gould KA, Pope CF, et al. Genome sequencing and characterization of an extensively drug-resistant sequence type 111 serotype O12 hospital outbreak strain of *Pseudomonas aeruginosa*. *Clin Microbiol Infect*. 2014 Oct;20(10):O609-18. doi: 10.1111/1469-0691.12528. PubMed PMID: 24422878.
98. Rizzo L, Manaia C, Merlin C, et al. Urban wastewater treatment plants as hotspots for antibiotic resistant bacteria and genes spread into the environment: a review. *Science of the total environment*. 2013;447:345-360.
99. Marti E, Variatza E, Balcazar JL. The role of aquatic ecosystems as reservoirs of antibiotic resistance. *Trends Microbiol*. 2014 2014/01/01;22(1):36-41. doi: <https://doi.org/10.1016/j.tim.2013.11.001>.
100. Aminov RI. Horizontal gene exchange in environmental microbiota. *Front Microbiol*. 2011;2:158-158. doi: 10.3389/fmicb.2011.00158. PubMed PMID: 21845185; eng.
101. Bruchmann J, Kirchen S, Schwartz T. Sub-inhibitory concentrations of antibiotics and wastewater influencing biofilm formation and gene expression of multi-resistant *Pseudomonas aeruginosa* wastewater isolates [journal article]. *Environmental Science and Pollution Research*. 2013 June 01;20(6):3539-3549. doi: 10.1007/s11356-013-1521-4.
102. Karkman A, Do TT, Walsh F, et al. Antibiotic-Resistance Genes in Waste Water. *Trends in Microbiology*. 2018 2018/03/01;26(3):220-228. doi: <https://doi.org/10.1016/j.tim.2017.09.005>.
103. Gunathilaka GU, Tahlan V, Mafiz AI, et al. Phages in urban wastewater have the potential to disseminate antibiotic resistance. *International Journal of Antimicrobial*


- Agents. 2017 2017/11/01/;50(5):678-683. doi: <https://doi.org/10.1016/j.ijantimicag.2017.08.013>.
104. Regmi C, Joshi B, Ray SK, et al. Understanding Mechanism of Photocatalytic Microbial Decontamination of Environmental Wastewater [Mini Review]. *Frontiers in Chemistry*. 2018 2018-February-28;6(33). doi: 10.3389/fchem.2018.00033. English.
105. Fujishima A, Zhang X. Titanium dioxide photocatalysis: present situation and future approaches. *Comptes Rendus Chimie*. 2006 2006/05/01/;9(5):750-760. doi: <https://doi.org/10.1016/j.crci.2005.02.055>.
106. Chong MN, Jin B, Chow CWK, et al. Recent developments in photocatalytic water treatment technology: A review. *Water Res*. 2010 2010/05/01/;44(10):2997-3027. doi: <https://doi.org/10.1016/j.watres.2010.02.039>.
107. Laxma Reddy PV, Kavitha B, Kumar Reddy PA, et al. TiO₂-based photocatalytic disinfection of microbes in aqueous media: A review. *Environmental Research*. 2017 2017/04/01/;154:296-303. doi: <https://doi.org/10.1016/j.envres.2017.01.018>.
108. Daneshvar N, Niaei A, Akbari S, et al. Photocatalytic disinfection of water polluted by *Pseudomonas aeruginosa*. *Global Nest J*. 2007;9(3):1-5.
109. Amezaga-Madrid P, Silveyra-Morales R, Cordoba-Fierro L, et al. TEM evidence of ultrastructural alteration on *Pseudomonas aeruginosa* by photocatalytic TiO₂ thin films. *J Photochem Photobiol B: Biol*. 2003;70(1):45-50.
110. Ratova M, Redfern J, Verran J, et al. Highly efficient photocatalytic bismuth oxide coatings and their antimicrobial properties under visible light irradiation. *Applied Catalysis B: Environmental*. 2018 2018/12/30/;239:223-232. doi: <https://doi.org/10.1016/j.apcatb.2018.08.020>.
111. Markowska-Szczupak A, Ulfig K, Morawski AW. The application of titanium dioxide for deactivation of bioparticulates: An overview. *Catal Today*. 2011 2011/07/01/;169(1):249-257. doi: <https://doi.org/10.1016/j.cattod.2010.11.055>.
112. Lawton LA, Robertson PKJ, Robertson RF, et al. The destruction of 2-methylisoborneol and geosmin using titanium dioxide photocatalysis. *Applied Catalysis B: Environmental*. 2003 2003/08/08/;44(1):9-13. doi: [https://doi.org/10.1016/S0926-3373\(03\)00005-5](https://doi.org/10.1016/S0926-3373(03)00005-5).
113. Murar M, Dhumale V. Antimicrobial potential of TiO₂ nanoparticles against MDR *Pseudomonas aeruginosa* AU - Arora, Bindu. *J Exp Nanosci*. 2015 2015/07/24/;10(11):819-827. doi: 10.1080/17458080.2014.902544.
114. Ryu H, Crittenden J, Abbaszadegan M. Photocatalytic inactivation of viruses using titanium dioxide nanoparticles and low-pressure UV light AU - Gerrity, Daniel. *Journal of Environmental Science and Health, Part A*. 2008 2008/07/29/;43(11):1261-1270. doi: 10.1080/10934520802177813.

1
2
3 Figure 1. Genomic diversity of *P. aeruginosa* clinical isolates. (A) Phylogenetic tree of *P.*
4 *aeruginosa* clinical isolates. Phylogenetically related ST groups with >7 members are shown
5 in different colors. The prevalent lineages constitute 52% of all strains analyzed. ST groups
6 that have only 1 member are delineated in grey, while black denotes phylogenetic lineages
7 with 2 to 6 members. The phylogenetic locations of the common laboratory strains PAO1,
8 PA14, and PA7 are marked in black for reference, while the 4 nonclinical isolates are
9 labelled with small black asterisks. Figure reprinted from Van Belkum et al. mBio. 2015 [36].
10
11

12
13 Figure 2. Summary of key virulence determinants and factors important for biofilm
14 formation produced by *Pseudomonas aeruginosa*. Reproduced from Diggle and Whiteley,
15 2019 [110].
16

17
18 Figure 3. The hierarchy quorum sensing network in *Pseudomonas aeruginosa*. Reproduced
19 from Lee and Zhang, Protein and Cell, 2014 [80].
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

A






Figure 3. Schematic representation of the four QS signaling networks in *P. aeruginosa* and their respective regulons. Arrows indicate a stimulatory effect. Perpendicular lines indicate an inhibitory effect.