

1
2
3
4 **Title:**
5

6 A patient-reported outcome measure of functional vision for children and young
7 people aged 8 to 18 years with visual impairment.
8

9 **Short title:**

10 Measuring functional vision of children/young people
11

12 **Authors:**

13
14 Alexandra O Robertson^{1,3}

15 email: alexandra.robertson.13@ucl.ac.uk

16 Valerija Tadić^{1,2,3}

17 email: v.tadic@greenwich.ac.uk

18 Mario Cortina-Borja¹

19 email: m.cortina@ucl.ac.uk

20 Jugnoo S Rahi^{1,3,4,5}

21 email: j.rahi@ucl.ac.uk

22 *For the Child Vision PROMs group**

23 *Members of the Child Vision PROM group are listed in the Acknowledgements
24
25

26 **Affiliations:**

27
28
29
30 ¹ Population, Policy and Practice Research & Teaching Department, UCL Great
31 Ormond Street Institute of Child Health, UK

32 ² School of Human Sciences, University of Greenwich

33 ³ Great Ormond Street Hospital NHS Foundation Trust, UK

34 ⁴ National Institute for Health Research (NIHR) Biomedical Research Centre at
35 Moorfields Eye Hospital NHS Foundation Trust and UCL Institute of Ophthalmology,
36 London, UK

37 ⁵ Ulverscroft Vision Research Group, UK
38
39

40 **Corresponding author (and address for reprints):**

41
42 Jugnoo S Rahi, UCL Great Ormond Street Institute of Child Health, Life Course
43 Epidemiology and Biostatistics Section, Population, Policy and Practice Research &
44 Teaching Department, 30 Guilford Street, London, WC1N 1EH, UK; Telephone: 44
45 (0)20 7905 2250; Email: j.rahi@ucl.ac.uk
46
47

48 **Supplemental Material:**

49 Supplementary Table 1 available at AJO.com
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 ABSTRACT
5

6 **Purpose:** To develop age-appropriate extensions of a patient-reported outcome
7 measure for capturing the functional impact of visual impairment on daily activities of
8 children and young people aged 8 up to 18 years.
9

10 **Design:** Questionnaire development and validation study.
11

12 **Setting:** Pediatric Ophthalmology departments at Great Ormond Street Hospital and
13 Moorfields Eye Hospital, and, in the final study phase, 20 further UK hospitals.
14

15 **Participants:** Children and young people (aged 6-19 years) with visual impairment
16 (acuity of the logarithm of the minimum angle of resolution (LogMAR) worse than
17 0.50 in the better eye) due to any cause but without significant non-ophthalmic
18 impairments.
19
20

21 **Methods:** We used our prototype FVQ_CYP for 10-15 year olds as the foundation.
22 Twenty-nine semi-structured interviews confirmed relevance of existing, and
23 identified new, age-specific items. Twenty-eight cognitive interviews captured
24 information regarding comprehensibility and format. The FVQ_Child (8-12 years) and
25 FVQ_Young Person (13-18 years), were evaluated with a national sample of 113
26 children and 96 young people using Rasch analysis.
27
28

29 **Results:** Issues emerging from interviews with children and young people were
30 largely congruent with those elicited originally with 10-15 year olds. The 28-item
31 FVQ_Child and 38-item FVQ_Young Person versions have goodness-of-fit statistics
32 within the interval 0.5, 1.5 and person separation values of 5.87 and 6.09
33 respectively. Twenty-four overlapping 'core' items enabled their calibration on the
34 same measurement scale. Correlations with acuity ($r = 0.47$) demonstrated construct
35 validity.
36
37

38 **Conclusions:** The FVQ_C and FVQ_Young Person are robust age-appropriate
39 versions of the FVQ_CYP which can be used cross-sectionally or
40 sequentially/longitudinally across the age-range of 8-18 years in clinical practice and
41 research.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

INTRODUCTION

Visual impairment (VI) affects a child's ability to perform everyday tasks and activities, with cumulative effects on their educational, social and occupational prospects, and engagement in daily life.^{1 2} In keeping with the international drive to use patient-reported outcome measures (PROMs)³ to assess the impact of eye conditions and any treatment undertaken, the ability to accurately assess the affected child's perspective of their functional vision (FV) i.e. vision for everyday tasks, would complement clinical (objective) measures.

However, until recently, age-appropriate measures of FV for children and young people with VI have been lacking. Recently, instruments comprising a single measure applicable to the whole age-range of 8-18 years^{4 5} have been reported but it is unclear whether their content is developmentally appropriate, given the significant differences in activities that are meaningful and relevant to children versus young people, for example an 8 year old versus an 18 year old, as well as the evolution of their abilities to self-assess and self-report.

In response to both the importance and the lack of age-appropriate, child-centred, psychometrically robust PROMs for use in Pediatric Ophthalmology⁶ we developed and used a child-centred approach to generate our 'foundation' PROM for capturing FV of children and young people with VI aged 10-15 years (the FVQ_CYP).⁷

We now report the development of age-specific extensions of this instrument to allow for use with a broader age-range of children and young people with VI. This work forms part of our broader program of development of pediatric PROMs, in which we have developed age-appropriate versions of a PROM assessing the complementary but distinct construct of vision-related quality of life (the VQoL_CYP).⁸⁻¹⁰

METHODS

This instrument development study was approved by the National Health Service (NHS) Research Ethics Committee for East of England, United Kingdom (UK) and followed tenets of the Declaration of Helsinki. Participants >16 years consented and those aged <16 years assented alongside their parents' consent.

Sample

Participants were recruited from two patient populations between September 2014 and May 2017, comprising those attending the Department of Ophthalmology at Great Ormond Street Hospital, and the Pediatric Glaucoma Service and Genetic Eye Disease Service at Moorfields Eye Hospital, London UK supplemented (in the final phase only) by patients attending 20 other hospitals across the UK (see Acknowledgements). Children and young people aged 6-19 years (with final age boundaries for the instrument versions determined empirically later) were eligible if they were visually impaired, severely visually impaired or blind (corrected acuity in the better eye of LogMAR 0.50 or worse or Snellen worse than 6/18 or additional visual defects causing VI) due to any disorder, but without any other significant non-ophthalmic impairment. By sampling across multiple sources nationally in the final phase, where the largest sample was needed, we ensured our sample was as

1
2
3
4 representative as possible of the UK population of children and young people with VI
5 with respect to ethnicity, socio-economic status, and disorder.
6

7 Procedures

8
9 Instrument development was undertaken in three standard phases using our
10 foundation FVQ_CYP for 10-15 year olds⁷ and its underpinning archived interview
11 data as the springboard for adaptation.
12

13 Phase 1: Item development and adaptation

14
15 Individual in-depth, interviews were conducted with children younger than 10 and
16 young people older than 15 years to investigate the relevance of issues covered by
17 the FVQ_CYP items (from the 10-15 year olds' instrument⁷) to those outside the age-
18 range of 10-15 years, and to identify any new age-specific issues. We used our
19 existing data from the development of the original FVQ_CYP, involving 32 interviews
20 with 10-15 year olds,⁷ as the foundation for data collection, and reached data
21 saturation after 12 interviews with children and 17 interviews with young people.
22 Interviews were transcribed and coded using NVivo10.¹¹ Qualitative analysis
23 revealed areas of overlap, discrepancy or omissions in the new data, compared to
24 the issues covered by the existing FVQ_CYP instrument. New, age-appropriate
25 items were developed to address any new issues not addressed in the foundation
26 FVQ_CYP. To ensure existing FVQ_CYP items were developmentally appropriate
27 for children younger than those for whom it was originally designed, participants <10
28 years completed the FVQ_CYP (10-15 years) with parental assistance. Feedback
29 informed the early draft of the FVQ_CYP version for younger children. This was not
30 considered necessary for participants older than 15 years, who were
31 developmentally well placed to comprehend the existing FVQ_CYP (10-15 years)
32 items.
33
34
35
36
37

38 Phase 2: Pre-testing

39
40 The upper and lower age boundary for each new age-appropriate FVQ instrument
41 version was developed empirically throughout Phase 2. To ensure the new draft
42 instrument versions would be comprehensible and age-appropriate to a broader age-
43 range, recruitment in this phase was focused on participants younger than 10 years
44 and older than 15 years. One-to-one cognitive interviews with 12 children aged 7-10
45 years and 16 young people aged 13-18 years were conducted. Items were evaluated
46 for importance, comprehensibility, difficulty and response format. The original
47 interviews with 10-15 year olds were re-read,⁷ and feedback from children and young
48 people, their parents, and study group consensus was used to determine the age
49 thresholds for the new instrument versions as 8-12 years and 13-18 years.
50
51
52

53 Phase 3: Piloting and validation

54
55 The age-appropriate instrument versions were piloted with a national sample (UK) of
56 113 children aged 8-12 years and 96 young people aged 13-18 years to confirm their
57 psychometric properties.
58
59
60
61
62
63
64
65

1
2
3
4 Participants received invitation letters, accompanied by consent/assent forms, child
5 and parent information sheets, and the age-appropriate instrument versions in large
6 print (including a link to an electronic version) and a postage-paid envelop for return
7 of the completed documents.
8

9
10 Data from the returned instrument versions were entered into IBM SPSS version
11 24,¹² and verified through double-checking, with no errors detected. Data from
12 participants with >25% of item responses, and items with >60% of participant
13 responses missing were excluded.¹³
14

15 Rasch analysis¹⁴ and the Andrich Rasch Rating Scale model defined the item
16 reduction. Criteria used to assess the appropriateness of the two instrument
17 versions^{13 15} are detailed in Table 2 and Figures 1 and 2. Prior to analysis, 1 – 4
18 responses were coded into a scale of 0 – 3.
19
20

21 *Calibrating the FVQ_Child and FVQ_Young Person versions*

22

23 We used the model resulting from equating both age-appropriate instrument versions
24 (as outlined by Linacre¹⁶) to ensure that they measure the same construct in children
25 and young people. This model utilizes the ‘core’ items common to both instrument
26 versions and provides continuity of measurement across the age-range of 8-18
27 years. Thus the instrument versions can be used in cross-sectional studies and also
28 at different time points with the same participants, to allow for longitudinal analysis.
29 In this transformation, all items are assumed to have equal importance, and
30 response categories are scaled accordingly to provide an equal value with uniform
31 increments between consecutive categories. A final differential item functioning (DIF)
32 analysis was conducted using these ‘core’ items common to both instrument
33 versions, to investigate whether the equated Rasch person measures from the two
34 age groups (8-12 and 13-18 years) were comparable.¹⁷
35
36
37

38 We assessed unidimensionality using infit and outfit statistics, following the criteria
39 described in Table 2.¹³ DIF statistics (Table 2) represent the effect size of the
40 difference between the two classifications of persons, in logits.¹⁸
41

42 FVQ total summary scores were calculated by adding item scores across the scale
43 and converted into Rasch person measures ranging from 0 (denoting lower difficulty
44 and excellent FVQ) to 100 (denoting greater difficulty and severely reduced FVQ).
45 This was done using the score-to-measure conversion tables for each version (Table
46 3). These conversion tables allow the derived measures to be compared between
47 the two age-appropriate versions regardless of the differences in the number and
48 wording of items.
49
50

51 For those participants with any missing items, Rasch person measures were imputed
52 applying a procedure which is consistent with item response theory.^{19 20} This
53 approach uses adjusted score-to-measure conversion tables derived from Table 3.
54
55

56 *Construct validity*

57

58 Construct validity, assessing the instrument’s ability to truly measure the underlying
59 latent construct, was assessed through Spearman’s rank correlation coefficients
60
61
62
63
64
65

1
2
3
4 between Rasch person measures on the FVQ_Child and FVQ_Young Person and
5 objectively measured visual acuity.
6

7 Rasch analysis was conducted using Winsteps 4.0.1.¹¹ and all other analyses using
8 SPSS.
9

10 **RESULTS**

11 Participants represented the overall “target” UK population of children and young
12 people with VI able to self-report (i.e. without additional significant impairment) in
13 terms of clinical and socio-demographic characteristics and ophthalmic diagnoses
14 (Table 1).^{7 9 21}
15
16

17 Phase 1: Item development and adaptation

18 The issues raised by children younger than 10 years and those older than 15 years
19 overlapped significantly with those addressed by the original FVQ_CYP instrument
20 for 10-15 year olds.⁷ Nevertheless, domain-pertinent issues in the original instrument
21 were not relevant to younger children and older participants reported engagement in
22 additional activities (e.g. attending parties, and using mobile phones) different to
23 those covered by the original FVQ_CYP.
24
25

26 The original FVQ_CYP instrument for 10-15 year olds has 36 items addressing
27 activities at home, school and leisure, restrictions and limitations, levels of
28 functioning, mobility, and communication. Of these, 28 were retained for the new
29 extension for children <10 years i.e. the FVQ_Child, and one new item capturing
30 outdoor/playground games was added. Thirty-one of the original 36 items were
31 retained following minor linguistic adaptations (e.g. references to ‘school’ were
32 changed to ‘school/college’) for the extension for those aged >15 years i.e. the
33 FVQ_Young Person. We added 7 items that drew on our foundation research and 2
34 entirely new items related to maintaining physical appearance and using a mobile
35 phone for social networking.
36
37

38 Item presentation was modified to calibrate the instrument versions by retaining a
39 consistent format and structure across them. All items were presented as a question
40 stem (*‘Because of my eyesight, I find...’*) followed by an activity (e.g. *‘Watching TV’*),
41 with four response options: *‘1: Very easy’*; *‘2: Easy’*; *‘3: A bit difficult’* (*‘Difficult’* in the
42 FVQ_Young Person); and *‘4: Very difficult or impossible’*.⁷ The prompt: *‘Remember
43 to tell us how things are for you when wearing your glasses (if you wear them), with
44 your low vision aids and other devices (if you use them for these activities) and with
45 the best lighting and contrast for you’* was inserted between items.
46
47

48 Phase 2: Pre-testing of the 29-item FVQ_Child and 40-item FVQ_Young Person

49 One item *‘Getting around outdoors by myself’* was divided into two items in both
50 instrument versions to specify context (*‘in daylight’* and *‘when it’s dark’*). Age-
51 boundaries for the extensions were re-adjusted as 8-12 years and 13-18 years
52 empirically, reflecting the minimum age for accurate self-reporting.²²
53
54

55 Phase 3: Piloting and validation

1
2
3
4 Four children and two young people were excluded from Phase 3 because they had
5 >25% missing data. These participants had visual acuity ranging from 0.48 to
6 perception of light only. In the remaining children and young people, missing data
7 per child (aged 8-12 years) was $\leq 7\%$, and $\leq 22\%$ among young people (aged 13-18
8 years). A Poisson regression model revealed a non-significant relationship between
9 the number of missing items and severity of visual impairment ($p = 0.351$).

10
11 Missing data per item was $\leq 20\%$ in the child dataset and $\leq 17\%$ for young people.

12
13 Following Rasch analysis, one item was removed from the FVQ_Child and 3 items
14 were removed from the FVQ_Young Person based on outfit MNSQ statistics and
15 notable DIF (see online Supplementary Table 1).

16 17 Calibrating the FVQ_Child and FVQ_Young Person instrument versions

18
19 Analysis of DIF between children and young people on the combined datasets for
20 the overlapping 'core' items revealed that the item '*Reading small writing such as*
21 *food packets, tickets, and labels*' was more difficult for children than young people.
22 Results from the preliminary item reduction stage were re-visited and this item was
23 removed from the FVQ_Child only, based on the finding that 57% of children (vs.
24 35.5% of young people) rated this item as '*Very difficult or impossible*', confirming an
25 age-related bias. All remaining overlapping 'core' items were productive for
26 measurement of FV in both instrument versions.
27
28
29

30
31 The final 28-item FVQ_Child and 38-item FVQ_Young Person contain 24
32 overlapping 'core' items and 4 and 14 age-specific items, respectfully (Table 2). Both
33 instrument versions showed fit statistics and DIF values within acceptable limits. Item
34 probability plots showed good ordering and acceptable distinction between 4
35 response categories (Figure 1), and targeting of items to respondents (Figure 2). The
36 FVQ_Child and FVQ_Young Person showed precision as indicated by the indices for
37 person separation (5.87 and 6.09, respectively).
38
39

40 Score-to-measure transformation

41
42 Rasch person measures from the FVQ_Child and FVQ_Young Person may be
43 compared on a linear scale ranging from 0 to 100. Table 3 shows the transformation
44 of scores into person measures which enable easy and precise scoring, and direct
45 comparison of scores from individuals of different ages, and scores over time.
46

47 Construct validity

48
49 In keeping with published criteria,¹³ Rasch person measures on the FVQ_Child and
50 FVQ_Young Person correlated positively with participants' latest recorded visual
51 acuity ($r = 0.48$, $p = <.001$ for FVQ_Child, $r = 0.43$, $p = <.001$ for FVQ_Young
52 Person, and $r = 0.46$, $p <.001$ for the combined FVQ_Child and FVQ_Young Person
53 datasets), indicating, as hypothesized, that lower FV is reported by children with
54 poorer acuity in both age-groups.
55
56

57 **DISCUSSION**

58
59
60
61
62
63
64
65

1
2
3
4 We report development of age/stage appropriate versions of a robust PROM
5 assessing the functional impact of VI on children and young people. The novel
6 equating approach we used to calibrate the two instrument versions means that
7 Rasch person measures from either version can be compared using one linear scale
8 representing FV, despite age-specific variation. This affords many advantages when
9 used in practice, namely that the instrument can be used cross-sectionally and
10 sequentially, with children and young people aged from 8 years up to 18 years, and
11 without loss of continuity of measurement as subjects get older by using an
12 alternative instrument. We provide log transformation tables, which can be used to
13 convert summary scores into Rasch person measures, which are also accompanied
14 by the model-based standard error of each measure, which should be used in future
15 clinical research.
16
17
18

19 Our research adhered to best practice via independent self-report from children and
20 young people themselves, through one-to-one individual interviews, expert
21 consultations, and provision of age-appropriate materials. This rigor is reflected in
22 the content, format and evidence of construct (convergent) validity of both instrument
23 versions. By deliberately isolating activities on which VI can impact, we have avoided
24 any conflation between FV and the psychosocial emotional impact of VI which is
25 captured instead in our corresponding vision-related quality of life instrument.⁸⁻¹⁰
26
27
28

29 The relatively small sample size of our study (reflecting the rarity of childhood VI) has
30 implications for Rasch analysis, particularly the stability of DIF analyses and item fit
31 statistics. We addressed this in the analysis of DIF by age, by grouping participants
32 by individual year groups to optimize use of the sample. We carefully considered the
33 trade-off between retaining meaningful items which are productive for measurement
34 and thus preserving content and face validity, versus removing those which did not fit
35 the 'perfect' measurement scale. The broader criteria we used for assessing item fit
36 reflects this.
37
38

39 Although FV is not formally defined in the extant literature, it bridges the gap
40 between health conditions and associated symptoms (i.e. reduced visual function)
41 and contextual factors (i.e. environmental and personal factors inherent to daily
42 activities) specified by the International Classification of Functioning, Disability and
43 Health.²³ We framed FV as the ability to complete meaningful daily activities in real
44 everyday environments. Consequently, our instrument captures activities performed
45 at home and in school, with age-appropriate items reflecting increasing
46 independence and responsibility with age.
47
48

49 Our instrument differs from some other current vision-specific PROMs which capture
50 some aspects of FV of children and young people, by being applicable to all/any
51 cause of VI versus a single eye condition^{24 25} and to an English speaking
52 population.²⁶ The most direct comparators are the Cardiff Visual Ability
53 Questionnaire for Children (CVAQC)⁴ and the LV Prasad-Functional Questionnaire
54 Second Version (LVP-FVQ II)⁵ but neither has age-appropriate versions capable of
55 capturing change in the nature of tasks of daily living over time. The recently
56 reported PedEyeQ²⁷ addresses age-specificity through separate instruments for
57
58
59
60
61
62
63
64
65

1
2
3
4 different age-groups but lacks the calibration required to allow for valid comparisons
5 of these measures.
6

7 The benefits of using PROMs within clinical practice include improvements in
8 patient-clinician communication such as advice and diagnoses given by health
9 professionals,²⁸ and increased 'patient centricity' within clinical care.²⁹ PROMs are
10 also valuable at a higher institutional level, with potential to trigger changes in clinical
11 practice and monitor the quality of healthcare provided.³⁰ The instrument versions we
12 have developed enhance these uses by also affording the opportunity to compare
13 scores meaningfully from individuals across the age-range of 8 up to 18 years whilst
14 maintaining specificity to differences between the two age-groups. This makes them
15 useful in assessments of key, age-related or vision-specific milestones or
16 interventions without the need for clinicians to use and interpret multiple instruments;
17 the latter a well-documented barrier to routine use of PROMs.^{31 32}

18 The age-boundaries for our instrument versions are empirically-based and echo
19 most child-centred, vision-specific PROMs.^{5 28 33} However, given the specific
20 developmental profile of the population of children and young people with VI, we
21 advocate tailoring the choice of version³⁴ to the patient's developmental needs rather
22 than just her/his age.
23

24 To ensure ability to self-report and focus on the impact of VI per se, we restricted our
25 participant population to those without additional impairments. Further work is
26 necessary to address the challenge of developing our FV measure to make it
27 appropriate for children/young people in whom VI may be one of a number of co-
28 existing impairments. Whilst parent or proxy reporting is not considered best practice
29 due to the potential for discordance between proxies and those affected i.e. risk of
30 misinterpreting the child's views,³⁵ this may nevertheless be required when complex
31 health conditions preclude self-reporting by children/young people. This may be the
32 way forward for our FVQ_CYP instrument as parents rate physical symptoms more
33 accurately than subjective well-being or quality of life.³⁶
34

35 Our child-centred and resource efficient approach has enabled development of a
36 robust age- and stage-appropriate PROM allowing children and young people to self-
37 assess and report on the functional impact of their VI. This instrument can be used
38 cross-sectionally (e.g. in population burden of disease studies) or sequentially (e.g.
39 moving from the FVQ_Child to the FVQ_Young Person over time in clinical trials) in
40 clinical practice and research to provide a deeper understanding and alternative
41 quantification of the impact of eye disease and its treatment than objective clinical
42 measures alone can afford.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 ACKNOWLEDGEMENTS/DISCLOSURES
5

6 Funding/Support:

7
8 This study was funded by a Fight for Sight Project Grant (1321/1322) and a UCL
9 GOS Institute of Child Health Clinical Health Research Trust PhD Studentship.
10

11 Financial Disclosures:

12
13 This work was undertaken at University College London (UCL) Institute of Child
14 Health (ICH)/Great Ormond Street Hospital and Moorfields Eye Hospital/UCL
15 Institute of Ophthalmology, both of which receive a proportion of funding from the
16 Department of Health's National Institute for Health Research (NIHR) Biomedical
17 Research Centres funding scheme. Members of the team are also supported by the
18 Ulverscroft Foundation. Professor Rahi is an NIHR Senior Investigator. The views
19 expressed in this article are those of the author(s) and not necessarily those of the
20 NHS, the NIHR, or the Department of Health.
21
22

23 Other Acknowledgements:

24
25 We acknowledge the contribution of the members of the Child Vision PROMs group
26 (Ameenat Lola Solebo, Phillippa Cumberland, Naomi Dale, Peng Tee Khaw, Gillian
27 Lewando Hundt, Alki Liasis, Anthony Moore and Alison Salt) and the study advisory
28 group (Corie Brown, Jackie Osborne, Paula Thomas, and Jude Thompson).
29
30

31 We thank the following UK hospitals and colleagues who helped with patient
32 identification and recruitment: East Lancashire Hospitals NHS Trust (May Mohan,
33 Matthew Milner and Heather Collier, on behalf of the Ophthalmology Team),
34 Southampton University Hospitals NHS Trust (Jay Self and Megan Ranger, on
35 behalf of the Ophthalmology Team), West Suffolk NHS Foundation Trust (Anthony
36 Vivian and Jen Bacon, on behalf of the Ophthalmology Team), Royal Cornwall
37 Hospitals NHS Trust, Hampshire Hospitals NHS Foundation Trust (Luke Clifford, on
38 behalf of the Ophthalmology Team), Countess of Chester Hospitals NHS Foundation
39 Trust (Jeremy Butcher on behalf of the Ophthalmology Team), Hinchingsbrook
40 Healthcare NHS Trust (Melanie Hingorani and Paula Thurnbull, on behalf of the
41 Ophthalmology Team), Mid Cheshire Hospitals NHS Foundation Trust (Simon
42 Walker and Sally Smith, on behalf of the Ophthalmology Team), University Hospitals
43 of North Midlands NHS Trust (Annie Joseph and Ruth Jones, on behalf of the
44 Ophthalmology Team), The Queen Elizabeth Hospital Kings Lynn NHS Foundation
45 Trust (Vineet Singh, on behalf of the Ophthalmology Team), Royal Devon & Exeter
46 NHS Foundation Trust (Anthony Quinn, on behalf of the Ophthalmology Team),
47 University Hospital of Wales, Cardiff and Vale University Health Board (Patrick Watts
48 and Tina McDonald, on behalf of the Ophthalmology Team), Birmingham Children's
49 Hospital NHS Foundation Trust (Joe Abbott, Manoj Parulekar and Laura Ramm, on
50 behalf of the Ophthalmology Team), Epsom & St. Helier University Hospitals NHS
51 Trust (Jane Leitch, on behalf of the Ophthalmology Team), Leeds Teaching
52 Hospitals NHS Trust (Vernon Long and Janice Hoole, on behalf of the
53 Ophthalmology Team), Portsmouth Hospitals NHS Trust (Kate Bolton, on behalf of
54 the Ophthalmology Team), Bristol Eye Hospital, University Hospitals Bristol NHS
55 Foundation Trust (Cathy Williams and Bekki Coles, on behalf of the Ophthalmology
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Team), Bradford Royal Infirmary, Bradford Teaching Hospitals NHS Foundation Trust (Rachel Pilling and Shegufta Farooq, on behalf of the Ophthalmology Team), Addenbrookes Hospital, Cambridge University Hospitals NHS Trust (Louise Allen, on behalf of the Ophthalmology Team), Manchester Royal Eye Hospital, Central Manchester University Hospitals NHS Foundation Trust (Jane Ashworth, on behalf of the Ophthalmology Team).

REFERENCES

1. D'Allura T. Enhancing the social interaction skills of preschoolers with visual impairments. *J Vis Impair Blind*. 2002;96(08):576-584.
2. Khadka J, Ryan B, Margrain TH, Woodhouse JM, Davies N. Listening to voices of children with a visual impairment: a focus group study. *Br J Vis Impair*. 2012;30(3):182-196.
3. Reeve BB, Wyrwich KW, Wu AW, et al. ISOQOL recommends minimum standards for patient-reported outcome measures used in patient-centered outcomes and comparative effectiveness research. *Qual Life Res*. 2013;22(8):1889-1905.
4. Khadka J, Ryan B, Margrain TH, Court H, Woodhouse JM. Development of the 25-item Cardiff Visual Ability Questionnaire for Children (CVAQC). *Br J Ophthalmol*. 2010;94(6):730-735.
5. Gothwal VK, Sumalini R, Bharani S, Reddy S, Bagga DK. The second version of the L. V. Prasad-functional vision questionnaire. *Optom Vis Sci*. 2012;89(11):1601-1610.
6. Tadić V, Hogan A, Sobti N, Knowles RL, Rahi JS. Patient-reported outcome measures (PROMs) in paediatric ophthalmology: a systematic review. *Br J Ophthalmol*. 2013;97:1369-1381.
7. Tadić V, Cooper A, Cumberland P, Lewando-Hundt G, Rahi JS. Development of the Functional Vision Questionnaire for Children and Young People with Visual Impairment: the FVQ_CYP. *Ophthalmology* 2013;120(12):2725-2732.
8. Rahi JS, Tadić V, Keeley S, Lewando-Hundt G. Capturing children and young people's perspectives to identify the content for a novel vision-related quality of life instrument. *Ophthalmology*. 2011;118(5):819-824.
9. Tadić V, Cooper A, Cumberland P, Lewando-Hundt G, Rahi JS. Measuring the quality of life of visually impaired children: first stage psychometric evaluation of the novel VQoL_CYP instrument. *PLoS One*. 2016;11(2):e0146225.
10. Tadić V, Robertson AO, Cortina-Borja M, Rahi JS. An age-and stage-appropriate patient-reported outcome measure of vision-related quality of life (VQoL) of children and young people with visual impairment. *Ophthalmology*. 2020;127(2):249-260.
11. Castleberry A. NVivo 10 [software program]. Version 10. QSR International. *Am J Pharm Educ*. 2014;7(1):25.
12. IBM SPSS Statistics. Version 24. IBM Corp: Armonk, New York.
13. Pesudovs K, Burr JM, Harley C, Elliott DB. The development, assessment, and selection of questionnaires. *Optom Vis Sci*. 2007;84(8):663-674.
14. Rasch G. *Some Probabilistic Models for Intelligence and Attainment Tests*. Chicago: University of Chicago Press; 1980.
15. Wright BD, Douglas GA. *Best Test Design and Self-Tailored Testing. Research Memorandum No 19*. Chicago: Statistical Laboratory, Department of Education, University of Chicago; 1975.
16. J.M. Linacre. A User's Guide to WINSTEPS MINISTEPS Rasch-Model Computer Programs [book on the internet]. Winsteps; 2012. Chapter 18.31: Equating

- and linking tests [cited 29 July 2019]; p. 567-573. Available at <https://www.winsteps.com/winman/equating.htm>.
17. Stocking ML, Lord FM. Developing a common metric in item response theory. *Appl Psychol Meas*. 1983;7(2):201-210.
 18. Dorans NJ, Kulick E. Differential item functioning on the Mini-Mental State Examination: an application of the Mantel-Haenszel and standardization procedures. *Med Care*. 2006;44(11):S107-S14.
 19. Linacre J. Missing data. <https://www.winsteps.com/winman/missingdata.htm> (accessed August 30, 2019).
 20. Fischer GH, Molenaar IW. *Rasch models: Foundations, Recent Developments, and Applications*. New York: Springer Science & Business Media; 2012.
 21. Rahi JS, Cable N. Severe visual impairment and blindness in children in the UK. *Lancet*. 2003;362(9393):1359-1365.
 22. Varni JW, Limbers CA, Burwinkle TM. How young can children reliably and validly self-report their health-related quality of life?: an analysis of 8,591 children across age subgroups with the PedsQL™ 4.0 Generic Core Scales. *Health Qual Life Outcomes*. 2007;5(1):1.
 23. World Health Organization. *International Classification of Functioning, Disability and Health: ICF*. Geneva: World Health Organization; 2001.
 24. Angeles-Han ST, Griffin KW, Lehman TJ, et al. The importance of visual function in the quality of life of children with uveitis. *J AAPOS*. 2010;14(2):163-168.
 25. Bokhary KA, Suttle C, Alotaibi AG, Stapleton F, Ying Boon M. Development and validation of the 21-item children's vision for living scale (CVLS) by Rasch analysis. *Clin Exp Optom*. 2013;96(6):566-576.
 26. Elsmann EBM, van Nispen RMA, van Rens G. Feasibility of the Participation and Activity Inventory for Children and Youth (PAI-CY) and Young Adults (PAI-YA) with a visual impairment: a pilot study. *Health Qual Life Outcomes*. 2017;15(1):98.
 27. Hatt SR, Leske DA, Castañeda YS, et al. Development of pediatric eye questionnaires for children with eye disease. *Am J Ophthalmol*. 2019;200:201-217.
 28. Valderas J, Kotzeva A, Espallargues M, et al. The impact of measuring patient-reported outcomes in clinical practice: a systematic review of the literature. *Qual Life Res*. 2008;17(2):179-193.
 29. Janssens A, Thompson Coon J, Rogers M, et al. A systematic review of generic multidimensional patient-reported outcome measures for children, part I: descriptive characteristics. *Value Health*. 2015;18(2):315-333.
 30. Robertson R, Wenzel L, Thompson J, Charles A. *Understanding NHS financial pressures. How are they affecting patient care?* London: The King's Fund; 2017.
 31. Batty MJ, Moldavsky M, Foroushani PS, et al. Implementing routine outcome measures in child and adolescent mental health services: from present to future practice. *Child Adolesc Ment Health*. 2013;18(2):82-87.
 32. Greenhalgh J, Long AF, Flynn R. The use of patient reported outcome measures in routine clinical practice: lack of impact or lack of theory? *Soc Sci Med*. 2005;60(4):833-843.

- 1
2
3
4 33. Angeles-Han ST, Griffin KW, Harrison MJ, et al. Development of a vision-related
5 quality of life instrument for children ages 8-18 years for use in juvenile
6 idiopathic arthritis-associated uveitis. *Arthritis Care Res.* 2011;63(9):1254-
7 1261.
8
9 34. Tadić V, Rahi JS. One size doesn't fit all: time to revisit patient-reported outcome
10 measures (PROMs) in paediatric ophthalmology? *Eye (Lond).*
11 2017;31(4):511.
12
13 35. Eiser C, Morse R. Can parents rate their child's health-related quality of life?
14 Results of a systematic review. *Qual Life Res* 2001;10:347-357.
15
16 36. Matza LS, Patrick DL, Riley AW, et al. Pediatric patient-reported outcome
17 instruments for research to support medical product labeling: report of the
18 ISPOR PRO good research practices for the assessment of children and
19 adolescents task force. *Value Health* 2013;16(4):461-479.
20
21 37. Linacre J. Optimizing rating scale category effectiveness. *J Appl Meas.*
22 2002;3(1):85-106.
23
24 38. Khadka J, Pesudovs K, McAlinden C, Vogel M, Kernt M, Hirneiss C.
25 Reengineering the glaucoma quality of life-15 questionnaire with rasch
26 analysis. *Invest Ophthalmol Vis Sci.* 2011;52(9):6971-6977.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 **FIGURE CAPTIONS**
5

6 Figure 1. Category probability curves for the 28-item FVQ Child (left), and 38-item
7 FVQ Young Person (right)
8

9 Figure 1: Category probability curves showing the probability of selecting response
10 categories across the scale of item difficulty for age-appropriate extensions of the
11 FVQ_CYP³⁷
12

13 Figure 2. Item-Person map for the 28-item FVQ Child (left), and 38-item
14 FVQ Young Person (right)
15

16 Figure 2: Item-person maps illustrating acceptable targeting of FVQ items (located
17 on the right hand side of the dashed line) to responders (located on the left side of
18 the dashed line and represented by X).³⁸ Participants with higher functional vision
19 and items with higher difficulty are at the bottom half of the map. M = mean; S = 1
20 standard deviation from the mean; T = 2 standard deviations from the mean.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

TABLES

Table 1. Demographic and clinical characteristics of participants in each phase of FVQ_CYP instrument adaptation.						
Demographic characteristic	Phase 1		Phase 2		Phase 3	
	Children (<i>n</i> = 12)	Young People (<i>n</i> = 17)	Children (<i>n</i> = 12)	Young People (<i>n</i> = 16)	Children (<i>n</i> = 113^a)	Young People (<i>n</i> = 96^b)
Age						
6	1 (8.3)	-	-	-	-	-
7	-	-	2 (16.7)	-	3 (2.65)	-
8	4 (33.3)	-	6 (50)	-	22 (19.47)	-
9	7 (58.3)	-	3 (25)	-	26 (23)	-
10	-	-	1 (8.3)	-	15 (13.27)	-
11	-	-	-	-	24 (21.24)	-
12	-	-	-	-	22 (19.47)	-
13	-	-	-	3 (18.75)	1 (0.88)	12 (12.5)
14	-	-	-	2 (12.5)	-	25 (26.04)
15	-	-	-	3 (18.75)	-	19 (19.79)
16	-	7 (41.18)	-	2 (12.5)	-	18 (18.75)
17	-	8 (47.06)	-	3 (18.75)	-	20 (20.83)
18	-	1 (5.88)	-	3 (18.75)	-	2 (2.08)
19	-	1 (5.88)	-	-	-	-
Gender						
Male	8 (66.7)	10 (58.82)	8 (66.7)	8 (50)	52 (46.02)	52 (54.17)
Female	4 (33.3)	7 (41.18)	4 (33.3)	8 (50)	61 (53.98)	44 (45.83)
Ethnicity						
White UK majority (White British)	8 (66.7)	10 (58.82)	5 (41.7)	11 (68.75)	62 (54.87)	62 (64.58)
White other (e.g. African, Polish, Turkish)	-	1 (5.88)	2 (16.7)	1 (6.25)	9 (7.96)	7 (7.29)
Black (British, African, Caribbean)	1 (8.3)	-	1 (8.3)	-	9 (7.96)	3 (3.13)
Asian (Indian, Bangladeshi, Pakistani)	2 (16.7)	3 (17.65)	2 (16.7)	4 (25)	25 (22.12)	12 (12.5)
Asian other (Arabic)	-	1 (5.88)	-	-	3 (2.65)	2 (2.08)
Chinese	-	-	-	-	-	-

Table 1. Demographic and clinical characteristics of participants in each phase of FVQ_CYP instrument adaptation.

Demographic characteristic	Phase 1		Phase 2		Phase 3	
	Children (<i>n</i> = 12)	Young People (<i>n</i> = 17)	Children (<i>n</i> = 12)	Young People (<i>n</i> = 16)	Children (<i>n</i> = 113 ^a)	Young People (<i>n</i> = 96 ^b)
Mixed	1 (8.3)	2 (11.76)	2 (16.7)	-	3 (2.65)	2 (2.08)
Missing	-	-	-	-	2 (1.77)	8 (8.33)
Severity of visual impairment						
LV: logMAR ≤0.46	-	1 (5.88)	-	-	5 (4.42)	1 (1.04)
VI1: logMAR 0.48-0.70	4 (33.3)	8 (47.06)	4 (33.3)	9 (56.25)	50 (44.25)	29 (30.21)
VI2: logMAR 0.72-1.00	5 (41.7)	3 (17.65)	3 (25)	5 (31.25)	40 (35.4)	37 (38.54)
SVI: logMAR 1.02-1.30	-	2 (11.76)	1 (8.3)	1 (6.25)	8 (7.08)	12 (12.5)
Blind: logMAR ≥1.32	3 (25)	3 (17.65)	4 (33.3)	1 (6.25)	10 (8.85)	17 (17.71)
Timing of onset of visual impairment						
Early (≤2 years)	12 (100)	15 (88.24)	12 (100)	10 (62.5)	99 (87.61)	79 (82.29)
Late	-	2 (11.76)	-	6 (37.5)	14 (12.39)	17 (17.71)
Nature of deterioration of visual impairment						
Stable	9 (75)	12 (70.59)	6 (50)	5 (31.25)	74 (65.49)	81 (84.38)
Progressive	3 (25)	5 (29.41)	6 (50)	11 (68.75)	39 (34.51)	15 (15.62)
Diagnosis by site of visual impairment^d						
Whole globe and anterior segment	-	1 (5.88)	1 (8.3)	1 (6.25)	2 (1.77)	3 (3.13)
Glaucoma, primary or secondary	1 (8.3)	-	3 (25)	-	10 (8.85)	10 (10.42)
Cornea (sclerocornea and corneal capacities)	-	-	-	1 (6.25)	2 (1.77)	2 (2.08)
Lens (cataract and aphakia)	1 (8.3)	-	1 (8.3)	2 (12.5)	14 (12.39)	9 (9.38)
Uvea	-	-	-	-	6 (5.31)	8 (8.33)
Retina	9 (75)	12 (70.59)	8 (66.67)	9 (56.25)	71 (62.83)	68 (70.83)
Optic nerve	1 (8.3)	3 (17.65)	1 (8.3)	3 (18.75)	13 (11.5)	6 (6.25)
Cerebral/visual pathways	1 (8.3)	-	-	1 (6.25)	5 (4.42)	9 (9.38)

Table 1. Demographic and clinical characteristics of participants in each phase of FVQ_CYP instrument adaptation.

Demographic characteristic	Phase 1		Phase 2		Phase 3	
	Children (<i>n</i> = 12)	Young People (<i>n</i> = 17)	Children (<i>n</i> = 12)	Young People (<i>n</i> = 16)	Children (<i>n</i> = 113 ^a)	Young People (<i>n</i> = 96 ^b)
Other (idiopathic nystagmus, high refractive error)	-	6 (35.29)	1 (8.3)	-	19 (16.81)	16 (16.67)
Index of multiple deprivation quintile rank						
1: most deprived	2 (16.7)	1 (5.88)	1 (8.3)	2 (12.5)	22 (19.47)	18 (18.75)
2	1 (8.3)	2 (11.76)	5 (41.7)	-	23 (20.35)	19 (19.79)
3	3 (25)	4 (23.53)	2 (16.7)	4 (25)	25 (22.12)	15 (15.62)
4	2 (16.7)	8 (47.06)	3 (25)	3 (18.75)	19 (16.81)	17 (17.71)
5: least deprived	4 (33.3)	2 (11.76)	1 (8.3)	7 (43.75)	21 (18.58)	27 (28.13)
Missing	-	-	-	-	3 (2.65) ^c	-

^a Four children excluded from analysis due to incomplete (more than 25% data missing) child data (e.g. parent proxy report provided instead).

^b Two young people excluded from analysis due to completely missing (*n*=1) young person data (e.g. parent proxy report provided instead) and failure to consent (*n*=1) to use of young person data.

^c Data missing due to postcode data not provided by the managing clinical team, as per local governance approval at the patient identification centre.

^d Does not add up to 100% because some children had visual impairment originating in multiple sites.

Table 2. Rasch Fit Statistics, item measure and differential item functioning (DIF) contrasts for the 28-item and 38-item age-appropriate FVQ instrument extensions, and DIF contrasts for the overlapping items (overlapping items shown in bold).

FVQ_Child	FVQ_Young Person	FVQ_Child					FVQ_Young Person					Core items
Item	Item	Item measure (logits)	Infit MNSQ ^a	Outfit MNSQ	DIF ^b contrast by age (logits)	DIF contrast by gender (logits)	Item measure (logits)	Infit MNSQ	Outfit MNSQ	DIF contrast by age (logits)	DIF contrast by gender (logits)	DIF contrasts by sample (i.e. children vs. young people)
Watching TV	Watching TV	0.31	0.94	0.89	-0.26	0.05	0.33	0.87	0.96	-0.22	0.19	-0.19
Playing video and computer games	Playing video and computer games	0.27	0.98	0.99	-0.19	-0.35	-0.16	1.04	1.08	-0.60	0.23	0.22
Playing other indoor games, such as board games or card games	Playing indoor games, such as board games or card games	0.60	0.76	0.72	0.34	0	0.26	0.80	0.88	-0.07	0.32	0.22
Playing outdoor games, such as tag or hide and seek		0.03	0.97	0.94	-0.23	-0.06						-
Using the computer at home to do my school work	Using the computer at home to do my homework	0.37	1.32	1.29	-0.54	-0.24	0.62	1.24	1.33	0.77	0.21	-0.39
	Reading food packets, tickets, labels or recipes						-1.30	0.78	0.73	-0.07	0.28	-
Doing household jobs, for example, tidying up my toys	Doing household chores, for example, washing up or tidying my bedroom	1.33	1.07	1.04	0.02	0.33	0.99	0.79	0.80	-0.08	-0.07	0.31
	Looking after my appearance, for example, doing my hair, shaving, or putting on make-up						0.62	0.95	0.94	0.31	-0.44	-

Table 2. Rasch Fit Statistics, item measure and differential item functioning (DIF) contrasts for the 28-item and 38-item age-appropriate FVQ instrument extensions, and DIF contrasts for the overlapping items (overlapping items shown in bold).

FVQ_Child		FVQ_Child					FVQ_Young Person					Core items
Item	Item	Item measure (logits)	Infit MNSQ ^a	Outfit MNSQ	DIF ^b contrast by age (logits)	DIF contrast by gender (logits)	Item measure (logits)	Infit MNSQ	Outfit MNSQ	DIF contrast by age (logits)	DIF contrast by gender (logits)	DIF contrasts by sample (i.e. children vs. young people)
	Making myself a snack at home						1.60	0.68	0.66	0.63	0.43	-
	Making myself a meal						0.37	0.92	0.90	0.84	0.34	-
	Finding objects I have dropped such as coins or glasses on a low contrast surface						-1.33	1.06	1.22	0	0.23	-
Using the computer in school lessons	Using the computer at school or college to do schoolwork/coursework	0.16	0.89	0.87	-0.19	-0.22	0.43	1.01	1.02	0.30	-0.09	-0.45
Reading small print worksheets and textbooks like dictionaries	Reading small print textbooks, worksheets and exam papers	-1.93	0.99	0.92	-0.08	0.50	-2.21	0.88	0.91	-0.18	-0.44	-0.15
Reading enlarged worksheets and textbooks like dictionaries		1.53	1.20	1.40	-0.18	-0.14						-
Drawing or painting		0.90	1.18	1.23	-0.32	0.72						-
Reading other people's handwriting	Reading other people's handwriting	-1.23	0.60	0.60	0.30	0	-1.59	0.85	0.83	-0.20	0.08	0
Seeing the board in the classroom	Seeing the board in the classroom when sitting at the	-1.38	1.11	1.02	-0.49	0.29	-1.21	1.06	1.00	-0.47	0.23	-0.54

Table 2. Rasch Fit Statistics, item measure and differential item functioning (DIF) contrasts for the 28-item and 38-item age-appropriate FVQ instrument extensions, and DIF contrasts for the overlapping items (overlapping items shown in bold).

FVQ_Child	FVQ_Young Person	FVQ_Child					FVQ_Young Person					Core items
Item	Item	Item measure (logits)	Infit MNSQ ^a	Outfit MNSQ	DIF ^b contrast by age (logits)	DIF contrast by gender (logits)	Item measure (logits)	Infit MNSQ	Outfit MNSQ	DIF contrast by age (logits)	DIF contrast by gender (logits)	DIF contrasts by sample (i.e. children vs. young people)
	front											
Recognising people, for example in school corridors	Recognising people, for example, in corridors at school/college or shops	-0.20	1.01	1.02	0.34	0.16	-0.89	1.31	1.35	-0.74	-0.10	0.41
Recognising other people's facial expressions	Recognising other people's facial expressions when they are close to me/at arm's length	0.25	1.06	1.02	0.40	0.31	0.16	1.34	1.27	-0.32	0.51	-0.11
Finding friends in the playground	Finding friends in crowded areas	-1.10	0.97	0.89	0.21	0	-1.77	0.90	1.17	0	-0.41	0.29
Doing maths in lessons	Doing maths	0.73	1.16	1.11	-0.24	-0.30	1.26	1.15	1.15	0.23	0.16	-0.56
Doing literacy in lessons		0.67	0.92	0.96	-0.21	-0.02						-
	Doing science						0.44	1.08	1.10	0.06	0.41	-
Doing PE	Doing sports at school/college	0.05	1.12	1.20	0	-0.52	-0.19	1.42	1.47	-0.19	-0.57	0
Keeping up with the teacher in lessons	Keeping up with the teacher or tutor in lessons	0.32	1.04	1.07	0	0	0.45	0.81	0.80	-0.14	0.41	-0.29
Keeping up with other children in lessons	Keeping up with other students in lessons	0.10	0.85	0.91	0.52	-0.10	0.51	0.71	0.71	-0.06	0	-0.59

Table 2. Rasch Fit Statistics, item measure and differential item functioning (DIF) contrasts for the 28-item and 38-item age-appropriate FVQ instrument extensions, and DIF contrasts for the overlapping items (overlapping items shown in bold).

FVQ_Child	FVQ_Young Person	FVQ_Child					FVQ_Young Person					Core items
Item	Item	Item measure (logits)	Infit MNSQ ^a	Outfit MNSQ	DIF ^b contrast by age (logits)	DIF contrast by gender (logits)	Item measure (logits)	Infit MNSQ	Outfit MNSQ	DIF contrast by age (logits)	DIF contrast by gender (logits)	DIF contrasts by sample (i.e. children vs. young people)
Getting around school without someone helping me	Getting around school/college by myself	1.82	1.19	1.02	-0.39	0	1.71	0.76	0.72	-0.19	-0.09	0.17
Playing team sports without special balls	Playing team sports, such as football, without adaptations such as special balls	-0.31	1.25	1.18	0.41	-0.35	-0.68	1.24	1.17	-0.52	-0.91	0.09
Seeing small balls when playing games like tennis or cricket	Seeing small balls when playing games, such as tennis or cricket	-1.10	1.05	1.00	0.28	0	-2.35	0.92	0.89	0.26	0.16	0.87
Seeing big moving objects, such as bicycles passing by	Seeing big moving objects, such as bikes passing, in daylight	0.59	0.73	0.74	0.44	0.14	0.36	0.81	0.79	0	-0.51	0.09
Getting around outdoors in daytime	Getting around outdoors e.g. shops or the park, by myself when it's daylight	0.79	0.76	0.74	0.05	-0.13	0.74	0.54	0.52	0.57	-0.41	-0.05
	Getting around outdoors e.g. shops or the park, by						-0.71	1.08	1.02	0.31	-0.28	-

Table 2. Rasch Fit Statistics, item measure and differential item functioning (DIF) contrasts for the 28-item and 38-item age-appropriate FVQ instrument extensions, and DIF contrasts for the overlapping items (overlapping items shown in bold).

FVQ_Child	FVQ_Young Person	FVQ_Child					FVQ_Young Person					Core items
Item	Item	Item measure (logits)	Infit MNSQ ^a	Outfit MNSQ	DIF ^b contrast by age (logits)	DIF contrast by gender (logits)	Item measure (logits)	Infit MNSQ	Outfit MNSQ	DIF contrast by age (logits)	DIF contrast by gender (logits)	DIF contrasts by sample (i.e. children vs. young people)
	myself when it's dark											
	Getting around in crowds by myself						-0.61	0.95	0.88	0.75	-0.53	-
	Finding my way around an unfamiliar house or a new building						-0.24	0.81	0.77	0.68	-0.44	-
Reading signs and posters at stations or shops	Reading signs and posters at stations or shops	-0.96	0.60	0.63	-0.07	0.10	-1.18	0.85	0.76	-0.23	-0.07	0.48
	Finding correct money to pay when shopping						0.75	0.97	1.00	0.09	0	-
Watching films in the cinema	Watching films in the cinema	1.04	1.01	0.95	0.35	0	0.69	0.74	0.72	0	-0.09	0.27
Watching shows at the theatre	Watching shows, such as plays, at the theatre	-0.26	1.09	1.07	-0.34	0.36	-0.65	0.98	1.00	-0.21	0.24	0.14
	Crossing the road by myself						0.28	0.97	0.95	0.46	-0.76	-
	Using public transport, such as trains, buses or the tube by myself						-0.22	1.02	1.02	0.40	-0.83	-
	Using a mobile phone to text people						1.34	1.27	1.15	0.15	0.70	-
	Using a mobile phone or tablet for						1.63	1.13	1.02	-0.21	-0.21	

Table 2. Rasch Fit Statistics, item measure and differential item functioning (DIF) contrasts for the 28-item and 38-item age-appropriate FVQ instrument extensions, and DIF contrasts for the overlapping items (overlapping items shown in bold).

FVQ_Child		FVQ_Child					FVQ_Young Person					Core items
Item	Item	Item measure (logits)	Infit MNSQ ^a	Outfit MNSQ	DIF ^b contrast by age (logits)	DIF contrast by gender (logits)	Item measure (logits)	Infit MNSQ	Outfit MNSQ	DIF contrast by age (logits)	DIF contrast by gender (logits)	DIF contrasts by sample (i.e. children vs. young people)
	social networking, for example, Facebook, Twitter or MySpace											

^a MNSQ = Mean-square standardized residual within the pre-defined interval (0.5, 1.5).¹⁰

^b DIF = Differential item functioning within a 1 logit threshold.^{11, 14}

Table 3a. Conversion table for transforming raw scores on the 28-item FVQ_Child version) into comparable Rasch person measures^a

Score	Measure	S.E. ^b	Score	Measure	S.E.	Score	Measure	S.E.
0	0.00	14.02	29	42.88	2.04	58	58.28	2.13
1	9.40	7.78	30	43.42	2.03	59	58.87	2.15
2	15.00	5.61	31	43.96	2.01	60	59.48	2.17
3	18.39	4.66	32	44.49	2.01	61	60.10	2.19
4	20.88	4.10	33	45.01	2.00	62	60.74	2.22
5	22.87	3.72	34	45.54	1.99	63	61.39	2.24
6	24.55	3.45	35	46.06	1.99	64	62.06	2.27
7	26.01	3.24	36	46.57	1.98	65	62.74	2.31
8	27.31	3.07	37	47.09	1.98	66	63.45	2.34
9	28.49	2.93	38	47.60	1.98	67	64.18	2.38
10	29.57	2.81	39	48.11	1.98	68	64.94	2.43
11	30.57	2.72	40	48.62	1.97	69	65.73	2.48
12	31.51	2.63	41	49.13	1.97	70	66.55	2.53
13	32.39	2.56	42	49.64	1.98	71	67.41	2.59
14	33.22	2.49	43	50.16	1.98	72	68.31	2.67
15	34.02	2.44	44	50.67	1.98	73	69.27	2.75
16	34.78	2.39	45	51.19	1.98	74	70.29	2.84
17	35.51	2.34	46	51.70	1.99	75	71.39	2.96
18	36.22	2.30	47	52.22	1.99	76	72.59	3.09
19	36.90	2.26	48	52.75	2.00	77	73.91	3.26
20	37.56	2.23	49	53.27	2.01	78	75.39	3.46
21	38.21	2.20	50	53.80	2.02	79	77.08	3.74
22	38.83	2.17	51	54.34	2.03	80	79.08	4.11
23	39.44	2.15	52	54.88	2.04	81	81.59	4.66
24	40.04	2.13	53	55.43	2.05	82	84.99	5.61
25	40.63	2.10	54	55.98	2.06	83	90.59	7.79
26	41.20	2.09	55	56.54	2.08	84	100.00	14.02
27	41.77	2.07	56	57.11	2.09			
28	42.33	2.05	57	57.69	2.11			

Table 3b. Conversion table for transforming raw scores on the 38-item FVQ_Young Person into comparable Rasch person measures^a

Score	Measure	S.E. ^b	Score	Measure	S.E.	Score	Measure	S.E.
0	0.00	12.49	39	43.46	1.70	78	59.18	1.71
1	8.41	6.96	40	43.88	1.69	79	59.61	1.72
2	13.45	5.03	41	44.30	1.68	80	60.05	1.73
3	16.53	4.19	42	44.71	1.68	81	60.49	1.74
4	18.80	3.70	43	45.13	1.67	82	60.94	1.75
5	20.63	3.37	44	45.54	1.66	83	61.39	1.76
6	22.18	3.12	45	45.94	1.66	84	61.85	1.77
7	23.52	2.94	46	46.35	1.66	85	62.32	1.79
8	24.73	2.79	47	46.75	1.65	86	62.79	1.80
9	25.82	2.66	48	47.15	1.65	87	63.27	1.82
10	26.82	2.56	49	47.55	1.64	88	63.76	1.83
11	27.75	2.47	50	47.95	1.64	89	64.26	1.85
12	28.62	2.39	51	48.34	1.64	90	64.77	1.87
13	29.44	2.32	52	48.74	1.64	91	65.29	1.89
14	30.21	2.26	53	49.13	1.64	92	65.82	1.92
15	30.95	2.21	54	49.53	1.63	93	66.37	1.94
16	31.65	2.16	55	49.92	1.63	94	66.93	1.97
17	32.33	2.12	56	50.31	1.63	95	67.51	2.00
18	32.97	2.08	57	50.70	1.63	96	68.11	2.03
19	33.60	2.04	58	51.09	1.63	97	68.73	2.07
20	34.20	2.01	59	51.49	1.63	98	69.37	2.11
21	34.79	1.98	60	51.88	1.63	99	70.03	2.15
22	35.36	1.95	61	52.27	1.63	100	70.73	2.20
23	35.91	1.93	62	52.67	1.64	101	71.46	2.26
24	36.45	1.90	63	53.06	1.64	102	72.24	2.32
25	36.98	1.88	64	53.45	1.64	103	73.06	2.40
26	37.49	1.86	65	53.85	1.64	104	73.93	2.48
27	38.00	1.84	66	54.25	1.64	105	74.88	2.59
28	38.49	1.82	67	54.95	1.65	106	75.91	2.71
29	38.98	1.81	68	55.05	1.65	107	77.04	2.86
30	39.45	1.79	69	55.45	1.65	108	78.32	3.04
31	39.92	1.78	70	55.85	1.66	109	79.79	3.29
32	40.38	1.77	71	56.26	1.66	110	81.54	3.62
33	40.84	1.75	72	56.67	1.67	111	83.73	4.12
34	41.29	1.74	73	57.08	1.67	112	86.71	4.97
35	41.73	1.73	74	57.49	1.68	113	91.66	6.91
36	42.17	1.72	75	57.91	1.69	114	100.00	12.47
37	42.60	1.71	76	58.33	1.69			
38	43.03	1.70	77	58.75	1.70			

^a Scores ranging from 1-4 must be re-scored into a scale of 0-3 before conversion.

^b Model-based standard error of the measure.

Figure 1. Category probability curves.
[Click here to download high resolution image](#)





Figure 2. Item-Person map
[Click here to download high resolution image](#)





Table of contents statement:

We report the development of age-specific extensions of the FVQ_CYP to allow for use with a broader age-range of children and young people with visual impairment. The FVQ_Child and FVQ_Young Person are psychometrically robust, age-appropriate versions of the FVQ_CYP, which can be used cross-sectionally or sequentially/longitudinally across the age-range of 8 up to 18 years in clinical practice and research.

Highlights

- Age-appropriate versions of a patient-reported outcome measure were developed.
- The FVQ_Child (8-12 years) and FVQ_Young Person (13-18 years) measure functional vision.
- Development comprised Rasch analysis and calibration on the same measurement scale.
- They can be used cross-sectionally or sequentially in practice and research.

SUPPLEMENTARY TABLE

Supplementary Table 1. Item reduction in Phase 3

Items removed – FVQ_Child		Items removed – FVQ_Young Person	
Item	Removal criteria	Item	Removal criteria
Reading small writing such as food packets or instructions for toys	Rasch – <i>removed during calibration because of DIF^a (more difficult for children (vs. young people))</i>		
		Reading enlarged textbooks, worksheets and exam papers	Rasch – <i>removed because of DIF by age (more difficult for older young people)</i>
		Drawing or painting	Rasch – <i>removed because of DIF by age (more difficult for older young people)</i>
		Doing English or literacy	Rasch – <i>removed because of DIF by gender (more difficult for males)</i>
Getting around outdoors when it is dark	Rasch – <i>removed due to item fit (OUTFIT MNSQ^b = 1.71)</i>		

^a DIF = Differential item functioning

^b MNSQ = Mean squared standardized residuals