



## LJMU Research Online

Yankelevich, V

**Notes on counter-orbiting globular clusters in the Milky Way**

<http://researchonline.ljmu.ac.uk/id/eprint/12887/>

### Article

**Citation** (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Yankelevich, V (2016) Notes on counter-orbiting globular clusters in the Milky Way. *Astronomical and Astrophysical Transactions*, 28 (4). ISSN 1055-6796**

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact [researchonline@ljmu.ac.uk](mailto:researchonline@ljmu.ac.uk)

<http://researchonline.ljmu.ac.uk/>

# Notes on counter-orbiting globular clusters in the Milky Way

V. Yankelevich\*

Southern Federal University, Rostov-on-Don, Russia

## Abstract

It is argued that Galactic globular clusters (GC) rotating retrograde may originate from prograde GCs that change their angular momentum due to gravitational perturbations from the Magellanic Cloud galaxies. It is shown that those galactic GCs with orbits near the Lagrange point of the system “Milky Way – Magellanic Clouds” can change the sign of their angular momentum in few Gyr time scale.

---

\*Email: vik-yank@yandex.ru

# 1 Introduction

The Galactic globular cluster system counts 150 clusters: 33 of them are recognized to orbit prograde, i.e. in the same direction as the halo stellar population of the Milky Way does, 15 of them are retrograde, i.e. counter-rotate the stellar halo, while for the rest the rotation direction is not identified yet [1, 2]. The origin of retrograde GCs is widely debated and remains totally controversial. A common understanding is that they are born outside the Milky Way and only relatively recently – a few Gyr ago – have been captured by the MW gravitation field on to confined orbits [3, 4, 5]. However a possibility that they could have formed in the MW and later on have changed their rotation under the gravitational influence from the Andromeda galaxy cannot be totally excluded (see discussion in [5]). In this model the captured clusters are to differ from the galactic ones in two respects: first, they seem to have higher angular momenta and lie predominantly on the peripheric regions of the Milky Way, and second, in general they are expected to form a separate population of clusters.

In Fig. 1 we show the distance-metallicity relation for those globular clusters with an identified orientation of their rotation from the Harris catalogue [1]. It is clearly seen that the distributions for prograde and retrograde are mostly similar and hardly can be distinguished: the retrograde clusters are spread lightly narrower and lie to a certain extent *within* the distribution of the prograde ones. A very vague anti-correlation between the metallicity and the distance for the two cluster populations does not look different. In addition, the most distant clusters are strikingly the progrades: *i*) the most distant prograde cluster orbits at around 100 kpc which is 3 times of the orbit of the most distant retrograde; and *ii*) the mean galactocentric radius for the prograde clusters  $\langle R \rangle = 11.76$  kpc, while for the retrograde clusters  $\langle R \rangle = 10.19$  kpc, which is in conflict to an intuitive expectation. Moreover, the most distant retrograde cluster has the galactocentric radius  $R \simeq 30$  kpc, while the most distant prograde one has  $R \simeq 100$  kpc – half an order higher. Crosses in Fig. 1 show globular clusters with uncertain direction of orbital motion, and the same behavior as demonstrated by the prograde and retrograde clusters are readily observed here. Overall, from this point of view the retrograde clusters do not seem to differ crucially from the other clusters. The only distinguishing feature in Fig. 1 is that the retrograde clusters are less spread in metallicities, which might be though attributed to individual evolutionary peculiarities.




Figure 1: The “distance-metallicity” relation for prograde GCs (open circles) and retrograde GCs (asterisks); crosses show clusters with unidentified rotation direction.

Neither differences between the progrades and retrogrades are seen in the distributions of their total (the sum of azimuthal and radial) velocities  $v_t = \sqrt{v_\theta^2 + v_r^2}$  as shown in Fig 2: both the populations show nearly equally wide distributions ranging from  $|v_\theta| \simeq 100 \text{ km s}^{-1}$  to  $|v_\theta| \simeq 400 \text{ km s}^{-1}$ , apart from one retrograde with  $v_t = 550 \text{ km s}^{-1}$  which is mostly due to the radial motion, nor in the other distributions, as for instance, the “mass-radius” and the “mass-metallicity” relations for both cluster subpopulations (Fig. 3a and 3b). Therefore, besides their reversed orbital motions the retrograde clusters are fairly similar to the whole population of galactic globular clusters and can hardly be distinguished from the other clusters.

In this brief note we demonstrate a principal possibility for galactic globular clusters to change the sign of their angular momentum under the action of the gravitational perturbations from the Magellanic Clouds.

## 2 Reversal of angular momentum of a globular cluster

Where the origin of retrograde globular clusters is concerned it is implicitly assumed that gravitational potential of the Milky Way is spherically (or




Figure 2: The distribution function of the modulus of total velocity  $v_t = \sqrt{v_r^2 + v_\theta^2}$  km s<sup>-1</sup> for the prograde (upper) and retrograde (lower) clusters; high velocities of the retrogrades are due to their high radial velocities.




Figure 3: The “mass-radius” (left) and “mass-metallicity” (right) relations for the progrades (open squares) and retrogrades (asterisk).

axially) symmetric, which implies conservation of the angular momentum of globular clusters. However, as soon as the orbits of globular clusters may extend up to distances comparable to the separation between the Milky Way and Magellanic Clouds, the non-axisymmetry of the potential and possible violation of the angular momentum conservation have to be noted and accounted for.

In order to demonstrate the possibility for globular clusters to reverse their angular momentum, in this paper we apply a simplistic dynamical model of a cluster moving in the joint gravitational field of the Milky Way and the Magellanic Clouds. For the Milky Way we assume the total mass  $M_{\text{halo}} = 7 \times 10^{11} M_{\odot}$ , the halo radius  $R_{\text{halo}} = 400$  kpc [4, 6, 7], and the Navarro-Frenk-White (NFW) density profile [8]

$$\rho(r) = \frac{\rho_0}{\frac{r}{h}(1 + \frac{r}{h})^2} \quad (1)$$

where  $h = 12.5$  kpc,  $\rho_0 = 1.06 \cdot 10^{-2} M_{\odot} \text{pc}^{-3}$ . The initial conditions for the positions and the velocities of globular clusters are assumed to correspond to Harris catalog. It is worth stressing that current estimates of the galactic mass are rather uncertain and may vary within factor 4–5 [7, 9, 10]. Depending on its exact value dynamics of the Magellanic Clouds may change dramatically from orbiting confined gravitationally to the Milky Way to move gravitationally unbound [11]. Our choice here corresponds to the gravitationally bound system “Milky Way – Magellanic Clouds” with the velocities  $v_x = 0$ ,  $v_y = -219$  km s $^{-1}$ ,  $v_z = 186$  km s $^{-1}$  for the LMC, and  $v_x = 0$ ,  $v_y = -174$  km s $^{-1}$ ,  $v_z = 173$  km s $^{-1}$  for the SMC [12], the accepted total

masses are [7]:  $M_{\text{LMC}} = 1.6 \cdot 10^{10} M_{\odot}$ , and  $M_{\text{SMC}} = 6.5 \cdot 10^9 M_{\odot}$ .

We start calculations of dynamics of a GC from a typical configuration shown in Fig. 4. In general, only those clusters experiencing sufficiently strong influence of the non-axisymmetry of the gravitational potential whose initial positions and velocities do lie in a restricted box. In other words, only the clusters occupying a restricted volume of the initial phase space are attracted in further evolution to the Magellanic Clouds. Therefore, a number of runs covering a set of initial conditions has to be checked in order to determine the trajectories which could, in principle, change considerably the angular momentum in the vicinity of the Magellanic Clouds.

We performed simulations by making use of the ordinary differential equation Runge-Kutta solver of 7-8 order. Each run covered 10 Gyr evolution with the accuracy  $10^{-8}$ .

In order to demonstrate the very possibility of a reversal of the angular momentum for globular clusters, first of all, we show here a typical example for a cluster starting from a box of the initial phase space with trajectories that attract to the Magellanic Clouds:  $y = 36$  kpc,  $z = 36$  kpc,  $v_0 = 212$  km s $^{-1}$ ; the trajectory is depicted in Fig. 4: in this particular example the cluster reverts its momentum in 7.45 Gyr. The corresponding evolution of the angular momentum is shown in Fig. 5 – the reversal occurs on a short time scale of  $\sim 1$  Myr.

Further, we demonstrate the quantity of GCs which potentially change angular momentum. A set of 3D 4 body simulations was performed. All the initial parameters of MW, LMC and SMC were the same as in the previous example. We took 10000 test particles for GCs, where the initial distance from the galactic center is distributed according with the law  $r \cdot e^{\frac{z}{\sigma}}$ . This law approximately coincides with the real distribution of known galactic clusters [1],[2]. The initial distribution of polar and azimuthal angles of GC is even. All the initial velocities except  $v_{tet}$  are distributed by the normal law with dispersion 130 km/s, for  $v_{tet}$  we used the absolute value of the normal distribution with dispersion 130 km/s. Thus all the GCs are prograde in the beginning. Each run covered 10 Gyr evolution with the accuracy  $10^{-8}$ . In the results there is about 3 – 4% of the GCs which become retrograde. Thus we demonstrate that the gravitational interactions in our system can change the direction of rotation of a GC though the number of such clusters is relatively small.




Figure 4: The trajectory of a GC moving in the joint field of the NFW density profile of the Milky Way and point-like potentials of LMC and SMC. Filled square and circles at around  $y = -40$  kpc and  $z = -40$  kpc correspond to the initial positions of the clobular cluster, and LMC and SMC galaxies, respectively; thin solid line is the trajectory of the GC, long dash line corresponds to the trajectory of LMC, while dash dotted line to the SMC. At  $t = 7.45$  Gyr the critical interaction of the GC with the orbiting each other SMC and LMC followed by a loop in the GC's trajectory is clearly seen.






Figure 5: The graph of total angular momentum of GC in the Navarro-Frenk-White density profile.

### 3 Conclusion

We have presented arguments favoring a new concept of the origin of the retrograde globular clusters. We have found that when a cluster starts from a selected volume of the initial phase space it falls into the area of attraction to the Magellanic Clouds, where strong non-axisymmetric violations of gravitational potential may change the cluster orbital angular momentum within the Hubble time. As a result, some of the clusters from this selected volume of the initial phase space may reverse the momentum, and turn from normal prograde rotation to the retrograde. We therefore argue that at least some of the retrograde clusters might have originated from the galactic population of globular clusters having been formed in the initial phase space with trajectories attractive to the Magellanic Clouds. Note, however, that this our scenario works only if the Magellanic Cloud galaxies are gravitationally confined to the Milky Way galaxy.

## 4 Acknowledgments

I thank prof. Yu. A. Shchekinov for formulation of the problem and guidance, prof. V. A. Marsakov for discussions and providing his processed Harris catalog, Douglas Heggie for useful discussions. I acknowledge support from the “Dynasty” Foundation, the Ministry of Education and Sciences of RF (project coded 14.A18.21.0787). Part of this work has been done while I was visiting the Henri Poincare Institute.

## References

- [1] Harris, W. E., 1996, *Astron. J*, 112, 1487
- [2] Borkova, T. A., Marsakov, V. A., 2000, *AstR*, 77, 750
- [3] van den Bergh, S., 1993, *AJ*, 105, 971
- [4] Boylan-Kolchin M., Springel V., White S., Jenkins A. , 2010, *MNRAS*, 406, 896
- [5] Kravtsov, V. V., 2001, *Astron. Astrophys. Transacti.*, 20, 1066
- [6] Deason A., Belokurov V., Evans N., An J., 2012, *MNRAS*, 424L, 44
- [7] Sofue, Y., 2012, *PASJ*, 64, 8
- [8] Navarro, J. F., Frenk, C. S., White, S. D. M., 1996, *ApJ*, 462, 563
- [9] McMillan, P. J., 2011, *MNRAS*, 414, 2446
- [10] Irrgang, A., Wilcox, B., Tucker, E., Schiefelbein, L., 2013, *A& A*, 549, A137
- [11] Besla, G., Kallivayalil, N., Hernquist, L., van der Marel, R. P., Cox, T. J., Keres, D., 2010, *ApJ*, 721L, 97
- [12] Brüns, C., <http://www.astro.uni-bonn.de/~rcbruens/simulationen/lmc/research.html>