

---

Poster

## Development of a reporter system for screening anti-biofilm activities



Ana Díaz Navarro, Aroa López Sánchez, Fernando Govantes Romero

Área de Microbiología/Centro Andaluz de Biología del Desarrollo, Universidad Pablo de Olavide, Ctra. De Utrera, Km 1, 41013 Sevilla

**Keywords:** c-di-GMP; Diguanylate cyclase; Biofilm formation

---

### ABSTRACT

Biofilm formation is responsible for increasing antibiotic tolerance in pathogenic bacteria. It is estimated that approximately 80% of chronic infections are associated with this phenomenon. Therefore, the search for therapeutic agents with specific biofilm targets has become of vital importance. One of the main strategies is the search for enzymes that degrade the different components of the extracellular matrix. However, since the composition of the matrix varies among the different microorganisms, an alternative would be to interfere with the signaling cascades that lead to the formation of the biofilm or stimulate its dispersion.

The production of the second intracellular di-cGMP messenger by diguanylate cyclases (DGCs) is a widely conserved process and has a central role in the transition between the planktonic life stage and the biofilm in both Gram positive and Gram negative bacteria, so that high levels promote biofilm formation while low levels induce biofilm dispersal. Molecules with the capacity to inhibit DGC activity have been identified by screening collections of chemical compounds that are capable of inhibiting biofilm formation, although no metagenomic search has been carried out to date.

The strategy of our project will be the construction of a *P. putida* indicator strain that will help us to recognize inhibitors of the DGC activity produced by clones from previously constructed environmental meta-libraries. To this end, our strain contains in the chromosome the gene that encodes PleD DGC under the control of the *Psal* promoter, which is induced by the presence of salicylate thanks to the NahR regulator and a *lacZ* transcriptional fusion of the *PparC* promoter of *P. putida*, which is repressed in the presence of di-cGMP.

In the presence of X-gal, a lawn of the indicator strain contained in top agar will be white, while in the case that any of the clones of the meta-library produces a compound capable of inhibiting DGC or interfering with di-cGMP signaling, a blue halo will appear around the colony. To test the operation of the screening system, once the indicator strain was constructed, the ability to form blue haloes was tested in the presence of sulfathiazole, a compound that has demonstrated its ability to inhibit di-cGMP synthesis.

### REFERENCES

Koo, H., Allan, R. N., Howlin, R. P., Stoodley, P., & Hall-Stoodley, L. (2017). Targeting microbial biofilms: current and prospective therapeutic strategies. *NATURE REVIEWS MICROBIOLOGY*, 15(12), 740.

Sambanthamoorthy, K., Luo, C., Pattabiraman, N., Feng, X., Koestler, B., Waters, C. M., & Palys, T. J. (2014). Identification of small molecules inhibiting diguanylate cyclases to control bacterial biofilm development. *BIOFOULING*, 30(1), 17-28.

Antoniani, D., Bocci, P., Maciąg, A., Raffaelli, N., & Landini, P. (2010). Monitoring of diguanylate cyclase activity and of cyclic-di-GMP biosynthesis by whole-cell assays suitable for high-throughput screening of biofilm inhibitors. *Applied microbiology and biotechnology*, 85(4), 1095-1104.