

Ratio Mathematica, 19, pp. 1-10

On properties of fuzzy subspaces of vector spaces

H. Hedayati

Department of Mathematics, Faculty of Basic Science,

Babol University of Technology, Babol, Iran

E-mail: h.hedayati@nit.ac.ir, hedayati143@yahoo.com

Abstract

In this paper, we introduce the notion of normal fuzzy subspace of vector spaces. By using it, we construct new fuzzy subspaces. We also show that, under certain conditions, a fuzzy subspace of a vector space is two-valued and takes 0 and 1.

Mathematics Subject Classification: 08A72

Keywords: vector space, fuzzy subspace, normal fuzzy subspace

1 Introduction

Zadeh in [17] introduced the notion of fuzzy set and started a generalized logic. After that reconsideration of mathematics concepts begun. Also there have been a number of generalizations of this fundamental concept. Fuzzy algebraic

structures play a prominent role in mathematics with wide applications in many other branches such as theoretical physics, computer sciences, control engineering, information sciences, coding theory, topological spaces, logic, set theory, group theory, groupoids, real analysis, measure theory etc (see [4], [5], [10] and [12]). In 1977, Katsaras and Liu [7] formulated and studied the concept of a fuzzy subspace of a vector space. Since then, a host of mathematicians are involved in extending the basic concepts and results from the theory of crisp vector spaces to the broader framework of the fuzzy setting. However, not all the results can be fuzzified. In [8], among other concepts and results, the fuzzy coset of a fuzzy subspace is defined and the algebraic nature of fuzzy subspaces under homomorphism is studied. In [9], the fuzzy basis and dimension of a fuzzy subspace are defined and studied. In [1] and [3], the fuzzy subspaces over fuzzy fields are discussed.

In this paper, some properties of fuzzy subspaces of vector spaces are investigated. Specially, some ways are created to construct new fuzzy subspaces from the old. Also the notion of normal fuzzy subspace of vector spaces is introduced. We see some normal fuzzy subspaces can be constructed by a fuzzy subspace. Finally, we show that, when a non-constant normal fuzzy subspace be a maximal in the partial ordered set of normal fuzzy subspaces of a vector space, then this fuzzy subspace is two-valued and takes the values 0 and 1.

2 Preliminaries

An abelian group $(\mathcal{V}, +)$ on a field F is called a *vector space* on F if there exists a map $\cdot : \mathcal{V} \times F \longrightarrow \mathcal{V}$ such that for all $x, y \in \mathcal{V}$ and $a, b \in F$ the following

conditions hold:

- (i) $1x = x$,
- (ii) $(ab)x = a(bx)$,
- (iii) $a(x + y) = ax + ay$,
- (v) $(a + b)x = ax + by$.

Also a non-empty subset \mathcal{W} of a vector space \mathcal{V} is called a *subspace*, if \mathcal{W} is a vector space on F .

Let X be an ordinary set. By a *fuzzy set* μ in X , we mean a function $\mu : X \longrightarrow [0, 1]$ with the grade of membership $\mu(x)$ for $x \in X$. If $t \in [0, 1)$, then $\mu_t = \{x \in X \mid \mu(x) \geq t\}$ is called a *level subset* of μ .

3 Fuzzy normal subspaces

In what follows, \mathcal{V} is a vector space on a field F , unless otherwise specified.

Definition 3.1. ([7], [11]) A fuzzy set μ of \mathcal{V} is called a *fuzzy subspace* of \mathcal{V} , if for all $x, y \in \mathcal{V}$ and $a \in F$ the following conditions hold:

- (i) $\mu(x + y) \geq \mu(x) \wedge \mu(y)$,
- (ii) $\mu(-x) \geq \mu(x)$,
- (iii) $\mu(ax) \geq \mu(x)$.

Clearly, if μ is a fuzzy subspace of \mathcal{V} , then $\mu(0) \geq \mu(x)$ for all $x \in \mathcal{V}$. Also, μ is a fuzzy subspace of \mathcal{V} if and only if μ_t is a subspace of \mathcal{V} for all $t \in [0, 1)$.

Lemma 3.2. *If μ is a fuzzy subspace of \mathcal{V} , then the set $\mathcal{V}_\mu = \{x \in \mathcal{V} \mid \mu(x) = \mu(0)\}$ is a subspace of \mathcal{V} .*

Proof. Let $x, y \in \mathcal{V}_\mu$. Then $\mu(x) = \mu(y) = \mu(0)$. Since μ is a fuzzy

subspace, it follows that

$$\mu(x - y) \geq \mu(x) \wedge \mu(y) = \mu(0) \wedge \mu(0) = \mu(0).$$

On the other hand $\mu(x - y) \leq \mu(0)$. Hence we have $\mu(x - y) = \mu(0)$ and so $x - y \in \mathcal{V}_\mu$. Also for any $x \in \mathcal{V}_\mu$ and $a \in F$, we get $\mu(ax) \geq \mu(x) = \mu(0)$. On the other hand $\mu(ax) \leq \mu(0)$. Hence, we obtain $\mu(ax) = \mu(0)$, which shows that $ax \in \mathcal{V}_\mu$. Consequently, the set \mathcal{V}_μ is a subspace of \mathcal{V} . \square

Definition 3.3. A fuzzy subspace of \mathcal{V} is said to be *normal* if there exists $x \in \mathcal{V}$ such that $\mu(x) = 1$. Note that if a fuzzy subspace of \mathcal{V} is normal, then $\mu(0) = 1$. Hence μ is a normal fuzzy subspace if and only if $\mu(0) = 1$.

Theorem 3.4. Let μ be a fuzzy subspace of \mathcal{V} and let $\tilde{\mu}$ be a fuzzy set in \mathcal{V} defined by $\tilde{\mu}(x) = \mu(x) + 1 - \mu(0)$ for all $x \in \mathcal{V}$. Then $\tilde{\mu}$ is a normal fuzzy subspace of \mathcal{V} containing μ .

Proof. Let $x, y \in \mathcal{V}$ and $a \in F$. Then

$$\tilde{\mu}(x - y) = \mu(x - y) + 1 - \mu(0) \geq (\mu(x) \wedge \mu(y)) + 1 - \mu(0) =$$

$$(\mu(x) + 1 - \mu(0)) \wedge (\mu(y) + 1 - \mu(0)) = \tilde{\mu}(x) \wedge \tilde{\mu}(y).$$

Also we have

$$\tilde{\mu}(ax) = \mu(ax) + 1 - \mu(0) \geq \mu(x) + 1 - \mu(0) = \tilde{\mu}(x).$$

Clearly, $\tilde{\mu}(0) = 1$ and $\mu \subseteq \tilde{\mu}$. This completes the proof. \square

Corollary 3.5. If μ is a fuzzy subspace of \mathcal{V} satisfying $\tilde{\mu}(x) = 0$ for some $x \in \mathcal{V}$, then $\mu(x) = 0$.

Lemma 3.6. *Let $\chi_{\mathcal{W}}$ be the characteristic function of a subset $\mathcal{W} \subseteq \mathcal{V}$. Then \mathcal{W} is a subspace of \mathcal{V} if and only if $\chi_{\mathcal{W}}$ is a fuzzy subspace of \mathcal{V} .*

Proof. It is directly followed from discussion after Definition 3.1. \square

Theorem 3.7. *For any subspace \mathcal{W} of \mathcal{V} , the characteristic function $\chi_{\mathcal{W}}$ is a normal fuzzy subspace of \mathcal{V} and $\mathcal{V}_{\chi_{\mathcal{W}}} = \mathcal{W}$.*

Proof. Straightforward. \square

Theorem 3.8. *A fuzzy subspace μ of \mathcal{V} is normal if and only if $\tilde{\mu} = \mu$.*

Proof. If $\tilde{\mu} = \mu$, then it is obvious that μ is a normal fuzzy subspace of \mathcal{V} . Assume that μ is a normal fuzzy subspace of \mathcal{V} and let $x \in \mathcal{V}$. Then $\tilde{\mu}(x) = \mu(x) + 1 - \mu(0) = \mu(x)$, and hence $\tilde{\mu} = \mu$. \square

Theorem 3.9. *If μ is a fuzzy subspace of \mathcal{V} , then $(\tilde{\tilde{\mu}}) = \tilde{\mu}$.*

Proof. Straightforward. \square

Theorem 3.10. *Let μ be a fuzzy subspace of \mathcal{V} . If there exists a fuzzy subspace ν of \mathcal{V} satisfying $\tilde{\nu} \subseteq \mu$, then μ is a normal fuzzy subspace of \mathcal{V} .*

Proof. Suppose there exists a fuzzy subspace ν of \mathcal{V} such that $\tilde{\nu} \subseteq \mu$. Then $1 = \tilde{\nu}(0) \leq \mu(0)$, and therefore $\mu(0) = 1$. \square

Corollary 3.11. *Let μ be a fuzzy subspace of \mathcal{V} . If there exists a fuzzy subspace ν of \mathcal{V} satisfying $\tilde{\nu} \subseteq \mu$, then $\tilde{\mu} = \mu$.*

Proof. It is immediately obtained from Theorem 3.10 and definition of $\tilde{\mu}$. \square

Theorem 3.12. *Let μ be a fuzzy subspace of \mathcal{V} and $f : [0, \mu(0)] \longrightarrow [0, 1]$*

be an increasing map. Define a fuzzy set $\mu^f : \mathcal{V} \longrightarrow [0, 1]$ by $\mu^f(x) = f(\mu(x))$ for all $x \in \mathcal{V}$. Then μ^f is a fuzzy subspace of \mathcal{V} . In particular, if $f(t) \geq t$ for all $t \in [0, \mu(0)]$ then $\mu \subseteq \mu^f$.

Proof. Let $x, y \in \mathcal{V}$. Then

$$\mu^f(x - y) = f(\mu(x - y)) \geq f(\mu(x) \wedge \mu(y)) =$$

$$f(\mu(x)) \wedge f(\mu(y)) = \mu^f(x) \wedge \mu^f(y).$$

Also if $a \in F$ and $x \in \mathcal{V}$, then $\mu^f(ax) = f(\mu(ax)) \geq f(\mu(x)) = \mu^f(x)$. Hence μ^f is a fuzzy subspace of \mathcal{V} . Assume that $f(t) \geq t$ for all $t \in [0, \mu(0)]$. Then $\mu^f(x) = f(\mu(x)) \geq \mu(x)$ for all $x \in \mathcal{V}$, which means $\mu \subseteq \mu^f$. \square

Theorem 3.13. *Let μ be a non-constant normal fuzzy subspace of \mathcal{V} , which is maximal in the partial ordered set of normal fuzzy subspaces of \mathcal{V} under fuzzy sets inclusion. Then μ is a two-valued fuzzy subspace and takes the values 0 and 1.*

Proof. We know $\mu(0) = 1$. Let $x \in \mathcal{V}$ be such that $\mu(x) \neq 1$. It is enough to show that $\mu(x) = 0$. Assume that there exists $x' \in \mathcal{V}$ such that $0 < \mu(x') < 1$. Define a fuzzy set $\nu : \mathcal{V} \longrightarrow [0, 1]$ by $\nu(x) = 1/2(\mu(x) + \mu(x'))$ for all $x \in \mathcal{V}$. Then clearly ν is well-defined. Let $x, y \in \mathcal{V}$. Then

$$\nu(x - y) = 1/2(\mu(x - y) + \mu(x')) \geq 1/2((\mu(x) \wedge \mu(y)) + \mu(x')) =$$

$$(1/2(\mu(x) + \mu(x'))) \wedge (1/2(\mu(y) + \mu(x'))) = \nu(x) \wedge \nu(y).$$

Also if $a \in F$ and $x \in \mathcal{V}$, then

$$\nu(ax) = 1/2(\mu(ax) + \mu(x')) \geq 1/2(\mu(x) + \mu(x')) = \nu(x).$$

Hence ν is a fuzzy subspace of \mathcal{V} . Now we have

$$\tilde{\nu}(x) = \nu(x) + 1 - \nu(0) =$$

$$1/2(\mu(x) + \mu(x')) + 1 - 1/2(\mu(0) + \mu(x')) = 1/2(\mu(x) + 1).$$

So $\tilde{\nu}(0) = 1/2(\mu(0) + 1) = 1$. Thus $\tilde{\nu}$ is a normal fuzzy subspace of \mathcal{V} . Also $\tilde{\nu}(0) = 1 > \tilde{\nu}(x') = 1/2(\mu(x') + 1) > \mu(x')$. We know that $\tilde{\nu}$ is non-constant. So by $\tilde{\nu}(x') > \mu(x')$, it follows that μ is not maximal, which is a contradiction. Therefore μ takes only the values 0 and 1. \square

Theorem 3.14. *Let μ be a fuzzy subspace of \mathcal{V} and let $\bar{\mu}$ be a fuzzy set in \mathcal{V} defined by $\bar{\mu}(x) = \mu(x)/\mu(0)$ for all $x \in \mathcal{V}$. Then $\bar{\mu}$ is a normal fuzzy subspace of \mathcal{V} containing μ .*

Proof. For any $x, y \in \mathcal{V}$, we have

$$\mu(x - y) = \mu(x - y)/\mu(0) \geq (1/\mu(0))(\mu(x) \wedge \mu(y)) =$$

$$(\mu(x)/\mu(0)) \wedge (\mu(y)/\mu(0)) = \bar{\mu}(x) \wedge \bar{\mu}(y).$$

Also if $a \in F$ and $x \in V$ we get

$$\bar{\mu}(ax) = (\mu(ax)/\mu(0)) \geq (\mu(x)/\mu(0)) = \bar{\mu}(x).$$

Hence $\bar{\mu}$ is a fuzzy subspace of \mathcal{V} . Clearly $\bar{\mu}(0) = 1$ and $\mu \subseteq \bar{\mu}$. \square

Corollary 3.15. *If μ is a fuzzy subspace of \mathcal{V} satisfying $\bar{\mu}(x) = 0$ for some $x \in \mathcal{V}$, then $\mu(x) = 0$.*

Proof. Obvious. \square

Theorem 3.16. *Let μ be a non-constant fuzzy subspace of \mathcal{V} such that $\tilde{\mu}$ is a maximal in the partial ordered set of normal fuzzy subspace of \mathcal{V} under fuzzy sets inclusion. Then*

- (1) μ is normal.
- (2) μ takes only the values 0 and 1.
- (3) $\chi_{\mathcal{V}_\mu} = \mu$.
- (4) \mathcal{V}_μ is a maximal subspace of \mathcal{V} .

Proof. Since μ is non-constant, so $\tilde{\mu}$ is non-constant maximal. Also $\tilde{\mu}$ is normal, which implies $\tilde{\mu}$ takes only values 0 and 1 by Theorem 3.13. . If $\tilde{\mu}(x) = 1$, then $\mu(x) = \mu(0)$ and if $\tilde{\mu}(x) = 0$, then $\mu(x) = \mu(0) - 1$. By Corollary 3.5, we have $\mu(x) = 0$ which implies $\mu(0) = 1$. Therefore μ is normal, and also $\tilde{\mu} = \mu$ by Theorem 3.8, which proves (1) and (2).

(3) Clearly $\chi_{\mathcal{V}_\mu} \subseteq \mu$ and $\chi_{\mathcal{V}_\mu}$ takes only the values 0 and 1. Let $x \in \mathcal{V}$ and $\mu(x) = 0$, then $\mu \subseteq \chi_{\mathcal{V}_\mu}$. If $\mu(x) = 1$ then $x \in \mathcal{V}_\mu$ and so $\chi_{\mathcal{V}_\mu}(x) = 1$. In any case $\mu \subseteq \chi_{\mathcal{V}_\mu}$.

(4) Since μ is non-constant, \mathcal{V}_μ is a proper subspace of \mathcal{V} . Let \mathcal{W} be a subspace of \mathcal{V} such that $\mathcal{V}_\mu \subseteq \mathcal{W}$. Then we obtain $\mu = \chi_{\mathcal{V}_\mu} \subseteq \chi_{\mathcal{W}}$. Since μ and $\chi_{\mathcal{W}}$ are normal and $\mu = \tilde{\mu}$ is maximal in the partial ordered set of normal fuzzy subspaces under fuzzy sets inclusion, we have $\mu = \chi_{\mathcal{W}}$ or $\chi_{\mathcal{W}}(x) = 1$ for all $x \in \mathcal{V}$, so $\mathcal{W} = \mathcal{V}$. If $\mu = \chi_{\mathcal{W}}$ then $\mathcal{V}_\mu = \mathcal{V}_{\chi_{\mathcal{W}}} = \mathcal{W}$ by Theorem 3.7 . Therefore \mathcal{V}_μ is a maximal subspace of \mathcal{V} . \square

References

- [1] **M. T. Abu osman**, " *On t -fuzzy subfield and t -fuzzy vector subspace*", Fuzzy Sets Syst 33 (1989), 111-117.
- [2] **M. T. Abu osman**, " *On fuzzy vector spaces via triangular norms*", Bull. Malaysian Math. Soc. 9(1) (1986), 33-42.
- [3] **R. Biswas**, " *Fuzzy fields and fuzzy linear spaces redefiend*", Fuzzy Sets Syst 33 (1989), 257-259.
- [4] **C. L. Chang**, " *Fuzzy topological spaces*", J. Math. Anal. Appl. 24 (1968), 182-190.
- [5] **P. Das**, " *Fuzzy vector spaces under triangular norms*", Fuzzy Sets Syst 25 (1988), 73-85.
- [6] **A. K. Katsaras**, " *Linear fuzzy neighborhood spaces*", Fuzzy Sets Syst 16 (1985), 25-40.
- [7] **A. K. Katsaras, D. B. Liu**, " *Fuzzy vector spaces and fuzzy topological spaces*", J. Math. Anal. Appl. 58 (1977), 135-146.
- [8] **R. Kumar**, " *Fuzzy vector spaces and fuzzy cosets*", Fuzzy Sets Syst 45 (1992), 109-116.
- [9] **R. Kumar**, " *On the dimension of a fuzzy subspace*", Fuzzy Sets Syst 54 (1993), 229-234.
- [10] **R. Lowen**, " *Convex fuzzy sets*", Fuzzy Sets Syst 3 (1980), 291-310.

- [11] **P. Lubczonok**, "Fuzzy vector spaces", Fuzzy Sets Syst 38 (1990), 329-343.
- [12] **P. P. Ming, Y. M. Ming**, "Fuzzy topology I, neighborhood structure of a fuzzy point and Moors-Smith convergence", J Math Anal 76 (1980), 571-599.
- [13] **S. Nanda**, "Fuzzy fields and fuzzy linear spaces", Fuzzy Sets Syst 19 (1986), 89-94.
- [14] **Y. Terao, N. Kitsunezaki**, "Fuzzy sets and linear mapping on vector spaces", Math. Japonica 39, No. 1(1994), 61-68.
- [15] **Y. Yandong**, "Finitely generated T -fuzzy linear spaces", Fuzzy Sets Syst 30 (1989), 69-81.
- [16] **Y. D. Yu**, "Fuzzy linear spaces redefined", Fuzzy Math. 4(3) (1984), 59-62.
- [17] **L. A. Zadeh**, "Fuzzy sets" Inform. Control, 8(1965), 338-353.