

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Determination of In Vitro Antiprotease, Antimicrobial, and Antibiofilm Activities of *Beta vulgaris* var. *cicla* against Multidrug-Resistant Strains of *Pseudomonas aeruginosa*

Hayet Edziri, Rim Nasri, Marwa Hamdi and Maha Mastouri

## Abstract

Antibiotic resistance of *Pseudomonas aeruginosa* causes many infectious diseases and it is great. So, the aim of the present work was to assess the antibacterial, antibiofilm activity of *Beta vulgaris* extracts against resistance bacteria *P. aeruginosa* that were clinically isolated and tested for their antiprotease potential. Result showed that methanol extract exhibited important antiprotease activity against Trypsin, Savinase, and digestive proteases of blue crab with percentage of inhibition of 94.66, 91.39, and 86.41%, respectively. It showed also important antibiofilm activities against multidrug-resistant *P. aeruginosa* with inhibition values upper than 80% with a concentration of 4MIC. Our investigation delivered that *Beta vulgaris* might be possible source of natural antienzymatic, antimicrobial, and antibiofilm agents.

**Keywords:** *Beta vulgaris*, antibacterial, antibiofilm, antiprotease, multidrug-resistant *P. aeruginosa*

## 1. Introduction

Many studies have demonstrated that vegetables play a significant role in human nutrition. They reduced risk of many chronic diseases, like diabetes, cardiovascular illnesses, and cancers [1–3] and Alzheimer's diseases [4, 5]. These advantageous properties of vegetables are due to the bioactive compositions known for their important antioxidant activities [6, 7].

Microbial contamination and the resistance of pathogenic bacteria to antibiotics are considered as major problems of public health [1, 8, 9].

*Pseudomonas aeruginosa* are bacteria that cause nosocomial infections. They are able to be resistant to a great number of antibiotics such as Carbapenems like imipenem [10, 11].

*Beta vulgaris* L. belongs to the *Amaranthaceae* family. Its juice had important biological properties like as antimicrobial, hemostatic, and anticancer [12, 13]. *Beta vulgaris* is categorized among the best vegetables with important antioxidant activity; in addition, many researches have showed that *Beta vulgaris* extracts had other important activities (anti-inflammatory, antiallergenic, antithrombotic, and anticoagulant) [14, 15].

The objective of this chapter was to investigate the antiprotease, antimicrobial, and antibiofilm activities of *Beta vulgaris var. cicla* against multidrug-resistant strains of *Pseudomonas aeruginosa*.

## 2. Materials and methods

### 2.1 Plant material

The fresh beetroots were bought from a market in Sousse (Tunisia) and the roots were identified and a voucher specimen was placed in our laboratory at the Faculty of Pharmacy (Monastir).

### 2.2 Preparation of aqueous extract of *Beta vulgaris*

About 200 ml of distilled water was added to 50 g of *Beta vulgaris* pieces. Then they were allowed to boil for 30 min. The extract was filtered using a Whatman paper. The filtrate was kept at  $-25^{\circ}\text{C}$ .

### 2.3 Preparation of methanolic extract of *Beta vulgaris*

Beetroots were washed and sliced into small pieces and then 200 ml of methanol was added to 100-g root in brown bottle for 3 days at room temperature, filtered through Whatman filter paper, and dried with rotavapor. Then the extract was kept at  $4^{\circ}\text{C}$ .

### 2.4 Total polyphenol content

The total phenolic content was tested by Folin-Ciocalteu method (Edziri et al.) [16]. The total polyphenols content is expressed as mg gallic acid equivalents (GAEs) per g of extract.

### 2.5 Total flavonoid content

The flavonoids content was tested by the method of Othmana et al. [17]. The result is expressed in mg quercetin equivalents (QEs) per g of extract.

### 2.6 Total tannin contents

Total tannin content in *Beta vulgaris* extracts was tested by using Folin-Denis reagent [18].

### 2.7 Total carotenoid content

Total carotenoids content of *Beta vulgaris* was determined by the ARNON method [19].

## 2.8 Antiprotease activity

The impact of *Beta vulgaris* extracts, at a concentration of 250 µg/ml, on several proteases' activity was evaluated. So, enzymes were pre-incubated with each extract for 30 min at 30°C. Then, the residual enzyme activity was evaluated according to the method of Georgé et al. [20] using casein as a substrate at the optimal pH and temperature for each enzyme: Purafect (pH 10.0; 50°C), Savinase (pH 10.0; 60°C), (pH 8.5; 50°C) and enzyme of blue crab (pH 8.0; 60°C), trypsin and chymotrypsin (pH 8.0; 37°C). The activity of the enzyme assayed in the absence of inhibitors was taken as 100%.

## 2.9 Antibacterial activity of *Beta vulgaris* extracts

### 2.9.1 Microdilution assay

Minimum inhibitory concentration (MIC) values were determined by a microdilution method as indicated by Edziri et al. [21]. The MIC was defined as the lowest concentration that inhibits the development of bacteria, after 24 h of incubation [22].

### 2.9.2 Antimicrobial activity

#### 2.9.2.1 Micro-well determination of MIC and MBC

The minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) were determined for all bacteria tested in this work by a microdilution method as described by [23]. The MIC was distinct as the lowest concentration of the extract to inhibit the development of the bacteria.

### 2.9.3 Inhibition of biofilm formation

The biofilm inhibition was tested against five multidrug-resistant strains of *Pseudomonas aeruginosa*, by microdilution technique according to Edziri et al. [21].

## 3. Results and discussion

### 3.1 Phytochemical screening

The proportions of the phenols, flavonoids, tannins, and carotenoids contents are summarized in **Table 1**. Methanolic extract of *Beta vulgaris* was the richest with phenols, flavonoids, tannins, and carotenoids as shown in **Table 1**.

### 3.2 Antiprotease, antibacterial and antibiofilm activities

The antiprotease activity of various *Beta vulgaris* extracts was tested at a concentration of 250 µg/ml. **Table 2** demonstrates that all extracts were able to decrease the protease activities by about 51.53–94.66%, suggesting that they were rich sources of the protease inhibitors. In fact, results revealed that the preincubation of Trypsin, Savinase, and digestive proteases of blue crab with methanol extract caused the loss of 94.66, 91.39, and 86.41% of their activity, respectively; however, a decrease of about 89.31 and 91.39% was observed after incubation of this extract with Purafect and chymotrypsin, respectively.

It is interesting to note that the proteolytic activities of Savinase® and Purafect®, commercial microbial proteases, were mostly inhibited by methanol extract by about 91.39 and 89.31%, respectively. In addition, it was efficient to reduce 94.66% of digestive trypsin activity. In addition aqueous extracts exhibit good antiprotease activity.

According to **Table 3**, the values of MIC for two extracts against multidrug-resistant *Pseudomonas aeruginosa* varied between 50 and 100 mg/ml, without difference between the two extracts of *Beta vulgaris*. Furthermore, the MBC values were of 250 mg/ml. The observed activity of *Beta vulgaris* roots may be attributed to the higher content of polyphenols, flavonoids, and tannin, which are known for their important antimicrobial activity.

The two extracts showed important antibiofilm activity (**Figures 1 and 2**). Furthermore, methanolic extract exhibited the greatest antibiofilm property against all resistant strains of PA with inhibition values greater than 80% at the concentration of 4MIC. In addition, aqueous extract inhibited the biofilm formation of PA greater than 50% with 2MIC. We can observe that there is not any difference between the tested strains of PA. Also methanol extract of *Beta*

	Aq	M
Total polyphenols (mg GAE/ g) extract	99.47 ± 0.45	134.55 ± 0.6
Total flavonoids (mg EQ/g) extract	1.29 ± 0.50	4.34 ± 0.02
Total tannin content(mg TA/g)	6.15 ± 1.4	7.5 ± 0.5
Carotenoids (mg/100 g FW)	2.1 ± 1.2	2.97 ± 0.4

GAE: gallic acid equivalent, CE: catchin equivalent, TA: tannic acid, Aq: aqueous extract, M: methanol extract.

**Table 1.**  
Phytochemical analysis of *Beta vulgaris* var. *cicla*.

Enzymes	M	Aq
Purafect	89.31 ± 1.13	60.70 ± 0.96
Savinase	91.39 ± 1.38	81.07 ± 0.43
Chymotrypsin	65.28 ± 0.47	51.53 ± 0.51
Trypsin	94.66 ± 0.32	87.42 ± 1.55
Digestive proteases of blue crab	86,41 ± 0.34	70.11 ± 0.61


Values are mean ± SD of three replicate analyses, Aq: aqueous extract, M: methanol extract.

**Table 2.**  
Antiprotease activity of *Beta vulgaris* L.


Strains extracts	P.S1		P.S2		P.S3		P.S4		P.S5	
	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC
Aq	100	250	100	250	100	250	100	250	100	250
M	50	250	100	250	100	250	100	250	100	250

Aq: aqueous extract, M: methanol extract, MIC and MBC are in mg/ml.

**Table 3.**  
Antipseudomonal activity of *Beta vulgaris* extracts.



**Figure 1.**  
 Antibiofilm activity of *Beta vulgaris* aqueous extracts.



**Figure 2.**  
 Antibiofilm activity of *Beta vulgaris* methanolic extracts.

*vulgaris* displayed a respectable antibiofilm property beside P.S5 with percentages of inhibition of 88.9% at a concentration of 4MIC. The present search was investigated for the first time on the antibiofilm capacity of *Bea vulgaris* against multidrug-resistant strains of *Pseudomonas aeruginosa*. On the other hand, the antibiofilm action is mostly due to the great amount of phenolic content, such as that of flavonoids and tannin known for their good biological activities [24, 25].

#### 4. Conclusion

From this study, we can see that *Beta vulgaris* showed good antiprotease, antibacterial, and antibiofilm activities against different resistant *Pseudomonas aeruginosa* strains. This study also showed that the utilization of this vegetable can lead to the inhibition of bacterial growth.

Furthermore, this vegetable can be used as a source of natural antienzymatic, antimicrobial, and antibiofilm agents. Research is in progress to identify and isolate the bioactive molecules and to test them in vivo.

IntechOpen

## Author details

Hayet Edziri<sup>1\*</sup>, Rim Nasri<sup>2</sup>, Marwa Hamdi<sup>2</sup> and Maha Mastouri<sup>1</sup>

1 Laboratoire des maladies transmissibles et des substances biologiquement actives, Faculté de Pharmacie, Monastir, Tunisie

2 Laboratory of Enzyme Engineering and Microbiology, National Engineering School of Sfax, University of Sfax, Sfax, Tunisia

\*Address all correspondence to: jaziri\_hayet@yahoo.fr

## IntechOpen

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

## References

- [1] Wruss J, Waldenberger G, Huemer S, Uygun P, Lanzerstorfer P, Müller U, et al. Compositional characteristics of commercial beetroot products and beetroot juice prepared from seven beetroot varieties grown in Upper Austria. *Journal of Food Composition and Analysis*. 2015;**42**:46-55
- [2] Guldiken B, Toydemir G, Nur Memis K, Okur S, Boyacioglu D, Capanoglu E. Home-processed red beetroot (*Beta vulgaris* L.) products: Changes in antioxidant properties and bioaccessibility. *International Journal of Molecular Sciences*. 2016;**17**:858
- [3] Biondo PBF, Boeing JS, Barizo RO, Souza NED, Matsushita M, Oliveira CCD, et al. Evaluation of beetroot (*Beta vulgaris* L.) leaves during its developmental stages: A chemical composition study. *Food Science and Technology (Campinas)*. 2014;**34**:94-101
- [4] Pai SR, D'Mello P. Stability evaluation of beetroot colour in various pharmaceutical matrices. *Indian Journal of Pharmaceutical Sciences*. 2004;**66**:696-699
- [5] Esatbeyoglu T, Wagner AE, Schini-Kerth VB, Rimbach G. Betanin—A food colorant with biological activity. *Molecular Nutrition & Food Research*. 2015;**59**:36-47
- [6] Sainath M, Kumar KS, Babu KA. Formulation and evaluation of herbal lipstick. *IJARMP*. 2016;**3**:2058-2060
- [7] Vinson JA, Hao Y, Su X, Zubik L. Phenol antioxidant quantity and quality in foods: Vegetables. *Journal of Agricultural and Food Chemistry*. 1998;**46**:3630-3634
- [8] Kähkönen MP, Hopia AI, Vuorela HJ, Rauha JP, Pihlaja K, Kujala TS. Antioxidant activity of plant extracts containing phenolic compounds. *Journal of Agricultural and Food Chemistry*. 1999;**47**:3954-3962
- [9] Vulić J, Čanadanović-Brunet J, Četković G, Tumbas V, Djilas S, Četojević-Simin D, et al. Antioxidant and cell growth activities of beet root pomace extracts. *Journal of Functional Foods*. 2012;**4**:670-678
- [10] Ehtesham-Gharaee M, Eshaghi A, Shojaee S, Asili J, Emami SA, Behravan J, et al. Protective effects of *Scutellaria lindbergii* root extract against oxidative-induced cell and DNA damage in mouse fibroblast-like cells. *Drug and Chemical Toxicology*. 2015;**38**:293-299
- [11] Edzir H, Ammar S, Souad L, Mahjoub MA, Mastouri M, Aouni M, et al. *In vitro* evaluation of antimicrobial, antioxidant activities of some Tunisian vegetables. *South African Journal of Botany*. 2012;**78**:252-256
- [12] El-Sabour MF, Abd E-S, Abo MA, Rizk M. Physiological and chemical responses of sunflower to the application of previous organic waste composts to sandy soil. *Environmental Management and Health*. 1997;**8**:128-132
- [13] Ormsbee MJ, Lox J, Arciero PJ. Beetroot juice and exercise performance. *Nutrition and Dietary Supplements*. 2013;**5**:27-35
- [14] Pandey KB, Rizvi SI. Plant polyphenols as dietary antioxidants in human health and disease. *Oxidative Medicine and Cellular Longevity*. 2009;**2**:270-278
- [15] Edziri H, Jaziri R, Chehab H, Verschaeve L, Flamini G, Boujnah D, et al. A comparative study on chemical composition, antibiofilm and biological activities of leaves extracts of four



Tunisian olive cultivars. *Heliyon*. 2019;**55**:E01604

[16] Saeed N, Khan M, Shabbir M. Antioxidant activity, total phenolic and total flavonoid contents of whole plant extracts *Torilis leptophylla* L. *BMC Complementary and Alternative Medicine*. 2012;**12**:221

[17] Othmana A, Ismaila A, Ghania NA, Adenan I. Antioxidant capacity and phenolic content of cocoa beans. *Food Chemistry*. 2007;**100**:1523-1530

[18] Edziri H, Mastouri M, Mahjoub A, Anthonissen R, Mertens B, Cammaerts S, et al. Toxic and mutagenic properties of extracts from Tunisian traditional medicinal plants investigated by the neutral red uptake, VITOTOX and alkaline comet assays. *South African Journal of Botany*. 2011;**77**:703-710

[19] Jeyanthi RL, Sharmila S, Das MP, Seshiah C. Extraction and purification of carotenoids from vegetables. *Journal of Chemical and Pharmaceutical Research*. 2014;**6**(4):594-598

[20] Georgé S, Brat P, Alter P, Amiot MJ. Rapid determination of polyphenols and vitamin C in plant-derived products. *Journal of Agricultural and Food Chemistry*. 2005;**53**:1370-1373

[21] Edziri H, Hechmi A, Chehab H, Luc V, Guido F, Dalenda B, et al. A comparative study on chemical composition, antibiofilm and biological activities of leaves extracts of four Tunisian olive cultivar. *Heliyon*. 2019;**5**:e01604

[22] Uttara B, Singh AV, Zamboni P, Mahajan RT. Oxidative stress and neurodegenerative diseases: A review of upstream and downstream antioxidant therapeutic options. *Current Neuropharmacology*. 2009;**7**:65-74

[23] Edziri H, Maha M, Samia A, Mata M, Gros P, Raida H, et al. Antimicrobial, antioxidant, and

antiviral activities of *Retama raetam* (Forssk) Webb flowers growing in Tunisia. *World Journal of Microbiology and Biotechnology*. 2008;**24**:2933-2940

[24] Clark G, Ting KN, Wiart C, Fry J. High correlation of 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging, ferric reducing activity potential and total phenolics content indicates redundancy in use of all three assays to screen for antioxidant activity of extracts of plants from the Malaysian rainforest. *Antioxidants*. 2013;**2**:1-10

[25] Dutta-Roy AK, Crosbie L, Gordon MJ. Effect of tomato extract on human platelet aggregation in vitro. *Platelets*. 2001;**12**:218-227