

Journal of Cataract & Refractive Surgery

Visual Function and Subjective Perception of Vision following bilateral implantation of monofocal and multifocal intraocular lenses: A Randomised Controlled Trial. --Manuscript Draft--

Manuscript Number:	JCRS-19-1154R2
Article Type:	Full Length Article
Section/Category:	Cataract
Keywords:	multifocal; Intraocular lens
Corresponding Author:	Phillip J Buckhurst, Ph.D. Plymouth, UNITED KINGDOM
First Author:	Elizabeth M. Law, MSc
Order of Authors:	Elizabeth M. Law, MSc Rajesh K. Aggarwal, BM, FRCOphth Hetal Buckhurst, PhD Hosam E. Kasaby, MBChB, FRCOphth Jonathan Marsden, PhD Gary Shum, PhD Phillip J Buckhurst, Ph.D.
Abstract:	<p>PURPOSE: Following implantation with Multifocal intraocular lenses (MIOLs) or monofocal intraocular lenses (IOLs), the study examines monocular and binocular visual function and patient reporting outcomes using a rigorous series of clinical assessments. Setting: BMI Southend Hospital, UK</p> <p>DESIGN: Prospective, randomised, double-masked clinical trial.</p> <p>METHODS: 100 subjects were randomised for bilateral implantation of either Bi-Flex 677MY MIOL or Bi-Flex 677AB IOL and were assessed at 3-6 months (V1) and 12-18 months (V2). Primary outcomes included distance, intermediate and near LogMAR visual acuities (VA) and defocus curve profile assessment. Secondary outcomes included reading speed, contrast sensitivity (CS) and the subjective perception of quality-of-vision.</p> <p>RESULTS: Uncorrected (MIOL 0.10 ± 0.09LogMAR; IOL 0.09 ± 0.11LogMAR) and best distance-corrected VA (MIOL 0.04 ± 0.06LogMAR; IOL 0.01 ± 0.07LogMAR) were comparable ($p > 0.05$). Unaided near VA (UNVA $p < 0.001$: MIOL 0.23 ± 0.13LogMAR; IOL 0.55 ± 0.20LogMAR) and distance-corrected near VA (DCNVA $p < 0.001$: MIOL 0.24 ± 0.13LogMAR; IOL 0.54 ± 0.17LogMAR) were significantly improved with MIOLs. There was no significant difference in distance-corrected intermediate VA (DCIVA $p = 0.431$: MIOL 0.38 ± 0.13; IOL 0.39 ± 0.13). Defocus curves demonstrated an increased range-of-focus amongst MIOLs (MIOL 4.14 ± 1.10D; IOL 2.57 ± 0.77D). Pelli-Robson CS was different at V1 ($p < 0.001$) but similar by V2 ($p = 0.059$). Overall satisfaction was high (>90%) in both groups for distance tasks whereas significantly different for near (MIOL 18.45 ± 16.53LogUnits; IOL 55.59 ± 22.52LogUnits).</p> <p>CONCLUSIONS : Unaided near visual acuity is demonstrably better with MIOLs and there was greater subjective satisfaction with their quality-of-near-vision. Halos reported by the MIOL group was significant compared to the IOL group, but did not show an adverse effect on overall satisfaction.</p>



UNIVERSITY OF
PLYMOUTH

Plymouth University
Room SF25
School of Health Professions
Plymouth
PL6 8BH
Tel: (+44) 1752 588884
Email:
phillip.buckhurst@plymouth.ac.uk
24th February 2020

Dear Prof Mamalis,

Thank you for reviewing of our manuscript. Please see below our responses to the comments highlighting our revisions that have been made to the manuscript

Reviewers' comments: Reviewer #1: The authors have responded to the various issues raised in my initial review and inserted comments accordingly in the article. Although this paper does not offer an original comparison (monofocal versus multifocal), it offers the advantage of a beautiful prospective methodology, which can be used as a reference in particular for the defence of multifocal optics before the authorities of the various ministries of health, eager for quality scientific evidence.

Thank you for your review

Editors comments: Would the space constraints be met if we made all of the tables (there are seven of them) online only but kept all of the figures? Yes, the tables could be made online only and the figures could all be kept as long as they are all one column.

Thanks for your answer to our query.

We have changed the figures in order for them to keep to a single column. To do this we have separated the Visit 1 and Visit 2 graphs and are only presenting the visit 2 data in the main manuscript. The Visit 1 graphs are now online only along with all the data within the tables (supplementary material).

Kind regards

Phill

Dr Phillip Buckhurst

Associate Professor | Associate Dean for Research | Associate Head of School (Research)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Visual Function and Subjective Perception of Vision following bilateral implantation of monofocal and multifocal intraocular lenses: A Randomised Controlled Trial.

Elizabeth M. Law, MSc,^{1,2} Rajesh K. Aggarwal, BM, FRCOphth,² Hetal Buckhurst, PhD,¹ Hosam E. Kasaby MBChB, FRCOphth,² Jonathan Marsden, PhD,¹ Gary Shum, PhD,¹ and Phillip J. Buckhurst, PhD.¹

¹University of Plymouth, School of Health Professions, Peninsula Allied Health Centre, Derriford Road, Plymouth, United Kingdom


²BMI Southend Hospital, Fairfax Drive, Westcliff on Sea, United Kingdom

Funding: The work was funded by Medicontur Medical Engineering (Zsámbék, Hungary). Medicontur had no role in the design or conduct of this research.

Financial Disclosure: No conflicting relationship exists for any author.

Running Head: Visual Function with MIOLs

Correspondence and reprint requests to Phillip J. Buckhurst, PhD, University of Plymouth, School of Health Professions, Peninsula Allied Health Centre, Derriford Road, Plymouth, PL6 8BH, United Kingdom. E-mail: Phillip.buckhurst@plymouth.ac.uk



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

ABSTRACT

PURPOSE:

Following implantation with Multifocal intraocular lenses (MIOLs) or monofocal intraocular lenses (IOLs), the study examines monocular and binocular visual function and patient reporting outcomes using a rigorous series of clinical assessments.


Setting: BMI Southend Hospital, UK

DESIGN: Prospective, randomised, double-masked clinical trial.

METHODS: 100 subjects were randomised for bilateral implantation of either Bi-Flex 677MY MIOL or Bi-Flex 677AB IOL and were assessed at 3-6 months (V1) and 12-18 months (V2). Primary outcomes included distance, intermediate and near LogMAR visual acuities (VA) and defocus curve profile assessment. Secondary outcomes included reading speed, contrast sensitivity (CS) and the subjective perception of quality-of-vision.

RESULTS: Uncorrected (MIOL 0.10 ± 0.09 LogMAR; IOL 0.09 ± 0.11 LogMAR) and best distance-corrected VA (MIOL 0.04 ± 0.06 LogMAR; IOL 0.01 ± 0.07 LogMAR) were comparable ($p > 0.05$). Unaided near VA (UNVA $p < 0.001$: MIOL 0.23 ± 0.13 LogMAR; IOL 0.55 ± 0.20 LogMAR) and distance-corrected near VA (DCNVA $p < 0.001$: MIOL 0.24 ± 0.13 LogMAR; IOL 0.54 ± 0.17 LogMAR) were significantly improved with MIOLs. There was no significant difference in distance-corrected intermediate VA (DCIVA $p = 0.431$: MIOL 0.38 ± 0.13 ; IOL 0.39 ± 0.13).

Defocus curves demonstrated an increased range-of-focus amongst MIOLs (MIOL 4.14 ± 1.10 D; IOL 2.57 ± 0.77 D). Pelli-Robson CS was different at V1 ($p < 0.001$) but similar by V2 ($p = 0.059$).



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65


Overall satisfaction was high (>90%) in both groups for distance tasks whereas significantly different for near (MIOL 18.45±16.53LogUnits; MIOL 55.59±22.52LogUnits).

CONCLUSIONS: Unaided near visual acuity is demonstrably better with MIOLs and there was greater subjective satisfaction with their quality-of-near-vision. Halos reported by the MIOL group was significant compared to the IOL group, but did not show an adverse effect on overall satisfaction.

Multifocal intraocular lenses (MIOLs) are widely considered the most reliable method of achieving spectacle independence following cataract surgery.¹⁻³ MIOLs distribute the light between distant and near focal points whereby the vergence of the incident light dictates which focal point is conjugate to the retinal plane.

The separation of these multiple focal points is determined by the addition power of the MIOL and to a lesser extent the biometry of the eye. High addition MIOLs (+4.00D or higher) are the zeitgeist of the designs used in the late 90s-early 2000s. Disadvantages of these early lenses included a close working distance and reduced intermediate vision. Moreover, the size of the dysphotopic phenomenon (commonly described as halo), associated with MIOLs, increases according to the addition power; these higher addition lenses generate larger haloes.^{4,5}

The light energy distribution between the retinal focal points created by a MIOL influences the overall quality of vision at different viewing distances. MIOLs that split light equally, create two focal points of comparative image quality. In contrast, distance dominant MIOLs allocate a higher



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65


percentage of light towards the distance retinal focal point and consequently near vision is relatively compromised. Conversely, the intensity of the halo is influenced by the light distribution relationship: the more distance dominant the lower the dysphotopic intensity.

In 2016, a Cochrane Review⁶ highlighted the need for robust randomised control trials examining the efficacy of MIOLs over monofocal intraocular lens (IOL) implantation and called for standardization of outcome measures in MIOL studies. The review concluded that it was unclear whether the achieved benefits of MIOL implantation i.e. greater near vision and increased spectacle independence, outweighed disadvantages such as reduced contrast sensitivity and increased dysphotopsia. Subsequently others have also highlighted the importance of patient reported outcomes in MIOLs.⁷ Despite these conclusions, in the subsequent three years there has only been a single RCT published comparing MIOLs with IOLs.⁸

The present study compared the efficacy of the Bi-Flex 677MY MIOL over its parent monofocal IOL using standardized methods for assessing both visual function and the subjective perception of the quality of vision.

METHODS

This study was a prospective, parallel double masked randomised clinical trial. The study protocol adheres to the Declaration of Helsinki and ethical approval was obtained prior to commencement of the trial. The study was registered with clinicaltrials.gov (NCT02338882) and written consent was obtained from all subjects. No modifications to the protocol or outcome measures were




1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

made during the study. The aim was to assess the IOLs using recognised methods that would provide rigour and establish a comprehensive method which could be utilised with all IOLs and allow easy comparison of results.

Patient Selection

Between September 2015 and May 2017, one hundred subjects were recruited from routine cataract clinics at the BMI Southend Hospital on a consecutive – if – eligible basis according to the inclusion/exclusion criteria ([Supplementary Table 1](#)). All subjects underwent initial examination by a consultant ophthalmic surgeon including dilated fundus examination; in the event of suspected macular pathology an OCT was carried out and if pathology was detected, the patient was excluded as per the study criterion. The anterior segments and ocular surface were also evaluated to confirm lack of pathology and minor ocular surface dryness was treated by commencement of ocular lubricants. Any ocular surface disease deemed moderate or marked resulted in exclusion. The allocation of IOLs was randomly designated and was masked to both the participant and the investigator conducting the post-operative study assessments. On enrolment, a study number was assigned to each subject. Using this study number, the allocation of lenses for all subjects was randomized in Microsoft Excel using blocked randomization with a 1:1 allocation ratio. Following allocation of the subject number, the unmasked surgeons and theatre staff accessed the randomization log and a series of sealed opaque envelopes that described which lenses were to be implanted (MIOL or IOL).

Surgical Technique



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65


All surgeries were performed by one of two experienced consultant ophthalmic surgeons (RA and HK) using small incision phacoemulsification. The same surgeon implanted both lenses for an individual subject. In each case, a 2.2mm clear corneal incision was located according to the steepest corneal meridian. The pre- and post-operative medication regime was the same regardless of surgeon. Second eye surgery occurred within 4 weeks of first eye surgery.

Masking

All post-operative study outcome measures were collected by a study investigator, who was masked to the allocation of study group. The subjects were also masked to their grouping allocation and were only informed of the type of lens implanted once they had completed the study. Post-operative slit lamp examination was performed by the unmasked consultant surgeon in order to maintain masking of the study investigator.

Intraocular Lenses

Each group had fifty subjects assigned. The Bi-Flex 677 AB is a single piece, aspheric aberration neutral IOL. The Bi-Flex MY MIOL has the same platform as the monofocal but the anterior surface has a 3mm apodized, diffractive central region with a near addition of 3.50D at the IOL plane ([Supplementary Table 2](#)). The Bi-Flex MY MIOL design is intended to provide distance dominance with greater mydriasis, thus maximizing contrast and minimizing halos when driving at night. Pupil miosis changes the light distribution relationship and results in a relatively equal split of light, hence, the Bi-Flex MY MIOL exploits the near miosis that occurs with reading. This type of MIOL was chosen for the study given that the unique aspect of the Bi-Flex MY is its low number of diffractive echelons (seven) which is theorized to improve the optical image quality of




1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

the resultant image. The identical platform and material of the two IOLs allowed unhindered assessment of the multifocality.

Primary Outcomes Measures

A masked investigator assessed the subjects at two study visits, 3-6 months (V1) and 12-18 months (V2) post-operatively. At each visit, monocular and binocular LogMAR acuities for unaided distance visual acuity (UDVA) and corrected distance visual acuity (CDVA) were measured using computerised test charts (Thomson Software Solutions Ltd) at 6m following the Bailey-Lovie principles and employing Sloan letters consistent with testing methods established by the Early Treatment Diabetic Retinopathy Study (ETDRS).⁹⁻¹³ Subjective refraction was conducted at 6m with a distance fixation target. The assessment of unaided near visual acuity (UNVA), distance corrected near visual acuity (DCNVA) and distance corrected intermediate visual acuity (DCIVA) utilised ETDRS charts for near (40cm) and intermediate (70cm) (Precision Vision) working distances respectively. To further assess intermediate and near vision at a range of distances, defocus profiles were plotted from -5.00D to 1.50D in 0.50D steps.¹⁴ The letters and defocus lenses were randomised between measures and subjects were prompted once using the phrase “can you read any more letters on the line below?”.¹⁵ All measures of visual acuity were performed with illuminance 120 cd/m² and luminance of 95 lux.

Secondary Outcome Measures




1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Contrast Sensitivity was assessed binocularly with the CSV-1000 (Precision Vision) calibrated to 2.4m and both monocularly and binocularly using Pelli-Robson charts at 6m (Thomson Software). Radner reading charts were used to assess reading speed at 40cm following the method outlined by Radner using a digital stopwatch.¹⁶ The subjective perception of vision was assessed using a quality of vision questionnaire¹⁷ and NAVQ.¹⁸ The Carl Zeiss Meditec Glare simulator was used to quantify the appearance of halos and glare. All secondary measures were assessed at V1 and V2. The same assessment room was used throughout the study and all secondary tests were carried out by the same masked investigator in photopic light conditions of illuminance 120cd/m² and luminance of 95 lux.

Statistical Analysis

The sample size for the study was calculated using G*power3 (University of Dusseldorf). Power calculations were based on a medium effect size ($f = 0.30$) based on *a-priori* matched paired *t test* design and a desired statistical power of 90% with an error probability of 0.05. Statistical analysis was performed using SPSS software, version 24 (IBM). All data were tested for normality using the Shapiro-Wilks test and visual examination of histogram plots. In all instances $p < 0.05$ was considered statistically significant. In order to evaluate effect size, Cohen's *d* was calculated, with $d > 0.2, 0.5$ and 0.8 corresponding to small, medium and large effect sizes, respectively.

A repeated measures ANOVA was used to establish similarity between right and left eye data for both monofocal and multifocal IOL data. No significant differences were found and as such only right eye data is presented.¹⁹ Where differences were found after repeated measures ANOVA, further pairwise tests were used to compare the monofocal and multifocal groups for all visual



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

acuity and contrast sensitivity measurements. Conversion of the NAVQ results to a Rasch score allowed significance to be determined with a Wilcoxon rank-sum test.

The Radner reading speed data was fitted with a non-linear regression (exponential rise to a maximum). Maximum reading speed (MRS) was defined as the asymptote of this curve and Critical print size (CPS) was calculated as the value for x (print size) when the reading speed was 95% of the MRS.

$$x = \frac{\text{Log}(1 - (y-c/a))}{b} \quad \text{equation 1}$$

Three methods were used to describe the defocus curves using the metrics published by Buckhurst et al.²⁰ After accounting for magnification of the defocus lenses, the direct comparison method determined significance at each level of acuity; a two way repeated measures ANOVA and pairwise comparison was used to determine if there was a significant difference between groups. Subsequently, fitting spline curves to the dataset allowed the calculation of the range-of-focus, determined using 0.3LogMAR as the threshold. Finally, the near, intermediate and distance areas of the curve were calculated using 0.3LogMAR as the upper limit.²⁰

RESULTS

Patient Demographics

Ninety subjects completed the study, one subject had a surgical complication (posterior capsular rupture) prior to IOL insertion and was thus excluded from the study. All subjects attended the

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

initial post-operative assessment with the consultant surgeon 3-4 weeks post-surgery, however nine subjects were lost to follow up thereafter, seven of these were excluded due to failure to attend one or both of their study visits despite repeated requests, one failed to attend due to ill health and the remaining subject was deceased (Figure 1). There were no adverse or serious adverse events reported in any subjects.

There were no significant differences in pre-operative measures between subjects in the monofocal IOL and MIOL groups, $p > 0.05$ in all instances ([Supplementary Table 3](#)).

Post-Operative Refraction

For all participants, manifest spherical equivalent (MSE) was calculated and astigmatism was analysed using the power vector method as described by Thibos.²¹ The effect of uncorrected astigmatism²² is known to be detrimental to outcomes and as such vector analysis was used to ensure that astigmatic effect was similar between groups. No significant differences were found between groups ($p > 0.05$) ([Supplementary Table 4](#)).

Visual Acuity

Significant differences were found for UNVA ($p < 0.01$) and DCNVA ($p < 0.01$) both monocularly and binocularly at V1 and V2. With near visual acuity being significantly better in the MIOL group.

No significant difference was found for intermediate vision (70cm) (Figure 2)([Supplementary Figure 1](#))([Supplementary Table 5](#)).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Defocus

A two-way repeated measure ANOVA was performed and a significant difference found ($F_{1,28} = 131.889$ $p < 0.001$). Pairwise comparisons identified that the differences were significant through the defocus range -2.00 to -5.00 ($p < 0.001$) at both visits, monocularly and binocularly (Figure 3)([Supplementary Figure 2](#)). Cohen's D effect size was calculated and remained > 1 throughout this range, thus categorized as a large effect size.

Defocus curves were also analysed using the area under the curve method as previously described.²⁰ MATLAB R2017b (The Mathworks Inc) curve fitting software was used to fit a spline curve to each data set. The same software was then used to calculate the area below the curve assuming $y = 0.3\text{LogMAR}$. The ranges were divided into distance (-0.5 to +0.5 defocus), intermediate (-0.5 to -2.0D defocus) and near (-2.0 to -4.0D defocus). A cut-off value of 0.3LogMAR was used as this is the UK, European and American binocular visual acuity driving standards.^{23,24}

Distance area was significantly greater in the monofocal group at Visit 1 but not at Visit 2, no difference was found in the intermediate area but the MIOL group showed a larger near area at both visits. In addition to the area metrics, range of focus was calculated as the dioptric range where VA was ≥ 0.3 LogMAR, by finding the roots of the spline curve fitted. The MIOL group had a significantly larger range of focus ($p < 0.001$) (Figure 4)([Supplementary Figure 2](#)) ([Supplementary Table 6](#)).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Reading Speed


There was significantly better critical print size (CPS) and reading acuity achieved in the MIOL group at V1 ($p < 0.001$) and V2 ($p < 0.001$). No significant difference in MRS was found at either visit ($p = 0.534$ V1 and $p = 0.555$ V2) (Figure 5) ([Supplementary Figure 4](#)).

Contrast Sensitivity

Monocular and binocular measures of contrast sensitivity with the Pelli-Robson charts showed a significant difference ($p < 0.001$) at Visit 1 with a large effect size demonstrated (Cohen's $d = 0.845$ and 1.031 respectively) ([Supplementary Figure 5](#)). However, at Visit 2, there was no significant difference between groups when tested binocularly ($p = 0.059$) (Figure 6).

Binocular contrast sensitivity, measured with the CSV-1000, was greater in the IOL group at visit 1 when measured at 3, 6 and 12cpd spatial frequencies ([Supplementary Figure 6](#)); this difference was only present for 12 and 18cpd at Visit 2 (Figure 7) ([Supplementary Table 7](#)).

Prior to visit 1 no subject underwent YAG capsulotomy whereas by visit 2; in the monofocal group one subject required YAG capsulotomy unilaterally and one bilaterally, and in the multifocal group one subject required YAG capsulotomy unilaterally and three required it bilaterally. No post-operative procedures were performed, for the correction of residual ametropia, on any of the subjects.




1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Questionnaire

75% of the MIOL group were completely spectacle independent compared to 6.7% of the monofocal group at Visit1. At Visit 2, 66.7% and 4.7% respectively remained completely spectacle independent ~~(Supplementary Figure 7).Figure 8a, 8b).~~

The type of spectacles worn in both groups was different post-operatively compared to pre-operatively with fewer subjects using bifocals or varifocals. Single vision near spectacles (reading only) were the most common refractive correction in both groups. A small proportion of subjects used spectacles for distance; this finding was consistent with the satisfaction results. In addition 2.5% of the MIOL group used varifocal spectacles post-operatively due to patient preference for varifocals rather than single vision reading spectacles and not due to a need for full time correction. Difficulty scores were low for everyday tasks such as driving and watching TV (Figure 9).

Overall satisfaction was high (> 90% of subjects) in both groups for distance tasks. Satisfaction was greater for the MIOL group at both intermediate and near (Figure 9a, ~~9b~~). Significant differences were found between groups for all near tasks (Figure 9~~b~~~~c~~, ~~9d~~) and at both visits the monofocal group reported significantly more difficulty using a VDU screen (Figure 9~~c~~~~e~~, ~~9f~~). However, satisfaction scores were similar for distance tasks such as driving and watching TV.



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65


Subjects were asked to rate the difficulty invoked in general night vision, and with glare, halos, starburst and ghost images (Figure 9~~g~~^h). Significant difference between groups were only evident for halos at both visits; MIOL scores were higher but still categorised as low difficulty (between 1 and 3 for all subjects).

The Zeiss (Carl Meditec Ltd) Glare simulator was used and subjects asked to adjust the settings in order to pictorially display halos/glare akin to those they observe at night. 77% of the MIOL group reported halos, compared to just 6% of the IOL group. Halo size and intensity was quantified using the simulator on a scale of 0 (no halo) to 100 (maximum). Results showed a significant difference in halo size reported in the MIOL group (Figure 10)([Supplementary Figure 9](#)).

The MIOL group had a significantly better NAVQ score, consistent with the greater spectacle independence achieved amongst participants in that group (Figure 11)([Supplementary Figure 10](#)).

DISCUSSION

The 2016 Cochrane review⁶ highlighted the need for the evaluation of MIOLs using a core set of standardised outcome measures and graded the current certainty of evidence for efficacy as very low to moderate. This RCT aimed to build on the evidence base by evaluating MIOLs using a comprehensive set of standard outcome measures. Participants were recruited from patients




1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

referred for cataract surgery under the UK NHS. As such the subjects did not attend expecting MIOL implantation and were not motivated for achieving spectacle independence which may in fact have biased the results towards spectacle dependence. Conversely, most existing studies of this nature are non-randomised and hence prone to bias towards spectacle independence in addition to influencing IOL selection.²⁵ In addition the mean age of the subjects in this study represent the oldest population of all of the IOL/MIOL RCTs and is the first where the subjects have a mean age greater than 75. As such, the results provide a generalizable dataset for an older patient base.

Near vision

Good uncorrected near vision is the primary motivation for MIOL implantation but assessing it requires a multifaceted approach. Previous studies have shown good near vision with bifocal IOLs, and improved satisfaction with near tasks and spectacle independence²⁵⁻²⁷ When compared with a monofocal IOL the present study demonstrated improved unaided and best distance corrected near vision with a MIOL. These results are further supported by the defocus curve analysis, via both the traditional direct comparison method and through the area and range of focus metrics.²⁰ Additionally the Radner reading charts showed significantly smaller critical print size was achieved whilst maintaining maximum reading speed in the MIOL group. The subjective perception of near vision was also enhanced in the MIOL group as evident via the observations of the two questionnaires used in this study, (QoV questionnaire¹⁷ and the previously validated NAVQ¹⁸); no differences in satisfaction scores were identified for the distance and intermediate vision.




1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

It must be noted, in most studies, including this study, an arbitrary reading distance of 40cm was used, this is likely to show optimum reading performance for an IOL that has an addition of +2.50D in the spectacle plane, however higher adds will have optimum acuity at a shorter focal length. Therefore, it is possible that maximum UNVA and DCNVA has not been recorded due to this imposed working distance.

Distance Vision

UDVA, CDVA and the direct comparison method of defocus curve analysis demonstrated no difference in vision at distance between the two lens types. Whilst the distance area-of-focus metric was greater at V1, by V2 both distance areas were similar. However, contrast sensitivity measurements were lower in the MIOL group at visit 1. This is consistent with the findings of other studies ^{26, 28-33} and is an expected finding with any RCT comparing MIOLs with IOLs. All MIOLs have a near focal point, which creates a myopic blur circle around the distance focal point; it is this blur that affects CS. The MIOL examined in the present study is designed to be distant dominant when viewing a distance object (provided a large pupil is present), this will reduce the intensity of the blur circle minimizing its impact on CS and preserving distance vision quality. By months 12-18 there was no significant difference in CS as measured on the Peli-Robson and at all but the low spatial frequencies on the CSV-1000. Given that there was no significant difference in distance visual acuity, and that the subjective satisfaction of distance vision was comparable, it is probable that the lens design has minimized the impact of the blur circle to the point whereby it is no longer of clinical significance.




1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Subjects implanted with MIOLs reported halos at both visits according to both the questionnaire data and glare simulator. This is to be expected as these halos are created by the defocus of the second focal point and are present with all MIOLs. The intensity of the halo is an important consideration with MIOL design. Theoretically distance dominant MIOL demonstrate lower halo intensities. The study MIOL incorporates a partially diffractive surface which is distance dominant with large pupil sizes and given that the perception of halos occurs mainly at night it is likely that the impact of halos on vision has been minimized: This may explain how, despite the presence of halos, overall satisfaction with distance vision was high (97%).

Intermediate Vision

Intermediate vision is relatively difficult to define and hence this study has used a variety of methods to assess visual function in this region. The intermediate area-of-focus metric defined by Buckhurst and colleagues²⁰ and used in this study evaluates vision quality between a defocus of -0.50 to -2.00D (corresponding to a working distance of approximately 0.50 to 2.00m). The intermediate area-of-focus results showed no significant difference between the MIOL and IOL; affirmed by the non-significant finding for intermediate vision using the ETDRS chart at 70cm. The Direct comparison method of defocus curve analysis demonstrated an improved visual acuity with a -2.00D of optical defocus corresponding with a distance of 50cm. This is similar to the findings of Hayashi³⁴ who found that an MIOL of +3.00D addition vision provided similar acuities to a monofocal IOL at distances of 1.0 and 0.7m whilst better acuities at 0.5 and 0.3m. Hitherto, the only study to have examined the Bi-Flex 677MY MIOL was a non-control cohort study on 25 subjects²⁷. Analogous to the present observations the investigators noted similar defocus curves



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65


with a peak in visual acuity at approximately -2.50D of defocus with a similar profile across the intermediate range. Comparability between the present study and this cohort study is limited as only mean defocus curve acuity values were reported and mean age of the cohort was over 10 years younger than that of the present study. Subsequent to the results of this study a revised version of this optic has been designed (the Liberty MIOL), that distributes light to the intermediate zone.

Interestingly, in the present study the perception of quality of vision for computer use was superior amongst the MIOL group; suggesting that improved acuity at 0.5m is sufficient to notice an improvement in vision for VDU use.

Spectacle independence

67% of the MIOL group were found to be entirely spectacle independent, whilst the remaining 33% of patients only wore glasses occasionally. This is a lower level of spectacle independence than has been recorded in previous studies.^{25,28,35,36} Motivation for spectacle independence is likely to be an important factor in these disparate observations; given that in the present study, participants attended for cataract removal rather than for a specific refractive outcome. Individuals with a prior motivation to be spectacle independent are more likely to tolerate near and intermediate blur and hence comparability between studies can be limited.

Only 5% of the monofocal group were found to be spectacle independent with 30% requiring constant correction and the remaining 65% occasionally wearing spectacles. A disparity between the type of spectacles worn was evident between groups, with 35% of subjects implanted with monofocal IOLs wearing either bifocals or varifocals post-operatively when compared to just 3%



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

of the MIOL group. It is important to note that overall satisfaction of distance vision was similar in both groups whilst satisfaction of near and intermediate vision was considerably greater in the MIOL group with 95% of subjects satisfied.

Unaided near visual acuity is demonstrably improved with the Bi-Flex MY IOL with greater spectacle independence. With regard to visual acuity measures, it must be noted that this study aimed to compare the MIOL and monofocal IOL using a standardised method, with specific lighting levels and working distances for near and intermediate. Limitations in visual performance due to halos, glare and reduction in contrast were evident amongst the MIOL group, and although statistically significant, they do not appear to limit the subject's visual function nor their perception of vision and overall satisfaction. Thus, the study concludes that the Bi-Flex MY multifocal IOL demonstrates efficacy for the correction of near and distance vision and is indicated when improved near vision/spectacle independence is required.


WHAT WAS KNOWN

Multifocal IOLs provide both distance and near vision whereas monofocal IOLs provide image quality at a single distance

Multifocal IOLs cause an increased prevalence of dysphotopsia and result in reduced retinal image contrast

A new Biconvex, aspheric, apodized, diffractive MIOL with a +3.50D add has been designed with a relatively low number of diffractive echelons aimed to improving the optical image quality of the resultant image.

WHAT THIS PAPER ADDS



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Distance visual acuity was comparable between the MIOL and IOL. Contrast sensitivity was reduced 3-6 months post-operatively whereas by months 12-18 were similar at all but low spatial frequencies.

Near vision was superior in the MIOL group and subjects in the MIOL group were more satisfied with the quality of vision at near and intermediate.

There was a statistically significant increase in the presence of dysphotopsia in the MIOL group, however, satisfaction with distance vision was high in both groups

REFERENCES


1. Alio JL, Kaymak H, Breyer D, Cochener B, Plaza-Puche AB. Quality of life related variables measured for three multifocal diffractive intraocular lenses: a prospective randomised clinical trial. *Clin Exp Ophthalmol*. 2018;46(4):380-388.
2. Alio JL, Plaza-Puche AB, Fernandez-Buenaga R, Pikkell J, Maldonado M. Multifocal intraocular lenses: An overview. *Surv Ophthalmol*. 2017;62(5):611-634.
3. Greenstein S, Pineda R, 2nd. The Quest for Spectacle Independence: A Comparison of Multifocal Intraocular Lens Implants and Pseudophakic Monovision for Patients with Presbyopia. *Semin Ophthalmol*. 2017;32(1):111-115.
4. Alba-Bueno F, Garzon N, Vega F, Poyales F, Millan MS. Patient-Perceived and Laboratory-Measured Halos Associated with Diffractive Bifocal and Trifocal Intraocular Lenses. *Curr Eye Res*. 2018;43(1):35-42.
5. Vega F, Alba-Bueno F, Millan MS, Varon C, Gil MA, Buil JA. Halo and Through-Focus Performance of Four Diffractive Multifocal Intraocular Lenses. *Invest Ophthalmol Vis Sci*. 2015;56(6):3967-3975.
6. de Silva SR, Evans JR, Kirthi V, Ziaei M, Leyland M. Multifocal versus monofocal intraocular lenses after cataract extraction. *Cochrane Database Syst Rev*. 2016;12:CD003169.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

7. Grzybowski A, Kanclerz P, Muzyka-Wozniak M. Methods for evaluating quality of life and vision in patients undergoing lens refractive surgery. *Graefes Arch Clin Exp Ophthalmol*. 2019.
8. Maxwell A, Holland E, Cibik L, et al. Clinical and patient-reported outcomes of bilateral implantation of a +2.5 diopter multifocal intraocular lens. *J Cataract Refract Surg*. 2017;43(1):29-41.
9. Ferris FL, 3rd, Kassoff A, Bresnick GH, Bailey I. New visual acuity charts for clinical research. *Am J Ophthalmol*. 1982;94(1):91-96.
10. Hazel CA, Elliott DB. The dependency of logMAR visual acuity measurements on chart design and scoring rule. *Optom Vis Sci*. 2002;79(12):788-792.
11. Shah N, Laidlaw DA, Brown G, Robson C. Effect of letter separation on computerised visual acuity measurements: comparison with the gold standard Early Treatment Diabetic Retinopathy Study (ETDRS) chart. *Ophthalmic Physiol Opt*. 2010;30(2):200-203.
12. Rosser DA, Murdoch IE, Fitzke FW, Laidlaw DA. Improving on ETDRS acuities: design and results for a computerised thresholding device. *Eye (Lond)*. 2003;17(6):701-706.
13. Williams MA, Moutray TN, Jackson AJ. Uniformity of visual acuity measures in published studies. *Invest Ophthalmol Vis Sci*. 2008;49(10):4321-4327.
14. Wolffsohn JS, Jinabhai AN, Kingsnorth A, et al. Exploring the optimum step size for defocus curves. *J Cataract Refract Surg*. 2013;39(6):873-880.
15. Gupta N, Wolffsohn JS, Naroo SA. Optimizing measurement of subjective amplitude of accommodation with defocus curves. *J Cataract Refract Surg*. 2008;34(8):1329-1338.
16. Radner W, Diendorfer G, Kainrath B, Kollmitzer C. The accuracy of reading speed measurement by stopwatch versus measurement with an automated computer program (rad-rd(c)). *Acta Ophthalmol*. 2017;95(2):211-216.
17. Law EM, Aggarwal RK, Kasaby H. Clinical outcomes with a new trifocal intraocular lens. *Eur J Ophthalmol*. 2014;24(4):501-508.
18. Buckhurst PJ, Wolffsohn JS, Gupta N, Naroo SA, Davies LN, Shah S. Development of a questionnaire to assess the relative subjective benefits of presbyopia correction. *J Cataract Refract Surg*. 2012;38(1):74-79.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

19. Ray WA, O'Day DM. Statistical analysis of multi-eye data in ophthalmic research. *Invest Ophthalmol Vis Sci.* 1985;26(8):1186-1188.
20. Buckhurst PJ, Wolffsohn JS, Naroo SA, et al. Multifocal intraocular lens differentiation using defocus curves. *Invest Ophthalmol Vis Sci.* 2012;53(7):3920-3926.
21. Thibos LN, Wheeler W, Horner D. Power vectors: an application of Fourier analysis to the description and statistical analysis of refractive error. *Optom Vis Sci.* 1997;74(6):367-375.
22. Wolffsohn JS, Bhogal G, Shah S. Effect of uncorrected astigmatism on vision. *J Cataract Refract Surg.* 2011;37(3):454-460.
23. Bron AM, Viswanathan AC, Thelen U, et al. International vision requirements for driver licensing and disability pensions: using a milestone approach in characterization of progressive eye disease. *Clin Ophthalmol.* 2010;4:1361-1369.
24. Rees GB. Vision standards for driving: what ophthalmologists need to know. *Eye (Lond).* 2015;29(6):719-720.
25. Cochener B, Arnould B, Viala M, Roborel de Climens A, Berdeaux G. Corrected and uncorrected near and distance vision with ReSTOR compared to monofocal intraocular lens implantation after cataract surgery: a pooled analysis. *Ophthalmologica.* 2009;223(2):128-135.
26. Ji J, Huang X, Fan X, Luo M. Visual performance of Acrysof ReSTOR compared with a monofocal intraocular lens following implantation in cataract surgery. *Exp Ther Med.* 2013;5(1):277-281.
27. Garcia-Bella J, Ventura-Abreu N, Morales-Fernandez L, et al. Visual outcomes after progressive apodized diffractive intraocular lens implantation. *Eur J Ophthalmol.* 2018;28(3):282-286.
28. Cillino S, Casuccio A, Di Pace F, et al. One-year outcomes with new-generation multifocal intraocular lenses. *Ophthalmology.* 2008;115(9):1508-1516.
29. Zhao G, Zhang J, Zhou Y, Hu L, Che C, Jiang N. Visual function after monocular implantation of apodized diffractive multifocal or single-piece monofocal intraocular lens Randomized prospective comparison. *J Cataract Refract Surg.* 2010;36(2):282-285.



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

30. Wilkins MR, Allan BD, Rubin GS, et al. Randomized trial of multifocal intraocular lenses versus monovision after bilateral cataract surgery. *Ophthalmology*. 2013;120(12):2449-2455 e2441.
31. Pedrotti E, Carones F, Aiello F, et al. Comparative analysis of visual outcomes with 4 intraocular lenses: Monofocal, multifocal, and extended range of vision. *J Cataract Refract Surg*. 2018;44(2):156-167.
32. Harman FE, Maling S, Kampougeris G, et al. Comparing the 1CU accommodative, multifocal, and monofocal intraocular lenses: a randomized trial. *Ophthalmology*. 2008;115(6):993-1001 e1002.
33. Kamlesh, Dadeya S, Kaushik S. Contrast sensitivity and depth of focus with aspheric multifocal versus conventional monofocal intraocular lens. *Can J Ophthalmol*. 2001;36(4):197-201.
34. Hayashi K, Manabe S, Hayashi H. Visual acuity from far to near and contrast sensitivity in eyes with a diffractive multifocal intraocular lens with a low addition power. *J Cataract Refract Surg*. 2009;35(12):2070-2076.
35. Baig R, T AC, Kukreja S, Shakil S, Ahmad K. Patients' satisfaction and spectacle independence after cataract surgery with multifocal intraocular lens implantation in a tertiary care hospital. *J Pak Med Assoc*. 2016;66(6):745-747.
36. Mendicute J, Kapp A, Levy P, et al. Evaluation of visual outcomes and patient satisfaction after implantation of a diffractive trifocal intraocular lens. *J Cataract Refract Surg*. 2016;42(2):203-210.


Supplementary Table 1: Inclusion/Exclusion Criteria

Supplementary Table 2: Characteristics of the Intraocular lenses

Supplementary Table 3: Patient Demographics

Supplementary Table 4: Refraction

Supplementary Table 5: Visual Acuity Results



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Supplementary Table 6: Area Under Defocus

Supplementary Table 7: CSV-1000

Figure 1: Trial Profile

Figure 2: ~~a) Visit 1 Monocular Visual Acuity, b) Visit 1 Binocular Visual Acuity, a)c) Visit 2 Monocular Visual Acuity, b)d) Visit 2 Binocular Visual Acuity.~~

Figure 3: ~~a) Visit 1 Monocular Defocus Curve, b) Visit 1 Binocular Defocus Curve, a)e) Visit 2 Monocular Defocus Curve, b)d) Visit 2 Binocular Defocus Curve~~


Figure 4: ~~a) Visit 1 Area under defocus curve, a)b) Visit 2 Area under defocus curve, c) Visit 1 Range of focus, b)d) Visit 2 Range of focus~~

Figure 5: ~~a) Visit 1 Maximum Reading Speed, b)a) Visit 2 Maximum Reading Speed, e) Visit 1 Reading Acuity, d)b) Visit 2 Reading Acuity, e) Visit 1 95% Critical Print Size f)c) Visit 2 95% Critical Print Size~~

Figure 6: ~~a) Visit 1 Monocular Contrast Sensitivity, b) Visit 1 Binocular Contrast Sensitivity c) Visit 2 Monocular Contrast Sensitivity, b)d) Visit 2 Binocular Contrast Sensitivity.~~

Figure 7: ~~a) Visit 1 CSV-1000 Contrast sensitivity, b) Visit 2 CSV-1000 Contrast sensitivity~~

Figure 8: Spectacle Wear ~~a) Visit 1 Frequency of wear, b)a) Visit 2 Frequency of wear, c) Visit 1 Type of spectacles, d, b) Visit 2 Type of spectacles, e c) Pre-Op spectacle wear~~



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 9: Quality of Vision Questionnaire results ~~a) Visit 1 Satisfaction~~, ~~ba) Visit 2 Satisfaction~~,
~~e) Visit 1 Near Tasks~~, ~~db) Visit 2 Near Tasks~~, ~~e) Visit 1 Everyday Tasks~~, ~~fc) Visit 2 Everyday Tasks~~,
~~g) Visit 1 Night Vision~~, ~~hd) Visit 2 Night Vision~~

Figure 10: Visit 2 Glare Simulator Scores ~~a) Visit 1~~ ~~b) Visit 2~~

Figure 11: Visit 2 NAVQ Scores ~~a) Visit 1~~ ~~b) Visit 2~~

Supplementary Figure 1: a) Visit 1 Monocular Visual Acuity, b) Visit 1 Binocular Visual Acuity

Supplementary Figure 2: a) Visit 1 Monocular Defocus Curve, b) Visit 1 Binocular Defocus Curve

Supplementary Figure 3: a) Visit 1 Area under defocus curve, b) Visit 1 Range of focus

Supplementary Figure 4: a) Visit 1 Maximum Reading Speed, b) Visit 1 Reading Acuity, c) Visit 1
95% Critical Print Size

Supplementary Figure 5: a) Visit 1 Monocular Contrast Sensitivity, b) Visit 1 Binocular Contrast
Sensitivity


Supplementary Figure 6: Visit 1 CSV-1000 Contrast sensitivity

Supplementary Figure 7: Spectacle Wear a) Visit 1 Frequency of wear, b) Visit 1 Type of
spectacles


Supplementary Figure 8: Quality of Vision Questionnaire results a) Visit 1 Satisfaction, b) Visit 1
Near Tasks c) Visit 1 Everyday Tasks, d) Visit 1 Night Vision

Supplementary Figure 9: Visit 1 Glare Simulator Scores

Supplementary Figure 10: Visit 1 NAVQ Score



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65



Formatted: Justified

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Visual Function and Subjective Perception of Vision following bilateral implantation of monofocal and multifocal intraocular lenses: A Randomised Controlled Trial.

Elizabeth M. Law, MSc,^{1,2} Rajesh K. Aggarwal, BM, FRCOphth,² Hetal Buckhurst, PhD,¹ Hosam E. Kasaby MBChB, FRCOphth,² Jonathan Marsden, PhD,¹ Gary Shum, PhD,¹ and Phillip J. Buckhurst, PhD.¹

¹University of Plymouth, School of Health Professions, Peninsula Allied Health Centre, Derriford Road, Plymouth, United Kingdom

²BMI Southend Hospital, Fairfax Drive, Westcliff on Sea, United Kingdom

Funding: The work was funded by MediconTur Medical Engineering (Zsámbék, Hungary). MediconTur had no role in the design or conduct of this research.

Financial Disclosure: No conflicting relationship exists for any author.

Running Head: Visual Function with MIOLs

Correspondence and reprint requests to Phillip J. Buckhurst, PhD, University of Plymouth, School of Health Professions, Peninsula Allied Health Centre, Derriford Road, Plymouth, PL6 8BH, United Kingdom. E-mail: Phillip.buckhurst@plymouth.ac.uk

1
2
3
4 **ABSTRACT**
5

6
7
8 **PURPOSE:**
9

10 Following implantation with Multifocal intraocular lenses (MIOLs) or monofocal intraocular
11 lenses (IOLs), the study examines monocular and binocular visual function and patient reporting
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Following implantation with Multifocal intraocular lenses (MIOLs) or monofocal intraocular lenses (IOLs), the study examines monocular and binocular visual function and patient reporting outcomes using a rigorous series of clinical assessments.

Setting: BMI Southend Hospital, UK

DESIGN: Prospective, randomised, double-masked clinical trial.

METHODS: 100 subjects were randomised for bilateral implantation of either Bi-Flex 677MY MIOL or Bi-Flex 677AB IOL and were assessed at 3-6 months (V1) and 12-18 months (V2). Primary outcomes included distance, intermediate and near LogMAR visual acuities (VA) and defocus curve profile assessment. Secondary outcomes included reading speed, contrast sensitivity (CS) and the subjective perception of quality-of-vision.

RESULTS: Uncorrected (MIOL 0.10 ± 0.09 LogMAR; IOL 0.09 ± 0.11 LogMAR) and best distance-corrected VA (MIOL 0.04 ± 0.06 LogMAR; IOL 0.01 ± 0.07 LogMAR) were comparable ($p > 0.05$). Unaided near VA (UNVA $p < 0.001$: MIOL 0.23 ± 0.13 LogMAR; IOL 0.55 ± 0.20 LogMAR) and distance-corrected near VA (DCNVA $p < 0.001$: MIOL 0.24 ± 0.13 LogMAR; IOL 0.54 ± 0.17 LogMAR) were significantly improved with MIOLs. There was no significant difference in distance-corrected intermediate VA (DCIVA $p = 0.431$: MIOL 0.38 ± 0.13 ; IOL 0.39 ± 0.13).

Defocus curves demonstrated an increased range-of-focus amongst MIOLs (MIOL 4.14 ± 1.10 D; IOL 2.57 ± 0.77 D). Pelli-Robson CS was different at V1 ($p < 0.001$) but similar by V2 ($p = 0.059$).

1
2
3
4 Overall satisfaction was high (>90%) in both groups for distance tasks whereas significantly
5
6 different for near (MIOL 18.45±16.53LogUnits; MIOL 55.59±22.52LogUnits).
7
8
9

10 **CONCLUSIONS:** Unaided near visual acuity is demonstrably better with MIOLs and there was
11
12 greater subjective satisfaction with their quality-of-near-vision. Halos reported by the MIOL
13
14 group was significant compared to the IOL group, but did not show an adverse effect on overall
15
16 satisfaction.
17
18
19
20
21
22
23
24
25
26

27 Multifocal intraocular lenses (MIOLs) are widely considered the most reliable method of
28
29 achieving spectacle independence following cataract surgery.¹⁻³ MIOLs distribute the light
30
31 between distant and near focal points whereby the vergence of the incident light dictates which
32
33 focal point is conjugate to the retinal plane.
34
35
36
37

38 The separation of these multiple focal points is determined by the addition power of the MIOL
39
40 and to a lesser extent the biometry of the eye. High addition MIOLs (+4.00D or higher) are the
41
42 zeitgeist of the designs used in the late 90s-early 2000s. Disadvantages of these early lenses
43
44 included a close working distance and reduced intermediate vision. Moreover, the size of the
45
46 dysphotopic phenomenon (commonly described as halo), associated with MIOLs, increases
47
48 according to the addition power; these higher addition lenses generate larger haloes.^{4,5}
49
50
51
52
53

54 The light energy distribution between the retinal focal points created by a MIOL influences the
55
56 overall quality of vision at different viewing distances. MIOLs that split light equally, create two
57
58 focal points of comparative image quality. In contrast, distance dominant MIOLs allocate a higher
59
60
61
62
63
64
65

1
2
3
4 percentage of light towards the distance retinal focal point and consequently near vision is
5
6 relatively compromised. Conversely, the intensity of the halo is influenced by the light
7
8 distribution relationship: the more distance dominant the lower the dysphotopic intensity.
9
10

11
12 In 2016, a Cochrane Review⁶ highlighted the need for robust randomised control trials examining
13
14 the efficacy of MIOLs over monofocal intraocular lens (IOL) implantation and called for
15
16 standardization of outcome measures in MIOL studies. The review concluded that it was unclear
17
18 whether the achieved benefits of MIOL implantation i.e. greater near vision and increased
19
20 spectacle independence, outweighed disadvantages such as reduced contrast sensitivity and
21
22 increased dysphotopsia. Subsequently others have also highlighted the importance of patient
23
24 reported outcomes in MIOLs.⁷ Despite these conclusions, in the subsequent three years there
25
26 has only been a single RCT published comparing MIOLs with IOLs.⁸
27
28
29
30
31
32
33

34 The present study compared the efficacy of the Bi-Flex 677MY MIOL over its parent monofocal
35
36 IOL using standardized methods for assessing both visual function and the subjective perception
37
38 of the quality of vision.
39
40
41
42
43
44
45

46 **METHODS**

47
48
49

50 This study was a prospective, parallel double masked randomised clinical trial. The study protocol
51
52 adheres to the Declaration of Helsinki and ethical approval was obtained prior to commencement
53
54 of the trial. The study was registered with clinicaltrials.gov (NCT02338882) and written consent
55
56 was obtained from all subjects. No modifications to the protocol or outcome measures were
57
58
59
60
61
62
63
64
65

1
2
3
4 made during the study. The aim was to assess the IOLs using recognised methods that would
5
6 provide rigour and establish a comprehensive method which could be utilised with all IOLs and
7
8 allow easy comparison of results.
9

10 11 12 **Patient Selection**

13
14
15
16 Between September 2015 and May 2017, one hundred subjects were recruited from routine
17
18 cataract clinics at the BMI Southend Hospital on a consecutive – if – eligible basis according to
19
20 the inclusion/exclusion criteria (Supplementary Table 1). All subjects underwent initial
21
22 examination by a consultant ophthalmic surgeon including dilated fundus examination; in the
23
24 event of suspected macular pathology an OCT was carried out and if pathology was detected, the
25
26 patient was excluded as per the study criterion. The anterior segments and ocular surface were
27
28 also evaluated to confirm lack of pathology and minor ocular surface dryness was treated by
29
30 commencement of ocular lubricants. Any ocular surface disease deemed moderate or marked
31
32 resulted in exclusion. The allocation of IOLs was randomly designated and was masked to both
33
34 the participant and the investigator conducting the post-operative study assessments. On
35
36 enrolment, a study number was assigned to each subject. Using this study number, the allocation
37
38 of lenses for all subjects was randomized in Microsoft Excel using blocked randomization with a
39
40 1:1 allocation ratio. Following allocation of the subject number, the unmasked surgeons and
41
42 theatre staff accessed the randomization log and a series of sealed opaque envelopes that
43
44 described which lenses were to be implanted (MIOL or IOL).
45
46
47
48
49
50
51
52
53
54
55
56
57

58 **Surgical Technique**

59
60
61
62
63
64
65

1
2
3
4 All surgeries were performed by one of two experienced consultant ophthalmic surgeons (RA and
5
6
7 HK) using small incision phacoemulsification. The same surgeon implanted both lenses for an
8
9 individual subject. In each case, a 2.2mm clear corneal incision was located according to the
10
11 steepest corneal meridian. The pre- and post-operative medication regime was the same
12
13 regardless of surgeon. Second eye surgery occurred within 4 weeks of first eye surgery.
14
15
16
17

18 **Masking**

19
20
21 All post-operative study outcome measures were collected by a study investigator, who was
22
23 masked to the allocation of study group. The subjects were also masked to their grouping
24
25 allocation and were only informed of the type of lens implanted once they had completed the
26
27 study. Post-operative slit lamp examination was performed by the unmasked consultant surgeon
28
29 in order to maintain masking of the study investigator.
30
31
32
33
34

35 **Intraocular Lenses**

36
37
38 Each group had fifty subjects assigned. The Bi-Flex 677 AB is a single piece, aspheric aberration
39
40 neutral IOL. The Bi-Flex MY MIOL has the same platform as the monofocal but the anterior
41
42 surface has a 3mm apodized, diffractive central region with a near addition of 3.50D at the IOL
43
44 plane (Supplementary Table 2). The Bi-Flex MY MIOL design is intended to provide distance
45
46 dominance with greater mydriasis, thus maximizing contrast and minimizing halos when driving
47
48 at night. Pupil miosis changes the light distribution relationship and results in a relatively equal
49
50 split of light, hence, the Bi-Flex MY MIOL exploits the near miosis that occurs with reading. This
51
52 type of MIOL was chosen for the study given that the unique aspect of the Bi-Flex MY is its low
53
54 number of diffractive echelons (seven) which is theorized to improve the optical image quality of
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 the resultant image. The identical platform and material of the two IOLs allowed unhindered
5
6
7 assessment of the multifocality.
8
9

10 **Primary Outcomes Measures**

11
12
13
14 A masked investigator assessed the subjects at two study visits, 3-6 months (V1) and 12-18
15
16 months (V2) post-operatively. At each visit, monocular and binocular LogMAR acuities for
17
18 unaided distance visual acuity (UDVA) and corrected distance visual acuity (CDVA) were
19
20 measured using computerised test charts (Thomson Software Solutions Ltd) at 6m following the
21
22 Bailey-Lovie principles and employing Sloan letters consistent with testing methods established
23
24 by the Early Treatment Diabetic Retinopathy Study (ETDRS).⁹⁻¹³ Subjective refraction was
25
26 conducted at 6m with a distance fixation target. The assessment of unaided near visual acuity
27
28 (UNVA), distance corrected near visual acuity (DCNVA) and distance corrected intermediate
29
30 visual acuity (DCIVA) utilised ETDRS charts for near (40cm) and intermediate (70cm) (Precision
31
32 Vision) working distances respectively. To further assess intermediate and near vision at a range
33
34 of distances, defocus profiles were plotted from -5.00D to 1.50D in 0.50D steps.¹⁴ The letters and
35
36 defocus lenses were randomised between measures and subjects were prompted once using the
37
38 phrase “can you read any more letters on the line below?”.¹⁵ All measures of visual acuity were
39
40 performed with illuminance 120 cd/m² and luminance of 95 lux.
41
42
43
44
45
46
47
48
49
50
51
52
53

54 **Secondary Outcome Measures**

55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 Contrast Sensitivity was assessed binocularly with the CSV-1000 (Precision Vision) calibrated to
5
6 2.4m and both monocularly and binocularly using Pelli-Robson charts at 6m (Thomson Software).
7
8 Radner reading charts were used to assess reading speed at 40cm following the method outlined
9
10 by Radner using a digital stopwatch.¹⁶ The subjective perception of vision was assessed using a
11
12 quality of vision questionnaire¹⁷ and NAVQ.¹⁸ The Carl Zeiss Meditec Glare simulator was used to
13
14 quantify the appearance of halos and glare. All secondary measures were assessed at V1 and V2.
15
16 The same assessment room was used throughout the study and all secondary tests were carried
17
18 out by the same masked investigator in photopic light conditions of illuminance 120cd/m² and
19
20 luminance of 95 lux.
21
22
23
24
25
26
27

28 **Statistical Analysis**

30
31
32 The sample size for the study was calculated using G*power3 (University of Dusseldorf). Power
33
34 calculations were based on a medium effect size ($f = 0.30$) based on *a-priori* matched paired *t* test
35
36 design and a desired statistical power of 90% with an error probability of 0.05. Statistical analysis
37
38 was performed using SPSS software, version 24 (IBM). All data were tested for normality using
39
40 the Shapiro-Wilks test and visual examination of histogram plots. In all instances $p < 0.05$ was
41
42 considered statistically significant. In order to evaluate effect size, Cohen's *d* was calculated, with
43
44 $d > 0.2$, 0.5 and 0.8 corresponding to small, medium and large effect sizes, respectively.
45
46
47
48
49

50
51 A repeated measures ANOVA was used to establish similarity between right and left eye data for
52
53 both monofocal and multifocal IOL data. No significant differences were found and as such only
54
55 right eye data is presented.¹⁹ Where differences were found after repeated measures ANOVA,
56
57 further pairwise tests were used to compare the monofocal and multifocal groups for all visual
58
59
60
61
62
63
64
65

1
2
3
4 acuity and contrast sensitivity measurements. Conversion of the NAVQ results to a Rasch score
5
6 allowed significance to be determined with a Wilcoxon rank-sum test.
7
8
9

10 The Radner reading speed data was fitted with a non-linear regression (exponential rise to a
11
12 maximum). Maximum reading speed (MRS) was defined as the asymptote of this curve and
13
14 Critical print size (CPS) was calculated as the value for x (print size) when the reading speed was
15
16
17
18 95% of the MRS.
19
20

$$x = \frac{\text{Log}(1 - (y-c/a))}{b} \quad \text{equation 1}$$

21
22
23
24
25
26

27 Three methods were used to describe the defocus curves using the metrics published by
28
29 Buckhurst et al.²⁰ After accounting for magnification of the defocus lenses, the direct comparison
30
31 method determined significance at each level of acuity; a two way repeated measures ANOVA
32
33 and pairwise comparison was used to determine if there was a significant difference between
34
35 groups. Subsequently, fitting spline curves to the dataset allowed the calculation of the range-of-
36
37 focus, determined using 0.3LogMAR as the threshold. Finally, the near, intermediate and distance
38
39 areas of the curve were calculated using 0.3LogMAR as the upper limit.²⁰
40
41
42
43
44
45
46
47
48

49 **RESULTS**

50 51 52 **Patient Demographics**

53
54
55
56 Ninety subjects completed the study, one subject had a surgical complication (posterior capsular
57
58 rupture) prior to IOL insertion and was thus excluded from the study. All subjects attended the
59
60
61
62
63
64
65

1
2
3
4 initial post-operative assessment with the consultant surgeon 3-4 weeks post-surgery, however
5
6
7 nine subjects were lost to follow up thereafter, seven of these were excluded due to failure to
8
9
10 attend one or both of their study visits despite repeated requests, one failed to attend due to ill
11
12 health and the remaining subject was deceased (Figure 1). There were no adverse or serious
13
14
15 adverse events reported in any subjects.
16
17

18 There were no significant differences in pre-operative measures between subjects in the
19
20
21 monofocal IOL and MIOL groups, $p > 0.05$ in all instances (Supplementary Table 3).
22
23
24
25
26

27 **Post-Operative Refraction**

28
29
30
31 For all participants, manifest spherical equivalent (MSE) was calculated and astigmatism was
32
33
34 analysed using the power vector method as described by Thibos.²¹ The effect of uncorrected
35
36
37 astigmatism²² is known to be detrimental to outcomes and as such vector analysis was used to
38
39
40 ensure that astigmatic effect was similar between groups. No significant differences were found
41
42
43 between groups ($p > 0.05$) (Supplementary Table 4).
44
45
46
47

48 **Visual Acuity**

49
50
51 Significant differences were found for UNVA ($p < 0.01$) and DCNVA ($p < 0.01$) both monocularly
52
53
54 and binocularly at V1 and V2. With near visual acuity being significantly better in the MIOL group.
55
56
57 No significant difference was found for intermediate vision (70cm) (Figure 2)(Supplementary
58
59
60
61
62
63
64
65 Figure 1)(Supplementary Table 5).

Defocus

A two-way repeated measure ANOVA was performed and a significant difference found ($F_{1,28} = 131.889$ $p < 0.001$). Pairwise comparisons identified that the differences were significant through the defocus range -2.00 to -5.00 ($p < 0.001$) at both visits, monocularly and binocularly (Figure 3)(Supplementary Figure 2). Cohen's D effect size was calculated and remained > 1 throughout this range, thus categorized as a large effect size.

Defocus curves were also analysed using the area under the curve method as previously described.²⁰ MATLAB R2017b (The Mathworks Inc) curve fitting software was used to fit a spline curve to each data set. The same software was then used to calculate the area below the curve assuming $y = 0.3\text{LogMAR}$. The ranges were divided into distance (-0.5 to +0.5 defocus), intermediate (-0.5 to -2.0D defocus) and near (-2.0 to -4.0D defocus). A cut-off value of 0.3LogMAR was used as this is the UK, European and American binocular visual acuity driving standards.^{23,24}

Distance area was significantly greater in the monofocal group at Visit 1 but not at Visit 2, no difference was found in the intermediate area but the MIOL group showed a larger near area at both visits. In addition to the area metrics, range of focus was calculated as the dioptric range where VA was ≥ 0.3 LogMAR, by finding the roots of the spline curve fitted. The MIOL group had a significantly larger range of focus ($p < 0.001$) (Figure 4)(Supplementary Figure 2)(Supplementary Table 6).

1
2
3
4 **Reading Speed**
5

6
7 There was significantly better critical print size (CPS) and reading acuity achieved in the MIOL
8 group at V1 ($p < 0.001$) and V2 ($p < 0.001$). No significant difference in MRS was found at either visit
9
10 ($p = 0.534$ V1 and $p = 0.555$ V2) (Figure 5) (Supplementary Figure 4).
11
12
13
14

15
16
17
18
19 **Contrast Sensitivity**
20

21
22
23
24
25
26 Monocular and binocular measures of contrast sensitivity with the Pelli-Robson charts showed
27
28 a significant difference ($p < 0.001$) at Visit 1 with a large effect size demonstrated (Cohen's $d =$
29
30 0.845 and 1.031 respectively) (Supplementary Figure 5). However, at Visit 2, there was no
31
32 significant difference between groups when tested binocularly ($p = 0.059$) (Figure 6).
33

34
35
36
37 Binocular contrast sensitivity, measured with the CSV-1000, was greater in the IOL group at visit
38
39 1 when measured at 3, 6 and 12cpd spatial frequencies (Supplementary Figure 6); this difference
40
41 was only present for 12 and 18cpd at Visit 2 (Figure 7)(Supplementary Table 7).
42
43
44

45
46
47
48 Prior to visit 1 no subject underwent YAG capsulotomy whereas by visit 2; in the monofocal
49
50 group one subject required YAG capsulotomy unilaterally and one bilaterally, and in the
51
52 multifocal group one subject required YAG capsulotomy unilaterally and three required it
53
54 bilaterally. No post-operative procedures were preformed, for the correction of residual
55
56 ametropia, on any of the subjects.
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7 **Questionnaire**
8
9

10 75% of the MIOL group were completely spectacle independent compared to 6.7% of the
11 monofocal group at Visit1. At Visit 2, 66.7% and 4.7% respectively remained completely spectacle
12 independent (Supplementary Figure 7).
13
14
15

16
17
18
19 The type of spectacles worn in both groups was different post-operatively compared to pre-
20 operatively with fewer subjects using bifocals or varifocals. Single vision near spectacles (reading
21 only) were the most common refractive correction in both groups. A small proportion of subjects
22 used spectacles for distance; this finding was consistent with the satisfaction results. In addition
23
24
25
26
27
28
29 2.5% of the MIOL group used varifocal spectacles post-operatively due to patient preference for
30 varifocals rather than single vision reading spectacles and not due to a need for full time
31 correction. Difficulty scores were low for everyday tasks such as driving and watching TV (Figure
32
33
34
35
36
37 9).
38
39
40
41
42
43

44 Overall satisfaction was high (> 90% of subjects) in both groups for distance tasks. Satisfaction
45 was greater for the MIOL group at both intermediate and near (Figure 9a). Significant differences
46 were found between groups for all near tasks (Figure 9b) and at both visits the monofocal group
47 reported significantly more difficulty using a VDU screen (Figure 9c). However, satisfaction scores
48
49
50
51
52
53
54
55 were similar for distance tasks such as driving and watching TV.
56
57
58
59
60
61
62
63
64
65

1
2
3
4 Subjects were asked to rate the difficulty invoked in general night vision, and with glare, halos,
5
6
7 starburst and ghost images (Figure 9d). Significant difference between groups were only evident
8
9
10 for halos at both visits; MIOL scores were higher but still categorised as low difficulty (between 1
11
12 and 3 for all subjects).
13
14

15
16
17
18 The Zeiss (Carl Meditec Ltd) Glare simulator was used and subjects asked to adjust the settings
19
20
21 in order to pictorially display halos/glare akin to those they observe at night. 77% of the MIOL
22
23 group reported halos, compared to just 6% of the IOL group. Halo size and intensity was
24
25 quantified using the simulator on a scale of 0 (no halo) to 100 (maximum). Results showed a
26
27 significant difference in halo size reported in the MIOL group (Figure 10)(Supplementary Figure
28
29
30
31 9).
32
33

34
35
36
37 The MIOL group had a significantly better NAVQ score, consistent with the greater spectacle
38
39 independence achieved amongst participants in that group (Figure 11)(Supplementary Figure
40
41
42 10).
43
44

45 46 47 48 **DISCUSSION** 49

50
51
52 The 2016 Cochrane review⁶ highlighted the need for the evaluation of MIOLs using a core set of
53
54 standardised outcome measures and graded the current certainty of evidence for efficacy as very
55
56 low to moderate. This RCT aimed to build on the evidence base by evaluating MIOLs using a
57
58 comprehensive set of standard outcome measures. Participants were recruited from patients
59
60
61
62
63
64
65

1
2
3
4 referred for cataract surgery under the UK NHS. As such the subjects did not attend expecting
5
6
7 MIOL implantation and were not motivated for achieving spectacle independence which may in
8
9
10 fact have biased the results towards spectacle dependence. Conversely, most existing studies of
11
12 this nature are non-randomised and hence prone to bias towards spectacle independence in
13
14 addition to influencing IOL selection.²⁵ In addition the mean age of the subjects in this study
15
16
17 represent the oldest population of all of the IOL/MIOL RCTs and is the first where the subjects
18
19
20 have a mean age greater than 75. As such, the results provide a generalizable dataset for an older
21
22
23 patient base.

24 25 26 **Near vision**

27
28
29 Good uncorrected near vision is the primary motivation for MIOL implantation but assessing it
30
31
32 requires a multifaceted approach. Previous studies have shown good near vision with bifocal
33
34 IOLs, and improved satisfaction with near tasks and spectacle independence²⁵⁻²⁷ When compared
35
36
37 with a monofocal IOL the present study demonstrated improved unaided and best distance
38
39
40 corrected near vision with a MIOL. These results are further supported by the defocus curve
41
42
43 analysis, via both the traditional direct comparison method and through the area and range of
44
45 focus metrics.²⁰ Additionally the Radner reading charts showed significantly smaller critical print
46
47
48 size was achieved whilst maintaining maximum reading speed in the MIOL group. The subjective
49
50
51 perception of near vision was also enhanced in the MIOL group as evident via the observations
52
53
54 of the two questionnaires used in this study, (QoV questionnaire¹⁷ and the previously validated
55
56
57 NAVQ¹⁸); no differences in satisfaction scores were identified for the distance and intermediate
58
59
60
61
62
63
64
65 vision.

1
2
3
4 It must be noted, in most studies, including this study, an arbitrary reading distance of 40cm was
5
6 used, this is likely to show optimum reading performance for an IOL that has an addition of
7
8 +2.50D in the spectacle plane, however higher adds will have optimum acuity at a shorter focal
9
10 length. Therefore, it is possible that maximum UNVA and DCNVA has not been recorded due to
11
12 this imposed working distance.
13
14
15
16
17

18 **Distance Vision**

19
20
21 UDVA, CDVA and the direct comparison method of defocus curve analysis demonstrated no
22
23 difference in vision at distance between the two lens types. Whilst the distance area-of-focus
24
25 metric was greater at V1, by V2 both distance areas were similar. However, contrast sensitivity
26
27 measurements were lower in the MIOL group at visit 1. This is consistent with the findings of
28
29 other studies ^{26, 28-33} and is an expected finding with any RCT comparing MIOLs with IOLs. All
30
31 MIOLs have a near focal point, which creates a myopic blur circle around the distance focal point;
32
33 it is this blur that affects CS. The MIOL examined in the present study is designed to be distant
34
35 dominant when viewing a distance object (provided a large pupil is present), this will reduce the
36
37 intensity of the blur circle minimizing its impact on CS and preserving distance vision quality. By
38
39 months 12-18 there was no significant difference in CS as measured on the Peli-Robson and at all
40
41 but the low spatial frequencies on the CSV-1000. Given that there was no significant difference
42
43 in distance visual acuity, and that the subjective satisfaction of distance vision was comparable,
44
45 it is probable that the lens design has minimized the impact of the blur circle to the point whereby
46
47 it is no longer of clinical significance.
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 Subjects implanted with MIOLs reported halos at both visits according to both the questionnaire
5
6 data and glare simulator. This is to be expected as these halos are created by the defocus of the
7
8 second focal point and are present with all MIOLs. The intensity of the halo is an important
9
10 consideration with MIOL design. Theoretically distance dominant MIOL demonstrate lower halo
11
12 intensities. The study MIOL incorporates a partially diffractive surface which is distance
13
14 dominant with large pupil sizes and given that the perception of halos occurs mainly at night it is
15
16 likely that the impact of halos on vision has been minimized: This may explain how, despite the
17
18 presence of halos, overall satisfaction with distance vision was high (97%).
19
20
21
22
23
24

25 **Intermediate Vision**

26
27
28
29 Intermediate vision is relatively difficult to define and hence this study has used a variety of
30
31 methods to assess visual function in this region. The intermediate area-of-focus metric defined
32
33 by Buckhurst and colleagues²⁰ and used in this study evaluates vision quality between a defocus
34
35 of -0.50 to -2.00D (corresponding to a working distance of approximately 0.50 to 2.00m). The
36
37 intermediate area-of-focus results showed no significant difference between the MIOL and IOL;
38
39 affirmed by the non-significant finding for intermediate vision using the ETDRS chart at 70cm.
40
41
42
43
44 The Direct comparison method of defocus curve analysis demonstrated an improved visual acuity
45
46 with a -2.00D of optical defocus corresponding with a distance of 50cm. This is similar to the
47
48 findings of Hayashi³⁴ who found that an MIOL of +3.00D addition vision provided similar acuities
49
50 to a monofocal IOL at distances of 1.0 and 0.7m whilst better acuities at 0.5 and 0.3m. Hitherto,
51
52 the only study to have examined the Bi-Flex 677MY MIOL was a non-control cohort study on 25
53
54 subjects²⁷. Analogous to the present observations the investigators noted similar defocus curves
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 with a peak in visual acuity at approximately -2.50D of defocus with a similar profile across the
5
6
7 intermediate range. Comparability between the present study and this cohort study is limited as
8
9
10 only mean defocus curve acuity values were reported and mean age of the cohort was over 10
11
12 years younger than that of the present study. Subsequent to the results of this study a revised
13
14 version of this optic has been designed (the Liberty MIOL), that distributes light to the
15
16
17 intermediate zone.
18

19
20
21 Interestingly, in the present study the perception of quality of vision for computer use was
22
23 superior amongst the MIOL group; suggesting that improved acuity at 0.5m is sufficient to notice
24
25 an improvement in in vision for VDU use.
26
27

28 29 **Spectacle independence**

30
31
32
33 67% of the MIOL group were found to be entirely spectacle independent, whilst the remaining
34
35 33% of patients only wore glasses occasionally. This is a lower level of spectacle independence
36
37 than has been recorded in previous studies.^{25,28,35,36} Motivation for spectacle independence is
38
39 likely to be an important factor in these disparate observations; given that in the present study,
40
41 participants attended for cataract removal rather than for a specific refractive outcome.
42
43 Individuals with a prior motivation to be spectacle independent are more likely to tolerate near
44
45 and intermediate blur and hence comparability between studies can be limited.
46
47
48
49

50
51
52 Only 5% of the monofocal group were found to be spectacle independent with 30% requiring
53
54 constant correction and the remaining 65% occasionally wearing spectacles. A disparity between
55
56 the type of spectacles worn was evident between groups, with 35% of subjects implanted with
57
58 monofocal IOLs wearing either bifocals or varifocals post-operatively when compared to just 3%
59
60
61
62
63
64
65

1
2
3
4 of the MIOL group. It is important to note that overall satisfaction of distance vision was similar
5
6
7 in both groups whilst satisfaction of near and intermediate vision was considerably greater in the
8
9
10 MIOL group with 95% of subjects satisfied.

11
12
13 Unaided near visual acuity is demonstrably improved with the Bi-Flex MY IOL with greater
14
15
16 spectacle independence. With regard to visual acuity measures, it must be noted that this study
17
18
19 aimed to compare the MIOL and monofocal IOL using a standardised method, with specific
20
21
22 lighting levels and working distances for near and intermediate. Limitations in visual performance
23
24
25 due to halos, glare and reduction in contrast were evident amongst the MIOL group, and although
26
27
28 statistically significant, they do not appear to limit the subject's visual function nor their
29
30
31 perception of vision and overall satisfaction. Thus, the study concludes that the Bi-Flex MY
32
33
34 multifocal IOL demonstrates efficacy for the correction of near and distance vision and is
35
36
37 indicated when improved near vision/spectacle independence is required.

38 **WHAT WAS KNOWN**

39
40 Multifocal IOLs provide both distance and near vision whereas monofocal IOLs provide image
41
42
43 quality at a single distance

44
45 Multifocal IOLs cause an increased prevalence of dysphotopsia and result in reduced retinal
46
47
48 image contrast

49
50 A new Biconvex, aspheric, apodized, diffractive MIOL with a +3.50D add has been designed with
51
52
53 a relatively low number of diffractive echelons aimed to improving the optical image quality of
54
55
56 the resultant image.

57 **WHAT THIS PAPER ADDS**

58
59
60
61
62
63
64
65

1
2
3
4 Distance visual acuity was comparable between the MIOL and IOL. Contrast sensitivity was
5 reduced 3-6 months post-operatively whereas by months 12-18 were similar at all but low spatial
6 frequencies.
7
8

9
10
11 Near vision was superior in the MIOL group and subjects in the MIOL group were more satisfied
12 with the quality of vision at near and intermediate.
13
14

15
16 There was a statistically significant increase in the presence of dysphotopisa in the MIOL group,
17 however, satisfaction with distance vision was high in both groups
18
19
20
21
22

23 REFERENCES

- 24
25
26 1. Alio JL, Kaymak H, Breyer D, Cochener B, Plaza-Puche AB. Quality of life related variables
27 measured for three multifocal diffractive intraocular lenses: a prospective randomised
28 clinical trial. *Clin Exp Ophthalmol*. 2018;46(4):380-388.
29
30
- 31
32 2. Alio JL, Plaza-Puche AB, Fernandez-Buenaga R, Pikkell J, Maldonado M. Multifocal
33 intraocular lenses: An overview. *Surv Ophthalmol*. 2017;62(5):611-634.
34
35
- 36 3. Greenstein S, Pineda R, 2nd. The Quest for Spectacle Independence: A Comparison of
37 Multifocal Intraocular Lens Implants and Pseudophakic Monovision for Patients with
38 Presbyopia. *Semin Ophthalmol*. 2017;32(1):111-115.
39
40
- 41 4. Alba-Bueno F, Garzon N, Vega F, Poyales F, Millan MS. Patient-Perceived and Laboratory-
42 Measured Halos Associated with Diffractive Bifocal and Trifocal Intraocular Lenses. *Curr*
43 *Eye Res*. 2018;43(1):35-42.
44
45
- 46 5. Vega F, Alba-Bueno F, Millan MS, Varon C, Gil MA, Buil JA. Halo and Through-Focus
47 Performance of Four Diffractive Multifocal Intraocular Lenses. *Invest Ophthalmol Vis Sci*.
48 2015;56(6):3967-3975.
49
50
51
- 52 6. de Silva SR, Evans JR, Kirthi V, Ziaei M, Leyland M. Multifocal versus monofocal intraocular
53 lenses after cataract extraction. *Cochrane Database Syst Rev*. 2016;12:CD003169.
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4 7. Grzybowski A, Kanclerz P, Muzyka-Wozniak M. Methods for evaluating quality of life and
5 vision in patients undergoing lens refractive surgery. *Graefes Arch Clin Exp Ophthalmol.*
6 2019.
7
8
- 9
10 8. Maxwell A, Holland E, Cibik L, et al. Clinical and patient-reported outcomes of bilateral
11 implantation of a +2.5 diopter multifocal intraocular lens. *J Cataract Refract Surg.*
12 2017;43(1):29-41.
13
14
- 15 9. Ferris FL, 3rd, Kassoff A, Bresnick GH, Bailey I. New visual acuity charts for clinical
16 research. *Am J Ophthalmol.* 1982;94(1):91-96.
17
18
- 19 10. Hazel CA, Elliott DB. The dependency of logMAR visual acuity measurements on chart
20 design and scoring rule. *Optom Vis Sci.* 2002;79(12):788-792.
21
22
- 23 11. Shah N, Laidlaw DA, Brown G, Robson C. Effect of letter separation on computerised visual
24 acuity measurements: comparison with the gold standard Early Treatment Diabetic
25 Retinopathy Study (ETDRS) chart. *Ophthalmic Physiol Opt.* 2010;30(2):200-203.
26
27
- 28 12. Rosser DA, Murdoch IE, Fitzke FW, Laidlaw DA. Improving on ETDRS acuities: design and
29 results for a computerised thresholding device. *Eye (Lond).* 2003;17(6):701-706.
30
31
- 32 13. Williams MA, Moutray TN, Jackson AJ. Uniformity of visual acuity measures in published
33 studies. *Invest Ophthalmol Vis Sci.* 2008;49(10):4321-4327.
34
35
- 36 14. Wolffsohn JS, Jinabhai AN, Kingsnorth A, et al. Exploring the optimum step size for defocus
37 curves. *J Cataract Refract Surg.* 2013;39(6):873-880.
38
39
- 40 15. Gupta N, Wolffsohn JS, Naroo SA. Optimizing measurement of subjective amplitude of
41 accommodation with defocus curves. *J Cataract Refract Surg.* 2008;34(8):1329-1338.
42
43
- 44 16. Radner W, Diendorfer G, Kainrath B, Kollmitzer C. The accuracy of reading speed
45 measurement by stopwatch versus measurement with an automated computer program
46 (rad-rd(c)). *Acta Ophthalmol.* 2017;95(2):211-216.
47
48
- 49 17. Law EM, Aggarwal RK, Kasaby H. Clinical outcomes with a new trifocal intraocular lens.
50 *Eur J Ophthalmol.* 2014;24(4):501-508.
51
52
- 53 18. Buckhurst PJ, Wolffsohn JS, Gupta N, Naroo SA, Davies LN, Shah S. Development of a
54 questionnaire to assess the relative subjective benefits of presbyopia correction. *J*
55 *Cataract Refract Surg.* 2012;38(1):74-79.
56
57
58
59
60
61
62
63
64
65

19. Ray WA, O'Day DM. Statistical analysis of multi-eye data in ophthalmic research. *Invest Ophthalmol Vis Sci.* 1985;26(8):1186-1188.
20. Buckhurst PJ, Wolffsohn JS, Naroo SA, et al. Multifocal intraocular lens differentiation using defocus curves. *Invest Ophthalmol Vis Sci.* 2012;53(7):3920-3926.
21. Thibos LN, Wheeler W, Horner D. Power vectors: an application of Fourier analysis to the description and statistical analysis of refractive error. *Optom Vis Sci.* 1997;74(6):367-375.
22. Wolffsohn JS, Bhogal G, Shah S. Effect of uncorrected astigmatism on vision. *J Cataract Refract Surg.* 2011;37(3):454-460.
23. Bron AM, Viswanathan AC, Thelen U, et al. International vision requirements for driver licensing and disability pensions: using a milestone approach in characterization of progressive eye disease. *Clin Ophthalmol.* 2010;4:1361-1369.
24. Rees GB. Vision standards for driving: what ophthalmologists need to know. *Eye (Lond).* 2015;29(6):719-720.
25. Cochener B, Arnould B, Viala M, Roborel de Climens A, Berdeaux G. Corrected and uncorrected near and distance vision with ReSTOR compared to monofocal intraocular lens implantation after cataract surgery: a pooled analysis. *Ophthalmologica.* 2009;223(2):128-135.
26. Ji J, Huang X, Fan X, Luo M. Visual performance of Acrysof ReSTOR compared with a monofocal intraocular lens following implantation in cataract surgery. *Exp Ther Med.* 2013;5(1):277-281.
27. Garcia-Bella J, Ventura-Abreu N, Morales-Fernandez L, et al. Visual outcomes after progressive apodized diffractive intraocular lens implantation. *Eur J Ophthalmol.* 2018;28(3):282-286.
28. Cillino S, Casuccio A, Di Pace F, et al. One-year outcomes with new-generation multifocal intraocular lenses. *Ophthalmology.* 2008;115(9):1508-1516.
29. Zhao G, Zhang J, Zhou Y, Hu L, Che C, Jiang N. Visual function after monocular implantation of apodized diffractive multifocal or single-piece monofocal intraocular lens Randomized prospective comparison. *J Cataract Refract Surg.* 2010;36(2):282-285.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
30. Wilkins MR, Allan BD, Rubin GS, et al. Randomized trial of multifocal intraocular lenses versus monovision after bilateral cataract surgery. *Ophthalmology*. 2013;120(12):2449-2455 e2441.
 31. Pedrotti E, Carones F, Aiello F, et al. Comparative analysis of visual outcomes with 4 intraocular lenses: Monofocal, multifocal, and extended range of vision. *J Cataract Refract Surg*. 2018;44(2):156-167.
 32. Harman FE, Maling S, Kampougeris G, et al. Comparing the 1CU accommodative, multifocal, and monofocal intraocular lenses: a randomized trial. *Ophthalmology*. 2008;115(6):993-1001 e1002.
 33. Kamlesh, Dadeya S, Kaushik S. Contrast sensitivity and depth of focus with aspheric multifocal versus conventional monofocal intraocular lens. *Can J Ophthalmol*. 2001;36(4):197-201.
 34. Hayashi K, Manabe S, Hayashi H. Visual acuity from far to near and contrast sensitivity in eyes with a diffractive multifocal intraocular lens with a low addition power. *J Cataract Refract Surg*. 2009;35(12):2070-2076.
 35. Baig R, T AC, Kukreja S, Shakil S, Ahmad K. Patients' satisfaction and spectacle independence after cataract surgery with multifocal intraocular lens implantation in a tertiary care hospital. *J Pak Med Assoc*. 2016;66(6):745-747.
 36. Mendicute J, Kapp A, Levy P, et al. Evaluation of visual outcomes and patient satisfaction after implantation of a diffractive trifocal intraocular lens. *J Cataract Refract Surg*. 2016;42(2):203-210.

45
46 **Supplementary Table 1:** Inclusion/Exclusion Criteria

47
48
49 **Supplementary Table 2:** Characteristics of the Intraocular lenses

50
51
52 **Supplementary Table 3:** Patient Demographics

53
54
55 **Supplementary Table 4:** Refraction

56
57
58 **Supplementary Table 5:** Visual Acuity Results

59
60
61
62
63
64
65

1
2
3
4 **Supplementary Table 6: Area Under Defocus**
5

6
7 **Supplementary Table 7: CSV-1000**
8
9

10 **Figure 1: Trial Profile**
11

12
13 **Figure 2: a) Visit 2 Monocular Visual Acuity, b) Visit 2 Binocular Visual Acuity**
14
15

16
17 **Figure 3: a) Visit 2 Monocular Defocus Curve, b) Visit 2 Binocular Defocus Curve**
18
19

20 **Figure 4: a) Visit 2 Area under defocus curve, b) Visit 2 Range of focus**
21
22

23
24 **Figure 5: a) Visit 2 Maximum Reading Speed, b) Visit 2 Reading Acuity, c) Visit 2 95% Critical Print**
25
26
27 Size
28

29
30 **Figure 6: a) Visit 2 Monocular Contrast Sensitivity, b) Visit 2 Binocular Contrast Sensitivity**
31
32

33 **Figure 7: Visit 2 CSV-1000 Contrast sensitivity**
34
35

36
37 **Figure 8: Spectacle Wear a) Visit 2 Frequency of wear, b) Visit 2 Type of spectacles, c) Pre-Op**
38
39 spectacle wear
40
41

42
43 **Figure 9: Quality of Vision Questionnaire results a) Visit 2 Satisfaction, b) Visit 2 Near Tasks, c)**
44
45 Visit 2 Everyday Tasks, d) Visit 2 Night Vision
46
47

48
49 **Figure 10: Visit 2 Glare Simulator Scores**
50
51

52 **Figure 11: Visit 2 NAVQ Scores**
53
54

55 **Supplementary Figure 1: a) Visit 1 Monocular Visual Acuity, b) Visit 1 Binocular Visual Acuity**
56
57

58
59 **Supplementary Figure 2: a) Visit 1 Monocular Defocus Curve, b) Visit 1 Binocular Defocus Curve**
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Supplementary Figure 3: a) Visit 1 Area under defocus curve, b) Visit 1 Range of focus

Supplementary Figure 4: a) Visit 1 Maximum Reading Speed, b) Visit 1 Reading Acuity, c) Visit 1
95% Critical Print Size

Supplementary Figure 5: a) Visit 1 Monocular Contrast Sensitivity, b) Visit 1 Binocular Contrast
Sensitivity

Supplementary Figure 6: Visit 1 CSV-1000 Contrast sensitivity


Supplementary Figure 7: Spectacle Wear a) Visit 1 Frequency of wear, b) Visit 1 Type of
spectacles


Supplementary Figure 8: Quality of Vision Questionnaire results a) Visit 1 Satisfaction, b) Visit 1
Near Tasks c) Visit 1 Everyday Tasks, d) Visit 1 Night Vision


Supplementary Figure 9: Visit 1 Glare Simulator Scores


Supplementary Figure 10: Visit 1 NAVQ Score


Randomised control trial comparing visual and subjective satisfaction with monofocal and multifocal intraocular lenses (MIOLs). Results demonstrated greater near vision and satisfaction with MIOLs and equivocal distance acuity and satisfaction




























—▽— Multifocal
—▲— Monofocal




■ Monofocal
▨ Multifocal




■ Monofocal
▨ Multifocal








Click here to access/download
Supplementary Material
Supplimentary Table 1.docx





Click here to access/download
Supplementary Material
Supplimentary Table 2.docx






Click here to access/download
Supplementary Material
Supplimentary Table 3.docx





Click here to access/download
Supplementary Material
Supplimentary Table 4.docx






Click here to access/download
Supplementary Material
Supplimentary Table 5.docx






Click here to access/download
Supplementary Material
Supplimentary Table 6.docx





Click here to access/download
Supplementary Material
Supplimentary Table 7.docx







Click here to access/download
Supplementary Material
Supplimentary Figure 2.TIF











Click here to access/download
Supplementary Material
Supplimentary Figure 6.TIF







Click here to access/download
Supplementary Material
Supplimentary Figure 8.TIF







Click here to access/download
Supplementary Material
Supplimentary Figure 10.TIF



Journal of Cataract and Refractive Surgery Authorship Contribution Form

The Journal of Cataract and Refractive Surgery follows the Uniform Requirements set out by the International Committee of Medical Journal Editors (www.icmje.org) for authorship. **To qualify for authorship, each author must make a substantial contribution to the intellectual content of the manuscript in each of the following categories:**

- Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; **AND**
- Drafting the work or revising it critically for important intellectual content; **AND**
- Final approval of the version to be published; **AND**
- Agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

The corresponding author is responsible for ensuring that all coauthors meet the requirements for authorship. Each author agrees that the corresponding author will be responsible for the submission of the manuscript to the Journal and any associated activities. By submitting this manuscript, each of the authors indicates that he or she has had full access to all data in this study and takes complete and public responsibility for the integrity of the data and the accuracy of the data analysis. The issue of authorship must be resolved before submission of the manuscript. **All authors must sign this form, confirming he or she has made the contributions listed in the chart below.**

Please list all the manuscript authors and their contribution in the Contribution Table below. This form should be submitted with the original submission.

Article Title: Visual Function and Subjective Perception of Vision following bilateral implantation of monofocal and multifocal intraocular lenses: A Randomised Controlled Trial

I have made substantive intellectual contributions to the content of this manuscript in the following areas:

Author Name	Concept and design	Data acquisition	Data analysis / interpretation	Drafting manuscript	Critical revision of manuscript	Statistical analysis	Securing funding	Admin, technical or material support	Supervision	Final approval
Elizabeth M. Law	Y	Y	Y	Y	Y	Y	Y	Y		Y
Rajesh K. Aggarwa	Y	Y	Y		Y		Y	Y		Y
Hetal Buckhurst	Y		Y	Y	Y	Y	Y	Y	Y	Y
Hosam E. Kasaby	Y	Y	Y		Y			Y		Y
Jonathan Marsden	Y				Y	Y			Y	Y
Gary Shum	Y		Y		Y				Y	Y
Phillip J. Buckhurst	Y		Y	Y	Y	Y	Y	Y	Y	Y

Other contributions:

(Please have all authors sign and date.)

1. Corresponding author Phillip J. Buckhurst

Signature of **corresponding** author: Phillip Buckhurst Digitally signed by Phillip Buckhurst
Date: 2019.11.20 22:43:20 Z Date Completed: 19/11/2019

2. Author Elizabeth M. Law

Signature of author: Elizabeth Law Digitally signed by Elizabeth Law
Date: 2019.11.20 20:10:44 Z Date Completed: 20/11/2019


3. Author Rajesh K. Aggarwal

Signature of author: Rajesh Aggarwal Digitally signed by Rajesh Aggarwal
Date: 2019.11.21 07:59:16 Z Date Completed: 20/11/2019

4. Author Hetal Buckhurst

Signature of author: Hetal Buckhurst Digitally signed by Hetal Buckhurst
Date: 2019.11.21 21:37:46 Z Date Completed: 20/11/2019


5. Author Hosam E. Kasaby

Signature of author:  Date Completed: 20/11/2019

6 Author Jonathan Marsden

Signature of author: Jonathan Marsden Digitally signed by Jonathan Marsden
Date: 2019.11.21 12:24:45 Z Date Completed: 20/11/2019

7. Author Gary Shum

Signature of author:  Digitally signed by Gary Shum
Date: 2019.11.21 14:31:17 Z Date Completed: 20/11/2019

8 Author _____

Signature of author: _____ Date Completed: _____

9. Author _____

Signature of author: _____ Date Completed: _____

10. Author _____

Signature of author: _____ Date Completed: _____

11. Author _____

Signature of author: _____ Date Completed: _____

12. Author _____

Signature of author: _____ Date Completed: _____

If additional author and signature fields are needed, please duplicate this form as needed.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.