

1 **Deficiency of GDP-L-galactose phosphorylase, an enzyme required for ascorbic acid**
2 **synthesis, reduces tomato fruit yield.**

3

4 Matías L Alegre^{1*}, Charlotte Steelheart^{1*}, Pierre Baldet², Christophe Rothan², Daniel
5 Just², Yoshihiro Okabe^{3,4}, Hiroshi Ezura^{3,4}, Nicholas Smirnoff⁵, Gustavo E Gergoff
6 Grozeff¹, Carlos G Bartoli¹

7

8 ¹ INFIVE, Facultades de Ciencias Agrarias y Forestales y Ciencias Naturales,
9 Universidad Nacional de La Plata-CCT CONICET La Plata, Argentina.

10 ² Institut National de la Recherche Agronomique (INRA), Université de Bordeaux,
11 UMR 1332 Biologie du Fruit et Pathologie, F-33140 Villenave d'Ornon, France.

12 ³ Faculty of Life and Environmental Sciences, University of Tsukuba, Tsukuba, Ibaraki
13 305-8572 Japan.

14 ⁴ Tsukuba Plant Innovation Research Center, University of Tsukuba, Tsukuba, Ibaraki
15 305-8572 Japan.

16 ⁵ Biosciences, College of Life and Environmental Sciences, University of Exeter, Exeter
17 EX4 4QD, UK.

18

19 *Equally contributed to this work.

20

21 **Corresponding author:**

22

Carlos G Bartoli

23

e-mail: carlos.bartoli@agro.unlp.edu.ar

24

Telephone number: 0054 221 4233698

25

26 **Main conclusion:**

27 Reduced GDP-L-galactose phosphorylase expression and deficiency of ascorbic acid
28 content lead to decreased fruit set and yield in tomato plants.

29

30 **Author Contribution Statement**

31 MA, MCS, GEGG, PB, DJ, OY, EH carried out the experiments and made the statistical
32 analysis of the data. PB, CR, NS and CGB wrote de manuscript. All authors read and
33 approved the manuscript.

34

35 **Abstract**

36 GDP-L-galactose phosphorylase (GGP) catalyzes the first step committed to
37 ascorbic acid synthesis. The participation of GDP-L-galactose phosphorylase and
38 ascorbate in tomato fruit production and quality was studied in this work using two
39 *S/GGP1* deficient EMS Micro-Tom mutants. The *S/GGP1* mutants display decreased
40 concentrations of ascorbate in roots, leaves, flowers and fruit. The initiation of anthesis
41 is delayed in *ggp1* plants but the number of flowers is similar to wild type. The number
42 of fruits is reduced in *ggp1* mutants with an increased individual weight. However, the
43 whole fruit biomass accumulation is reduced in both mutant lines. Fruits of the *ggp1*
44 plants produce more ethylene and show higher firmness and soluble solids content than
45 the wild type after the breaker stage. Leaf CO₂ uptake decreases about 50 % in both
46 *ggp1* mutants at saturating light conditions; however, O₂ production in an enriched CO₂
47 atmosphere is only 19 % higher in wild type leaves. Leaf conductance that is largely
48 reduced in both mutants may be the main limitation for photosynthesis. Sink-source
49 assays and hormone concentration were measured to determine restrictions to fruit
50 yield. Manipulation of leaf area/fruit number relationship demonstrates that the number
51 of fruits and not the provision of photoassimilates from the source restricts biomass
52 accumulation in the *ggp1* lines. The lower gibberellins concentration measured in the
53 flowers would contribute to the lower fruit set, thus impacting in tomato yield. Taken as
54 a whole these results demonstrate that ascorbate biosynthetic pathway critically
55 participates in tomato development and fruit production.

56

57 **Keywords:** antioxidant, ascorbate, fruit, GGP, ripening, tomato, yield.

58 **Introduction**

59

60 Ascorbate is one of the most abundant compounds in plants and there is great
61 interest in its multiple functions as an antioxidant and enzyme cofactor (Foyer and
62 Noctor 2011; Smirnov 2018). It is synthesized *via* GDP-mannose and GDP-L-galactose
63 and the first enzyme in this pathway considered to be specific to ascorbate synthesis is
64 GDP-L-galactose phosphorylase (Dowdle et al. 2007; Laing et al. 2007, Linster et al.
65 2007). It is encoded by paralogues in various species, including arabidopsis (*VTC2* and
66 *VTC5*). Double *vtc2 vtc5* mutants which are unable to make ascorbate are not viable but
67 can be rescued by ascorbate supplementation (Dowdle et al. 2007; Lim et al. 2016). A
68 range of other *vtc* mutants and transgenic plants in different parts of the ascorbate
69 biosynthesis pathway with 10-20% of wild type ascorbate concentrations grow
70 relatively normally but exhibit various subtle developmental changes, increased
71 sensitivity to environmental stresses and increased basal resistance to pathogens (Pavet
72 et al. 2005; Barth et al. 2006; Senn et al. 2016; Caviglia et al. 2018; Plumb et al. 2018).
73 Therefore, it is apparent that relatively severe decreases in ascorbate still enable its
74 essential functions while higher concentrations must be assumed to be beneficial.
75 Notably, high light intensity increases ascorbate concentration in leaves, associated with
76 its role in removal of hydrogen peroxide and in photoprotection (Asada 1999; Bartoli et
77 al. 2006). GDP-L-galactose phosphorylase expression is strongly controlled by light and
78 repressed by high ascorbate in part *via* a conserved upstream open reading frame
79 (uORF) in the 5'-UTR. This, along with over-expression experiments, strongly supports
80 its role in controlling ascorbate biosynthesis (Dowdle et al. 2007; Gao et al. 2011;
81 Yoshimura et al. 2014; Laing et al. 2015; Macknight et al. 2017; Li et al. 2018).

82 Tomatoes are of interest as a source of ascorbate in the diet and control of its
83 synthesis and functions have been investigated by altering expression of various
84 biosynthesis genes (Alhaghdow et al. 2007; Gilbert et al. 2009; Gilbert et al. 2016). The
85 availability of tomato plants with GDP-L-galactose phosphorylase deficiency enables the
86 role of this enzyme and ascorbate in fruit production and quality (Baldet et al. 2013).
87 Two *GGP* genes encode GDP-L-galactose phosphorylase in tomato with complementary
88 function, and *SIGGP1* is about thousand time more expressed than *SIGGP2* (Massot et
89 al., 2012). The *Slggp1* mutant, although expressing *GGP2*, had low ascorbate

90 concentration in its leaves. When this *Slggp1* mutant was submitted to high irradiance
91 conditions chlorophyll bleaching was observed (Baldet et al. 2013). In addition,
92 transformed tomato with decreased GDP-L-galactose phosphorylase expression display
93 increased damage when exposed to chilling (Wang et al. 2013; Yang et al. 2017). These
94 results provide evidence for the increased susceptibility of GDP-L-galactose
95 phosphorylase deficient tomato plants to stress. However, studies focused in
96 modifications at the level of the fruit have not been done yet. The work focuses on the
97 effects of GDP-L-galactose phosphorylase expression and associated ascorbate
98 deficiency on tomato fruit yield and quality.
99

100 **Material and Methods**

101

102 The experiments were carried out with *Solanum lycopersicum* L cv Micro-Tom
103 plants with two lines deficient in expression of the *GGP1* gene encoding GDP-L-
104 galactose phosphorylase. The two EMS mutant Micro-Tom lines used here, GGP-5261
105 and GGP-49C12, were respectively from the NBRP-Tomato population (Tsukuba-
106 Japan) and TILLING-Tomato collection (Bordeaux-France). They are truncation and
107 splice junction mutants respectively (Baldet et al. 2013). Plants were grown
108 hydroponically in an air-conditioned greenhouse during spring and summer seasons
109 under an irradiance of 700 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ at midday and temperatures average of
110 25 ± 2 and 20 ± 3 °C during the day and night, respectively. The final harvest was taken
111 when half mutant plants were bearing the first fully red ripe fruit. Each experiment
112 included 10 plants of each genotype and the harvest was taken when half mutant plants
113 were bearing the first fully red ripe fruit.

114 Exogenous treatment of ascorbic acid was initiated when seedlings were twenty
115 days-old and finished after the development of the fourth inflorescence. Each plant
116 received 1 mL of 20 mM ascorbic acid solution (including 0.01% tween 20 as a
117 surfactant) wetting all above ground organs (Mainly in the adaxial side of the leaves).
118 The treatment was repeated four times a week.

119

120 **Sink-source experiment**

121 A sink-source experiment was performed leaving only 4 inflorescences on each
122 plant. The treatments consisted of plants with one (1F) or two (2F) tomatoes per
123 inflorescence, two levels of leaf pruning (–L and =L, for 50 and 75 % of leaf area
124 removal, respectively) and a control without organ removal (leaving only 4
125 inflorescences). Three independent experiments were carried out including at least five
126 plants per treatment for each genotype (i.e. 75 plants for each experiment).
127 Measurements were made when half mutant plants got at least one red fruit.

128

129 **Ascorbate determination**

130 Ascorbate was determined with a HPLC system (Shimadzu LC-10Atvp solvent
131 delivery module and Shimadzu UV-Vis SPD-10Avp detector) as previously described

132 (Bartoli et al. 2006). Root, leaf, flower and fruit tissues were ground in 6 % (v/v)
133 trifluoroacetic acid, centrifuged at 13000xg for 5 min and supernatants used for the
134 measurements. Total AA was determined after the treatment of an aliquot with 5 mM
135 dithiothreitol (DTT). Oxidized AA was calculated as the difference between samples
136 with or without DTT.

137

138 **Photosynthesis measurements**

139 CO₂ assimilation was measured in fully developed leaves with an infrared gas
140 analyzer (PLC 6, Cirrus-2 PPSsystems) at saturating irradiance (1200 μmol photon m⁻² s⁻¹,
141 *A_{max}*). In addition, photosynthesis was measured as O₂ evolution under saturating
142 irradiance and CO₂ concentration (*P_{pot}*). Leaf discs were placed in a gas tight chamber
143 equipped with a Clark type electrode (Hansatech, UK). Saturating CO₂ atmosphere was
144 generated including a mate imbibed with 1 M NaHCO₃ (Walker 1987). Photosynthetic
145 electron transport rate was determined with a modulated chlorophyll fluorescence
146 system (FMS-2, Hansatech Instruments Ltd., Norfolk, UK) and calculated according to
147 Genty et al. (1989). Chlorophyll fluorescence quenching analysis was carried out with a
148 CF Imager (Technologica Ltd., Colchester, UK) as described by Lim et al. (2016) and
149 the chlorophyll fluorescence parameters calculated according to Baker (2008). Leaf
150 temperature was measured with a thermographic camera (FLUKE Ti 400) with an
151 emissivity of 0.95. Measurements were taken to well watered plants exposed at 700
152 μmol photons m⁻² s⁻¹ during midday inside the greenhouse.

153

154 **Ripening parameters**

155 Fruit ethylene production was measured with a gas chromatograph system (Konik,
156 KNK-3000-HGRC) including an alumina column and a flame ionization detector as
157 previously reported (Bartoli et al. 1996). Firmness was measured in detached tomato
158 fruit with a texture analyzer (TA.XT.PLUS, Micro Systems TM Goldalming, Surrey,
159 UK) using a 2.5 mm diameter flat probe. The measurements were obtained by fruit
160 deformation for a distance of 0.5 mm at 0.25 mm s⁻¹ and 5.9 g trigger force. The
161 maximum force was recorded and results expressed in force g. Total soluble solids were
162 measured as previously described (Gergoff et al. 2016).

163

164 **Hormone determination**

165 Plant hormones were measured in whole flower tissues sampled at anthesis and
166 collected from several plants of each independent experiment. About 100 mg of
167 lyophilized tissues were added with 1% (v/v) AcH (40mg ml⁻¹) and ²H₅-indoleacetic
168 acid (IAA), ²H₂-gibberellin (GA), ²H₂-GA4, ²H₂-GA8, ²H₅-zeatin and ²H₆-ABA
169 (OlChemIm Ltd., Olomouc, Czech Republic) as internal standards. The aqueous
170 solution was partitioned 3 times with ethyl acetate at pH 3. Organic fractions were
171 combined, evaporated and then resuspended in methanol for hormone determination by
172 liquid chromatography-mass spectrometry with electrospray ionization (Waters Corp.,
173 New York, NY, USA). An Alliance 2695 (Separation Module, Waters, USA)
174 quaternary pump equipped with a Restek C18 column (2.1 x 100mm) (Restek, USA)
175 was used to analyze the samples. A binary solvent system used for elution consisted of
176 0.2% (v/v) acetic acid in H₂O and methanol. MS/MS assays were done with a
177 Micromass Quatro Ultima TM mass spectrometer (Micromass, Manchester City, UK)
178 as described by Masciarelli et al. (2014).

179

180 **Results**

181

182 **Growth, development and photosynthesis in GDP-L-galactose phosphorylase**
183 **mutants.**

184

185 The two GDP-L-galactose phosphorylase mutants had 34-50% of wild type
186 ascorbate in their leaves, roots, flowers and fruit (Fig. 1a-d). Ascorbate concentration
187 was lower in roots and the proportion of DHA was much higher than in leaves, flowers
188 and fruit. Exogenous ascorbate increased its concentration in leaves of the mutants by
189 66 to 79 % but not in wild type plants (Table 1).

190 The vegetative biomass and leaf area at harvest were similar in all genotypes
191 (Table 2). However, the mutants both allocated less biomass to roots (15% compared to
192 23% in wild type) and more to stems (20% compared to 13% in wild type). Therefore,
193 the mutants were visibly different since they had larger internodes (Supplementary Fig.
194 S1). The fruit fresh weight per plant was decreased in both mutants but the individual
195 fruit weight was increased (Table 2). The mutants had a similar number of flowers but
196 decreased fruit setting, and consequently, lower number of tomatoes than those of the
197 wild type. Exogenous ascorbate supplementation increased fruit setting in both mutants
198 but not in wild type plants (Table 1). In addition, anthesis was delayed in both mutants.
199 Impairment of fruit setting may be the consequence of altered hormone concentrations.
200 Consequently, the concentrations of various hormones in flowers at anthesis were
201 measured (Table 3). All three gibberellins measured were decreased in the mutants but
202 indole acetic acid (IAA), zeatin and ABA showed no differences.

203 The CO₂ assimilation and electron transport rates (ETR) of the mutants were 50
204 and 90 %, of wild type respectively (Fig. 2a, b). In addition, photosynthetic O₂
205 production under a saturating CO₂ atmosphere, was 19 % higher in wild type than in the
206 mutants (Fig 2c). Consistent with the small effect on ETR and O₂ assimilation under
207 saturating CO₂, there was no significant effect on photochemical quenching (qP), PSII
208 maximum operating quantum efficiency (Fv'/Fm') and PSII operating efficiency
209 (Fq'/Fm'; φPSII) (Fig. 3). Non photochemical quenching (NPQ), which is impaired in
210 ascorbate deficient arabidopsis mutants (Müller-Moule et al. 2002) was not affected in
211 the mutants. Stomatal conductance was 60 % higher in wild type than in the mutants

212 (Fig 2d). Consequently, leaf temperature measured by infra-red thermal imaging was
213 1.8 to 2.5°C higher in the mutants (Supplementary Fig. S2).

214

215 **The effect of GDP-L-galactose phosphorylase mutation on fruit quality**

216

217 Fruit firmness decreased along ripening but both GDP-L-galactose phosphorylase
218 mutants kept higher firmness than wild type at the red stage (Fig. 4a). Ethylene
219 production was higher in the mutant lines than wild type fruit at the breaker and red
220 stages (Fig. 4b). Soluble solids content (Brix°) increased during fruit ripening and was
221 also higher in mutants compared with wild type at breaker stage (Fig. 4c). Other
222 ripening parameters such as pH and titratable acidity show no differences between
223 genotypes (Data not shown).

224

225 **The effect of source-sink manipulation on fruit yield characteristics**

226

227 To investigate the relative importance of source or sink limitations on fruit yield
228 various fruit or leaf pruning treatments were performed (Table 4). Fruit production was
229 not modified in wild type or mutant plants after leaf removal. However, when plants
230 were limited to two tomatoes per inflorescence (i.e. eight tomatoes per plant) the
231 mutants kept their fruit weight per plant as in control treatment but it drastically
232 decreased in all genotypes when only one fruit per inflorescence is maintained. For an
233 easier interpretation of these results a principal component analysis is shown in Figure
234 5. It shows a contrasting relationship between different traits such as fruit number and
235 yield *versus* individual fruit fresh weight. Component 1 distinguishes treatments
236 independently of the genotypes and component 2 the wild type from mutants (more
237 markedly for fruit removal treatment).

238

239

240 Discussion

241

242 Arabidopsis *vtc2* mutants, which are knockouts of one of the two genes encoding
243 GDP-L-galactose phosphorylase, have ~20% of wild type ascorbate (Dowdle et al. 2007;
244 Barth et al. 2010) and have small decreases in rosette leaf area and biomass, the extent
245 most likely being sensitive to growth conditions (Lim et al. 2016; Caviglia et al. 2018;
246 Plumb et al. 2018). Here, the tomato GDP-L-galactose phosphorylase mutants, which
247 have 34-50% of wild type ascorbate in all organs, are similarly unaffected in total
248 vegetative biomass but do have reduced allocation to roots. The reason for the effect on
249 root growth is not evident but could be associated with re-allocation to shoots to
250 compensate for decreased CO₂ assimilation. Effects of the GDP-L-galactose
251 phosphorylase mutations are more evident in reproductive development, the mutants
252 being slower to anthesis and having markedly decreased fruit set and number. However,
253 early flowering has been reported in various ascorbate deficient Arabidopsis mutants
254 (Kotchoni et al. 2009). Whatever the delaying or accelerating effect, these results
255 together suggest that the plant ascorbic acid concentration is linked to the regulation of
256 time to flowering. This effect may be a species- but also an environmental-dependent
257 phenomenon. The decrease in fruit number is associated with increased fruit size in the
258 mutants. Larger fruit size is most easily explained by decreased competition between
259 fruits for assimilate, since the fruit removal experiment increased individual fruit weight
260 in all genotypes. Critically, ascorbate supplementation increased fruit set in the mutants,
261 suggesting that ascorbate is important for this process. This observation also suggests
262 that assimilate limitation (Ruan et al. 2012) may be only partially responsible for
263 reduced fruit set in this case. To further investigate the cause of reduced fruit set, the
264 effect of GDP-L-galactose phosphorylase mutations on flower hormones was
265 investigated. GA1, GA4 and GA8 were decreased in the mutants. The chemical
266 inhibition of gibberellin synthesis in tomato reduces fruit setting and this effect is
267 reversed by exogenous application of the hormone (Serrani et al. 2007). Therefore, the
268 low GA concentration in the flowers of the mutants could contribute to low fruit setting.
269 Ethylene treatment decreases fruit set in tomato, possibly by inhibiting GA synthesis
270 (Shinozaki et al. 2015). Arabidopsis *vtc2* mutants have increased ethylene production

271 (Caviglia et al. 2018). Therefore, if ascorbate deficiency also increases ethylene in
272 tomato, it is possible that this could be the cause of decreased GA and fruit setting.

273

274 The large decrease in CO₂ assimilation in the mutants was not matched by a large
275 decrease in biomass, although overall fruit yield was significantly less (~72% of wild
276 type). As noted above this could be partly attributed to lower fruit set. Measuring
277 photosynthesis under high CO₂ and by chlorophyll fluorescence showed that the
278 mutations do not limit photosynthetic capacity, but the limitation is most likely caused
279 by partial stomatal closure. Since the stomatal conductance was measured around
280 midday, it is possible that over the course of the light period, stomata are more open in
281 morning/evening, thus minimizing an overall effect on assimilation. Furthermore,
282 decreasing GDP-L-galactose phosphorylase expression in tomato in another study also
283 decreased ascorbate content by 50% while having no effect on CO₂ assimilation under
284 non-stressed conditions (Wang et al. 2013). Similarly, *Arabidopsis vtc2-1* has a similar
285 CO₂ assimilation rate to wild type (Senn et al. 2016). Therefore, studies to date show
286 that ascorbate deficiency (at least to 20% of wild type), has minimal effect on
287 photosynthesis and vegetative biomass.

288 The source-sink experiment demonstrates that the number of fruit is crucial to
289 determine fruit yield in tomato and that the higher abortion of fruit observed in *ggp1*
290 mutants does not seem to be the consequence of an insufficient provision of photo-
291 assimilates by the leaves. Similarly, Tanaka and Fujita (1974) demonstrate in assays
292 partially removing fruit and leaves that source is not limiting yield in tomato and also
293 that fruit size has a limited flexibility. These authors conclude that unraveling the
294 processes establishing fruit number is an important task. In this context, the present
295 work, using GDP-L-galactose phosphorylase deficient mutants and especially exogenous
296 ascorbic acid supplementation, demonstrates that ascorbic acid concentration is an
297 important factor in establishing fruit number and consequently for the improvement of
298 tomato yield.

299

300 Ascorbic acid constitutes an important nutritional attribute of fruit (Lee and Kader
301 2000). However, the impact of decreased ascorbate concentration on other fruit
302 characteristics has scarcely been addressed. Previous reports on tomato mutants in

303 enzymes earlier in the mannose/L-galactose biosynthetic pathway (GDP-mannose
304 pyrophosphorylase and GDP-mannose-3',5-epimerase) show effects on fruit
305 development and quality (Gilbert et al. 2009; Zhang et al. 2013; Gilbert et al. 2016) but
306 these may be caused by effects on cell wall composition as well as ascorbate deficiency
307 itself. Decreased GDP-L-galactose phosphorylase activity in the current experiments is
308 likely to affect ascorbate synthesis more specifically. Among effects observed here in
309 the fruit of GDP-L-galactose phosphorylase mutants is an increase in ethylene
310 production. *Vtc2* arabidopsis plants also produce more ethylene (Caviglia et al. 2018).
311 Ethylene production is associated with a stimulation of cell wall degrading enzymes
312 during fruit ripening (Osorio et al. 2011). However, the results observed in GDP-L-
313 galactose phosphorylase mutants suggest that other processes prevail, leading to
314 increased fruit firmness. For example, changes in cuticles may also delay fruit softening
315 (Saladié et al. 2007). The modifications observed in GDP-L-galactose phosphorylase
316 tomato lines (including increased soluble solids) suggest that ascorbate influences some
317 aspects of fruit quality.

318

319 **Conclusions**

320 Tomato mutants in the ascorbate biosynthesis the *GGPI* isoform of GDP-L-
321 galactose phosphorylase contain 34-50 % of wild type ascorbate. The results suggest
322 that this decrease in GDP-L-galactose phosphorylase expression and ascorbate
323 concentration influence tomato fruit set, possibly *via* decreased GA concentration, final
324 fruit size and some aspects of quality. A higher flower abortion is directly caused by
325 ascorbate deficiency. Fruit size increases in compensation, but not sufficiently to
326 prevent a decrease in total fruit biomass. Supply of assimilates for biomass and fruit
327 production is unlikely to be a major limiting factor.

328

329 **Acknowledgments**

330 We are really grateful to Dr Eduardo Tambussi for his valuable discussion of the
331 results, Dr Gabriela Cano for her technical assistance and Dr Diego Fanello for the
332 statistical analysis. GGG and CGB are researchers at the Consejo Nacional de
333 Investigaciones Científicas y Técnicas (CONICET, Argentina). MA and CS are fellows
334 at CONICET and Agencia Nacional de Promoción Científica y Técnica (ANPCyT,
335 Argentina), respectively. This work was supported by grants PICT 2015-0103
336 (ANPCyT) and A258 and A322 (Universidad Nacional de La Plata, Argentina).

337

338 **Literature**

339 Alhaghdow M, Mounet F, Gilbert L, Nunes-Nesi A, Garcia V, Just D, Petit J,
340 Beauvoit B, Fernie AR, Rothan C, Baldet P (2007) Silencing of the mitochondrial
341 ascorbate synthesizing enzyme L-galactono-1,4-lactone dehydrogenase affects plant and
342 fruit development in tomato. *Plant Physiol* 145:1408-1422.

343 Asada K (1999) The water-water cycle in chloroplasts: Scavenging of active
344 oxygens and dissipation of excess photons. *Annu Rev Plant Phys* 50:601-39.

345 Baker NR (2008) Chlorophyll fluorescence: a probe for photosynthesis *in vivo*.
346 *Annu Rev Plant Biol* 59: 89-113.

347 Baldet P, Bres C, Okabe Y, Mauxion JP, Just D, Bournonville C, Ferrand C, Mori
348 K, Ezure H, Rothan C (2013) Investigating the role of vitamin C in tomato through
349 TILLING identification of ascorbate-deficient tomato mutants. *Plant Biotechnol*
350 30:308-314.

351 Barth C, De Tullio M, Conklin PL (2006) The role of ascorbic acid in the control
352 of flowering time and the onset of senescence. *J Exp Bot* 57:1657-1665.

353 Barth C, Gouzd ZA, Steele HP, Imperio RM (2010) A mutation in GDP-mannose
354 pyrophosphorylase causes condition hypersensitivity to ammonium, resulting in
355 *Arabidopsis* root growth inhibition, altered metabolism, and hormone homeostasis. *J*
356 *Exp Bot* 61:379–394.

357 Bartoli CG, Simontachi M, Montaldi E, Puntarulo S (1996) Oxidative stress,
358 antioxidant capacity and ethylene production during ageing of cut carnation (*Dianthus*
359 *caryophyllus*) petals. *J Exp Bot* 47:595-601.

360 Bartoli CG, Yu J, Gómez F, Fernández L, McIntosh L, Foyer CH (2006) Inter-
361 relationships between light and respiration in the control of ascorbic acid synthesis and
362 accumulation in *Arabidopsis thaliana* leaves. *J Exp Bot* 57:1621-1631.

363 Caviglia M, Mazorra Morales LM, Concellón A, Gergoff Grozeff GE, Wilson M,
364 Foyer CH, Bartoli CG (2018) Ethylene signaling triggered by low concentrations of
365 ascorbic acid regulates biomass accumulation in *Arabidopsis thaliana*. *Free Radical Bio*
366 *Med* 122:130-136.

367 Dowdle J, Ishikawa T, Gatzek S, Rolinski S, Smirnoff N (2007) Two genes in
368 *Arabidopsis thaliana* encoding GDP-L-galactose phosphorylase are required for
369 ascorbate biosynthesis and seedling viability. *Plant J* 52:673-689.

370 Foyer CH, Noctor CH (2011) Ascorbate and glutathione: The heart of the redox
371 hub. *Plant Physiol* 155:2-18.

372 Gao Y, Badejo AA, Shibata H, Sawa Y, Maruta T, Shigeoka S, Page M, Smirnoff
373 N, Ishikawa T (2011) Expression analysis of the VTC2 and VTC5 genes encoding
374 GDP-L-Galactose Phosphorylase, an enzyme involved in ascorbate biosynthesis, in
375 *Arabidopsis thaliana*. *Biosci Biotechnol Biochem* 75:1783-1788.

376 Genty B, Briantais JM, Baker NR (1989) The relationship between the quantum
377 yield of photosynthetic electron transport and quenching of chlorophyll fluorescence.
378 *Biochim Biophys Acta* 990:87-92.

379 Gergoff Grozeff GE, Senn ME, Alegre M, Chaves AR, Bartoli CG (2016)
380 Nocturnal low irradiance pulses improve fruit yield and lycopene concentration in
381 tomato. *Sci Hortic-Amsterdam* 203:47-52.

382 Gilbert L, Alhagdow M, Nunes-Nesi A, Quemener B, Guillon F, Bouchet B,
383 Faurobert M, Gouble B, Page D, Garcia V, Petit J, Stevens R, Causse M, Fernie AR,
384 Lahaye M, Rothan C and Baldet P. (2009) GDP-D-mannose epimerase (GME) plays a
385 key role at the intersection of ascorbate and non cellulosic cell wall biosynthesis in
386 tomato. *Plant J* 60, 499-508.

387 Gilbert L, Dumont M, Ferrand C, Bournonville C, Monier A, Jorly J, Lemaire-
388 Chamley M, Mori K, Atienza I, Hernould M, Stevens R, Lehner A, Mollet JC, Rothan
389 C, Lerouge P, Baldet P (2016) Two tomato GDP-D-mannose epimerase isoforms
390 involved in ascorbate biosynthesis play specific roles in cell wall biosynthesis and
391 development. *J Exp Bot* 67:4767-4777.

392 Kotchoni SO, Larrimore KE, Mukherjee M, Kempinski CF, Barth C (2009)
393 Alterations in the endogenous ascorbic acid content affect flowering time in
394 *Arabidopsis*. *Plant Physiol* 149, 803–815.

395 Laing WA, Wright MA, Cooney J, Bulley SM (2007) The missing step of the L-
396 galactose pathway of ascorbate biosynthesis in plants, an L-galactose guanyltransferase,
397 increases leaf ascorbate content. *P Natl Acad Sci USA* 104: 9534-9539.

398 Laing WA, Martinez-Sanchez M, Wright MA, Bulley SM, Brewster D, Dare AP,
399 Rassam M, Wang D, Storey R, Macknight RC, Hellens RP (2015) An Upstream Open
400 Reading Frame Is Essential for Feedback Regulation of Ascorbate Biosynthesis in
401 *Arabidopsis*. *Plant Cell* 27:772-786.

402 Lee SK, Kader AA (2000) Preharvest and postharvest factors influencing vitamin
403 C content of horticultural crops. *Postharvest Biol Tech* 20:207-220.

404 Li T, Yang X, Yu Y, Si X, Zhai X, Zhang H, Dong W, Gao C, Xu C (2018)
405 Domestication of wild tomato is accelerated by genome editing. *Nat Biotechnol*
406 36:1160-1163.

407 Lim B, Smirnoff N, Cobbett CS, Golz JF (2016) Ascorbate-deficient *vtc2* mutants
408 in *Arabidopsis* do not exhibit decreased growth. *Front Plant Sci* 7:1025-1030.

409 Linster C, Gomez TA, Christensen KC, Adler LN, Young BD, Brenner C, Clarke
410 SG (2007) *Arabidopsis* VTC2 encodes a GDP-L-galactose phosphorylase, the last
411 unknown enzyme in the Smirnoff-Wheeler pathway of ascorbic acid in plants. *J Biol*
412 *Chem* 282:18879-18885.

413 Macknight RC, Laing WA, Bulley SM, Broad RC, Johnson AAT, Hellens RP
414 (2017) Increasing ascorbate levels in crops to enhance human nutrition and plant abiotic
415 stress tolerance. *Curr Opin Biotech* 44:153–160.

416 Masciarelli O, Llanes A, Luna V (2014) A new PGPR co-inoculated with
417 *Bradyrhizobium japonicum* enhances soybean nodulation. *Microbiol Res* 169:609-615.

418 Massot C, Stevens R, Génard M, Longuenesse JJ, Gautier H (2012) Light affects
419 ascorbate content and ascorbate-related gene expression in tomato leaves more than in
420 fruits. *Planta* 235:153-163.

421 Müller-Moulé P, Conklin PL, Niyogi KK (2002) Ascorbate deficiency can limit
422 violaxanthin de-epoxidase activity in vivo. *Plant Physiol* 128:970–977.

423 Osorio S, Alba R, Damasceno CMB, Lopez-Casado G, Lohse M, Zanor MI,
424 Tohge T, Usadel B, Rose JKC, Fei Z, Giovannoni JJ, Fernie AR (2011) Systems
425 biology of tomato fruit development: Combined transcript, protein, and metabolite
426 analysis of tomato transcription factor (*nor*, *rin*) and ethylene receptor (*Nr*) mutants
427 reveals novel regulatory interactions. *Plant Physiol* 157:405-425.

428 Pavet V, Olmos E, Kiddle G, Mowla S, Kumar S, Antoniw J, Alvarez ME, Foyer
429 CH (2005) Ascorbic acid deficiency activates cell death and disease resistance
430 responses in *Arabidopsis*. *Plant Physiol* 139:1291-1303.

431 Plumb W, Townsend AJ, Rasool B, Alomrani S, Razak N, Karpinska B, Ruban
432 AV, Foyer CH (2018) Ascorbate-mediated regulation of growth, photoprotection, and
433 photoinhibition in *Arabidopsis thaliana*. *J Exp Bot* 69:2823-2835.

434 Ruan Y-L, Patrick JW, Bouzayen M, Osorio S, Fernie AR (2012) Molecular
435 regulation of seed and fruit set. *Trends Plant Sci* 17:656-665.

436 Saladié M, Matas AJ, Isaacson T, Jenks MA, Goodwin SM, Niklas KJ, Xiaolin R,
437 Labavitch JM, Shackel KA, Fernie AR, Lytovchenko A, O'Neill MA, Watkins CB,
438 Rose JKC (2007) A reevaluation of the key factors that influence tomato fruit softening
439 and integrity. *Plant Physiol* 144:1012-1028.

440 Serrani JC, Sanjuán R, Ruiz-Rivero O, Fos M, García-Martínez JL (2007)
441 Gibberellin regulation of fruit set and growth in tomato. *Plant Physiol* 145:246-257.

442 Senn ME, Gergoff Grozeff GE, Alegre ML, Barrile F, De Tullio MC, Bartoli CG
443 (2016) Effect of mitochondrial ascorbic acid synthesis on photosynthesis. *Plant Physiol*
444 *Biochem* 104:29-35.

445 Shinozaki Y, Hao S, Kojima M, Sakakibara H, Ozeki-Iida Y, Zheng Y, Fei Z,
446 Zhong S, Giovannoni JJ, Rose JKC, Okabe Y, Heta Y, Ezura H, Ariizumi T (2015)
447 Ethylene suppresses tomato (*Solanum lycopersicum*) fruit set through modification of
448 gibberellin metabolism. *Plant J* 83:237-251.

449 Smirnoff N (2018) Ascorbic acid metabolism and functions: A comparison of
450 plants and mammals. *Free Radical Bio Med* 122:116-129.

451 Tanaka A, Fujita K (1974). Nutrio-physiological studies on the tomato plant IV.
452 Source-sink relationship and structure of the source-sink unit. *Soil Sci Plant Nutr*
453 20:305-315.

454 Walker D (1987) The use of the oxygen electrode and fluorescence probes in
455 simple measurements of photosynthesis. Oxygraphics Limited, University of Sheffield
456 Print Unit, Sheffield, UK pp203.

457 Wang L-Y, Li D, Deng Y-S, Lv W, Meng Q-W (2013) Antisense-mediated
458 depletion of tomato GDP-L-galactosephosphorylase increases susceptibility to chilling
459 stress. *J Plant Physiol* 170:303-314.

460 Yang D-Y, Li M, Ma N-N, Yang X-H, Meng Q-W (2017) Tomato *SIGGP-LIKE*
461 gene participates in plant responses to chilling stress and pathogenic infection. *Plant*
462 *Physiol Biochem* 112:218-226.

463 Yoshimura K, Nakane T, Kume S, Shiomi Y, Maruta T, Ishikawa T, Shigeoka S
464 (2014) Transient expression analysis revealed the importance of VTC2 expression level

465 in light/dark regulation of ascorbate biosynthesis in Arabidopsis. *Biosci Biotech Bioch*
466 78:60-66.

467 Zhang C, Ouyang B, Yang C, Zhang X, Liu H, Zhang Y, Zhang J, Li H, Ye Z
468 (2013) Reducing AsA leads to leaf lesion and defence response in knock-down of the
469 AsA biosynthetic enzyme GDP-D-Mannose pyrophosphorylase gene in tomato plant.
470 *PLoS ONE* 8(4): e61987.

471

472 **Legends to figures**

473

474 Figure 1. Ascorbate concentrations in leaves (A), roots (B), flowers (C) and fruit (D) of
475 *ggp1* mutants and wild type plants grown under greenhouse conditions. The values were
476 obtained from at least three independent experiments and expressed as the means \pm S.D.
477 (ANOVA, $P < 0.05$). Lower and upper case letters indicate statistical differences between
478 genotypes and stages, respectively.

479

480 Figure 2. Leaf gas exchange measurements in *ggp1* mutants and wild type plants as CO₂
481 uptake (A), ETR (B), O₂ production at saturating CO₂ (C) and stomatal conductance (D)
482 grown under greenhouse conditions. The values were obtained from at least three
483 independent experiments and expressed as the means \pm S.D. (ANOVA, $P < 0.05$). Lower
484 and upper case letters indicate statistical differences between genotypes.

485

486 Figure 3. Photosynthetic characteristics of *ggp1* mutants and wild type plants measured
487 by chlorophyll fluorescence imaging. The plants were grown in 16 h day, 21C day (150
488 $\mu\text{mol m}^{-2} \text{s}^{-1}$)/19C dark, 55 RH day/50 RH dark for 4 weeks before transferring to an
489 irradiance of 500 $\mu\text{mol m}^{-2} \text{s}^{-1}$ for 4 days. NPQ, non-photochemical quenching; qP,
490 photosystem II efficiency factor; F_q'/F_m' , PSII operating efficiency; F_v'/F_m' , PSII
491 maximum efficiency. The values were obtained from three independent experiments
492 and expressed as the means \pm S.D.

493 Figure 4. Firmness (A), ethylene production (B) and soluble solids content (C) in fruit
494 taken from *ggp1* mutants and wild type plants grown under greenhouse conditions at
495 green mature, breaker and red stages. The values were obtained from at least three
496 independent experiments and expressed as the means \pm S.D. (ANOVA, $P < 0.05$). Lower
497 and upper case letters indicate statistical differences between genotypes and stages,
498 respectively.

499

500 Figure 5. Bi-plot of the first and second principal components for tomato development
501 traits. Each parameter is represented by dark continuous lines and genotype-treatment
502 combination by symbols (\diamond). The treatments consisted in control plants that were

503 cultivated leaving only 4 trusses (C); limited number of fruit where each plant kept 4 or
504 8 fruits (1F and 2F respectively); and two levels of defoliation (-L and =L).

505 Table 1. Effect of 20 mM ascorbate supplement in ascorbate content and fruit setting in
 506 tomato wt and *GGP* deficient plants. One mL of ascorbate solution was sprayed on each plant
 507 four times a week.

Treatment	Leaf ascorbate content $\mu\text{mol g}^{-1}\text{FW}$	Fruit setting %
Wt	10.0 ± 0.6	62.7 ± 10.3
Wt + AA	11.3 ± 0.33	57.3 ± 10.9
GGP 5261	5.0 ± 0.6	25.0 ± 8.6
GGP 5261 + AA	$8.3 \pm 0.9^*$	$38.3 \pm 6.4^*$
GGP P49 C12	4.3 ± 0.3	29.7 ± 1.2
GGP P49 C12 + AA	$7.7 \pm 1.2^*$	$40.0 \pm 1.5^*$

508 Values are means \pm S.D. (n = 3). * Indicates statistical differences with non-treated plants
 509 (ANOVA, $P < 0.05$).

510

511

512 Table 2. Yield and vegetative parameters in wt and *ggpl* mutants tomato plants. Measurements were done when half mutant plants reached
 513 one mature fruit.

	Fruit yield	Individual	n° fruit	n° flowers	Fruit set	Days to	Vegetative	Leaf area	Leaf	Stem	Root
		fruit weight	per plant	per plant		anthesis	biomass		biomass	biomass	biomass
	(g FWplant ⁻¹)	(g FW)			(%)		(g FW)	(cm ²)	(g DW)	(g DW)	(g DW)
Wild type	109.9±15.9 A	2.52±1.82 A	40.16±9.15 A	50.1±10.7 A	74.0±8.1 A	53.06±4.7 A	58.06±15.3 A	853.2±165 A	3.5±0.90 A	0.7±0.12 A	1.23±0.32 A
GGP 5261	88.31±7.5 B	3.29±2.27 B	24.0±5.67 B	54.4±9.8 A	42.5±10.2 B	59.3±6.5 B	59.6±8.8 A	887.8±236.7 A	3.6±0.85 A	1.1±0.24 B	0.77±0.16 B
GGP P49C12	70.7±14.4 B	4.08±2.36 C	19.25±5.04 B	45.4±37 A	42.6±12.7 B	58.3±5.24 B	56.7±12.68 A	906.0±116.1 A	3.7±0.27 A	1.2±0.21 B	0.91±0.14 B

514 The values were obtained from three independent experiments and expressed as means ± S.D. (ANOVA, *P*<0.05). Letters indicates
 515 statistical differences between genotypes.

516

517

518 Table 3. Concentrations of several hormones in flowers of *ggp1* mutants and wild type tomato plants.
 519 Samples were taken at anthesis combining flowers from different plants.

	GA1	GA4	GA8	AIA	Zeatin	ABA
	$\mu\text{g g}^{-1}\text{DW}$					
Wt	2.85±0.15 A	19.5±1.9 A	1.59±0.08 A	1.13±0.07 A	1.26±0.08 A	16.2±0.6 A
GGP5261	1.77±0.06 B	9.3±0.3 B	0.34±0.03 B	1.11±0.03 A	1.44±0.2 A	16.0±1.1 A
GGPP49C12	2.20±0.11 B	7.7±0.4 B	0.28±0.02 B	1.06±0.02 A	1.14±0.03 A	18.1±1.1 A

520 Data are shown as means \pm S.D. from 3 independent experiments. Letters indicates statistical
 521 differences between genotypes (ANOVA, $P < 0.05$).
 522

523 Table 4. Biomass accumulation in different organs of wt and *GGPI* deficient tomato plants limiting the size of source and sink
524 tissues. The treatments consisted in control plants that were cultivated leaving only 4 trusses (C); limited number of fruit where
525 each plant kept 4 or 8 fruits (1F and 2F respectively); and two levels of defoliation (-L and =L). * Flowers were cut to establish
526 a limited number of fruits.

	Fruit yield g FWfr pl ⁻¹	Individual FW g FW fr ⁻¹	Fruit number fr pl ⁻¹	Flower number fl pl ⁻¹	Leaf g DW pl ⁻¹	Stem g DW pl ⁻¹	Root g DW pl ⁻¹	Total vegetative g DW pl ⁻¹
1F wt	18,7±3,8 B	3,60±0,31 B	4,0±0	*	2,51±0,2 A	1,96±0,3 A	1,97±0,5 B	6,44±1,0 A
2F wt	30,4±2,3 C	4,49±0,16 B	8,0±0	*	2,38±0,1 A	1,89±0,1 A	1,75±0,2 B	6,01±0,4 A
C wt	59,0±2,6 A	2,63±0,36 A	23,3±1,6 A	29,3±2,3 A	2,39±0,2 A	1,85±0,1 AC	1,65±0,2 AB	5,89±0,6 A
-L wt	59,1±2,7 A	2,36±0,23 A	21,7±0,75 A	29,2±0,9 A	1,32±0,09 B	1,41±0,03 BC	1,25±0,14 AB	3,97±0,2 B
=L wt	56,7±0,5 A	2,48±0,29 A	21,5±1,3 A	29,7±1,6 A	1,15±0,10 B	1,27±0,04 B	0,90±0,09 B	3,32±0,1 B
1F GGP 5261	20,9±2,3 B	4,98±1,02 B	4,0±0	*	1,75±0,1 A	1,82±0,1 A	0,99±0,2 A	4,57±0,2 AB
2F GGP 5261	39,1±2,30 A	5,47±0,41 B	8,0 0	*	2,03±0,26 A	1,73±0,23 A	1,09±0,15 A	4,85±0,6 A
C GGP 5261	51,8±1,7 A	3,75±0,22 A	14,7±0,8 A	29,3 ±4 A	2,05±0,1 A	1,41±0,2 A	1,0±0,1 A	4,6 ± 0,2 AB
-L GGP 5261	48,2±2,7 A	2,97±0,82 A	17,0±1,4 A	27,0±0,5 A	1,42±0,13 AB	1,52± 0,14 A	0,95±0,12 A	3,90±0,4 AB
=L GGP 5261	46,2±6,5 A	3,27±0,58 A	13,7±1,8 A	26,7±2,3 A	0,89±0,14 B	1,45± 0,22 A	0,72±0,13 A	3,06±0,5 B
1F GGP P49	26,3±2,8 B	5,76±0,59 B	4,0±0	*	1,89±0,2 A	1,72±0,2 A	1,44±0,2 A	5,07±0,6 A
2F GGP P49	43,1±5,4 A	6,73±0,75 B	8,0±0	*	1,78±0,15 A	1,57±0,16 A	1,07±0,08 AB	4,43±0,3 AB
C GGP P49	49,6±5,3 A	3,69±0,60 A	11,7±2,9 A	29,0±3,0 A	1,94±0,1 A	1,59±0,2 A	1,09±0,1 AB	4,65±0,4 AB
-L GGP P49	55,4±2,1 A	3,62±0,48 A	16,2±1,4 A	30,7±3,6 A	1,30±0,08 AB	1,73±0,08 A	1,02 0,04 AB	4,04±0,1 AB
=L GGP P49	49,2±2,7 A	3,70±0,57 A	15,4± 2,4 A	33,4±4,2 A	0,96±0,05 B	1,44±0,13 A	0,76±0,09 B	3,16±0,2 B

527 The values were obtained from three independent experiments and expressed as means ± S.D. (ANOVA, $P<0.05$). Letters
528 indicate significant differences between treatments for the same genotype.

529