



The influence of age and sex on cerebrovascular reactivity and ventilatory response to hypercapnia in children and adults

Christine M Tallon<sup>1</sup>, Alan R Barker<sup>2</sup>, Daniela Nowak-Flück<sup>1</sup>, Philip N Ainslie<sup>1</sup>, Ali M McManus<sup>1</sup>

<sup>1</sup>Centre for Heart, Lung, and Vascular Health, School of Health and Exercise Sciences, University of British Columbia, Kelowna, Canada

<sup>2</sup>Children's Health and Exercise Research Centre, Sport and Health Sciences, College of Life and Environmental Sciences, University of Exeter, Exeter, United Kingdom

*Running heading:* Influence of age and sex on hypercapnic reactivity

*Keywords:* Cerebrovascular reactivity, middle cerebral artery, ventilation, chemoreflex, children

*Total word count:* 7620

*Total number of references:* 48

*Corresponding author:* Christine Tallon

Centre for Heart, Lung, and Vascular Health,

School of Health and Exercise Sciences,

University of British Columbia,

1147 Research Road – ART 360,

Kelowna BC V1V 1Y3

Canada

*Tel:* +1 250 807 9873

*Email:* [christinemtallon@gmail.com](mailto:christinemtallon@gmail.com)

**Subject Area: Cardiovascular Control**

This is an Accepted Article that has been peer-reviewed and approved for publication in the Experimental Physiology, but has yet to undergo copy-editing and proof correction. Please cite this article as an Accepted Article; [doi: 10.1113/EP088293](https://doi.org/10.1113/EP088293).

This article is protected by copyright. All rights reserved.

Accepted Article

## **New Findings**

### **What is the central question of this study?**

This study investigated intracranial cerebrovascular and ventilatory reactivity to 6% CO<sub>2</sub> in children and adults and explored dynamic ventilatory and cerebrovascular onset responses.

### **What is the main finding and its importance?**

We show cerebrovascular reactivity is similar in children and adults, but the intracranial blood velocity onset response was markedly attenuated in children. Sex differences were apparent, with greater increases in intracranial blood velocity in females, and lower ventilatory reactivity in adult females. Our study confirms the importance of investigating dynamic onset responses when assessing the influence of development on cerebrovascular regulation.

**Abstract**

The purpose of this study was to compare the integrated intracranial cerebrovascular reactivity (CVR) and hypercapnic ventilatory response (HCVR) between children and adults, as well as explore the dynamic response of the middle cerebral artery mean velocity ( $MCA_v$ ). Children ( $n = 20$ ;  $9.9 \pm 0.7$  years) and adults ( $n = 21$ ;  $24.4 \pm 2.0$  years) completed assessment of CVR over 240s using a fixed concentration of inspired  $CO_2$  ( $F_iCO_2$ , 0.06). Baseline  $MCA_v$  was higher in the adult females compared to the males ( $p \leq .05$ ).  $MCA_v$  was greater in female children compared to male children ( $p \leq .05$ ), and in female adults compared to male adults ( $p \leq .05$ ) with hypercapnia. Relative CVR was similar in children and adults ( $3.71 \pm 1.06$  vs.  $4.12 \pm 1.32$  %/mmHg;  $p = .098$ ), with absolute CVR higher in adult females than males ( $3.27 \pm .86$  vs.  $2.53 \pm .70$  cm/s/mmHg;  $p \leq .001$ ). Likewise, HCVR did not differ between the children and adults ( $1.89 \pm 1.00$  vs.  $1.77 \pm 1.34$  L/min/mmHg;  $p = .597$ ), but was lower in adult females than males ( $1.815 \pm .37$  vs.  $2.33 \pm 1.66$  L/min/mmHg;  $p \leq .05$ ). The heart rate response to hypercapnia was greater in children than adults ( $p = .001$ ). A mono-exponential regression model was used to characterize the dynamic onset, consisting of a delay term, amplitude and time constant ( $\tau$ ). The results revealed that  $MCA_v \tau$  was faster in adults than in children ( $34 \pm 18$  vs.  $.74 \pm 28$  s;  $p = .001$ ). Our study provides new insight into the impact of age and sex on CVR and the dynamic response of the  $MCA_v$  to hypercapnia.

*Key words:* Adults; cerebrovascular reactivity; children; hypercapnia; middle cerebral artery

## Introduction

The partial pressure of arterial carbon dioxide ( $\text{PaCO}_2$ ) plays the primary role in the regulation of cerebral blood flow (CBF) during wakefulness (Hoiland *et al.*, 2019). It is well known that elevations or reductions in  $\text{PaCO}_2$  result in vasodilation or constriction, respectively, of the entire cerebrovascular tree from the extra- and intracranial arteries, to the pial vessels [reviewed extensively here (Willie *et al.*, 2014)]. This vasomotor mechanism is termed cerebrovascular reactivity (CVR) and provides a marker of the dynamic regulation of CBF and cerebrovascular reserve capacity (Fierstra *et al.*, 2013; Willie *et al.*, 2014).

The extensive neuronal development that occurs during childhood is reflected in a cerebral metabolic demand that is about twice that of adults (Vandekar *et al.*, 2017). Meeting this metabolic demand requires tight regulation of CBF. Although CVR has been extensively documented in adults [reviewed in: (Hoiland *et al.*, 2019)], far less is known about this response in healthy children and adolescents (Brouwers *et al.*, 1990; Leung *et al.*, 2016). There is, however, emerging evidence that age, sex and maturity may impact CBF during childhood and adolescence. Developmental trajectories reported from a large-scale ( $n=922$ ) study of 8 to 22 year olds reveal that resting cerebral perfusion (assessed from magnetic resonance imaging (MRI) arterial spin-labelling) peaks between the ages of 8 to 10 years, with values approximately 30% higher than adults; these CBF elevations are commensurate with the higher cerebral metabolic needs of this age group (Satterthwaite *et al.*, 2014). Resting CBF declines from early to late puberty in boys; however, in girls, declines are only evident in early puberty, with increases in CBF mid to late puberty (Vandekar *et al.*, 2017). When intracranial blood velocity of the anterior and posterior cerebral circulation was compared between girls and boys (4-8 years), middle and basilar cerebral artery velocities were higher in the girls compared to the boys (Tontisirin *et al.*, 2007). In contrast, cerebral autoregulation, indexed from changes in cerebral blood velocity in response to changes in blood pressure, did not differ by sex (Tontisirin *et al.*, 2007). Sex differences have also been noted in women compared to

men, with higher cerebral blood velocities reported in women (Vriens *et al.*, 1989; Marinoni *et al.*, 1998).

Leung *et al.* (2016) have reported the development of grey and white matter CVR in a group (n=34) between 9 to 30 years of age. Here, CVR was lowest in the youngest children, with increases of ~33% and 50% in grey and white matter CVR respectively between the ages of 9 to 15 years. These differences declined with age and, in comparison to young adults, children had grey and white matter CVR values ~20% and 30% lower, respectively. Higher levels of resting cerebral perfusion noted in the younger child may limit the capacity for further vasodilation, as such a lower cerebrovascular reserve, and may account for the blunted CVR observed by Leung *et al.* (2016). Higher levels of resting cerebral blood velocity have been observed in girls compared to boys (Tontisirin *et al.*, 2007). However, whether there are sex differences in CVR in the healthy child is currently unknown. It is also possible that the blunted CVR of children compared to adults may relate to a heightened hypercapnic ventilatory response (HCVR; the change in VE in response to hypercapnia) that has been reported in children (Gratas-Delamarche *et al.*, 1993; Marcus *et al.*, 1994). It is thought that the elevated ventilatory sensitivity is due to the activation of the central chemoreflex increasing the drive to breathe in order to 'blow off' the CO<sub>2</sub> [reviewed in (Ainslie & Duffin, 2009)]. In adults, there is some evidence that a blunted CVR is reflected in elevations in the ventilatory sensitivity to CO<sub>2</sub> (Xie *et al.*, 2006); however, this is not a universal finding (Hoiland *et al.*, 2015). Onset, or dynamic responses, to hypercapnia have also been used to interrogate the integration of cerebrovascular and ventilatory responses to hypercapnia. At least in adults, close coupling of middle cerebral artery blood velocity (MCA<sub>v</sub>) and PaCO<sub>2</sub> onset responses, in comparison to a slower onset in ventilation (V<sub>E</sub>), may infer cerebrovascular reactivity to CO<sub>2</sub> modulates the central ventilatory chemoreflex (Ogoh *et al.*, 2008; Ainslie & Duffin, 2009).

The integrated CBF and ventilatory response to changes in end-tidal CO<sub>2</sub> (P<sub>ET</sub>CO<sub>2</sub>) has yet to be defined in children and has implications for understanding the control of V<sub>E</sub> and CBF during

development. If an interaction exists between the chemoreceptor regulation of breathing and brain blood flow, changes in response to hypercapnia would be anticipated to be developmentally-dependent. Therefore, the purpose of this study was to compare intracranial CVR (indexed from the  $MCA_v$ ) between children and adults and explore the dynamic response of  $MCA_v$ ,  $V_E$  and  $P_{ET}CO_2$  by characterizing the kinetic onset responses to hypercapnia. Our primary hypotheses were (i) in response to hypercapnia, a blunted CVR in children would be reflective of an increased HCVR  $CO_2$  when compared to adults; (ii) the dynamic response of  $V_E$  to an increase in hypercapnia would be faster in children compared to adults, whereas the dynamic response of the  $MCA_v$  response to hypercapnia would be slower in children compared to adults; and (iii) females will have greater baseline  $MCA_v$  than males in both children and adults.

## Methods

### Ethical Approval

All experimental procedures and the protocols were approved by the University of British Columbia Clinical Research Ethics Board (H16-01281) and the study conformed to the standards set by the *Declaration of Helsinki*, except for registration in a database. Written informed consent was obtained from the adults and the parents of the children. Additionally, written and oral informed assent was obtained from each child.

### Study design

Twenty-four children (13 females, 11 males; aged 8.3 to 10.8 years) and 26 young adults (14 females, 12 males; aged 20.8 to 27.8 years) participated in this experimental study. Adult participants were recruited from the University of British Columbia Okanagan and children were recruited from a local elementary school.

Following baseline screening, participants visited the laboratory on one occasion. Per recommendations (Ainslie & Duffin, 2009), participants were asked to refrain from eating high fat foods, consuming caffeine or alcohol, or exercising strenuously 24 hours prior to testing. Additionally, room temperature and time of day were held constant for all participants, and no visual stimulation was allowed during the protocol. Upon arrival, anthropometric measures were completed and the participants were instrumented. After instrumentation, the protocol consisted of a 10-minute supine rest period breathing room air, and four-minutes of hypercapnia in the same supine position. Prior work in adults has demonstrated a steady-state response to hypercapnia using a fixed CO<sub>2</sub> concentration is achieved between 2 to 3 minutes (Fisher, 2016), therefore we chose a fixed concentration of 6% inspired CO<sub>2</sub> (F<sub>I</sub>CO<sub>2</sub>) over 4 minutes to quantify the responses of MCA<sub>V</sub> and V<sub>E</sub> to hypercapnia in children and adults. During hypercapnia, 6% CO<sub>2</sub> was administered in 21% oxygen (balance nitrogen). A 3-way Hans Rudolph valve allowed inspiratory gases to be switched from room air to the 6% CO<sub>2</sub> mixture (using a 20L Douglas bag). Throughout the protocol MCA<sub>V</sub>, P<sub>ET</sub>-CO<sub>2</sub>, end-tidal O<sub>2</sub> (P<sub>ET</sub>O<sub>2</sub>), heart rate (HR), blood pressure (BP), V<sub>E</sub> and its components tidal volume (V<sub>T</sub>) and respiratory rate (R<sub>R</sub>) were assessed continuously.

### Measures

*Anthropometrics and maturation.* Body mass and stature were measured barefoot in light clothing to the nearest 0.1 kg and 0.1 cm respectively. Physical maturation was assessed from predicted age at peak height velocity (aPHV), using the Mirwald equation (Mirwald *et al.*, 2002) and from parental report of Tanner stage for pubic hair and genitalia (Rasmussen *et al.*, 2015). All children were >-1 year from aPHV (mean aPHV: -2.3 ± 0.9 years). Fourteen of the 20 children were Tanner stage 1 and 6 were Tanner stage 2 (3 boys, 3 girls). Tanner stage 1 is defined by the lack of secondary sexual characteristics i.e., no pubic hair, breast or genitalia development and considered pre-pubertal. Tanner stage 2 marks the onset of secondary sex characteristics i.e., sparse pubic hair, breast buds

or onset of male genitalia growth and is considered early-pubertal (Marshall & Tanner, 1969; Marshall & Tanner, 1970).

*Cardiorespiratory measures:* Beat-by-beat BP was assessed using a Finometer Pro (Finapres Medical Systems). HR was assessed using a 3-lead electrocardiogram (ECG; ADInstruments BioAmp ML132).  $V_E$ ,  $R_R$ ,  $V_T$ ,  $P_{ET}CO_2$  and  $P_{ET}O_2$  were assessed breath by breath, using a metabolic cart (Oxycon Pro, Carefusion, USA). The metabolic cart was calibrated prior to each test using gases of a known concentration, in addition to calibration of volume. Respiratory variables were interpolated in second-by-second intervals and time-aligned with  $MCA_v$ , HR, and BP. We excluded data if there was relative hypocapnia at baseline – this was defined as  $P_{ET}CO_2 > 2$  standard deviations (SD) below the child or adult mean.

*Cerebrovascular measures:*  $MCA_v$  (right side) was assessed using a 2-MHz transcranial Doppler (TCD) ultrasound (Spencer Technologies). The TCD probe was secured via attachment to a headpiece (child-adapted and adult M600 bilateral head frame; Spencer Technologies). MCA insonation was performed through the trans-temporal window following previously described guidelines (Willie *et al.*, 2011). Our between-day coefficient of variation for resting  $MCA_v$  insonation is 4.7%.

## Data Processing

### *Baseline and steady-state responses to hypercapnia.*

Baseline values were calculated from 120 seconds of the supine rest. Response values were taken in the final 30s of the four-minute test and differences from baseline to hypercapnia ( $\Delta$ ) was calculated for HR, MAP,  $MCA_v$ ,  $V_E$ ,  $V_T$ ,  $R_R$ ,  $P_{ET}CO_2$  and  $P_{ET}O_2$ .

Cerebrovascular resistance index (CVRi) was calculated as MAP divided by  $MCA_v$  (mmHg/cm/s) at baseline and during the last 30 seconds of hypercapnia.



There is no standardized analytical approach for the calculation of CVR (Willie *et al.*, 2012; Skow *et al.*, 2013); therefore, CVR was calculated both in absolute (cm/s/mmHg) and relative (%/mmHg) terms:

$$\text{Absolute CVR} = \frac{(\text{response } MCA_V - \text{baseline } MCA_V)}{(\text{response } P_{ET}CO_2 - \text{baseline } P_{ET}CO_2)}$$

$$\text{Relative CVR} = \frac{\left( \frac{(\text{response } MCA_V - \text{baseline } MCA_V)}{\text{baseline } MCA_V} \right) * 100}{(\text{response } P_{ET}CO_2 - \text{baseline } P_{ET}CO_2)}$$

HCVR (L/min/mmHg) was calculated as:

$$\text{HCVR} = \frac{(\text{response } V_E - \text{baseline } V_E)}{(\text{response } P_{ET}CO_2 - \text{baseline } P_{ET}CO_2)}$$

We also examined MAP (mmHg/mmHg) and HR (beats/min/mmHg) reactivity to CO<sub>2</sub> to account for the impact changes in F<sub>I</sub>CO<sub>2</sub> may have on cardiovascular function:

$$\text{MAP reactivity} = \frac{(\text{response } MAP - \text{baseline } MAP)}{(\text{response } P_{ET}CO_2 - \text{baseline } P_{ET}CO_2)}$$

$$\text{HR reactivity} = \frac{(\text{response } HR - \text{baseline } HR)}{(\text{response } P_{ET}CO_2 - \text{baseline } P_{ET}CO_2)}$$

#### *Dynamic onset responses to hypercapnia.*

Previous studies investigating dynamic cerebrovascular responses to various stimuli including hypercapnia and exercise, have utilized monoexponential modelling, providing a robust fit for MCA<sub>V</sub>, V<sub>E</sub> and P<sub>ET</sub>CO<sub>2</sub> responses (Ogoh *et al.*, 2009; Billinger *et al.*, 2017). Likewise, we conducted exploratory analyses of the kinetic response of MCA<sub>V</sub>, V<sub>E</sub> and P<sub>ET</sub>CO<sub>2</sub> to hypercapnia using an iterative least-squares non-linear regression procedure, with corresponding 95% confidence intervals (95% CIs) to establish the precision of the estimate (GraphPad Prism Version 8.0.2, GraphPad Software,

San Diego, CA). The  $MCA_v$ ,  $V_E$  and  $P_{ET}CO_2$  onset responses for each participant were modelled using a monoexponential function including a delay term:

$$y(t) = y_0 + \Delta_A(1 - e^{-(t-TD)/\tau})$$

where  $y(t)$ ,  $y_0$ ,  $\Delta_A$ ,  $TD$  and  $\tau$  are the response at a given time, the baseline value, the baseline corrected change in amplitude from baseline to asymptote, the time delay and the time-constant of the response, respectively. Each participant's response was modelled from the onset of the 6%  $CO_2$  stimulus ( $t = 0$  s). Goodness of fit ( $r^2 > 0.50$ ) and normality of residuals were used to determine model acceptability.

### Statistical Analysis

Statistical analyses were performed using SPSS (version 25, SPSS; Chicago IL). Normality was checked and verified using the Shapiro-Wilk test normality test for all variables. Two-way analyses of variance (ANOVA) were used to compare baseline and responses to hypercapnia by age and sex. T-tests were used to deconstruct the ANOVA main and interaction effects. One-way ANOVA was used to compare onset kinetic responses by age. Pearson product moment correlation coefficients were computed to assess the relationship between CVR, HCVR, HR and MAP reactivity. Statistical significance for main effects and interactions was set *a priori* at  $p \leq 0.05$ . Data are presented as mean  $\pm$  SD unless stated otherwise.

### Results

Of the 24 children and 26 adults recruited, four children and five adults were excluded from analyses. Three children were excluded who had a  $P_{ET}CO_2 < 31.2$  mmHg at baseline and the MCA could not be insonated in one child. Of the five adults who were excluded from the analysis, two were excluded due to low  $P_{ET}CO_2 (< 34.7$  mmHg) at baseline. The MCA could not be insonated in two adults and the signal acquisition for  $V_E$  was too poor for data to be processed in one adult. The data presented are from 20 children (11 females, 9 males) and 21 adults (10 females, 11 males). Age of

the children was  $9.9 \pm 0.7$  years (stature:  $1.42 \pm .07$  m; body mass:  $33.4 \pm 6.2$  kg), while the age of the adults was  $24.4 \pm 2.0$  years (height:  $1.72 \pm .06$  m; body mass:  $72.9 \pm 10.4$  kg).

*Baseline and steady-state responses to hypercapnia.*

Absolute values for all variables (HR, MAP,  $MCA_v$ , CVRi,  $V_E$ ,  $V_T$ ,  $R_R$ ,  $P_{ET}CO_2$ , and  $P_{ET}O_2$ ) at baseline and during the last 30s of 4 minutes of  $CO_2$  administration are presented in Table 1. There were significant child-adult differences at baseline in HR, MAP,  $MCA_v$ , CVRi,  $V_T$ ,  $R_R$ ,  $P_{ET}CO_2$ , and  $P_{ET}O_2$ . An age difference remained for the  $\Delta$  response to hypercapnia only for HR, CVRi,  $V_T$  and  $R_R$  (see Table 1). The  $\Delta$ HR response to hypercapnia was significantly greater in children compared to adults, while the CVRi decline was much greater in the adults compared to children. There was no difference in  $V_E$  at baseline or in response to hypercapnia, but children achieved a similar  $V_E$  at all time points with a smaller  $V_T$  and faster  $R_R$ .

A main effect for sex was apparent for  $MCA_v$ , MAP, CVRi,  $V_E$ ,  $P_{ET}CO_2$ , and  $P_{ET}O_2$  (see Table 1). The  $MCA_v$  values were higher in adult females compared to males at baseline ( $p \leq .05$ ) and in the final 30s of hypercapnia ( $p \leq .05$ ). Baseline  $MCA_v$  values were similar in girls and boys, but greater in the final 30s of hypercapnia in the girls ( $p \leq .05$ ).

< Insert Table 1 here >

There was a main effect for sex for the MAP response to hypercapnia, with follow up analyses revealing a greater MAP response to hypercapnia in adult females compared to males ( $p \leq 0.05$ ), but not children. The CVRi response to hypercapnia also differed by sex in the adults only, decreasing more in the females than the males ( $p \leq 0.05$ ).

Ventilatory responses to hypercapnia (absolute and  $\Delta V_E$ ) were greater in adult males compared to females ( $p \leq 0.05$ ), but not in children. All males had a higher baseline  $P_{ET}CO_2$ , and a smaller  $\Delta P_{ET}CO_2$

in response to hypercapnia ( $p \leq .05$ ), whereas all females had a higher baseline  $P_{ET}O_2$ , but a smaller hypercapnic  $\Delta P_{ET}O_2$  ( $p \leq .05$ ).

Absolute and relative CVR are presented in Figure 1. There was no main effect for age in absolute CVR (see Figure 1, panel A and B), with values of  $3.38 \pm 0.89$  cm/s/mmHg in children and  $2.89 \pm 0.85$  cm/s/mmHg in adults. A main effect of sex was apparent ( $p = .04$ ), with follow-up analyses indicated absolute CVR was greater ( $p \leq .05$ ) in females ( $3.27 \pm .86$ ) than males ( $2.53 \pm .70$ ). When expressed in relative terms, no main effect for age or sex were found (see Figure 1, panel C and D), with values of  $3.71 \pm 1.17$  %/mmHg in children and  $4.12 \pm 1.32$  %/mmHg in adults.

< Insert Figure 1 here >

The HCVR, MAP and HR reactivity are presented in Figure 2. There was no main effect for age for absolute HCVR (see Figure 2, panel A). A main effect for sex was present (see Figure 2, panel B), with follow-up analyses indicating a higher ventilatory reactivity in adult males ( $2.33 \pm 1.66$  L/min/mmHg) compared to females ( $1.15 \pm 0.37$  L/min/mmHg;  $p \leq .05$ ), but not in children (males:  $2.32 \pm 1.21$  L/min/mmHg; females  $1.53 \pm 0.67$  v;  $p = .07$ ).

There was no main effect for age or sex for absolute MAP reactivity (see Figure 2, panel C and D).

There was however, a main effect for age for absolute HR reactivity (see Figure 2, panel E), but not sex. HR reactivity was 1.05 beats/min/mmHg higher in children than in adults.

< Insert Figure 2 here >

There were no significant relationships between CVR, HCVR, MAP or HR reactivity in children or adults, when expressed in either absolute or relative terms.

*Dynamic onset responses to hypercapnia.*

The estimated kinetic responses at the onset of hypercapnia are shown in Table 2. Of the 20 children and 21 adults, acceptable models for  $MCA_V$  and  $P_{ET}CO_2$  were determined in 14 children and 16 adults. Acceptable model fit for  $V_E$  was only achieved in 8 children and 10 adults. Given the smaller sample size of the dynamic responses, only a main effect for age was explored using one-way ANOVAs.

< Insert Table 2 here >

A typical  $MCA_V$  response profile for a child and an adult at the onset of hypercapnia is shown in Figure 3. One-way ANOVA analyses revealed significant main effects for age for  $MCA_V$  TD and  $\tau$  (see Table 2), but not  $\Delta_A$ . Children responded with a  $\tau$  that was on average 42 s slower than adults. The  $MCA_V$  TD was longer in children, as was the  $P_{ET}CO_2$  TD. There were no age differences in any of the  $V_E$  onset parameters.

< Insert Figure 3 here >

$MCA_V$  and  $V_E$   $\tau$  values did not differ significantly in children ( $p=.383$ ), but  $MCA_V$   $\tau$  was considerably slower than  $P_{ET}CO_2$   $\tau$  (mean difference 43 s;  $p \leq .001$ ). In contrast,  $MCA_V$   $\tau$  was on average 59 s faster than  $V_E$   $\tau$  in adults ( $p \leq .001$ ), with no significant differences between  $MCA_V$  and  $P_{ET}CO_2$   $\tau$  values ( $p=.299$ ).

## Discussion

This study is the first to report the impact of age and sex on CVR, HCVR and the dynamic onset response of  $MCA_V$ ,  $V_E$  and  $P_{ET}CO_2$  to hypercapnia in children and adults. In contrast to our primary hypothesis, we did not find a blunted CVR in children, or child-adult differences in HCVR. We do, however, observe a markedly attenuated  $MCA_V$   $\tau$  in children compared to adults. In partial keeping with our final hypothesis, we did find sex differences in  $MCA_V$  at baseline, but only in adults, with

higher  $MCA_v$  values in females. Sex differences in the  $MCA_v$  response to hypercapnia were however apparent in both children and adults, with a greater  $MCA_v$  response to 6% $CO_2$  in females compared to males.

#### *Child to adult differences in reactivity to $CO_2$*

The CVR of 3.8 %/mmHg we report in 9-10-year-old children was similar to the adult CVR of 4.1 %/mmHg. These values are similar to the average adult TCD derived  $MCA_v$  CVR of between 3 to 4 %/mmHg (Hoiland *et al.*, 2019) and would suggest that cerebrovascular reserve is similar in healthy children and adults. Leung *et al.* (2016) estimated white and grey matter CVR in children to be about 20 to 30% lower compared to adults of similar ages to those in our study, concluding this may indicate a reduced cerebrovascular reserve in the child. However, previous work by Leung *et al.* (2016) did not account for physical maturation and only 3 boys and 4 girls were under the age of 12 years, limiting conclusions regarding child-adult differences in CVR. Given the influence age and maturation have on CBF, our study adds to a scant literature on the development of CVR. Comparison of these two studies is limited because of methodological differences in the approaches used; however, the methodologies used and the impact they may have on the interpretation of the findings are important and discussed in more detail under '*Methodological Considerations*' section.

We hypothesized that HCVR would be amplified in children because of the striking developmental differences in the chemoreceptive responsivity to  $CO_2$  previously demonstrated. We failed to find developmental differences in HCVR or a relationship between CVR and HCVR. This study was designed to explore the potential effect of CVR on the central ventilatory  $CO_2$  chemoreflex, but the rise in  $V_E$  with hypercapnia resulted in elevations in  $P_{ET}O_2$  in both children and adults. The failure to maintain an iso-oxic hypercapnic stimulus is a limitation given increases in  $P_{ET}CO_2$  and  $P_{ET}O_2$  induce peripheral and central respiratory chemoreflex loops. An alternative approach is the use of hyperoxic gas administration to blunt the peripheral chemoreceptor response and thus isolate the central  $V_E$  chemoreflex. Although the evidence is limited, the impact of hyperoxic hypercapnia is not

uniform in children and adults. Using a step protocol comprising a single breath of 15% CO<sub>2</sub> in 85% O<sub>2</sub>, Gozal *et al.* (1994) demonstrate the slope of the ventilatory response to step hypercapnic stimuli ( $V_E$  vs  $P_{ET}CO_2$ ) was 0.507 in children compared to 0.182 in adults. Further work is needed that explores the impact of iso-oxic and hyperoxic administration of CO<sub>2</sub>, as well as modified re-breathing techniques (Keir *et al.*, 2019) to further probe the possible interplay between CVR and HCVR in children. Inclusion of the posterior cerebral circulation will be also important, given the central chemoreceptors are located near the ventral surface of the medulla and disparate blood flow regulation has been shown in the brainstem and cortex of adults (Willie *et al.*, 2012; Skow *et al.*, 2013).

An intriguing finding was the markedly slower  $MCA_V$  kinetic response in children compared to adults. The model fit was acceptable with normality residuals and mean  $r^2$  of .78, .71 and .91 for  $MCA_V$ ,  $V_E$  and  $P_{ET}CO_2$  respectively. In adults, similar to prior work (Ogoh *et al.*, 2009),  $MCA_V \tau$  was tightly coupled to  $P_{ET}CO_2 \tau$ , with  $V_E$  in lag, supporting the premise that CVR influences the central ventilatory CO<sub>2</sub> chemoreflex. In contrast, in children this does not appear to be the case. The  $MCA_V \tau$  was 42 seconds slower than  $P_{ET}CO_2 \tau$ , and similar to  $V_E \tau$ . These disparate findings suggest developmentally distinct regulatory mechanisms, which are discussed below.

#### *Potential Mechanism(s)*

Increasing  $F_iCO_2$  has been shown to increase MAP and thereby cause increases in cerebral perfusion pressure (Battisti-Charbonney *et al.*, 2011). We found an 8-9% increase in MAP with the increased  $F_iCO_2$  in both children and adults, a similar MAP reactivity to hypercapnia, no relationship between CVR and MAP reactivity, but a smaller decline in CVR<sub>i</sub> with hypercapnia in children. It is possible that child-adult differences in autonomic regulation and/or cerebral autoregulation during hypercapnia exist and influence the onset response of  $MCA_V$ , but further investigations are needed to substantiate this proposition. Alternatively, sex differences in the MAP response may have impacted

the MAP reactivity–CVR relationship since cerebral autoregulation is attenuated with hypercapnia and in females (Edgell *et al.*, 2012; Perry *et al.*, 2014; Labrecque *et al.*, 2019).

A notable difference between the children and adults in the current study was the elevated HR in children with the increased  $F_i\text{CO}_2$ . It is possible that this relates to altered sympathetic or parasympathetic nervous activity with hypercapnia, but this is yet to be determined in children. Of note, the fraction of cardiac output (CO) delivered to the brain is highest during childhood, declining into adulthood (Wu *et al.*, 2016). Total cerebral blood flow to aortic outflow ratio is about 10% higher in 8-10-year-old children compared to adults (Wu *et al.*, 2016). As result of the increased cerebral oxygen consumption at this age (Vandekar *et al.*, 2017), it is possible that the increased HR reflects the role CO plays in meeting increased cerebrovascular demands in the child. This possibility, however, needs to be confirmed in future studies.

In adults, there is evidence that transitory increases in  $F_i\text{CO}_2$  causes shear mediated vasodilation in the conduit extracranial vasculature, independent of changes in MAP, cardiac output or HR that often accompany sustained hypercapnia (Hoiland *et al.*, 2017). At least in adults, dilation of the MCA has been shown to occur with elevations in  $P_{\text{ET}}\text{CO}_2 >10$  mmHg (Coverdale *et al.*, 2014). Half of our participants had an increase in  $P_{\text{ET}}\text{CO}_2$  between 10-12 mmHg and it is possible this caused vasodilation of the MCA. If in fact hypercapnic hyperemia caused shear-mediated vasodilation in our participants, we are likely underestimating the change in cerebral perfusion. However, previous research has shown only increases of  $P_{\text{ET}}\text{CO}_2$  approximately 15 mmHg led to significant dilation of the MCA (Verbree *et al.*, 2014), in which case it is likely no significant dilation occurred in the present study (see discussion within ‘*Methodological Considerations*’ section). Future work will need to properly document the impact of hypercapnia on the cerebral vasculature vasomotion in children using volumetric techniques to help discern whether dilation occurs and if so whether hypercapnic hyperemia is shear mediated in this population.

#### *Sex Differences in $\text{MCA}_v$*

This article is protected by copyright. All rights reserved.



Women were found to have a significantly higher baseline  $MCA_v$  than men, however, baseline  $MCA_v$  was similar in girls and boys (see Table 1). The sex difference found in adults confirms our hypothesis and aligns with previous literature where significant sex differences were found in  $MCA_v$  of adults at rest (Vriens *et al.*, 1989; Marinoni *et al.*, 1998). However, the similarity between the girls ( $94.1 \pm 9.5$  cm/s) and boys ( $90.5 \pm 9.8$  cm/s) included in this data set is contrary to the tertiary hypothesis of the present study. The age of the sample population may provide a reasonable explanation for this finding. Using transcranial Doppler, Tontisirin *et al.* (2007) report higher levels of resting  $MCA_v$  in girls ( $99 \pm 11$  cm/s) compared to boys ( $92 \pm 13$  cm/s) in 48 children between the age of 4 and 8 years old. However, prior research reports no sex difference in baseline  $MCA_v$  in 76 children between any of the four age categories: 2-4 years; 5-9 years; 10-14 years; 15-19 years (Brouwers *et al.*, 1990). As the present study includes a range of children 8 to 11 years of age, an older population than those included by Tontisirin *et al.* (2007), yet within range of those included by Brouwers *et al.* (1990), it is reasonable that our results mirror the latter. However, future longitudinal work is needed to fully understand the influence of age on sex differences in resting intracranial blood velocities.

A number of reasons have been proposed for the elevated cerebral blood velocity response to hypercapnia in all females, including sex hormones, hemoglobin, cerebral metabolism, and shear stress (Vavilala *et al.*, 2002). Within the present study there was a lack of control for menstrual phase in the adult females. However, our findings are similar to others who detected sex differences in CVR without controlling for menstrual phase (Minhas *et al.*, 2018), but estrogen is known to impact vasodilation (Krause *et al.*, 2006) and we do not account for this. Additionally, estrogen has been linked to greater carotid artery compliance and distensibility in women compared to men (Marlatt *et al.*, 2013). Albeit within the peripheral vasculature, there is evidence of sex differences in vascular smooth muscle in children who were classified as Tanner stage 2 (McCue *et al.*, 2012). Despite only six of our participants being Tanner stage 2, the remaining 14 were Tanner stage 1, suggests other factors alongside sex hormones may account for the sex differences noted.

It is also important to consider whether sex differences in cerebral metabolic rate (CMR) account for the an elevated hypercapnic  $MCA_v$  response in the girls. In an early study of CMR using a modified Kety Schmidt procedure, Kennedy and Sokoloff (1957) report higher CMRs in children up to 11 years of age compared to adults. Tontisirin *et al.* (2007) also report an elevated  $MCA_v$  in girls compared to boys. In contrast, a recent investigation of sex differences in CMR in a large cohort of children and adolescents, showed a lower global CMR (expressed as a % of whole body resting metabolic rate) and lower cerebral glucose consumption in girls in comparison to boys (Vandekar *et al.*, 2017). This latter study highlighted the increased CMR in children compared to adults, alongside increased cerebral perfusion in the younger participants. The higher baseline perfusion and lower CVRi we note in the children compared to the adults is most likely an outcome of the higher CMR in the child.

#### *Methodological considerations*

How the  $CO_2$  challenge is delivered is important given the impact this has on the time required to reach a steady state cerebrovascular response and the (in)ability to target end-tidal gases and standardize the end-tidal to arterial  $CO_2$  gradient [reviewed in (Hoiland *et al.*, 2019)]. The present study implemented elevations in  $F_I CO_2$  to experimentally manipulate  $PaCO_2$ . The benefit of this approach being it allows interactions between CVR and  $V_E$  to be explored since the  $P_{ET}CO_2$  to  $PaCO_2$  gradients are able to vary (Hoiland *et al.*, 2019). Leung *et al.* (2016) used prospective targeting of  $P_{ET}CO_2$ , alternating 60s of normoxic isocapnia [ $P_{ET}CO_2$  of 40 mmHg] with 45s of normoxic hypercapnia [ $P_{ET}CO_2$  of 45 mmHg]). Previous work in adults shows the time to steady state differs substantially between these two methods, with steady state attained during the prospective targeting technique within 1-15s, compared to the 1 to 2 minutes required when using a fixed concentration  $F_I CO_2$  (Mark *et al.*, 2010; Fisher, 2016). Our dynamic onset data reveal a distinctly slower  $MCA_v$  onset in children compared to adults, and it will be valuable to understand differences in steady state between the  $F_I CO_2$  and prospective targeting of  $P_{ET}CO_2$  in children. Of note, we were only able to fit the monoexponential model to the ventilatory data in 8 children and 10 adults. The

ventilatory response to CO<sub>2</sub> usually requires more than four minutes to attain steady-state and it is likely our protocol was of insufficient length for the kinetic modelling of V<sub>E</sub> (Ogoh *et al.*, 2009).

While using a fixed F<sub>I</sub>CO<sub>2</sub> does allow the interactions between CVR and V<sub>E</sub> to unfold, it does not permit precise control of the change in P<sub>ET</sub>CO<sub>2</sub> (or P<sub>ET</sub>O<sub>2</sub>) and shifts in the PaCO<sub>2</sub> to P<sub>ET</sub>CO<sub>2</sub> gradient can result (Hoiland *et al.*, 2019). It is important to note that resting P<sub>ET</sub>CO<sub>2</sub> closely reflects PaCO<sub>2</sub> in both children and adults and is an acceptable, non-invasive surrogate (Ohuchi *et al.*, 1999). The baseline P<sub>ET</sub>CO<sub>2</sub> and P<sub>ET</sub>O<sub>2</sub> values we report for children (Table 1: 36.6±2.7 / 98.5±4.3 mmHg) are similar to those found in previous published work from our group (Ellis *et al.*, 2017) and others (MacLean *et al.*, 2016) reporting P<sub>ET</sub>CO<sub>2</sub> 36.1±3.8 and P<sub>ET</sub>O<sub>2</sub> 102.2±10.1 mmHg in healthy 9-12-year-olds. Hypercapnia can increase the physiological deadspace to tidal volume ratio that can inflate P<sub>ET</sub>CO<sub>2</sub> to a greater extent than PaCO<sub>2</sub> and modify the MCA<sub>V</sub> response. Interestingly, the response of P<sub>ET</sub>CO<sub>2</sub> and P<sub>ET</sub>O<sub>2</sub> to four minutes of increased F<sub>I</sub>CO<sub>2</sub> in our study were similar between the children and adults. Although inter-individual variability in the PaCO<sub>2</sub> to P<sub>ET</sub>CO<sub>2</sub> gradient is unavoidable using the fixed F<sub>I</sub>CO<sub>2</sub> approach, this does not appear to have resulted in differences in the P<sub>ET</sub>CO<sub>2</sub> response to hypercapnia between the children and adults in this study. It is worth noting also, similar to our previous findings (Ellis *et al.*, 2017), the pattern of response in MCA<sub>V</sub> in children did not mirror the response pattern in P<sub>ET</sub>CO<sub>2</sub>; however, the changes in P<sub>ET</sub>CO<sub>2</sub> and MCA<sub>V</sub> seems to be aligned in adults. In order to properly discern the trajectory of change in CVR throughout childhood from methodology it will be important for future work to carefully consider the impact the experimental approach to delivery of CO<sub>2</sub>, as well as the technique used to assess CBF or velocity has on the CVR response.

The implementation of TCD to assess of cerebral blood velocity has both limitations and benefits. TCD does not account for vessel diameter, only providing an assessment of blood velocity not blood flow (Hoiland *et al.*, 2019). As CO<sub>2</sub> is a known vasodilator, without a measure of vessel diameter the assessment of the change in CBF cannot be conducted and therefore the change in cerebral perfusion could be underestimated (Ainslie & Hoiland, 2014). Literature on the vasomotion of the

MCA with hypercapnia is convoluted, although it has been suggested that there is subtle vasodilation of the MCA with hypercapnia (Hoiland & Ainslie, 2016). However, the degree of the increase in  $P_{ET}CO_2$  matters. For example, in a MRI study Verbree *et al.* (2014), reported significant vasodilation of the MCA after an increase in  $P_{ET}CO_2$  from  $36.9 \pm 3.8$  mmHg at baseline to  $51.0 \pm 4.5$  mmHg in adults. However, no significant change in MCA diameter occurred with an increase from baseline values to  $45.0 \pm 3.8$  mmHg (Verbree *et al.*, 2014), suggesting the magnitude of the change above baseline values influences the extent of vasomotion of the MCA. It is also unknown how the vasculature within the child reacts to hypercapnia and whether the vasomotion of the MCA changes in the same pattern as the adults. In the current study, the increase in  $P_{ET}CO_2$  averaged  $10.2 \pm 1.9$  mmHg within all participant values; however,  $P_{ET}CO_2$  values of  $49.7 \pm 2.0$  mmHg were reached in the adults (compared to  $46.3 \pm 1.8$  mmHg in the children) due to their slightly higher resting  $P_{ET}CO_2$  at baseline. It would seem reasonable that it is the delta change – rather than absolute value – in  $PaCO_2$  that might influence the MCA diameter; therefore, both adults and children would have had the same vasomotion stimulus. Despite the noted limitations of both the fixed  $F_iCO_2$  experimental approach and TCD for measurement of cerebral blood velocity, they do provide robust and commonly used methods for assessing cerebrovascular reactivity that are applicable to a paediatric population (Hoiland *et al.*, 2019).

We explored the link between cerebral blood velocity and  $V_E$  by insonating the MCA. The MCA is one of the two most commonly insonated vessels, the other being posterior cerebral artery, these vessels supply blood to the anterior and posterior regions of the brain, respectively. If a change (either increase or decrease) in  $V_E$  were expected to influence cerebral blood velocity through an interaction with the respiratory centers (e.g. via an influence on local metabolism), this would be expected to manifest within the *posterior* not *anterior* circulation. Therefore, insonation solely of the MCA as per this study, may not fully capture potential relationships between ventilatory and cerebrovascular control. At least in adults, studies have reported no differences in CVR to

hypercapnia between posterior and anterior circulations (Willie *et al.*, 2012; Hoiland *et al.*, 2015); however, if any regional differences are present in children should be considered in future studies.

### Conclusion

These are the first findings to provide insight into the dynamic onset response of the MCA<sub>v</sub> to hypercapnia in children compared to adults and to explore the impact sex may have on cerebrovascular reactivity in the child. We show a similar CVR, but markedly attenuated MCA<sub>v</sub> onset response in the child, that may relate to differences in cerebral perfusion pressure, autonomic control or cerebral metabolism. Identifying which factors account for age, sex, and maturation differences in the MCA<sub>v</sub> response to hypercapnia is challenging, but has relevant application to clinical settings when interpreting cerebral hemodynamic abnormalities in children. Future work will require careful consideration of the impact of the methodological approach employed, examination of regional differences in CVR in children and a longitudinal design to fully understand the development of cerebrovascular regulation.

### References

- Ainslie PN & Duffin J (2009). Integration of cerebrovascular CO<sub>2</sub> reactivity and chemoreflex control of breathing: mechanisms of regulation, measurement, and interpretation. *American Journal of Physiology Regulatory, Integrative and Comparative Physiology* **296**, R1473-1495.
- Ainslie PN & Hoiland RL (2014). Transcranial Doppler ultrasound: valid, invalid, or both? *Journal of Applied Physiology* **117**, 1081-1083.
- Battisti-Charbonney A, Fisher J & Duffin J (2011). The cerebrovascular response to carbon dioxide in humans. *The Journal of Physiology* **589**, 3039-3048.
- Billinger SA, Craig JC, Kwapiszeski SJ, Sisante JV, Vidoni ED, Maletsky R & Poole DC (2017). Dynamics of middle cerebral artery blood flow velocity during moderate-intensity exercise. *Journal of Applied Physiology* **122**, 1125-1133.

Brouwers PJAM, Vriens EM, Musbach M, Wieneke GH & van Huffelen AC (1990). Transcranial pulsed doppler measurements of blood flow velocity in the middle cerebral artery: reference values at rest and during hyperventilation in healthy children and adolescents in relation to age and sex. *Ultrasound in Medicine and Biology* **16**, 1-8.

Coverdale NS, Gati JS, Opalevych O, Perrotta A & Shoemaker JK (2014). Cerebral blood flow velocity underestimates cerebral blood flow during modest hypercapnia and hypocapnia. *Journal of Applied Physiology* **117**, 1090-1096.

Edgell H, Robertson AD & Hughson RL (2012). Hemodynamics and brain blood flow during posture change in younger women and postmenopausal women compared with age-matched men. *J Appl Physiol (1985)* **112**, 1482-1493.

Ellis LA, Ainslie PN, Armstrong VA, Morris LE, Simair RG, Sletten NR, Tallon CM & McManus AM (2017). Anterior cerebral blood velocity and end-tidal CO<sub>2</sub> responses to exercise differ in children and adults. *American Journal of Physiology Heart and Circulatory Physiology* **312**, H1195-H1202.

Fierstra J, Sobczyk O, Battisti-Charbonney A, Mandell DM, Poublanc J, Crawley AP, Mikulis DJ, Duffin J & Fisher JA (2013). Measuring cerebrovascular reactivity: what stimulus to use? *The Journal of Physiology* **591**, 5809-5821.

Fisher JA (2016). The CO<sub>2</sub> stimulus for cerebrovascular reactivity: Fixing inspired concentrations vs. targeting end-tidal partial pressures. *Journal of Cerebral Blood Flow & Metabolism* **36**, 1004-1011.

Gozal D, Arens R, Omlin KJ, Marcus CL & Keens TG (1994). Maturational differences in step vs. ramp hypoxic and hypercapnic ventilatory responses. *J Appl Physiol* **76**, 1968-1975.

Gratas-Delamarche A, Mercier J, Ramonatxo M, Dasonville J & Préfaut C (1993). Ventilatory response of prepubertal boys and adults to carbon dioxide at rest and during exercise. *European Journal of Applied Physiology* **66**, 25-30.

- Hoiland RL & Ainslie PN (2016). CrossTalk proposal: The middle cerebral artery diameter does change during alterations in arterial blood gases and blood pressure. *The Journal of Physiology* **594**, 4073-4075.
- Hoiland RL, Ainslie PN, Wildfong KW, Smith KJ, Bain AR, Willie CK, Foster G, Monteleone B & Day TA (2015). Indomethacin-induced impairment of regional cerebrovascular reactivity: implications for respiratory control. *The Journal of Physiology* **593**, 1291-1306.
- Hoiland RL, Fisher JA & Ainslie PN (2019). Regulation of the Cerebral Circulation by Arterial Carbon Dioxide. *Comprehensive Physiology* **9**, 1101-1154.
- Hoiland RL, Smith KJ, Carter HH, Lewis NCS, Tymko MM, Wildfong KW, Bain AR, Green DJ & Ainslie PN (2017). Shear-mediated dilation of the internal carotid artery occurs independent of hypercapnia. *American Journal of Physiology Heart and Circulatory Physiology* **313**, H24-H31.
- Keir DA, Duffin J, Millar PJ & Floras JS (2019). Simultaneous assessment of central and peripheral chemoreflex regulation of muscle sympathetic nerve activity and ventilation in healthy young men. *J Physiol* **597**, 3281-3296.
- Kennedy C & Sokoloff L (1957). An adaptation of the nitrous oxide method to the study of the cerebral circulation in children; normal values for cerebral blood flow and cerebral metabolic rate in childhood. *The Journal of Clinical Investigation* **36**, 1130-1137.
- Krause DN, Duckles SP & Pelligrino DA (2006). Influence of sex steroid hormones on cerebrovascular function. *Journal of Applied Physiology* **101**, 1252-1261.
- Labrecque L, Rahimaly K, Imhoff S, Paquette M, Le Blanc O, Malenfant S, Drapeau A, Smirl JD, Bailey DM & Brassard P (2019). Dynamic cerebral autoregulation is attenuated in young fit women. *Physiol Rep* **7**, e13984.
- Leung J, Kosinski PD, Croal PL & Kassner A (2016). Developmental trajectories of cerebrovascular reactivity in healthy children and young adults assessed with magnetic resonance imaging. *The Journal of Physiology* **594**, 2681-2689.

- MacLean JE, DeHaan K, Fuhr D, Hariharan S, Kamstra B, Henderson L, Adatia I, Majaesic C, Lovering AT, Thompson RB, Nicholas D, Thebaud B & Stickland MK (2016). Altered breathing mechanics and ventilatory response during exercise in children born extremely preterm. *Thorax* **71**, 1012-1019.
- Marcus C, Glomb W, Basinski D, Ward SD & Keens T (1994). Development of hypercapnic and hypoxic ventilatory responses from childhood to adulthood. *Journal of Applied Physiology* **76**, 314-320.
- Marinoni M, Ginanneschi A, Inzitari D, Mugnai S & Amaducci L (1998). Sex-related differences in human cerebral hemodynamics. *Acta Neurologica Scandinavica* **97**, 324-327.
- Mark CI, Slessarev M, Ito S, Han J, Fisher JA & Pike GB (2010). Precise control of end-tidal carbon dioxide and oxygen improves BOLD and ASL cerebrovascular reactivity measures. *Magnetic Resonance in Medicine* **64**, 749-756.
- Marlatt KL, Kelly AS, Steinberger J & Dengel DR (2013). The influence of gender on carotid artery compliance and distensibility in children and adults. *Journal of Clinical Ultrasound* **41**, 340-346.
- Marshall WA & Tanner JM (1969). Variations in pattern of pubertal changes in girls. *Archives of Disease in Childhood* **44**, 291-303.
- Marshall WA & Tanner JM (1970). Variations in the pattern of pubertal changes in boys. *Archives of Disease in Childhood* **45**, 13-23.
- McCue MC, Marlatt KL, Kelly AS, Steinberger J & Dengel DR (2012). Evaluation of gender differences in endothelium-independent dilation using peripheral arterial tonometry. *Clinical Physiology and Functional Imaging* **32**, 94-98.
- Minhas JS, Panerai RB & Robinson TG (2018). Sex differences in cerebral haemodynamics across the physiological range of PaCO<sub>2</sub>. *Physiological Measurement* **39**, 105009-105018.



- Mirwald RL, Baxter-Jones ADG, Bailey DA & Beunen GP (2002). An assessment of maturity from anthropometric measurements. *Medicine & Science in Sports & Exercise* **34**, 689-694.
- Ogoh S, Ainslie PN & Miyamoto T (2009). Onset responses of ventilation and cerebral blood flow to hypercapnia in humans: rest and exercise. *Journal of Applied Physiology* **106**, 880-886.
- Ogoh S, Hayashi N, Inagaki M, Ainslie PN & Miyamoto T (2008). Interaction between the ventilatory and cerebrovascular responses to hypo- and hypercapnia at rest and during exercise. *J Physiol* **586**, 4327-4338.
- Ohuchi H, Kato Y, Tasato H, Arakaki Y & Kamiya T (1999). Ventilatory Response and Arterial Blood Gases during Exercise in Children. *Pediatric Research* **45**, 389-396.
- Perry BG, Lucas SJ, Thomas KN, Cochrane DJ & Mundel T (2014). The effect of hypercapnia on static cerebral autoregulation. *Physiol Rep* **2**.
- Rasmussen AR, Wohlfahrt-Veje C, Tefre de Renzy-Martin K, Hagen CP, Tinggaard J, Mouritsen A, Mieritz MG & Main KM (2015). Validity of self-assessment of pubertal maturation. *Pediatrics* **135**, 86-93.
- Satterthwaite TD, Shinohara RT, Wolf DH, Hopson RD, Elliott MA, Vandekar SN, Ruparel K, Calkins ME, Roalf DR, Gennatas ED, Jackson C, Erus G, Prabhakaran K, Davatzikos C, Detre JA, Hakonarson H, Gur RC & Gur RE (2014). Impact of puberty on the evolution of cerebral perfusion during adolescence. *Proceedings of the National Academy of Sciences of the United States of America* **111**, 8643-8648.
- Skow RJ, MacKay CM, Tymko MM, Willie CK, Smith KJ, Ainslie PN & Day TA (2013). Differential cerebrovascular CO<sub>2</sub> reactivity in anterior and posterior cerebral circulations. *Respir Physiol Neurobiol* **189**, 76-86.
- Tontisirin N, Muangman SL, Suz P, Pihoker C, Fisk D, Moore A, Lam AM & Vavilala MS (2007). Early childhood gender differences in anterior and posterior cerebral blood flow velocity and autoregulation. *Pediatrics* **119**, e610-615.

Vandekar SN, Shou H, Satterthwaite TD, Shinohara RT, Merikangas AK, Roalf DR, Ruparel K, Rosen A, Gennatas ED, Elliott MA, Davatzikos C, Gur RC, Gur RE & Detre JA (2017). Sex differences in estimated brain metabolism in relation to body growth through adolescence. *Journal of Cerebral Blood Flow & Metabolism* **39**, 524-535.

Vavilala MS, Lee LA & Lam AM (2002). Cerebral blood flow and vascular physiology. *Anesthesiology Clinics of North America* **20**, 247-264.

Verbree J, Bronzwaer AS, Ghariq E, Versluis MJ, Daemen MJ, van Buchem MA, Dahan A, van Lieshout JJ & van Osch MJ (2014). Assessment of middle cerebral artery diameter during hypocapnia and hypercapnia in humans using ultra-high-field MRI. *Journal of Applied Physiology* **117**, 1084-1089.

Vriens EM, Kraaier V, Musbach M, Wieneke GH & van Huffelen AC (1989). Transcranial pulsed Doppler measurements of blood velocity in the middle cerebral artery: reference values at rest and during hyperventilation in healthy volunteers in relation to age and sex *Ultrasound in Medicine and Biology* **15**, 1-8.

Willie CK, Colino FL, Bailey DM, Tzeng YC, Binsted G, Jones LW, Haykowsky MJ, Bellapart J, Ogoh S, Smith KJ, Smirl JD, Day TA, Lucas SJ, Eller LK & Ainslie PN (2011). Utility of transcranial Doppler ultrasound for the integrative assessment of cerebrovascular function. *Journal of Neuroscience Methods* **196**, 221-237.

Willie CK, Macleod DB, Shaw AD, Smith KJ, Tzeng YC, Eves ND, Ikeda K, Graham J, Lewis NC, Day TA & Ainslie PN (2012). Regional brain blood flow in man during acute changes in arterial blood gases. *The Journal of Physiology* **590**, 3261-3275.

Willie CK, Tzeng YC, Fisher JA & Ainslie PN (2014). Integrative regulation of human brain blood flow. *The Journal of Physiology* **592**, 841-859.

Wu C, Honarmand AR, Schnell S, Kuhn R, Schoeneman SE, Ansari SA, Carr J, Markl M & Shaibani A (2016). Age-Related Changes of Normal Cerebral and Cardiac Blood Flow in Children and Adults Aged 7 Months to 61 Years. *J Am Heart Assoc* **5**.

Xie A, Skatrud JB, Morgan B, Chenuel B, Khayat R, Reichmuth K, Lin J & Dempsey JA (2006). Influence of cerebrovascular function on the hypercapnic ventilatory response in healthy humans. *The Journal of Physiology* **577**, 319-329.

#### **Additional Information**

#### **Disclosure/ Conflict of Interest**

The authors declared no potential conflicts of interest.

#### **Author Contribution**

CT, DNF, PA & AM conceived the study design. CT, DNF & AM were involved with data collection. CM, AB, PA & AM were involved in data analysis and interpretation. CM, PA & AM drafted the manuscript. All authors critically reviewed and approved the final version of the manuscript and agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. All persons designated as authors qualify for authorship, and all those who qualify for authorship are listed.

#### **Sources of Funding**

This work was supported by Natural Sciences and Engineering Research Council Discovery Grant (AMM: 201503647) and a Canadian Foundation for Innovation Grant (AMM).

#### **Acknowledgements**

The authors are extremely grateful to the elementary school staff for facilitating this project and would like to thank the students for participating.

## Tables

**Table 1.** Ventilatory, cardiovascular, and cerebrovascular data at baseline and in the final 30s of hypercapnia in children and adults and by sex

		CHILDREN			ADULTS			ANOVA MAIN EFFECTS		
		Female	Male	All Children	Female	Male	All Adults	Age	Sex	Age*Sex
		(n = 11)	(n = 9)	(n = 20)	(n = 10)	(n = 11)	(n = 21)	<i>p value</i>	<i>p value</i>	<i>p value</i>
HR (beats/min)	BL	78 ± 8	78 ± 7	78 ± 8	69 ± 15	62 ± 12	65 ± 14	<b>.001</b>	.377	.292
	Hypercapnia	93 ± 10	92 ± 6	92 ± 8	73 ± 14	66 ± 8	69 ± 12	<b>.001</b>	.189	.427
	Δ	16 ± 6	13 ± 6	14 ± 6	4 ± 4	5 ± 6	4 ± 5	<b>.001</b>	.526	.492
MAP (mmHg)	BL	78 ± 10	78 ± 10	78 ± 9.5	88 ± 12	82 ± 9	85 ± 11	<b>.040</b>	.270	.384
	Hypercapnia	87 ± 10	83 ± 12	85 ± 11	97 ± 13	87 ± 10	92 ± 12	<b>.034</b>	<b>.048</b>	.403
	Δ	8 ± 6	5 ± 6	7 ± 6	9 ± 6	6 ± 7	7 ± 6	.687	.079	.957
MCA <sub>v</sub> (cm/s)	BL	94.1 ± 9.5	90.5 ± 9.8	92.7 ± 9.5	79.6 ± 15.4	64.0 ± 7.4	71.4 ± 14.1	<b>.001</b>	<b>.008</b>	.086

	Hypercapnia	129.9 ± 11.9	118.1 ± 13.3	125.6 ± 12.7	115.8 ± 14.8	88.2 ± 11.6	101.7 ± 19.4	<b>.001</b>	<b>.001</b>	.053
	Δ	35.8 ± 8.0	27.6 ± 7.7	32.9 ± 9.6	36.2 ± 9.7	24.2 ± 6.2	29.9 ± 10.0	.548	<b>.001</b>	.456
CVRI (mmHg/cm/s)	BL	0.84 ± 0.16	0.86 ± 0.12	0.85 ± 0.14	1.17 ± 0.22	1.29 ± 0.19	1.24 ± 0.21	<b>.001</b>	.143	.252
	Hypercapnia	0.67 ± 0.08	0.70 ± 0.08	0.68 ± 0.08	0.70 ± 0.22	.88 ± 0.24	0.80 ± 0.25	<b>.001</b>	<b>.033</b>	.138
	Δ	-0.24 ± 0.17	-0.16 ± 0.07	-0.17 ± 0.09	-0.48 ± 0.28	-0.41 ± 0.15	-0.44 ± 0.22	.394	.478	.698
V <sub>E</sub> (L/min)	BL	8.0 ± 1.5	7.7 ± 1.3	7.9 ± 1.3	8.2 ± 1.6	8.8 ± 1.9	8.5 ± 1.8	.196	.637	.366
	Hypercapnia	22.9 ± 5.4	27.0 ± 7.7	24.8 ± 6.7	20.7 ± 4.1	29.3 ± 11.7	25.3 ± 9.8	.978	<b>.014</b>	.369
	Δ	15.0 ± 5.2	19.3 ± 8.3	16.9 ± 7.0	12.6 ± 3.4	20.5 ± 11.4	16.7 ± 9.3	.812	<b>.016</b>	.468
V <sub>T</sub> (L)	BL	0.42 ± 0.10	0.48 ± 0.15	0.45 ± 0.10	0.72 ± 0.07	0.91 ± 0.31	0.82 ± 0.25	<b>.001</b>	<b>.037</b>	.253
	Hypercapnia	0.88 ± 0.14	0.94 ± 0.21	0.91 ± 0.17	1.41 ± 0.18	1.87 ± 0.41	1.66 ± 0.40	<b>.001</b>	<b>.003</b>	<b>.021</b>
	Δ	0.46 ± 0.16	0.46 ± 0.23	0.46 ± 0.19	0.69 ± 0.21	.96 ± 0.32	0.84 ± 0.31	<b>.001</b>	.084	.087
R <sub>R</sub> (bpm)	BL	20 ± 4	18 ± 4	19 ± 4	12 ± 3	11 ± 3	11 ± 3	<b>.001</b>	.142	.461
	Hypercapnia	28 ± 9	32 ± 11	30 ± 9	16 ± 3	16 ± 6	16 ± 5	<b>.001</b>	.373	.532
									.110	

	$\Delta$	$8 \pm 7$	$14 \pm 11$	$11 \pm 10$	$4 \pm 3$	$6 \pm 6$	$5 \pm 5$	<b>.010</b>		.325
P <sub>ET</sub> CO <sub>2</sub> (mmHg)	BL	$36.0 \pm 2.7$	$37.4 \pm 1.4$	$36.6 \pm 2.7$	$38.2 \pm 2.0$	$40.2 \pm 2.2$	$39.3 \pm 2.3$	<b>.001</b>	<b>.017</b>	.676
	Hypercapnia	$46.4 \pm 1.8$	$46.2 \pm 1.9$	$46.3 \pm 1.8$	$49.24 \pm 1.8$	$50.1 \pm 2.2$	$49.7 \pm 2.0$	<b>.001</b>	.641	.485
	$\Delta$	$10.4 \pm 2.3$	$8.8 \pm 1.5$	$9.7 \pm 2.1$	$11.1 \pm 1.1$	$9.9 \pm 2.0$	$10.5 \pm 1.7$	.121	<b>.016</b>	.801
P <sub>ET</sub> O <sub>2</sub> (mmHg)	BL	$100.9 \pm 4.1$	$96.6 \pm 2.3$	$98.5 \pm 4.3$	$96.1 \pm 5.3$	$92.8 \pm 4.9$	$94.4 \pm 5.4$	<b>.009</b>	<b>.004</b>	.482
	Hypercapnia	$132.9 \pm 4.3$	$133.0 \pm 5.6$	$132.9 \pm 4.3$	$125.7 \pm 4.6$	$127.2 \pm 5.4$	$126.5 \pm 4.6$	<b>.001</b>	.567	.610
	$\Delta$	$32.0 \pm 3.0$	$37.4 \pm 5.7$	$34.4 \pm 5.1$	$29.6 \pm 4.0$	$34.5 \pm 6.4$	$32.1 \pm 5.8$	.094	<b>.002</b>	.870


Values are means  $\pm$  SD. HR, heart rate; MAP, mean arterial blood pressure; CVRi, cerebrovascular resistance index; MCA<sub>v</sub>, middle cerebral artery mean velocity; V<sub>E</sub>, ventilation; V<sub>T</sub>, tidal volume; R<sub>R</sub>, respiratory rate; P<sub>ET</sub>CO<sub>2</sub>, end-tidal carbon dioxide; P<sub>ET</sub>O<sub>2</sub>, end-tidal oxygen; BL, baseline;  $\Delta$ , absolute change between the pre-stimulus baseline to the last 30s of hypercapnia.

**Table 2.** MCA<sub>V</sub>, P<sub>ET</sub>CO<sub>2</sub> and V<sub>E</sub> kinetic responses at the onset of hypercapnia in children and adults.

		<b>Children</b>	<b>Adults</b>	<b>ANOVA</b>
		<i>(n=14)</i>	<i>(n=16)</i>	<b>Main effect for age</b>
				<i>(p value)</i>
MCA <sub>V</sub>	Δ <sub>A</sub> (cm/s)	35.2 ± 8.1	29.2 ± 8.5	.081
	TD (s)	11 ± 5	5 ± 4	<b>.023</b>
	τ (s)	74 ± 28	34 ± 18	<b>.001</b>
P <sub>ET</sub> CO <sub>2</sub>	Δ <sub>A</sub> (mmHg/s)	10.7 ± 1.6	9.7 ± 2.3	.321
	TD (s)	8 ± 2	4 ± 2	<b>.006</b>
	τ (s)	31 ± 11	27 ± 16	.622
		<i>n=8</i>	<i>n=10</i>	
V <sub>E</sub>	Δ <sub>A</sub> (L/min)	16.6 ± 6.0	26.9 ± 14.3	.127
	TD (s)	11 ± 8	13 ± 7	.652
	τ (s)	102 ± 40	98 ± 24	.891


Values are means ± SD. MCA<sub>V</sub>, middle cerebral artery mean velocity; P<sub>ET</sub>CO<sub>2</sub>, end-tidal carbon dioxide; V<sub>E</sub>, ventilation. Δ<sub>A</sub> delta amplitude, TD time delay, τ time constant.

Figure Legends




**Figure 1.** Relative and absolute cerebrovascular reactivity (CVR) to hypercapnia by age and by sex. *A*, absolute CVR in children and adults. *B*, absolute CVR in males and females. *C*, relative CVR in children and adults. *D*, relative CVR in males and females. *P* values indicate the main effects for age and sex from ANOVA analyses.





**Figure 2.** Hypercapnic ventilatory and cardiovascular reactivity by age and by sex. *A* hypercapnic ventilatory reactivity (HCVR) in children and adults. *B*, HCVR in males and females. *C*, mean arterial pressure (MAP) reactivity in children and adults. *D*, MAP reactivity in males and females. *E*, heart rate (HR) reactivity in children and adults. *F*, HR reactivity in males and females. *P* values indicate the main effects for age and sex from ANOVA analyses.



**Figure 3.** Dynamic onset response of  $MCA_v$  in representative child and adult subjects. The *continuous black and grey lines* represent the fitted monoexponential function in the adults and children respectively. The *vertical dotted line* signifies the onset of the  $CO_2$  stimulus.