



Functional Outcome and Quality of Life After Hypoglossal-Facial Jump Nerve Suture

Gerd Fabian Volk^{1,2}, Maren Geitner^{1,2}, Katharina Geißler^{1,2}, Jovanna Thielker^{1,2}, Ashraf Raslan^{1,2,3,†}, Oliver Mothes⁴, Christian Dobel^{1,2} and Orlando Guntinas-Lichius^{1,2*}

¹ Department of Otorhinolaryngology, Jena University Hospital, Jena, Germany, ² Facial Nerve Center Jena, Jena University Hospital, Jena, Germany, ³ Department of Otorhinolaryngology, Assiut University Hospital, Assiut, Egypt, ⁴ Department of Computer Science, Friedrich Schiller University, Jena, Germany

Background: To evaluate the face-specific quality of life after hypoglossal-facial jump nerve suture for patients with long-term facial paralysis.

Methods: A single-center retrospective cohort study was performed. Forty-one adults (46% women; median age: 55 years) received a hypoglossal-facial jump nerve suture. Sunnybrook and eFACE grading was performed before surgery and at a median time of 42 months after surgery. The Facial Clinimetric Evaluation (FaCE) survey and the Facial Disability Index (FDI) were used to quantify face-specific quality of life after surgery.

Results: Hypoglossal-facial jump nerve suture was successful in all cases without tongue dysfunction. After surgery, the median FaCE Total score was 60 and the median FDI Total score was 76.3. Most Sunnybrook and eFACE grading subscores improved significantly after surgery. Younger age was the only consistent independent predictor for better FaCE outcome. Additional upper eyelid weight loading further improved the FaCE Eye comfort subscore. Sunnybrook grading showed a better correlation to FaCE assessment than the eFACE. Neither Sunnybrook nor eFACE grading correlated to the FDI assessment.

Conclusion: The hypoglossal-facial jump nerve suture is a good option for nerve transfer to reanimate the facial muscles to improve facial motor function and face-specific quality of life.

Keywords: facial nerve, hypoglossal nerve, facial paralysis, facial muscles, patient reported outcome measure, quality of life

OPEN ACCESS

Edited by:

Haralampos Gouveris,
Johannes Gutenberg University
Mainz, Germany

Reviewed by:

Georgios Psychogios,
University Hospital of Ioannina, Greece
Marcin Szymanski,
Medical University of Lublin, Poland

*Correspondence:

Orlando Guntinas-Lichius
orlando.guntinas@med.uni-jena.de

† Present address:

Ashraf Raslan,
Department of Otorhinolaryngology,
Ilm-Kreis-Kliniken, Arnstadt, Germany

Specialty section:

This article was submitted to
Otorhinolaryngology - Head and Neck
Surgery,
a section of the journal
Frontiers in Surgery

Received: 08 January 2020

Accepted: 02 March 2020

Published: 19 March 2020

Citation:

Volk GF, Geitner M, Geißler K, Thielker J, Raslan A, Mothes O, Dobel C and Guntinas-Lichius O (2020) Functional Outcome and Quality of Life After Hypoglossal-Facial Jump Nerve Suture. *Front. Surg.* 7:11. doi: 10.3389/fsurg.2020.00011

INTRODUCTION

Facial nerve reconstruction after complex damage of the peripheral facial nerve, especially if the central stump of the facial nerve is not available for nerve suture, or after longer denervation time, still is a challenge. The hypoglossal-facial jump nerve suture (i.e., a side-to-end nerve suture of the incised hypoglossal nerve to a nerve graft that is then sutured end-to-end to the distal facial nerve) is a well-established standard procedure for facial nerve reanimation for patients with permanent facial nerve paralysis (1–4). The jump technique has overcome the until then popular classical hypoglossal-facial nerve suture with transposition of the complete hypoglossal nerve and end-to-end nerve suture directly to the facial nerve. The complete transection of the

hypoglossal nerve unavoidably resulted in homolateral paralysis and hemitongue atrophy. Due to the tongue dysfunction many patients complained of permanent swallowing and speech problems (2). These problems are not seen after the jump technique with preservation of the function of the hypoglossal nerve.

Nevertheless, although hypoglossal-facial jump nerve suture and other techniques are well-established, clinical outcome research still is challenging due to a lack of standardization of outcome measurements after facial reanimation surgery (5–7). The Sunnybrook Facial Grading Scale seems to be the most robust grading scale out of the classical physician-based grading systems (8, 9). It became also popular in the last years to use the eFACE, an electronic and digitally graded facial measurement scale, as a modern alternative for physician-based facial grading (10). Further on, patient-reported outcome measures (PROMs) like the Facial Clinimetric Evaluation (FaCE) survey or the Facial Disability Index (FDI) as face-specific quality of life measures are essential to assess the facial disability in a holistic manner (11, 12).

Therefore, the present retrospective clinical study was performed to analyze systematically the outcome of hypoglossal-facial jump nerve suture using a standardized set of outcome measures including the Sunnybrook Facial Grading Scale, the eFACE, the FaCE, and the FDI.

MATERIALS AND METHODS

Study Design and Setting

This retrospective observational study included all patients who received a hypoglossal-facial jump nerve suture at the Department of Otorhinolaryngology, Jena University Hospital, Jena, Germany, between 2007 and 2017. This period was chosen to allow a sufficient follow-up for all patients. The local institutional review board approved the study protocol for this retrospective data analysis. In general, a facial nerve reconstruction technique was chosen if electromyography (EMG) confirmed a complete denervation of facial musculature or of parts of the facial musculature. A hypoglossal-facial jump nerve suture was proposed to the patient in different scenarios: First, the procedure was indicated if an early reconstruction within 12 months after onset of the lesion was possible, but the facial nerve stump proximal to the lesion was not available, or if the proximal stump was available but the defect was that large that a combined approach was needed (3, 4, 13). Second, the procedure was indication in cases of early reconstruction within 12 months after lesion, but the lesion site was far proximal, especially if proximal to the tympanic segment of the facial nerve. Finally, the procedure was indicated if late reconstruction later than 12 months after onset of the palsy was pursued. Denervation time longer than 24 months alone was no contraindication for hypoglossal-facial jump nerve suture. Nevertheless, the patients were informed that reconstruction results later than 12 months after lesion are might be worse than in patients who underwent surgery with a denervation time of <12 months. In any case, visibility of facial musculature was confirmed by facial muscle ultrasonography.

If present, almost all mimic muscles can be visualized with sonography. Details are presented elsewhere (14). Furthermore, the viability and contractibility of facial muscles was checked by direct surface electrostimulation of the muscles using a electrostimulation device (Paresestim, Krauth+Timmermann, Hamburg, Germany).

A standard hypoglossal-facial jump nerve suture technique was performed by the help of an operation microscope and 10/0 microsurgical suture material (2, 13). In all cases, the greater auricular nerve was used as interpositional nerve graft. Briefly, the hypoglossal-nerve was incised to one third. A side-to end nerve suture was performed between the incised hypoglossal nerve and the greater auricular nerve. In case of the standard hypoglossal-facial jump nerve suture, the other stump of the greater auricular nerve was sutured end-to-end to the main trunk of the peripheral extratemporal facial nerve. In cases with denervation of only parts of the peripheral facial nerve, the distal stump of the greater auricular nerve was sutured end-to-end only to the affected facial nerve branches. In a few cases of residual but functionally insufficient activity of the complete peripheral facial nerve, this distal suture to the facial nerve was performed side-to side, i.e., only the epineurium of the peripheral main trunk of the facial nerve was only incised and the distal stump of the greater auricular nerve was sutured end-to-side to the facial nerve.

The patients had follow-up visits every 3 months the first year after surgery, and every 6 months beginning with the second year. Patients' charts were reviewed for demographic characteristics, patients' history, and surgical techniques. Preoperative and postoperative evaluation included facial nerve grading using the Sunnybrook grading (8). The Sunnybrook Facial Grading Scale is a regional weighted system that rates three subscores: resting symmetry, the degree of voluntary facial muscle movement, involuntary muscle contraction (synkinesis). The three subscores are used to calculate a composite score (0 = total paralysis; 100 = normal function). An addition, the electronic facial assessment scale, eFACE, was used (15). The 16-item eFACE uses physician-graded visual analog scales and consists of 5 static, 7 dynamic, and 4 synkinesis items. The items are used to calculate static, dynamic, and synkinesis subscores, a total score, and three zonal domain scores, i.e., a periocular, midface and smile, and a lower face and neck score. Most scores range from 0 (worst) to 100 (best). Four items regarding static characteristics and items regarding dynamic characteristics relate to the nasolabial fold are rated on a scale of 0 (complete flaccidity) via 100 (balanced aesthetic appearance) to 200 (worst imaginable hypertonia). Two patient-reported outcome measures (PROMs), the Facial Clinimetric Evaluation (FaCE) scale, and the Facial Disability Index (FDI), were used (11, 12). The FaCE scores range from 0 (worst) to 100 (best). The FDI is divided into two domains and includes physical function and social/well-being function. The physical function scale is scored from –25 (worst) to 100 (best), while the social/well-being function scores from 0 (worst) to 100 (best). If several preoperative evaluations (Sunnybrook, FaCE, and FDI) were available, the evaluation closest before to surgery was selected. Out of the regular postoperative evaluations during the follow-up, the last postoperative evaluation was used.

Statistics

All outcome variables were analyzed with IBM SPSS software for medical statistics (Version 25; IBM, New York). Data are presented as frequencies or mean ± standard deviation (SD) if not otherwise indicated. Spearman's correlation was used for univariate correlation analysis. Preoperative and postoperative Sunnybrook gradings were compared with the non-parametric Wilcoxon test for paired data. Linear regression analyses including parameters from univariate analysis and $p < 0.01$ were performed to evaluate predictors for postsurgical FaCE subscores. The significance level was set at $p < 0.05$.

RESULTS

Baseline and Surgical Characteristics

Forty-one (41) patients (46% women; median age: 55 years) were included. Details on the characteristics of the patients are summarized in **Supplementary Table 1**. A classical hypoglossal-jump nerve suture was performed in most cases (68%). The variations of the suture technique are presented in **Table 1**. The median time between lesion and facial nerve reanimation surgery was 14 months (range: 0–307). All cases showed reinnervation of the facial muscles as controlled by needle EMG. A functional deficit of the tongue did not occur in any case. Thirty-three out of 41 cases (80.5%) also received an upper eyelid weight the day of hypoglossal-jump nerve suture. About half of the cases received further small surgical interventions later on during the follow-up period.

Functional Outcome and Quality of Life

The median time to the evaluation of the facial nerve function and quality of life was 42 months. The results of the Quality of life measurements using the FaCE and FDI after facial nerve reanimation with hypoglossal-facial jump nerve suture are presented in detail in **Table 2**. Out of the FaCE subscores, median values better than 70 points (best score: 100 points) were reached for FaCE Facial comfort (75.0 points), FaCE Oral function (75.0 points), and FaCE Social function (87.5 points). Using the FDI, more than 70 points were reached only for the FDI Social/well-being function subscore (82.0 points). The median FDI Physical function subscore was 65.0 points.

The functional assessment with the Sunnybrook grading system showed a significant improvement for the Symmetry of voluntary movement subscore ($p < 0.0001$), a higher (worse) Synkinesis subscore ($p = 0.009$), and improvement the Composite score ($p < 0.0001$) after hypoglossal-facial jump nerve suture compared to the preoperative function. The Resting symmetry subscore was not improved ($p = 0.345$; **Figure 1**; **Supplementary Table 2**).

The functional grading with the eFACE revealed an improvement of the Static subscore ($p < 0.0001$), the Dynamic subscore ($p = 0.017$), and a lower (worse) Synkinesis subscore ($p < 0.0001$). As a result, the eFACE total score was not significantly changed ($p = 0.065$). From the eFACE zonal scores, the Periocular score and the Midface and smile score were improved (both $p < 0.0001$), whereas the Lower face and neck

TABLE 1 | Surgical characteristics of the patients ($N = 41$).


Parameter	Absolute (N)	Relative (%)
Hypoglossal-facial-jump nerve suture		
side-to-end of main trunk of facial nerve	28	68.3
side-to-side of main trunk of facial nerve	4	9.8
side-to-end of cervicofacial main branch as part of a combined approach)	5	12.2
side-to-end of cervicofacial main branch	1	2.4
side-to-end of marginal mandibular branch	2	4.8
side-to-end of zygomatic branch	1	2.4
Brow lift	6	14.6
Upper eyelid weight	33	80.5
Upper eyelid plasty	3	7.3
Lower eyelid plasty	20	48.8
Sling plasty of the corner of the mouth	3	7.3
	Mean ± SD	Median, range
Denervation time, months	28.5 ± 51.2	14.0, 0–307
Time to first electrophysiological signs of reinnervation, months	5.6 ± 3.3	5.0, 2.0–14.0
Follow-up after surgery, months	39.5 ± 30.5	31.5, 6–134

SD, standard deviation.

TABLE 2 | Quality of life after hypoglossal-facial jump nerve suture ($N = 41$).

Parameter	Mean ± SD	Median, range
FaCE		
FaCE Facial movement	25.0 ± 22.0	16.6, 0–75
FaCE Facial comfort	67.3 ± 25.2	75.0, 17–100
FaCE Oral function	68.52 ± 28.7	75.0, 13–100
FaCE Eye comfort	40.3 ± 28.9	37.5, 0–100
FaCE Lacrimal control	52.8 ± 32.8	50.0, 0–100
FaCE Social function	75.2 ± 26.5	87.5, 6–100
FaCE Total score	56.5 ± 17.6	60.0, 23–83
FDI		
FDI Physical function	67.9 ± 17.5	65.0, 35–100
FDI Social/well-being function	75.1 ± 16.8	82.0, 40–96
FDI Total*	65.5 ± 15.5	76.3, 38–96
	Mean ± SD	Median, range
Interval between surgery to postoperative evaluation, months	48.9 ± 39.6	42.0, 6–131

SD, standard deviation; *mean score of FDI Physical function and FDI Social/well-being function.




score was decreased ($p = 0.014$) after facial reanimation surgery (Figure 2; Supplementary Table 3).

There was a good correlation between the Sunnybrook composite score after facial reanimation and the FaCE Total score ($\rho = 0.450$; $p = 0.014$; Figure 3). There was a moderate correlation between the eFACE Total score and the FaCE Total score after surgery ($\rho = 0.373$; $p = 0.056$; Figure 3). The correlation between the FaCE and FDI subscores among each other is shown in Supplementary Table 4. The ratings of the FaCE subscores were more independent among each other than the FDI subscores. The correlation between the Sunnybrook and eFACE subscores among each other is shown in Supplementary Table 5. Only some of the Sunnybrook and eFACE subscores correlated to each other.

The Supplementary Tables 6–9 contain the univariate correlation analyses for association between patients' and surgical characteristics and outcome in FaCE, FDI, Sunnybrook, and eFACE grading, respectively. Younger age and sometimes a higher Sunnybrook composite score were correlated to higher


FaCE subscores (all $p < 0.05$). Additional upper eyelid weight implantation was associated to a higher FaCE Eye comfort subscore ($p = 0.014$). Additional upper eyelid weight loading and a higher eFACE total score was associated to a higher FDI Physical function subscore ($p = 0.047$ and $p < 0.0001$, respectively). The final Sunnybrook composite score and also the final eFACE total score did not show any associations to patients' and surgical characteristics (all $p > 0.05$). Finally, the multivariate linear logistic regression analyses are presented in Table 3. Younger age (beta = -0.388 ; 95% confidence interval [CI] = -0.036 to -0.739) and a higher final Sunnybrook composite score (beta = 0.479 ; CI = 0.053 to 0.904) were independent predictors for a higher (better) FaCE Facial movement subscore. Younger age was also an independent predictor for higher FaCE Facial comfort subscore (beta = -0.543 ; CI = -1.004 to -0.083), FaCE Oral function subscore (beta = -0.565 ; CI = -1.131 to -0.0001), FaCE Eye comfort subscore (beta = -0.754 ; CI = -1.215 to -0.293), FaCE Lacrimal control subscore (beta = -1.119 ; CI = -1.651 to



-0.586), and a higher FaCE total score (beta = -0.335; CI = -0.633 to -0.038). Additionally, upper lid loading was an independent predictors for a higher FaCE Eye comfort subscore (beta = 29.038; CI = 8.995 to 49.081).

DISCUSSION

The hypoglossal-facial jump nerve suture is part of the standard repertoire of facial nerve repair techniques. Several case series



have shown that the jump technique is safe in regard of the preservation of the tongue function and leads to satisfactory functional results (1, 2, 4, 16–18). It is important to differentiate the jump technique with one third incision and normally side-to-end suture to a nerve graft from the splitting technique. Here, the hypoglossal nerve is split along its length and the split segment is transposed for end-to-end suture to the facial nerve. The splitting technique risks more injury to hypoglossal axons and results in mild to moderate hemitongue atrophy because many interweaving axons are transected (1, 19, 20).

Nevertheless, classical assessments of quality of life or with PROMs after hypoglossal-facial jump nerve suture are sparse. Lin et al. used apart from a Sunnybrook grading also the FDI

and also the short form SF-36 quality of life questionnaire to analyze several types of facial nerve repair (21). Fourteen patients with hypoglossal-facial jump nerve suture with or without interpositional graft and a Sunnybrook Composite score of 35.6–39.2 points revealed 67.1–70.1 points in the FDI Physical function subscore and 66.3–77.3 points in the Social/well-being function subscore. Due to the variability in patients' characteristics, it is difficult to compare this study with the present study, but the results FDI are in the same range, whereas functional outcome measuring with Sunnybrook grading are better in the present study. General quality of life measured with the SF-36 was highest after hypoglossal-facial jump nerve suture in the study by Lin et al. (21). General health values in the range of the

TABLE 3 | Linear regression analysis for independent predictors of the FaCE and FDI subscores.

Measure	beta	95% CI lower	95% CI upper	Stand.* beta	p**
FaCE Facial movement; $R^2 = 0.684$; $p = 0.001$					
Side (right = 0; left = 1)	14.377	-0.381	29.136	0.330	0.056
Age	-0.388	-0.036	-0.739	-0.333	0.032
Sunnybrook, composite score, after facial reanimation surgery	0.479	0.053	0.904	0.380	0.029
FaCE Facial comfort; $R^2 = 0.659$; $p = 0.011$					
Age	-0.543	-1.004	-0.083	-0.440	0.023
Interval between onset of palsy to surgery (denervation time)	0.051	-0.089	0.192	0.137	0.456
Sunnybrook, composite score, after facial reanimation surgery	0.453	-0.045	0.951	0.341	0.072
eFACE, total score, after facial reanimation surgery	0.325	-0.324	0.974	0.194	0.310
FaCE Oral function; $R^2 = 0.367$; $p = 0.050$					
Age	-0.565	-1.131	0.000	-0.367	0.050
FaCE Eye comfort; $R^2 = 0.504$; $p < 0.0001$					
Gender (female = 0; male = 1)	14.614	-2.508	31.736	0.254	0.091
Age	-0.754	-1.215	-0.293	-0.476	0.002
Additional upper eyelid weight (no = 0; yes = 1)	29.038	8.995	49.081	0.433	0.006
FaCE Lacrimal control; $R^2 = 0.646$; $p < 0.0001$					
Age	-1.119	-1.651	-0.586	-0.646	<0.0001
FaCE Social function					
Not applicable					
FaCE total score; $R^2 = 0.652$; $p = 0.013$					
Side (right = 0; left = 1)	2.949	-9.754	15.652	0.088	0.635
Age	-0.335	-0.633	-0.038	-0.383	0.029
Sunnybrook, composite score, after facial reanimation surgery	0.303	-0.091	0.696	0.321	0.125
eFACE, total score, after facial reanimation surgery	0.364	-0.086	0.814	0.306	0.108
FDI Physical function; $R^2 = 0.409$; $p = 0.111$					
Additional upper eyelid weight (no = 0; yes = 1)	14.560	-0.381	29.501	0.375	0.056
eFACE, total score, after facial reanimation surgery	0.185	-0.296	0.666	0.148	0.435
FDI Social/well-being function					
Not applicable					
FDI Total score					
Not applicable					

*Standardized beta; **p-values < 0.05 in bold.

normal population were reached after facial nerve repair and confirmed by another study (22). The SF-36 was not applied in the present study. It has been shown that the correlation of SF-36 and FDI or FaCE results is moderate (23, 24). Furthermore, function of individual facial regions seems not equally important for estimating facial palsy-related quality of life. The ability to smile seems to be of greatest importance (25). Therefore, one has to be careful with direct comparisons of the results of the different tools. Better facial motor function is highly correlated with better quality of life. This has also been shown by others and in general for patients with facial palsy (22, 24, 26). Furthermore, also other studies have revealed that age is negatively correlated with functional and quality of life outcome after facial nerve repair (22).

The present study is not without limitations: The retrospective design includes an uncontrolled selection bias. Although one of the largest series on hypoglossal-facial jump nerve suture, the sample size was not large. This limited the possibilities

for multivariate analyses. Strengths are the standardized use of reliable functional outcome measures (Sunnybrook and eFACE) and face-specific quality of life measures (FaCE and FDI). Another advantage is the long median follow-up of 31.5 months allowing a definite estimation of the final results.

What is lacking is a universal objective tool to measure the outcome after facial nerve repair (5). We could identify only one study using objective measure to evaluate the outcome after hypoglossal-facial jump nerve suture (20). Kochhar et al. used the facial asymmetry index (FAI) and the MEEI smile and oral Facegram software (27). Kochhar et al. revealed an objective reduction of facial asymmetry and improved lip excursions for smiling after hypoglossal-facial jump nerve suture. Recently, we have introduced an automated tool based on a machine learning approach for objective facial action coding and another tool for objective grading of patients with facial palsy (28, 29). An important next task will be to apply these objective tools

also for evaluation of the patients after hypoglossal-facial jump nerve suture.

CONCLUSION

This retrospective cohort study on 41 patients with hypoglossal-facial jump nerve suture confirms that this cross-nerve technique is a well-deserved part of the standard armamentarium for facial nerve repair. This was confirmed not only by standard facial grading but also by PROMs allowing a better inclusion of the patient's perspective. With good patient selection, this surgery achieves a very high overall success rate. The present study demonstrates a functionally relevant improvement for facial motor function after surgery in line with a good face-specific quality of life.

DATA AVAILABILITY STATEMENT

All datasets generated for this study are included in the article/**Supplementary Material**.

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by the ethics committee of the Jena University Hospital.

REFERENCES

- May M, Sobol SM, Mester SJ. Hypoglossal-facial nerve interpositional-jump graft for facial reanimation without tongue atrophy. *Otolaryngol Head Neck Surg.* (1991) 104:818–25. doi: 10.1177/019459989110400609
- Manni JJ, Beurskens CH, van de Velde C, Stokroos RJ. Reanimation of the paralyzed face by indirect hypoglossal-facial nerve anastomosis. *Am J Surg.* (2001) 182:268–73. doi: 10.1016/S0002-9610(01)00715-2
- Volk GF, Pantel M, Guntinas-Lichius O. Modern concepts in facial nerve reconstruction. *Head Face Med.* (2011) 6:25. doi: 10.1186/1746-160X-6-25
- Volk GF, Pantel M, Streppel M, Guntinas-Lichius O. Reconstruction of complex peripheral facial nerve defects by a combined approach using facial nerve interpositional graft and hypoglossal-facial jump nerve suture. *Laryngoscope.* (2011) 121:2402–5. doi: 10.1002/lary.22357
- Hadlock TA, Urban LS. Toward a universal, automated facial measurement tool in facial reanimation. *Arch Facial Plast Surg.* (2012) 14:277–82. doi: 10.1001/archfacial.2012.111
- Bhama P, Gliklich RE, Weinberg JS, Hadlock TA, Lindsay RW. Optimizing total facial nerve patient management for effective clinical outcomes research. *JAMA Facial Plast Surg.* (2014) 16:9–14. doi: 10.1001/jamafacial.2013.1382
- Hadlock T. Standard outcome measures in facial paralysis: getting on the same page. *JAMA Facial Plast Surg.* (2016) 18:85–6. doi: 10.1001/jamafacial.2015.2095
- Ross BG, Fradet G, Nedzelski JM. Development of a sensitive clinical facial grading system. *Otolaryngol Head Neck Surg.* (1996) 114:380–6. doi: 10.1016/S0194-5998(96)70206-1
- Fattah A, Gurusinghe A, Gavilan J, Hadlock TA, Marcus JR, Marres H, et al. Facial nerve grading scales: systematic review of the literature and suggestion for uniformity. *Plast Reconstr Surg.* (2015) 135:569–79. doi: 10.1097/PRS.0000000000000905
- Gaudin RA, Robinson M, Banks CA, Baiungo J, Jowett N, Hadlock TA. Emerging vs time-tested methods of facial grading among patients with facial paralysis. *JAMA Facial Plast Surg.* (2016) 18:251–7. doi: 10.1001/jamafacial.2016.0025
- VanSwearingen JM, Brach JS. Validation of a treatment-based classification system for individuals with facial neuromotor disorders. *Phys Ther.* (1998) 78:678–89. doi: 10.1093/ptj/78.7.678
- Kahn JB, Gliklich RE, Boyev KP, Stewart MG, Metson RB, McKenna MJ. Validation of a patient-graded instrument for facial nerve paralysis: the FaCE scale. *Laryngoscope.* (2001) 111:387–98. doi: 10.1097/00005537-200103000-00005
- Guntinas-Lichius O, Finkensieper M. Facial nerve reconstruction. In: Guntinas-Lichius O, Schaitkin B, editors. *Facial Nerve Disorders and Diseases: Diagnosis and Management.* Stuttgart: Thieme (2016). doi: 10.1055/b-0036-140496
- Volk GF, Pohlmann M, Sauer M, Finkensieper M, Guntinas-Lichius O. Quantitative ultrasonography of facial muscles in patients with chronic facial palsy. *Muscle Nerve.* (2014) 50:358–65. doi: 10.1002/mu.s.24154
- Banks CA, Bhama PK, Park J, Hadlock CR, Hadlock TA. Clinician-Graded electronic facial paralysis assessment: the eFACE. *Plast Reconstr Surg.* (2015) 136:223e–30e. doi: 10.1097/PRS.0000000000001447
- Atlas MD, Lowinger DS. A new technique for hypoglossal-facial nerve repair. *Laryngoscope.* (1997) 107:984–91. doi: 10.1097/00005537-199707000-00028
- Guntinas-Lichius O, Streppel M, Stennert E. Postoperative functional evaluation of different reanimation techniques for facial nerve repair. *Am J Surg.* (2006) 191:61–7. doi: 10.1016/j.amjsurg.2005.05.054
- Flores LP. Surgical results of the hypoglossal-facial nerve jump graft technique. *Acta Neurochir.* (2007) 149:1205–10. discussion: 1210. doi: 10.1007/s00701-007-1412-x
- Arai H, Sato K, Yanai A. Hemihypoglossal-facial nerve anastomosis in treating unilateral facial palsy after acoustic neuroma resection. *J Neurosurg.* (1995) 82:51–4. doi: 10.3171/jns.1995.82.1.0051
- Kochhar A, Albathi M, Sharon JD, Ishii LE, Byrne P, Boahene KD. Transposition of the intratemporal facial to hypoglossal nerve for reanimation

AUTHOR CONTRIBUTIONS

OG-L and GV: protocol development. MG, OM, KG, JT, and AR: acquisition of the analyzed data. OG-L, GV, and CD: data analysis. All authors: critical revision of work, and final approval of the version to be published. All authors complied with all aspects of the work. They ensure that questions related to the accuracy of the work are adequately discussed and solved.

FUNDING

AR was supported by a DAAD program German-Egyptian Research Long-Term Scholarship. OM acknowledges support by a Deutsche Forschungsgemeinschaft (DFG) grant DE 735/15-1 and OG-L by DFG grant GU 463/12-1.

ACKNOWLEDGMENTS

We thank Anish Raj for his help to collect the data and for the review of the Sunnybrook and eFACE gradings.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fsurg.2020.00011/full#supplementary-material>

- of the paralyzed face: the VII to XII Transposition Technique. *JAMA Facial Plast Surg.* (2016) 18:370–8. doi: 10.1001/jamafacial.2016.0514
21. Lin V, Jacobson M, Dorion J, Chen J, Nedzelski J. Global assessment of outcomes after varying reinnervation techniques for patients with facial paralysis subsequent to acoustic neuroma excision. *Otol Neurotol.* (2009) 30:408–13. doi: 10.1097/MAO.0b013e31819a8e26
 22. Guntinas-Lichius O, Straesser A, Streppel M. Quality of life after facial nerve repair. *Laryngoscope.* (2007) 117:421–6. doi: 10.1097/MLG.0b013e31802d83df
 23. Marsk E, Hammarstedt-Nordenvall L, Engstrom M, Jonsson L, Hultcrantz M. Validation of a Swedish version of the facial disability index (FDI) and the facial clinimetric evaluation (FaCE) scale. *Acta Otolaryngol.* (2013) 133:662–9. doi: 10.3109/00016489.2013.766924
 24. Volk GF, Granitzka T, Kreysa H, Klingner CM, Guntinas-Lichius O. Nonmotor disabilities in patients with facial palsy measured by patient-reported outcome measures. *Laryngoscope.* (2016) 126:1516–23. doi: 10.1002/lary.25695
 25. van Veen MM, Tavares-Brito J, van Veen BM, Dusseldorp JR, Werker PMN, Dijkstra PU, et al. association of regional facial dysfunction with facial palsy-related quality of life. *JAMA Facial Plast Surg.* (2019) 21:32–7. doi: 10.1001/jamafacial.2018.0804
 26. Volk GF, Granitzka T, Kreysa H, Klingner CM, Guntinas-Lichius O. Initial severity of motor and non-motor disabilities in patients with facial palsy: an assessment using patient-reported outcome measures. *Eur Arch Otorhinolaryngol.* (2017) 274:45–52. doi: 10.1007/s00405-016-4018-1
 27. Bray D, Henstrom DK, Cheney ML, Hadlock TA. Assessing outcomes in facial reanimation: evaluation and validation of the SMILE system for measuring lip excursion during smiling. *Arch Facial Plast Surg.* (2010) 12:352–4. doi: 10.1001/archfacial.2010.69
 28. Haase D, Minnigerode L, Volk GF, Denzler J, Guntinas-Lichius O. Automated and objective action coding of facial expressions in patients with acute facial palsy. *Eur Arch Otorhinolaryngol.* (2015) 272:1259–67. doi: 10.1007/s00405-014-3385-8
 29. Mothes O, Modersohn L, Volk GF, Klingner C, Witte OW, Schlattmann P, et al. Automated objective and marker-free facial grading using photographs of patients with facial palsy. *Eur Arch Otorhinolaryngol.* (2019) 276:3335–43. doi: 10.1007/s00405-019-05647-7

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Volk, Geitner, Geißler, Thielker, Raslan, Mothes, Dobel and Guntinas-Lichius. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.