

Accepted Manuscript

Lineage/species-specific expansion of the Mx gene family in teleosts: Differential expression and modulation of nine Mx genes in rainbow trout *Oncorhynchus mykiss*

Tingyu Wang, Fuguo Liu, Guangming Tian, Christopher J. Secombes, Tiehui Wang



PII: S1050-4648(19)30541-8

DOI: <https://doi.org/10.1016/j.fsi.2019.04.303>

Reference: YFSIM 6127

To appear in: *Fish and Shellfish Immunology*

Received Date: 22 February 2019

Revised Date: 28 April 2019

Accepted Date: 29 April 2019

Please cite this article as: Wang T, Liu F, Tian G, Secombes CJ, Wang T, Lineage/species-specific expansion of the Mx gene family in teleosts: Differential expression and modulation of nine Mx genes in rainbow trout *Oncorhynchus mykiss*, *Fish and Shellfish Immunology* (2019), doi: <https://doi.org/10.1016/j.fsi.2019.04.303>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21

Lineage/species-specific expansion of the Mx gene family in teleosts: Differential expression and modulation of nine Mx genes in rainbow trout *Oncorhynchus mykiss*

Tingyu Wang¹, Fuguo Liu¹, Guangming Tian¹, Christopher J Secombes^{1*}, Tiehui Wang^{1*}

^a Scottish Fish Immunology Research Centre, School of Biological Sciences, University of Aberdeen, Aberdeen AB24 2TZ, UK

***Corresponding authors:**

Dr. Tiehui Wang, E-mail: t.h.wang@abdn.ac.uk

Dr. Christopher J. Secombes, E-mail: c.secombes@abdn.ac.uk

22 **Abstract**

23

24 Myxovirus resistance (Mx) proteins are interferon (IFN)-inducible Dynamin-like GTPases, which play an
25 important role in antiviral immunity. Three Mx genes (Mx1-3) have been cloned previously in rainbow trout.
26 In this study, an additional six Mx genes were cloned that reside in four chromosomal loci. Further
27 bioinformatics analysis suggests the presence of three teleost Mx groups (TMG) each with a characteristic
28 gene organisation. Salmonid Mx belong to TMG1 and TMG2. The increased salmonid Mx gene copies are
29 due mainly to local gene duplications that happened before and after salmonid speciation, in a lineage/species
30 specific manner. Trout Mx molecules have been diversified in the loop 1 and 4 regions, and in the nuclear
31 localisation signal in loop 4. The trout Mx genes were shown to be differentially expressed in tissues, with
32 high levels of expression of TMG1 (Mx1-4) in blood and TMG2 (Mx5-9) in intestine. The expression of the
33 majority of the trout Mx genes was induced by poly IC *in vitro* and *in vivo*, and increased during development.
34 In addition, induction by antiviral (IFN) and proinflammatory cytokines was studied, and showed that type I
35 IFN, IFN γ and IL-1 β can induce Mx gene expression in an Mx gene-, cytokine- and cell line-dependent
36 manner. These results show that salmonids possess a large number Mx genes as well as complex regulatory
37 pathways, which may contribute to their success in an anadromous life style.

38

39 **Key words:** Rainbow trout, Mx, anti-viral defence, evolution, gene expression, modulation, type I interferon,
40 IFN γ , IL-1 β

41

42

43 1. Introduction

44 Mx (myxovirus resistance) proteins are interferon (IFN)-inducible Dynamin-like GTPases, with an important
45 role in antiviral immunity [1-2]. They are members of a family of large GTPases, and share an N-terminal
46 GTPase domain, a middle domain (MD), and a C-terminal GTPase effector domain (GED). The GTPase
47 domain is the most conserved part that consists of a tripartite GTP-binding motif (GDXXSGKS, DLPG, and
48 TKPD) and a dynamin signature (LPRXXGXXTR). The MD is important for oligomerization and viral target
49 recognition, whilst the GED has a conserved C-terminal leucine zipper that folds back to join the N-terminal
50 GTP-binding domain to establish the enzymatically active center of Mx proteins [1-3]. Mx proteins form
51 tetramers in solution that oligomerize further into large filaments and rings [3], with both GTPase activity and
52 oligomerization required for antiviral immunity.

53 A prototype Mx gene has been found in amphioxus, containing the N-terminal GTPase domain [4]. Typical
54 Mx genes are found in all vertebrate groups. The first evidence of Mx genes in fish started with the isolation
55 of an Mx genomic DNA fragment in perch (*Perca fluviatilis*) in 1989 [5]. The first full-length characterisation
56 of Mx genes was reported in rainbow trout *Oncorhynchus mykiss*, that has three Mx genes (Mx1-3) [6-7].
57 Subsequently Mx genes have been characterised in many fish species, with 1-9 genes present [4, 8-21].
58 However, some fish species such as the Gadiformes have lost their Mx genes [22]. The role of fish Mx
59 proteins in antiviral defence has been established in a few species, such as Japanese flounder *Paralichthys*
60 *olivaceus*, Atlantic salmon *Salmo salar* and grass carp *Ctenopharyngodon Idella* [23-26].

61 The multiple copies of mammalian Mx are closely linked and arise from local gene duplications [2]. How
62 multiple fish Mx genes have evolved is currently unclear [4,16]. A recent publication has shown that there are
63 nine Mx genes present on three chromosomes (Ch) in Atlantic salmon with Mx1-3 on Ch12, Mx4-8 on Ch25,
64 and Mx9 on Ch9 [27]. The origin of multiple copies of Mx genes on the same chromosome, that are linked
65 closely and share high sequence identities, is likely to also be via local gene duplications. However, due to the
66 third teleost-wide whole genome duplication (3R WGD) and the salmonid 4R WGD, many genes with single
67 copy in mammals are present as four copies on four chromosomes in salmonids [28]. Thus it is possible there
68 could be a fourth chromosome that harbours Mx genes in salmonids, and if discovered this may shed light on
69 how the different Mx-bearing chromosomes evolved in salmonids.

70 Mammalian Mx gene expression is induced by type I and type III IFNs but not by type II IFN γ or other
71 proinflammatory cytokines [29-31]. Interestingly, the diversified repertoire of Atlantic salmon Mx genes
72 appear to show some differential responsiveness to type I and II IFNs, with those on Ch12 being highly
73 induced by type I IFNs and those on Chr25 being more strongly induced by IFN γ than by type I IFN [27].
74 This finding is very interesting and raises the question as to whether a diversified Mx repertoire may also be
75 responsive to other cytokines released during innate antiviral defence, and remains to be examined. Hence, in
76 this study we aimed to shed light on Mx gene evolution in actinopterygian fish, in an attempt to establish a

77 better model of their evolution, and to establish whether the increased Mx copy number in salmonids has
78 allowed neo-functionalisation giving a broader responsiveness to a variety of cytokines. We first identified
79 and cloned an additional six Mx genes in rainbow trout, and found that all salmonids with a mapped genome
80 have four chromosomes harbouring Mx genes. We identified three groups of Mx genes present in teleosts in a
81 lineage-specific manner, with some (Ostariophysi) having all three groups, some having two groups
82 (Protacanthopterygii, including salmonids) but the percomorphs possessing only a single group. We next
83 investigated the expression of the nine trout Mx family members individually. We found that the trout Mx
84 genes are differentially expressed constitutively in tissues, that they increase during development, are induced
85 *in vivo* by poly IC, and are modulated *in vitro* by type I and type II IFNs, and by other proinflammatory
86 cytokine in a gene-, cytokine- and cell line-specific manner.

87

ACCEPTED MANUSCRIPT

88 2. Materials and methods

89 2.1. Rainbow trout

90 Healthy rainbow trout (~40 g) were purchased from the Mill of Elrich Trout Fishery (Aberdeenshire,
91 Scotland, UK). The fish were fed twice a day with a commercial diet (EWOS) and maintained in 1-m-
92 diameter fibreglass tanks with recirculating freshwater at 14°C at the Scottish Fish Immunology Research
93 Centre, University of Aberdeen, UK. Head kidney (HK) swabs were taken routinely and showed no bacterial
94 presence [32]. Fish were given at least two weeks for acclimation prior to use and ranged in size from 100-140
95 g when experiments were performed. All the experiments described comply with the Guidelines of the
96 European Union Council (2010/63/EU) for the use of laboratory animals, and were carried out under UK
97 Home Office project licence PPL 60/4013, approved by the ethics committee at the University of Aberdeen.

98 2.2. Identification, cloning and sequence analysis of Mx cDNA in rainbow trout

99 Three Mx genes (Mx1-3) are known in rainbow trout [6-7]. To identify additional Mx genes in this
100 species, we searched the recently released rainbow trout reference genome (GCF_002163495.1)
101 using TBLASTN [33] with the known trout Mx genes as query, resulting in the identification of four
102 genomic loci (Chromosomes (Ch)3, 11, 17 and 24) that harbour Mx genes. The Mx genes were then
103 predicted as described previously [34-35]. In addition, potential exons in untranslated regions (UTR)
104 were predicted by using trout RNA-seq datasets (SRP108798) through aligning to the reference
105 genome. Primers (supplementary **Table S1**) were subsequently designed in the predicted 5'- and 3'-
106 UTR for PCR cloning of the complete coding region of each predicted Mx gene. The general cloning
107 and sequence analysis was as described previously [34-35]. The nucleotide sequences generated
108 were assembled and analysed with the AlignIR programme (LI-COR, Inc.). Homology search was
109 performed using the BLAST program (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) [33] and the gene
110 organization was predicted using the Spidey program at NCBI. Protein prediction was undertaken
111 using software at the ExPASy Molecular Biology Server (<http://www.expasy.org/tools>) [36].
112 Multiple sequence alignments were generated using CLUSTALW [37]. Amino acid sequence
113 identity/similarity comparison was performed using the scoring matrix BLOSUM62 within the
114 MatGAT program, with a gap open penalty of 10 and gap extension penalty of 1 [38].

115 2.3. Analysis of Mx genes in other salmonids

116 The Mx genes in other salmonids were predicted/analysed using recently released genomes of
117 Atlantic salmon (*Salmo salar*, Atlantic, acc. no. GCF_000233375.1), chinook salmon (*Oncorhynchus*
118 *tshawytscha*, Chinook, acc. no. GCF_002872995.1), coho salmon (*Oncorhynchus kisutch*, Coho, acc.

119 no. GCF_002021735.1), and Arctic charr (*Salvelinus alpinus*, Charr, acc. no. GCF_002910315.2).
120 Each Mx aa and nucleotide sequence was mapped to chromosomes/scaffolds. Similarly, Mx genes
121 were analysed in the pike (*Esox lucius*) reference genome (acc. no. GCF_000721915.3), the closest
122 relative of salmonids that has not undergone the salmonid 4R WGD and that has a sequenced
123 genome. The aa sequences were used for phylogenetic tree analysis using MEGA7.0 software [39]
124 based on aa multiple alignments generated by CLUSTALW. The evolutionary distances were
125 computed using the JTT matrix-based method. A neighbour-joining phylogenetic tree was
126 constructed using pair-wise deletion option.

127 **2.4. Evolutionary analysis of teleost Mx family**

128 Mx genes/proteins were analysis at NCBI from selected teleost fish, including species known to
129 possess multiple Mx genes. The naming of Mx genes/proteins followed those already published [4,
130 16, 40-41] or simple Mx with an acc. no. For phylogenetic tree analysis, Mx protein sequences were
131 extracted from one holostean species, spotted gar (*Lepisosteus oculatus*, Lepisosteiformes) that is an
132 early actinopterygian fish species without the 3R WGD, twenty-one teleosts and three mammals
133 (human *Homo sapiens*, mouse *Mus musculus* and cow *Bos taurus*) as an outgroup. The teleost
134 species included an elopomorph, European eel (*Anguilla anguilla*, Anguilliformes), five Ostariophysi
135 (Otophysi) species including three Cypriniformes fish (common carp *Cyprinus carpio*, goldfish
136 *Carassius auratus* and zebrafish *Danio rerio*), channel catfish (*Ictalurus punctatus*, Siluriformes),
137 and Mexican tetra or blind cave fish (*Astyanax mexicanus*, Characiformes), five protacanthopterygii
138 (the salmonids and pike described above), and ten percomorphs including two Pleuronectiformes
139 (turbot *Scophthalmus maximus*, and olive flounder *Paralichthys olivaceus*), Atlantic killifish
140 (*Fundulus heteroclitus*, Cyprinodontiformes), stickleback (*Gasterosteus aculeatus*,
141 Gasterosteiformes), Medaka (*Oryzias latipes*, Beloniformes), fugu (*Takifugu rubripes*,
142 Tetraodontiformes), Nile tilapia (*Oreochromis niloticus*, Cichliformes) and three Perciforme fish
143 (gilt-head sea bream *Sparus aurata*, orange-spotted grouper *Epinephelus coioides*, and the Asian sea
144 bass *Lates calcarifer*). A neighbour joining phylogenetic tree was constructed as above. Synteny
145 analysis was performed using the Genomicus program [42] or with information extracted from
146 reference genome sequence at NCBI.

147 **2.5. Real-time PCR analysis of gene expression**

148 Specific primers for each Mx gene were carefully designed based on a multiple cDNA sequence alignment to
149 ensure that at least one primer was isoform specific, and one primer crosses an intron to prevent genomic
150 DNA amplification. The primers for qPCR analysis of Mx genes and other cytokine genes are detailed in

151 Table S1 and S2, respectively. Total RNA preparation, cDNA synthesis and qPCR analysis were as described
152 previously [43]. The expression of each gene was first normalized to that of the house keeping gene
153 elongation factor-1 α (EF-1 α). To directly compare the expression level of the different Mx paralogues, a
154 reference was constructed using equal molar amounts of PCR product from each gene, including EF-1 α .

155 **2.6. Tissue distribution of rainbow trout Mx gene family**

156 Six healthy rainbow trout (~140 g) were killed and seventeen tissues (blood, thymus, gills, scales,
157 skin, muscle, tail fins, adipose fin, brain, adipose tissue, spleen, liver, heart, intestine, gonad, head
158 kidney (HK) and caudal kidney) were collected and processed as described previously [34-35]. The
159 relative expression level of Mx genes in each sample was normalized against the expression level of
160 EF-1 α and expressed as arbitrary units (AU) where 1 AU = the expression level of EF-1 α /1,000,000.

161 **2.7. Ontogeny of the expression of the Mx gene family**

162 To investigate if the expression of Mx is correlated to immune capacity in early life, the ontogeny of the
163 expression of Mx genes was examined. Archived samples from a previous experiment were used in this study
164 as detailed in Wang et al. [44]. Briefly, eyed eggs, immediate post-hatch fry, pre-first feeding (Pre-feeding)
165 fry at the stage of full disappearance of the yolk sac, and fry 3 weeks following first feeding were sampled and
166 cDNA prepared. Six samples for each developmental stage were prepared. The qPCR quantification of gene
167 expression was as described above.

168 **2.8. Production of recombinant trout type I IFN α**

169 The cDNA sequence encoding the mature peptide of trout IFN α was amplified from a poly IC
170 stimulated cDNA sample using the primers IFN α F (TGTGACTGGATCCGACACCAT) and IFN α R
171 (GTACATCTGTGCTGCAAGGATATCC). The amplified product was cloned to a pTriEx vector
172 (Novagen) as described previously [45]. Sequence analysis of the construct used for recombinant
173 protein production revealed that it encodes a His-tag (MAHHHHHHHG) at the N-terminus
174 followed by the 152 aa mature peptide identical to XP_021480273. Thus, the recombinant trout IFN α
175 was 163 aa with a calculated molecular weight of 19.5 kDa and a theoretical pI of 9.17. A sequence
176 confirmed plasmid was transformed into BL21 Star (DE3) competent cells (Invitrogen). The protein
177 was produced, purified under denaturing conditions, refolded, and quantified as described previously
178 [34,43,45]. The refolding buffer was phosphate buffered saline (PBS, pH7.4, Sigma, UK) containing
179 10% glycerol, 0.5 M arginine monohydrochloride, and 5 mM 2-mercaptoethanol (2-ME). The
180 purified protein was buffer changed using a centrifugal concentrator (10 kDa cutoff, Thermo
181 Scientific). The storage buffer was PBS (pH7.4) containing 10% glycerol, 2 mM EDTA, 10 mM

182 arginine monohydrochloride, 10 mM glutamine, and 5 mM 2-ME. After sterilization with a 0.2 µm
183 filter, the recombinant protein was aliquoted and stored at -80°C ready for bioactivity analysis.

184 **2.9. Stimulation of cell lines with PAMPs and recombinant cytokines**

185 Three trout cell lines, a macrophage-like cell line RTS-11 from spleen [46], a fibroblast-like cell line RTG-2
186 from gonad [47], and an epithelial-like cell line RTGill from gills [48] were used for *in vitro* stimulation. All
187 the cells were maintained at 20°C in Leibovitz medium (L-15) supplemented with 100 U/ml penicillin and
188 100 µg/ml streptomycin (P/S), and 10% (for RTG-2 and RTGill cell lines) or 30% (for RTS-11 cells) foetal
189 bovine serum (FBS). The cells were seeded at 1x10⁶ cells/ml (RTS-11) or 0.5x10⁶ cells/ml (RTG-2 and
190 RTGill) in L-15 containing 10% FCS at 2 ml/well in 12-well cell culture plates overnight before stimulation.

191 RTS-11 cells were first stimulated with pathogen-associated molecular patterns (PAMPs), the bacterial cell
192 wall component lipopolysaccharide (LPS, from *E. coli* strain 055:B5, Sigma) and the viral dsRNA mimic
193 polyinosinic: polycytidylic acid (poly IC, Sigma). The stimulants were added to the cells at 25 µg/ml for LPS
194 and 50 µg/ml for poly IC, or medium alone as control. The treatments were terminated by dissolving the cells
195 in TRI reagent (Sigma, UK) 4 h, 8 h and 24 h post-stimulation. Total RNA isolation and gene expression
196 analysis was as described above.

197 The RTS-11 cells were then stimulated with five trout recombinant cytokines, IFNγ (20 ng/ml) [49], IFNα (25
198 ng/ml) prepared above, IL-1β (25 ng/ml) [50], IL-6 (100 ng/ml) [51] and TNFα (50 ng/ml) [52], or medium
199 alone as control. The treatments were terminated at 4 h, 8 h and 24 h and gene expression analysed as above.

200 Finally, RTG-2 and RTGill were stimulated with IFNγ (20 ng/ml) and IFNα (25 ng/ml) for 4 h and gene
201 expression analysed as above.

202 **2.10. Modulation of Mx gene expression *in vivo* by poly IC**

203 Poly I:C (Sigma, UK) was dissolved at 10 mg/ml in sterile cell culture-grade water, stored at -80 °C and
204 diluted to 5 mg/ml in PBS before intraperitoneal (ip) injection. Trout (~100 g, N=24) were injected
205 intraperitoneally (ip) with 1 mg poly IC in 0.2 ml of PBS, or the same amount of PBS as control. Six fish from
206 each group were killed at 6 h and 24 h post injection, and spleen, HK, gills and intestine were collected for
207 gene expression analysis as described previously [53]. The time points chosen were based on past studies of
208 the rapid PAMP response *in vivo* in rainbow trout [54]. The expression was expressed as AU after
209 normalisation with EF-1α, where 1 AU = the average expression level in control fish at 6 h in each tissue.

210 **2.11. Statistical analysis**

211 The data were statistically analyzed using the SPSS Statistics package 24 (SPSS Inc., Chicago, Illinois). The
212 analysis of real-time PCR data was as described previously (43). To improve the normality of data, real-time

213 quantitative PCR measurements were scaled, with the lowest expression level in a data set defined as 1, and
214 log₂ transformed. One way-analysis of variance (ANOVA) and the LSD post hoc test were used to analyse
215 the gene expression data, with $P \leq 0.05$ between treatment and control groups considered significant.

216

217

218 **3. Results**

219 **3.1. Identification, cloning and sequence analysis of Mx gene family in rainbow trout**

220 In addition to the known Mx1-3 in rainbow trout, six additional Mx genes (Mx4-9) have been identified and
221 cloned in this study (Supplementary Figs. S1-S6, acc. nos. MK301134-MK301139). Mx4, as with Mx1-3, was
222 located on Ch17 and was located between Mx2 and Mx3. Mx5-6, Mx7-8 and Mx9 were located on Ch3, Ch11
223 and Ch24, respectively (**Table 1**).

224 Each trout Mx cDNA sequence had a complete open reading frame that encoded for 635, 614, 606, 613, 608,
225 and 640 aa for Mx4-9, respectively. Each trout Mx had a N-terminal dynamin GTPase domain, and a C
226 terminal GTPase effector domain, that were well conserved as shown in a multiple alignment of the 9 trout
227 and two human Mx proteins (**Fig. 1**). The tripartite GTP-binding motif (GDXXSGKS, DLPG, and TKPD) in
228 all trout Mx were identical to human MxA and MxB. The dynamin signature (LPRXXGXXTR), and the
229 leucine residues that form leucine zipper folds in the GTPase effector domain, were also conserved (**Fig. 1**).
230 The middle domain and the GTPase effector domain of Mx fold into a four-helical bundle that constitutes a
231 stalk that mediates oligomerization and transmits conformational changes from the G domain to the target
232 structure [55]. The regions forming the helix, and loops L2 and L3 were all conserved. However, relatively
233 large differences were present in loops L1 that connects the N and C-terminal of the helix $\alpha 1$ and introduces a
234 kink, and L4 that connects the helix $\alpha 3$ and $\alpha 4$ (**Fig. 1**). Potential nuclear localisation signals (KKRKR) are
235 present in trout Mx2 and Mx4 in L4, where a lysine motif (KKKK) is also present in human MxA that
236 contribute to membrane association of MxA [2].

237 **3.2. Sequence analysis of Mx family in salmonids**

238 Nine Mx genes (Mx1-9) have been described recently in Atlantic salmon [27] that map to three chromosomes
239 (Ch9, 12 and 25, **Table 1**). In addition, a partial sequence for Atlantic Mx10 (XP_013998960) has been
240 mapped to Ch15 (**Table 1**). At least 6 Mx genes each in chinook salmon and coho salmon, and 10 Mx genes
241 in Arctic charr could be identified and mapped to chromosomes or scaffolds (**Table 1**). Partial sequences for
242 three pike Mx genes were also found, with Mx1 on Ch17 and Mx2-3 on Ch13 (**Table 1**).

243 A phylogenetic tree constructed using all the known salmonid Mx and the three pike Mx protein sequences
244 showed that the salmonid Mx family clustered into four separate clades (**Fig. 2A**). Moreover, salmonid Mx
245 genes are located on four cognate chromosomes, at least in rainbow trout, Atlantic, chinook and coho salmon,
246 in which their genome sequences have been mapped to chromosomes. The Mx genes on the same
247 chromosome are grouped together (**Fig. 2B**), as seen also in Atlantic salmon [27], suggesting that multiple
248 genes on the same chromosome originate from local gene duplication events in each species. Thus there are
249 four salmonid Mx groups (SMG, **Fig. 2**). Pike Mx1 was grouped with SMG1, whilst pike Mx2 and Mx3,
250 which are linked on Ch13, were grouped with SMG2. SMG1 consisted of Mx1-4 of trout, Mx1-3 of Atlantic,
251 chinook and coho salmon, and Mx1-2 of charr. SMG2 consisted of trout Mx5-6, Atlantic salmon Mx4-8,
252 chinook and coho salmon Mx4, and charr Mx3-8. SMG3 contained trout Mx7-8, Atlantic salmon Mx10,
253 chinook and coho salmon Mx5 and char Mx9-10. SMG4 had trout and Atlantic salmon Mx9, and chinook and
254 coho salmon Mx6 (**Fig. 2**).

255 It is notable that trout Mx1/3 and Mx2/4, along with their cognate salmonid Mx molecules formed two
256 separate branches with high bootstrap support in SMG1 group (**Fig. 2A**), suggesting that the existence of these
257 genes, or their ancestral gene predates salmonid speciation. A similar situation was also observed with trout
258 Mx7 and Mx8 in SMG3 group (**Fig. 2A**). Although more Mx genes might still be found, the data for
259 salmonids with an advanced (sequenced) genome suggests that the distinct numbers of Mx genes in SMG1-3
260 are due to species-specific independent local gene duplication or deletion events after salmonid speciation.

261 In agreement with four SMGs in the phylogenetic tree, the Mx aa sequences within each SMG share high aa
262 identities (**Table 2**). In SMG1, trout Mx1-4 share high aa sequence identities between each other (86.3-98.4%)
263 in similar range to SMG1 Mx from different salmonids (83.4-98.2%), but have relatively low identities to Mx
264 from SMG2 (43.8-47.0%), SMG3 (41.5-48.1%) and SMG4 (42.1-48.1) (**Table 2**). Similarly, Mx sequences
265 share high identities within SMG2 (82.7-93.6), SMG3 (57.8-93.8%) and SMG4 (80.2-97.0%). However, the
266 identities of Mx between SMGs are similarly low (41.5-51.6%) with the exception of Mx molecules in SMG2
267 and SMG3 that share moderate 53.4-71.2% aa identities (**Table 2**). Furthermore, the Mx bearing
268 chromosomes in rainbow trout (Ch3, 11, 17 and 24) and Atlantic salmon (Ch9, 12, 15 and 25) do not share
269 syntenic origins [56-57]. These data suggest that the four Mx-bearing chromosomes do not appear to originate
270 from the salmonid 4R WGD.

271 3.3. Phylogenetic tree analysis of Mx in vertebrates

272 To understand how the four SMGs evolved, we analysed the Mx gene family in other vertebrates with a focus
273 on teleost Mx genes. Three Mx genes, Mx1-2 on Ch17 and Mx3 on Ch3, are present in spotted gar, an early
274 Actinopterygian (Holostei) that has not undergone the 3R WGD that may represent an ancestral state [4, 27].
275 A neighbour-joining phylogenetic tree was constructed based on a multiple alignment of Mx proteins from
276 selected mammalian and teleost fish species. In agreement with previous studies, mammalian Mx form an

277 independent group separate from all teleost Mx molecules (**Fig. 3**). Three teleost Mx groups (TMG) can be
278 identified, with a gar Mx at the root of each clade. TMG1 contained gar Mx1, salmonid SMG1 and Mx from
279 all the major teleost groups (**Fig. 3**). TMG2 contained gar Mx3, salmonid SMG2, SMG3 and SMG4, and Mx
280 molecules from European eel, zebrafish, goldfish, Mexican tetra and catfish. TMG3 contained gar Mx2, and
281 Mx from Cypriniformes (zebrafish, common carp and goldfish) and Characiformes (tetra) (**Fig. 3**). This
282 phylogenetic tree may suggest that the 3R WGD duplicated 6 Mx genes (from the 3 ancestral Mx genes
283 present in gar) that have subsequently undergone lineage specific deletion, with Cypriniformes and
284 Characiformes species retaining a copy of each of the duplicates, protacanthopterygii such as the salmonids
285 and pike retained two whilst in the majority of teleosts, the percomorphs only one is present. Within a species
286 the numbers of Mx genes might be increased again by local gene duplication.

287 **3.4. Synteny analysis of Mx locus in vertebrates**

288 Despite much analysis, the evolutionary relationship of Mx genes in different vertebrates is still unclear [4,16].
289 In the present study we performed a synteny analysis using the most advanced genomes available. Pike Ch17
290 (Mx1) and trout Ch17 (Mx1-4), and pike Ch13 (Mx2-3) and trout Ch11 (Mx7-8) share a considerable syntenic
291 relationship (**Fig. 4**). However, trout Ch3 (Mx5-6) and Ch24 (Mx9) share no clear syntenic relationships to
292 pike Mx loci, but have a good relationship instead with gar Ch17 (Mx1-2) and Ch3 (Mx3), respectively (**Fig.**
293 **4**).

294 Interestingly, the Gar Mx3 (Ch3) locus also has considerable synteny with zebrafish Ch15 (MxF) and Ch25
295 (MxD, G1 and G2), in addition to the trout Mx9 locus, and all the Mx residing in these loci belong to TMG2.
296 Furthermore, zebrafish Ch15 and 25 combined share a perfect match syntenically to gar Ch3, suggesting a
297 break of the ancestral gar-like derived chromosome in zebrafish. The gar Mx3 locus also shares synteny with
298 the tetrapod Mx locus, eg. human Ch21 (MxA and MxB) (**Fig. 4**), as also reported by Robertsen et al. [27].
299 This suggests that a gar Mx3-like ancestral locus gave rise to the teleost Mx group loci in zebrafish and
300 salmonids, and led to the tetrapod Mx locus.

301 The gar Mx1-2 locus shares synteny to tetra Ch12 (Mx1-7) and the combined tetra Ch19 (Mx8) and scaffold
302 NW_019172839 (Mx9) of tetra, indicating the retention of two 3R derived Mx loci in this species. Similarly,
303 the gar Mx1-2 locus shares synteny to both zebrafish Ch1 (MxA/B) and Ch9 (MxC/E) (**Fig. 4**). It is notable
304 that the tetra Ch12 Mx locus has Mx genes that belong to TMG1 (Mx1) as well as TMG2 (Mx2-7), whilst the
305 zebrafish TMG2 resides in Ch15 and 25 derived from gar Ch3. Taken as a whole, two ancestral gar-like Mx
306 loci gave rise to the current teleost Mx loci in a lineage-specific manner.

307 **3.5. Gene organisation analysis of Mx genes in vertebrates**

308 To shed more light on the evolution of the three teleost Mx groups, we analysed the gene organisation of all
309 trout Mx genes in comparison with Mx genes from other teleosts and humans. All exon-intron boundaries of

310 trout Mx genes conformed to the consensus sequences (GT/AG). In TMG1, trout Mx1-4 genes all had a 12
311 coding exon/11 intron structure with identical intron phase. The coding region of exons were identical with
312 the exception of exon 8 and 11 (**Fig. 5**). A similar gene organisation of the coding region was observed with
313 other TMG1 genes, eg. zebrafish MxA and MxB that are on the same chromosome, tetra Mx1 and gar Mx1,
314 with the exception of a non-coding exon in the 5'UTR of the gar and tetra Mx gene in this group (**Fig. 5**).

315 Trout Mx5-9 belong to TMG2. They all had 13 coding exons with the coding regions of exons well conserved,
316 with the exception of the first and the last two exons (**Fig. 5**). Compared to TMG1 genes, the last eleven intron
317 phases were identical to that of TMG1 genes. The main difference in gene organisation was an extra N-
318 terminal coding exon that brought a phase II intron in Mx5-9 that was missing in Mx1-4. This gene
319 organisation was conserved in other TMG2 Mx genes except gar Mx3, that shared the same gene organisation
320 with TMG1 Mx genes (**Fig. 5**).

321 Human MxA and MxB also had the same 13 coding exon structure as TMG2 but with the first intron in phase
322 0 (**Fig. 5**). Interestingly, some TMG3 Mx genes had the same gene organisation as in humans (zebrafish MxE
323 and gar Mx2), and others (eg. zebrafish MxC and tetra Mx8) had the same as in TMG1 (**Fig. 5**). Some of
324 TMG3 Mx genes have lost the last third exon.

325 In general, the exon size and intron phase in the regions encoding for the N-terminal GTPase domain, the
326 middle domain and the C-terminal GTPase effector domain are well conserved. The noticeable variations in
327 size were the 5th last exon that encodes the L1 loop, and the second last exon that encodes for the L4 loop (**Fig.**
328 **5**).

329 **3.6. The expression of rainbow trout Mx family in tissues and cell lines**

330 The expression of each trout Mx gene was comparatively studied using gene specific primers and
331 serial dilutions of references, and expressed as arbitrary units (AU) relative to EF-1 α expression.
332 Thus the AU of the relative expression is on an equal molar basis. The expression of paralogues on
333 the same chromosome was grouped together, and the tissues were ordered according to the average
334 expression level of Mx1 (**Fig. 6**). The expression level of Mx1-4 on Ch7 was medium (AU =100 to
335 1,000) to high (AU > 1,000) across tissues (**Fig. 6A**). Mx5 and Mx6 expression was detectable in all
336 the seventeen tissues but at low levels (AU <100). The exceptions were Mx5 in intestine that was at
337 a high level, and Mx5 in thymus, gills, adipose fin, tail fins, spleen, scales and gonad, and Mx6 in
338 intestine and gills that was at medium expression levels (**Fig. 6B**). Mx8 expression was also
339 detectable in all tissues examined but at low levels except for high level expression in intestine and
340 medium level expression in thymus, brain and gonad. Mx7 expression was undetectable in head/
341 caudal kidney and tail fins, but had high level expression in intestine, medium level in thymus, with

342 low levels in other tissues (**Fig. 6C**). Mx9 expression was also high in intestine but low or
343 undetectable in other tissues (**Fig. 6D**).

344 Each Mx gene was differentially expressed across different tissues. In the same tissue, the majority
345 of Mx genes had varying expression levels, as shown by the ratio of different genes and paired-
346 samples T tests (**Table S3**). In general, the expression of the Mx1-4 and Mx5-9 was more similar
347 within each group than between groups. It is noteworthy that blood expressed highest levels of Mx1-
348 4 but low levels of Mx5-9. In contrast, intestine expressed highest levels of Mx5-9 genes amongst
349 the tissues examined (**Fig. 6**).

350 The constitutive expression of the trout Mx gene family was also examined in three trout cell lines.
351 The macrophage-like cell line RTS-11 expressed all the Mx genes at low level (**Fig. 6**). The
352 fibroblast-like cell line RTG-2 and epithelial-like cell line RTGill expressed medium levels of Mx2
353 and Mx3, and low levels of other Mx genes but Mx5 and Mx8 in RTG-2 and Mx9 in RTGill were
354 not detectable (AU < 1).

355 **3.7. Transcript expression of Mx gene family during developmental stages**

356 The high levels of expression of Mx gene family members in blood and intestine suggest an important role in
357 immune defence. We next examined the expression of these genes in eyed-eggs, immediately post-hatch fry,
358 pre-first feeding fry or fry 3 weeks after first feeding, which represent a critical period when the fish encounter
359 potential pathogens from the environment and food [44]. The expression levels of all Mx genes were
360 maintained from eyed-eggs till post-hatch. The expression of Mx1, Mx8 and Mx9 was increased in pre-
361 feeding fry and maintained at the same levels in post-feeding fry (**Fig. 7**). Mx5 expression was low in eyed-
362 eggs and post-hatch fry but increased significantly in pre-feeding fry and increased further in post-feeding fry.
363 The expression of Mx2 and Mx5 was only increased in post-feeding fry whilst that of Mx3, Mx4 and Mx6
364 was unchanged across the different developmental stages (**Fig. 7**).

365 **3.8. Modulation of the expression of trout Mx and proinflammatory cytokine genes *in vivo* by poly IC**

366 Poly IC, a known strong inducer of Mx expression, was used to investigate its ability to modulate Mx
367 expression *in vivo*. The expression of Mx genes was examined in two major systemic lymphoid tissues, the
368 spleen and HK, and two mucosa-associated lymphoid tissues, the gills and intestine. The expression of Mx1-4
369 was induced in all four tissues at both 6 h and 24 h post poly IC injection, with the exception of Mx1 and Mx3
370 in intestine at 6 h (**Fig. 8A-D**). As seen *in vitro*, poly IC did not increase Mx9 expression *in vivo* (**Fig. 8I**). The
371 induction of other Mx (5-8) genes was time- and tissue-dependent (**Fig. 8E-H**). In the spleen, poly IC
372 increased Mx5 and Mx7 expression at 24 h and Mx6 and Mx8 expression at both time points. In the HK, poly
373 IC increased Mx5 and Mx8 expression at both time points, and Mx6 expression at 24 h, but had no effect on

374 Mx7 expression. In the gills, poly IC increased the expression of Mx6-8 at 24 h, and Mx5 at both time points.
375 In the intestine that has high constitutive expression of Mx5-8, poly IC increased Mx5 expression at both time
376 points and Mx7 expression at 6 h, but had no effects on Mx6 and Mx8 (**Fig. 8E-H**). In summary, poly IC was
377 also a strong inducer of Mx gene expression *in vivo* with highest induction seen at 24 h post-injection (**Fig. 8**).

378 In addition to inducing Mx gene expression, poly IC also induced the expression of many pro-inflammatory
379 cytokines, including IL-1 β 1-2, IL-6, IL-8, TNF α 1-2, IFN α 1, IFN γ and CXCL11, at least at one time point, in
380 all the four tissues examined (**Fig. 9**). In contrast to the later (at 24 h) peak induction of Mx gene expression,
381 poly IC induced an early (6 h) induction of the majority of the proinflammatory cytokines studied (eg IL-1 β 1-
382 2, IL-6, IL-8, TNF α 2 and IFN α 1) (**Fig. 9**).

383 **3.9. Modulation of trout Mx expression in RTS-11 cells by PAMPs**

384 The expression of all trout Mx genes was detectable in the macrophage cell line RTS-11 (**Fig. 6**). Thus we
385 examined the modulation of trout Mx gene family members first in this cell line using poly IC and LPS,
386 classical viral and bacterial PAMPs. Poly IC was a strong inducer of Mx gene expression. It significantly
387 induced the expression of Mx2, Mx3 and Mx4 from 4 h to 24 h, that of Mx1, Mx5 and Mx6 from 8 h to 24 h,
388 and that of Mx7 and Mx8 at 8 h, but had little effect on Mx9 expression (**Fig. 10**). As expected, LPS had only
389 minor effects on Mx gene expression; it induced a small upregulation of Mx4 at 8 h and Mx5 at 24 h, and a
390 small downregulation of Mx1 at 24 h and Mx2 at 4 h (**Fig. 10**).

391 **3.10. Modulation of Mx expression by proinflammatory cytokines in RTS-11 cells**

392 The early peak induction of proinflammatory cytokine expression and late peak induction of Mx genes may
393 suggest that poly IC can induce Mx indirectly via proinflammatory cytokines as well as by virus sensing
394 pathways. Indeed, IFN γ has been shown recently to modulate some of the Mx isoforms in Atlantic salmon
395 [27]. Hence, the possibility of modulation of Mx gene expression by IFN α 1, IFN γ , IL-1 β , IL-6 and TNF α was
396 studied using RTS-11 cells. Mx9 expression was refractory to all the cytokines (**Fig. 11I**). However, the
397 expression modulation of the other Mx genes was cytokine-specific. IFN α induced the expression of Mx1-4
398 and Mx6 from 4 h to 24 h, Mx5 at 4 h and 24 h, and Mx7 at 8 h, but had no effects on Mx8 (**Fig. 11**). IFN γ
399 induced the expression of Mx2-6 from 4 h to 24 h, Mx7-8 at 24 h, but decreased Mx1 expression at 24 h (**Fig.**
400 **11A-H**). IL-1 β induced the expression of Mx3-4 and Mx8 from 4 h to 24 h, Mx2 and Mx6 at 4 h and 8 h, Mx5
401 at 24 h, but had no effects on the expression of Mx1 and Mx7 (**Fig. 11A-H**). IL-6 increased the expression of
402 Mx3 at 8 h and 24 h, Mx5 at 24 h, and Mx6 at 4 h, but decreased the expression of Mx 1 and Mx4 at 24 h, and
403 Mx8 at 8 h and 24 h. It had no effects on Mx2 and Mx7 (**Fig. 11A-H**). TNF α induced the expression of Mx2
404 and Mx4 at 8 h, Mx3 from 4 h to 24 h, Mx5 at 24 h, but decreased Mx1 expression at 24 h and Mx5
405 expression at 8 h. It had no effects on Mx6-8 (**Fig. 11A-H**). It is noteworthy that IFN α is a strong inducer of
406 the expression of Mx1-4 and Mx7, IFN γ is a strong inducer of Mx5-6 expression and IL-1 β a strong inducer

407 of Mx8 (**Fig. 11A-H**). In conclusion, all the Mx genes except Mx9 can be modulated by multiple antiviral and
408 proinflammatory cytokines in an Mx- and cytokine-dependent manner.

409 **3.11. Modulation of Mx expression by type I and type II IFNs in RTG-2 and RTGill cell lines**

410 The cytokine-dependent Mx modulation may also be cell-type dependent. Thus, the IFN modulated Mx gene
411 expression was further studied in the fibroblast-like cell line RTG-2 and the epithelial like cell line RTGill. In
412 RTG-2 cells, both IFN γ and IFN α induced the expression of Mx1-6, with similar potency for Mx5 and Mx6.
413 However, IFN α was more potent for Mx1-4 (**Fig. 12**). In RTGill, both IFN γ and IFN α induced the expression
414 of Mx2 and Mx3, with IFN γ more potent for Mx2, but had no effects on Mx1 and Mx4 (**Fig. 12A-D**). Only
415 IFN γ but not IFN α induced the expression of Mx5 and Mx6 in RTGill cells (**Fig. 12E-F**). The expression of
416 Mx7-9 was low and refractory (data not shown).

417

418 **4. Discussion**

419 This study reveals that at least 9 active Mx genes are present in the rainbow trout genome, the same number as
420 reported recently in Atlantic salmon [40]. However, in this study we show that there are in fact 4 Mx loci
421 present in salmonids and that the number of Mx genes at each locus differs between these two species at 3 of
422 these loci. Multiple Mx genes are also present in other salmonids at four chromosomal loci. The salmonid Mx
423 genes at the same genomic locus share high sequence identities within and between species, suggesting they
424 arose from local gene duplication events. It seems that local Mx gene duplication/gene loss is common with
425 some duplication events likely to have happened before salmonid speciation, eg. duplication of Mx1/3 and
426 Mx2/4 in SMG1, and Mx7 and Mx8 in SMG3, but others after salmonid speciation, eg. Atlantic salmon Mx4-
427 8 in SMG2. The four Mx bearing chromosomal loci could have arisen from the 3R and 4R WGDs as seen
428 with other genes when mammals have one and salmonids have 4 [28]. However, sequence homology, synteny
429 and phylogenetic tree analysis do not clearly support this, and past models [4] do not adequately explain their
430 evolutionary path in bony fish.

431 Multiple Mx genes (up to 10) can be found in many teleost species. Our phylogenetic tree analysis indicates
432 that three TMG exist. TMG1 are present in different teleost lineages, but TMG2 and TMG3 are found in only
433 more basal teleosts [57]. Each TMG has a unique gene organisation in terms of coding exon number and the
434 first intron phase. For example, whilst TMG1 has a 12 coding exon structure with the first intron in phase I,
435 TMG2 has 13 coding exons with the first intron in phase II, and TMG3 has either 13 coding exons with the
436 first intron in phase 0 (as seen with mammalian Mx genes) or the same organisation as in TMG1. Interestingly,
437 the spotted gar possesses three Mx genes, with one present in each TMG.

438 Although the four cognate Mx chromosomal loci between salmonids are well conserved, no clear syntenic
439 conservation have been observed in trout and other salmonid species between the four Mx loci. However, a
440 syntenic relationship between the two gar Mx loci and those in zebrafish/ tetra is apparent. For example, the
441 Gar Mx1-2 locus and tetra Mx loci on Ch12 and Ch19, that harbour Mx genes in all the three teleost Mx
442 groups. However, the zebrafish cognate Mx loci of gar Mx1-2 only have Mx genes that belong to TMG1 and
443 TMG3, and the zebrafish TMG2 locus shares synteny to the gar Mx3 locus. The two gar Mx loci also share
444 apparent synteny with two of the trout Mx loci whilst the other two show a syntenic relationship with the two
445 pike Mx loci. This complex syntenic relationship may suggest that the current Mx genes in 3R or 4R teleosts
446 may have arisen from the three Mx genes present at two chromosomal loci as seen in spotted gar, with the 3R
447 duplicated Mx loci retained/lost in a lineage specific manner (**Fig. 13**). This model differs from that in Qi et al.
448 [4] in taking into account the number of loci present in actinopterygians as well as Mx copy number.

449 The increased copy number of Mx genes seen in many teleosts may confer increased expression level and
450 hence heightened antiviral defence. The duplicated copies may also acquire novel sequence properties that
451 confer anti-viral specificity and efficiency. The nine trout Mx genes have considerable variation in the
452 nucleotide sequence coding for the L1 and L4 loops in the stalk, as seen in the multiple aa alignments and
453 their gene organisation. Both L1 and L4 are at the surface of the stalk [3] that can interact with surrounding
454 proteins and may be involved in interaction with viral components. L4 of mammalian Mx is a critical
455 determinant of viral substrate specificity [58-59]. The diversification of these regions might have been driven
456 by past virus exposure and life history traits of different species. For example, zebrafish and tetra have short
457 life cycles but live in diverse changing water environments. Their survival depends heavily on innate
458 immunity against viral pathogens. Salmonids survive successfully in both fresh and marine waters, and may
459 encounter a larger virus repertoire compared to species living in only fresh water or marine water. Hence, the
460 increase of copy number and types of Mx genes in these species may confer a fitness advantage.

461 Mx antiviral effects depend on where the Mx protein is present. Thus, the mouse Mx1 protein which is
462 localized in the nucleus mainly inhibits orthomyxoviruses that replicate in the nucleus, whereas mouse Mx2 is
463 confined to the cytoplasm and inhibits viruses with an exclusively cytoplasmic replication phase [60]. There is
464 a potential NLS in the L4 of some salmonid Mx proteins eg. trout Mx2, and Mx4, but not Mx1 and Mx3. This
465 NLS may indeed be functional as trout Mx2 is found in the nucleus, and Mx1/ Mx3 in the cytoplasm [7]. This
466 suggests the nuclear presence of trout Mx4. Taken as a whole, salmonids, such as rainbow trout are equipped
467 with a battery of diversified Mx genes with their protein products present in the cytoplasm and nucleus to
468 protect themselves from viral attack during their life cycle.

469 Investigation of Mx isoform expression will help understand their functional roles. Although multiple Mx
470 genes have been identified in several teleost species, a comparative expression study in healthy fish at the
471 individual gene level is lacking [17, 27]. Our results show that the nine trout Mx genes were differentially
472 expressed across different tissues and cell lines, as outlined below, suggesting a level of neofunctionalisation

473 of trout Mx paralogues through variation in expression patterns. The high levels of Mx1-4 transcript in blood
474 and Mx5-9 in intestine is of particular interest. Many different viruses can infect hosts via the intestine to
475 cause acute infectious gastroenteritis, or get access to the blood by physical breaches (wounds) or during
476 viremia [61]. So preventing their spread at these sites is a good antiviral strategy. The differential expression
477 of Mx genes in the cell lines may suggest that specific cell types preferentially express a particular Mx gene or
478 a set of Mx genes to defend against potential cell type-tropic viruses. However, the three cell lines examined
479 expressed relatively low levels of Mx genes compared to the tissues analysed, perhaps due to the need for
480 humoral factors present *in vivo* to maintain high level Mx gene expression.

481 The expression of Mx genes was also studied during development, and several were increased in pre-feeding
482 and post-feeding fry (eg Mx1, Mx2, Mx5, Mx7-9). First feeding is a critical stage in the life of a fish, when
483 potential food borne viruses are met directly for the first time and when the adaptive immune system has not
484 fully developed. Indeed, it was the genes preferentially expressed in the intestine in adults that were increased
485 in the post-hatch fry.

486 Next we studied whether the Mx genes could be modulated by PAMPS or cytokines. In agreement with Mx
487 induction in other species, poly IC was a strong inducer of trout Mx1-8 gene expression *in vitro* and *in vivo*,
488 with Mx9 more refractory. Although the induction patterns *in vivo* were gene- and tissue-dependent, highest
489 expression was seen at 24 h with most of the Mx genes. Injection of poly IC also induced the expression of
490 proinflammatory cytokines, such as IL-1 β , IL-6 and TNF α as well as type I and type II IFNs. In contrast to the
491 late peak of induction of Mx gene expression, poly IC caused an early peak of expression in the cytokines
492 studied. Therefore, the later peak in Mx expression could be influenced by such molecules. To test this
493 hypothesis, we stimulated RTS-11 cells with these cytokines to see if they could modulate Mx expression.
494 Seven of the nine Mx genes were induced by type I IFN α and type II IFN γ , and six were induced by IL-1 β . In
495 contrast, IL-6 and TNF α had only minor effects on Mx expression. This cytokine mediated induction was
496 gene-dependent. IFN α was a strong inducer of Mx1-4 and Mx6-7. Past studies have shown Mx1-3 to be
497 modulated by type I IFNs, and so it was no surprise that Mx4 as an additional SMG1/TMG1 member was also
498 induced. Mx6 and Mx7 on the other hand are TMG2 genes. Studies with two other cell lines confirmed the
499 induction of Mx6 by type I IFN as well as a small induction of Mx5 (as seen in RTS11 cells) but Mx7 was not
500 expressed in these cells. IFN γ was a strong inducer of Mx5-6, although some induction of Mx1-4 was also
501 seen in the different cell lines. Trout Mx5 and Mx6 are on the same locus as Mx4-8 in Atlantic salmon, that
502 were also responsive to IFN γ [27], and are SMG2/TMG2 genes. The SMG3 (Mx7-8) and SMG4 (Mx9) genes
503 did not show this responsiveness. IL-1 β was able to induce Mx8 (SMG3) in RTS-11 cells although some
504 induction of Mx2-4 and Mx5-6 was also seen, suggesting a broader responsiveness across SMGs. In common
505 with salmon, no induction of Mx9 was found with these PAMPS/cytokines and its role, if any, in antiviral
506 defence remains to be elucidated.

507 Such findings contrast with mammalian Mx genes that are strictly induced by type I and type III IFNs but are
508 not induced by IFN γ or other proinflammatory cytokines [29-31]. Salmonids possess multiple type I (IFN α -f)
509 and type II (IFN γ 1-2 and IFN γ rels) IFNs, but type III IFN has not been identified in any fish species [62]. In
510 addition to the induction of Mx genes by type I and type II IFNs, this study confirms that some
511 proinflammatory cytokines also influence Mx expression in fish. IL-1 β in particular has a clear impact on Mx
512 gene expression in trout and was the only cytokine that induced Mx8 expression. Thus, it is apparent that
513 cytokines other than IFNs can have a role in antiviral defence.

514 **Conclusions:**

515 Up to 10 Mx genes are present in salmonids that reside in four chromosomal loci. Three teleost Mx groups
516 (TMG) can be identified with characteristic gene organisations, each with a spotted gar Mx gene at the root in
517 the phylogenetic tree. Synteny analysis suggests that the current Mx genes in 3R or 4R teleosts may be
518 evolved from the three Mx genes present at two chromosomal loci in spotted gar, with the 3R duplicated Mx
519 loci retained/lost in a lineage specific manner. Salmonid Mx belong to TMG1 and TMG2. The increased
520 salmonid Mx gene copies are due to local gene duplications that have happened before and after salmonid
521 speciation in a lineage/species specific manner. Salmonids are equipped with a diversified battery of Mx
522 genes, with their protein products present in both cytoplasmic and nuclear locations to protect against viral
523 attack during their life in freshwater and seawater.

524 Trout Mx genes are differentially expressed in tissues with high levels of expression of TMG1 (Mx1-4) in
525 blood and TMG2 (Mx5-9) in the intestine. The expression of most of the trout Mx genes was induced by poly
526 IC (*in vitro* and *in vivo*), and increased during early developmental stages. In addition to induction by type I
527 IFN, IFN γ and IL-1 β also induced Mx expression in rainbow trout and are cytokines that are highly modulated
528 by viral infection. These results show that salmonids possess a large number Mx genes as well as complex
529 regulatory pathways to induce Mx gene expression for antiviral defence, which may contribute to their
530 success in an anadromous life style.

531

532 **5. Acknowledgements**

533 This work was partially funded by the Biotechnology and Biological Sciences Research Council (BBSRC,
534 BB/R008442/1). Tingyu Wang was supported by the Ministry of Science and Technology, Republic of China
535 (Taiwan) (MOST 107-2917-I-564-019). Fuguo Liu was supported by a Newton International Fellowship
536 funded by the Academy of Medical Sciences, UK (AMS, NIF004\1036). Guangming Tian was supported
537 financially by the State Scholarship Fund organised by the China Scholarship Council (201808420042).

538

539 **6. References**

- 540 [1] Haller O, Staeheli P, Schwemmler M, Kochs G. Mx GTPases: dynamin-like antiviral machines of innate
541 immunity. *Trends Microbiol.* 2015;23:154-63.
- 542 [2] Verhelst J, Hulpiau P, Saelens X. Mx proteins: antiviral gatekeepers that restrain the uninvited. *Microbiol*
543 *Mol Biol Rev.* 2013;77:551-66.
- 544 [3] Haller O, Gao S, von der Malsburg A, Daumke O, Kochs G. Dynamin-like MxA GTPase: structural
545 insights into oligomerization and implications for antiviral activity. *J Biol Chem.* 2010;285:28419-24.
- 546 [4] Qi F, Yang A, Ambreen S, Bai X, Hou Y, Lu X. Birth and death of Mx genes and the presence/absence of
547 genes regulating Mx transcription are correlated with the diversity of anti-pathogenicity in vertebrate species.
548 *Mol Genet Genomics.* 2019;294:121-133.
- 549 [5] Staeheli P, Yu YX, Grob R, Haller O. 1989. A double-stranded RNA-inducible fish gene homologous to
550 the murine influenza virus resistance gene Mx. *Mol. Cell. Biol.* 9:3117-21.
- 551 [6] Trobridge GD, Leong JA. Characterization of a rainbow trout Mx gene. *J Interferon Cytokine Res.*
552 1995;15:691-702.
- 553 [7] Trobridge GD, Chiou PP, Leong JA. Cloning of the rainbow trout (*Oncorhynchus mykiss*) Mx2 and Mx3
554 cDNAs and characterization of trout Mx protein expression in salmon cells. *J Virol.* 1997;71:5304-11.
- 555 [8] Robertsen B, Trobridge G, Leong JA. Molecular cloning of double-stranded RNA inducible Mx genes
556 from Atlantic salmon (*Salmo salar* L.). *Dev Comp Immunol.* 1997;21:397-412.
- 557 [9] Jensen V, Robertsen B. Cloning of an Mx cDNA from Atlantic halibut (*Hippoglossus hippoglossus*) and
558 characterization of Mx mRNA expression in response to double-stranded RNA or infectious pancreatic
559 necrosis virus. *J Interferon Cytokine Res.* 2000;20:701-10.
- 560 [10] Lee JY, Hirono I, Aoki T. Cloning and analysis of expression of Mx cDNA in Japanese flounder,
561 *Paralichthys olivaceus*. *Dev Comp Immunol.* 2000;24:407-15.
- 562 [11] Altmann SM, Mellon MT, Johnson MC, Paw BH, Trede NS, Zon LI, Kim CH. Cloning and
563 characterization of an Mx gene and its corresponding promoter from the zebrafish, *Danio rerio*. *Dev Comp*
564 *Immunol.* 2004;28:295-306.
- 565 [12] Zhang YB, Li Q, Gui JF. Differential expression of two *Carassius auratus* Mx genes in cultured CAB
566 cells induced by grass carp hemorrhage virus and interferon. *Immunogenetics.* 2004;56:68-75.
- 567 [13] Plant KP, Thune RL. Cloning and characterisation of a channel catfish (*Ictalurus punctatus*) Mx gene.
568 *Fish Shellfish Immunol.* 2004;16:391-405.
- 569 [14] Wu YC, Chi SC. Cloning and analysis of antiviral activity of a barramundi (*Lates calcarifer*) Mx gene.
570 *Fish Shellfish Immunol.* 2007;23:97-108.
- 571 [15] Zenke K, Kim KH. Molecular cloning and expression analysis of three Mx isoforms of rock bream,
572 *Oplegnathus fasciatus*. *Fish Shellfish Immunol.* 2009;26:599-605.
- 573 [16] Li G, Zhang J, Sun Y, Wang H, Wang Y. The evolutionarily dynamic IFN-inducible GTPase proteins
574 play conserved immune functions in vertebrates and cephalochordates. *Mol Biol Evol.* 2009;26:1619-30.
- 575 [17] Huang B, Huang WS, Nie P. Characterization of four Mx isoforms in the European eel, *Anguilla anguilla*.
576 *Fish Shellfish Immunol.* 2013;35:1048-54.

- 577 [18] Fernández-Trujillo MA, García-Rosado E, Alonso MC, Castro D, Álvarez MC, Béjar J. Mx1, Mx2 and
578 Mx3 proteins from the gilthead seabream (*Sparus aurata*) show in vitro antiviral activity against RNA and
579 DNA viruses. *Mol Immunol.* 2013;56:630-6.
- 580 [19] Alvarez-Torres D, Bejar J, Collet B, Alonso MC, Garcia-Rosado E. Structural and functional
581 characterization of the Senegalese sole (*Solea senegalensis*) Mx promoter. *Fish Shellfish Immunol.*
582 2013;35:1642-8.
- 583 [20] Novel P, Fernández-Trujillo MA, Gallardo-Gálvez JB, Cano I, Manchado M, Buonocore F, Randelli E,
584 Scapigliati G, Alvarez MC, Béjar J. Two Mx genes identified in European sea bass (*Dicentrarchus labrax*)
585 respond differently to VNNV infection. *Vet Immunol Immunopathol.* 2013;153:240-8.
- 586 [21] Saxena A, Belwal K, Chauhan A, Pande A. Interferon induced Mx protein from Indian snow trout
587 *Schizothorax richardsonii* (Gray) lacks critical functional features unlike its mammalian homologues. *Comput*
588 *Biol Chem.* 2018;73:31-40.
- 589 [22] Solbakken MH, Rise ML, Jakobsen KS, Jentoft S. Successive Losses of Central Immune Genes
590 Characterize the Gadiformes' Alternate Immunity. *Genome Biol Evol.* 2016;8:3508-3515.
- 591 [23] Caipang CM, Hirono I, Aoki T. In vitro inhibition of fish rhabdoviruses by Japanese flounder,
592 *Paralichthys olivaceus* Mx. *Virology.* 2003; 317:373–82
- 593 [24] Larsen R, Rokenes TP, Robertsen B. Inhibition of infectious pancreatic necrosis virus replication by
594 Atlantic salmon Mx1 protein. *J. Virol.* 2004; 78:7938–44
- 595 [25] Su J, Yang C, Zhu Z, Wang Y, Jang S, Liao L. Enhanced grass carp reovirus resistance of Mx-transgenic
596 rare minnow (*Gobiocypris rarus*). *Fish Shellfish Immunol.* 2009;26:828-35.
- 597 [26] Peng L, Yang C, Su J. Protective roles of grass carp *Ctenopharyngodon Idella* Mx isoforms against grass
598 carp reovirus. *PLoS One.* 2012;7:e52142.
- 599 [27] Robertsen B, Greiner-Tollersrud L, Jørgensen LG. Analysis of the Atlantic salmon genome reveals a
600 cluster of Mx genes that respond more strongly to IFN gamma than to type I IFN. *Dev Comp Immunol.*
601 2019;90:80-89.
- 602 [28] Husain M, Bird S, van Zwieten R, Secombes CJ, Wang T. Cloning of the IL-1 β gene and IL-1 β
603 pseudogene in salmonids uncovers a second type of IL-1 β gene in teleost fish. *Dev Comp Immunol.*
604 2012;38:431-46.
- 605 [29] Mordstein M, Neugebauer E, Ditt V, Jessen B, Rieger T, et al. 2010. Lambda interferon renders epithelial
606 cells of the respiratory and gastrointestinal tracts resistant to viral infections. *J. Virol.* 84:5670–77
- 607 [30] Holzinger D, Jorns C, Stertz S, Boisson-Dupuis S, Thimme R, et al. 2007. Induction of MxA gene
608 expression by influenza A virus requires type I or type III interferon signaling. *J. Virol.* 81:7776–85
- 609 [31] Simon A, Fah J, Haller O, Staeheli P. 1991. Interferon-regulated Mx genes are not responsive to
610 interleukin-1, tumor necrosis factor, and other cytokines. *J. Virol.* 65:968–71
- 611 [32] Wangkahart E, Secombes CJ, Wang T. Dissecting the immune pathways stimulated following injection
612 vaccination of rainbow trout (*Oncorhynchus mykiss*) against enteric redmouth disease (ERM). *Fish Shellfish*
613 *Immunol.* 2019;85:18-30.
- 614 [33] Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol*
615 1990; 215:403–10.

- 616 [34] Wang T, Hu Y, Wangkahart E, Liu F, Wang A, Zahran E, Maisey KR, Liu M, Xu Q, Imarai M,
617 Secombes CJ. Interleukin (IL)-2 Is a Key Regulator of T Helper 1 and T Helper 2 Cytokine Expression in Fish:
618 Functional Characterization of Two Divergent IL2 Paralogs in Salmonids. *Front Immunol.* 2018;9:1683.
- 619 [35] Wang B, Wangkahart E, Secombes CJ, Wang T. Insights into the evolution of the suppressors of cytokine
620 signalling (SOCS) gene family in vertebrates. *Mol Biol Evol.* 2019 36:393-411.
- 621 [36] Gasteiger E, Hoogland C, Gattiker A, Duvaud S, Wilkins MR, Appel RD, et al. Protein identification and
622 analysis tools on the ExPASy server. In: Walker JM, editor. *The Proteomics Protocols Handbook*. New York:
623 Humana Press; (2005). p. 571–607.
- 624 [37] Chenna R, Sugawara H, Koike T, Lopez R, Gibson TJ, Higgins DG, Thompson JD. Multiple sequence
625 alignment with the Clustal series of programs. *Nucleic Acids Res.* 2003; 31:3497-500.
- 626 [38] Campanella JJ, Bitincka L, Smalley J. MatGAT: an application that generates similarity/identity matrices
627 using protein or DNA sequences. *BMC Bioinformatics.* 2003; 4:29.
- 628 [39] Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for
629 Bigger Datasets. *Mol Biol Evol.* 2016; 33:1870-4.
- 630 [40] Staeheli P, Haller O, Boll W, Lindenmann J, Weissmann C. 1986. Mx protein: constitutive expression in
631 3T3 cells transformed with cloned Mx cDNA confers selective resistance to influenza virus. *Cell* 44:147–58
- 632 [41] Staeheli P, Sutcliffe JG. Identification of a second interferon-regulated murine Mx gene. *Mol. Cell. Biol.*
633 1988; 8:4524–28
- 634 [42] Louis A, Muffato M, Roest Crollius H. Genomicus: five genome browsers for comparative genomics in
635 eukaryota. *Nucleic Acids Res.* 2013;41(Database issue):D700-5.
- 636 [43] Wang T, Diaz-Rosales P, Costa MM, Campbell S, Snow M, Collet B, Martin SA, Secombes CJ.
637 Functional characterization of a nonmammalian IL-21: rainbow trout *Oncorhynchus mykiss* IL-21 upregulates
638 the expression of the Th cell signature cytokines IFN-gamma, IL-10, and IL-22. *J Immunol.* 2011;186:708-21.
- 639 [44] Wang T, Monte MM, Huang W, Boudinot P, Martin SA, Secombes CJ. Identification of two FoxP3
640 genes in rainbow trout (*Oncorhynchus mykiss*) with differential induction patterns. *Mol Immunol.*
641 2010;47:2563-74.
- 642 [45] Wangkahart E, Scott C, Secombes CJ, Wang T. Re-examination of the rainbow trout (*Oncorhynchus*
643 *mykiss*) immune response to flagellin: *Yersinia ruckeri* flagellin is a potent activator of acute phase proteins,
644 anti-microbial peptides and pro-inflammatory cytokines in vitro. *Dev Comp Immunol.* 2016;57:75-87.
- 645 [46] Ganassin RC, Bols NC. Development of a monocyte/macrophage-like cell line, RTS11, from rainbow
646 trout spleen. *Fish Shellfish Immunol.* 1998; 8:457–476
- 647 [47] Wolf K, Quimby MC. Established eurythermic lines of fish cells in vitro. *Science* 1962; 135:1065
- 648 [48] Schirmer K, Chan AG, Greenberg BM, Dixon DG, Bols NC. Ability of 16 priority PAHs to be
649 photocytotoxic to a cell line from the rainbow trout gill. *Toxicology.* 1998; 127:143–155
- 650 [49] Wang T, Huang W, Costa MM, Martin SA, Secombes CJ. Two copies of the genes encoding the subunits
651 of putative interleukin (IL)-4/IL-13 receptors, IL-4R α , IL-13R α 1 and IL-13R α 2, have been identified in
652 rainbow trout (*Oncorhynchus mykiss*) and have complex patterns of expression and modulation.
653 *Immunogenetics.* 2011;63:235-53.
- 654 [50] Hong S, Zou J, Crampe M, Peddie S, Scapigliati G, Bols N, Cunningham C, Secombes CJ. The
655 production and bioactivity of rainbow trout (*Oncorhynchus mykiss*) recombinant IL-1 beta. *Vet Immunol*
656 *Immunopathol.* 2001;81:1-14.

- 657 [51] Costa MM, Maehr T, Diaz-Rosales P, Secombes CJ, Wang T. Bioactivity studies of rainbow trout
658 (*Oncorhynchus mykiss*) interleukin-6: effects on macrophage growth and antimicrobial peptide gene
659 expression. *Mol Immunol.* 2011;48:1903-16.
- 660 [52] Hong S, Li R, Xu Q, Secombes CJ, Wang T. Two types of TNF- α exist in teleost fish: phylogeny,
661 expression, and bioactivity analysis of type-II TNF- α 3 in rainbow trout *Oncorhynchus mykiss*. *J Immunol.*
662 2013;191:5959-72.
- 663 [53] Husain M, Martin SA, Wang T. Identification and characterisation of the IL-27 p28 subunits in fish:
664 Cloning and comparative expression analysis of two p28 paralogues in Atlantic salmon *Salmo salar*. *Fish*
665 *Shellfish Immunol.* 2014. 41:102-12.
- 666 [54] Wangkahart E, Secombes CJ, Wang T. Studies on the Use of Flagellin as an Immunostimulant and
667 Vaccine Adjuvant in Fish Aquaculture. *Front Immunol.* 2019. 9:3054.
- 668 [55] Gao S, von der Malsburg A, Paeschke S, Behlke J, Haller O, Kochs G, Daumke O. Structural basis of
669 oligomerization in the stalk region of dynamin-like MxA. *Nature.* 2010. 465(7297):502-6.
- 670 [56] Berthelot C, Brunet F, Chalopin D, Juanchich A, Bernard M, Noël B, Bento P, Da Silva C, Labadie K,
671 Alberti A, Aury JM, Louis A, Dehais P, Bardou P, Montfort J, Klopp C, Cabau C, Gaspin C, Thorgaard GH,
672 Boussaha M, Quillet E, Guyomard R, Galiana D, Bobe J, Volff JN, Genêt C, Wincker P, Jaillon O, Roest
673 Crollius H, Guiguen Y. The rainbow trout genome provides novel insights into evolution after whole-genome
674 duplication in vertebrates. *Nat Commun.* 2014. 5:3657.
- 675 [57] Lien S, Koop BF, Sandve SR, Miller JR, Kent MP, Nome T, Hvidsten TR, Leong JS, Minkley DR, Zimin
676 A, Grammes F, Grove H, Gjuvsland A, Walenz B, Hermansen RA, von Schalburg K, Rondeau EB, Di
677 Genova A, Samy JK, Olav Vik J, Vigeland MD, Caler L, Grimholt U, Jentoft S, Våge DI, de Jong P, Moen T,
678 Baranski M, Palti Y, Smith DR, Yorke JA, Nederbragt AJ, Tooming-Klunderud A, Jakobsen KS, Jiang X, Fan
679 D, Hu Y, Liberles DA, Vidal R, Iturra P, Jones SJ, Jonassen I, Maass A, Omholt SW, Davidson WS. The
680 Atlantic salmon genome provides insights into rediploidization. *Nature.* 2016; 533(7602):200-5.
- 681 [58] Staeheli P, Haller O. Human MX2/MxB: a Potent Interferon-Induced Postentry Inhibitor of
682 Herpesviruses and HIV-1. *J Virol.* 2018; 92(24). pii:e00709-18.
- 683 [59] Goujon C, Moncorgé O, Bauby H, Doyle T, Barclay WS, Malim MH. Transfer of the amino-terminal
684 nuclear envelope targeting domain of human MX2 converts MX1 into an HIV-1 resistance factor. *J Virol.*
685 2014; 88:9017-26.
- 686 [60] Zurcher T, Pavlovic J, Staeheli P. Mouse Mx2 protein inhibits vesicular stomatitis virus but not influenza
687 virus. *Virology.* 1992; 187:796-800
- 688 [61] Stuempfig ND, Seroy J. *Viral Gastroenteritis.* 2019; StatPearls
- 689 [62] Secombes CJ, Zou J. Evolution of Interferons and Interferon Receptors. *Front Immunol.* 2017; 8:209.

691 **Table 1. Summary of Mx gene family in salmonids and pike**

692

Common name	Gene	Chromosome	Location	Genebank accession number (mRNA/protein)
Rainbow trout	Mx1	NC_035093.1 (Ch17)	54,895,743→54,907,262	NM_001171901.1
Rainbow trout	Mx2	NC_035093.1 (Ch17)	54,827,385→54,839,851	NM_001124751.1
Rainbow trout	Mx3	NC_035093.1 (Ch17)	54,879,332→54,883,908	XM_021569609.1
Rainbow trout	Mx4	NC_035093.1 (Ch17)	54,848,974→54,863,639	MK301134
Rainbow trout	Mx5	NC_035079.1 (Ch3)	82,015,213→81,992,259	MK301135
Rainbow trout	Mx6	NC_035079.1 (Ch3)	82,029,332→82,045,373	MK301136
Rainbow trout	Mx7	NC_035087.1 (Ch11)	76,228,384→76,215,926	MK301137
Rainbow trout	Mx8	NC_035087.1 (Ch11)	76,240,419→76,420,666	MK301138
Rainbow trout	Mx9	NC_035100.1 (Ch24)	21,286,705→21,268,999	MK301139
Atlantic salmon	Mx1	NC_027311(Ch12)	66,798,275→66,829,177	NM_001123690/NP_001117162
Atlantic salmon	Mx2	NC_027311(Ch12)	66,776,028→66,803,979	NM_001139918/NP_001133390
Atlantic salmon	Mx3	NC_027311(Ch12)	66,816,288→66,829,177	NM_001123675/NP_001117147
Atlantic salmon	Mx4	NC_027324(Ch25)	47,088,993→47,121,652	XM_014174614/XP_014030089
Atlantic salmon	Mx5	NC_027324(Ch25)	47,228,437→47,217,827	XM_014174615/XP_014030090
Atlantic salmon	Mx6	NC_027324(Ch25)	47,161,992→47,139,132	XM_014174616/XP_014030091
Atlantic salmon	Mx7	NC_027324(Ch25)	47,193,272→47,175,785	XM_014174617/XP_014030092
Atlantic salmon	Mx8	NC_027324(Ch25)	47,243,602→47,262,616	XM_014174618/XP_014030093
Atlantic salmon	Mx9	NC_027308(Ch9)	117,838,750→117,853,816	XM_014214722/XP_014070197
Atlantic salmon	Mx10	NC_027314(Ch15)	5,299,091→5,292,439	XM_014143485/XP_013998960
Chinook salmon	Mx1	NW_020142590	72,571→83,518	XM_024415949/XP_024271717
Chinook salmon	Mx2	NW_020142590	17,032→35,480	XM_024415950/XP_024271718
Chinook salmon	Mx3	NW_020142590	56,817→83,518	XM_024415946/XP_024271714
Chinook salmon	Mx4	NW_020133776	172→6,566	XM_024410424/XP_024266192
Chinook salmon	Mx5	NC_037108(Ch12)	2,001,377→2,013,759	XM_024438118/XP_024293886
Chinook salmon	Mx6	NC_037110(Ch14)	41,090,466→41,103,747	XM_024445373/XP_024301141
Coho salmon	Mx1	NC_034174(Ch1)	46,664,816→46,675,808	LOC109896993
Coho salmon	Mx2	NC_034174(Ch1)	46,587,607→46,621,678	XM_020468497/XP_020324086
Coho salmon	Mx3	NC_034174(Ch1)	46,651,400→46,656,202	XM_020491468/XP_020347057
Coho salmon	Mx4	NW_018090236	57,121→68,549	GDQG01031501/ /Q6PW23
Coho salmon	Mx5	NC_034181(Ch8)	66,940,031→66,941,209	XM_020488627/XP_020344216
Coho salmon	Mx6	NC_034191(Ch18)	53,794,477→53,822,277	XM_020508491/XP_020364080
Arctic charr	Mx1	NC_036838(Ch1)	44,797,136→44,802,044	XM_023993827/XP_023849595
Arctic charr	Mx2	NC_036838(Ch1)	44,762,649→44,772,138	XM_023993825/XP_023849593
Arctic charr	Mx3	NW_019943275	202,142→225,309	XM_024139811/XP_023995579
Arctic charr	Mx4	NW_019943275	93,732→108,405	XM_024139809/XP_023995577
Arctic charr	Mx5	NW_019943275	87,444→170,634	XM_024139810/XP_023995578
Arctic charr	Mx6	NW_019945020	48,231→54,383	XM_024143207/XP_023998975
Arctic charr	Mx7	NW_019945020	11,050→28,797	XM_024143206/XP_023998974
Arctic charr	Mx8	NW_019946381	2,678→17,616	XM_024144359/XP_024000127
Arctic charr	Mx9	NW_019942645	359,971→369,223	XM_024136430/XP_023992198
Arctic charr	Mx10	NW_019942645	378,229→388,058	XM_024136431/XP_023992199
Pike	Mx1	NC_025984(Ch17)	25,113,053→25,132,843	XM_013138351/XP_012993805
Pike	Mx2	NC_025980(Ch13)	25,725,708→25,740,882	ENSELUG00000023570/ENSELUT00000043341
Pike	Mx3	NC_025980(Ch13)	25,694,138→25,702,004	ENSELUG00000023626/ENSELUT00000036437

693

694

695 **Table 2. Comparison of trout Mx aa sequence identities to Mx from other salmonids, spotted gar and**
 696 **mammals.** The amino acid number for each sequence is also shown. Only full-length aa sequences were
 697 included in the analysis.

698

		No. of aa	SMG1				SMG2		SMG3		SMG4
			Trout Mx1	Trout Mx2	Trout Mx3	Trout Mx4	Trout Mx5	Trout Mx6	Trout Mx7	Trout Mx8	Trout Mx9
SMG1	Trout Mx1	621	100.0	86.5	96.3	86.5	46.2	46.3	47.9	46.4	47.7
	Trout-Mx2	635	86.5	100.0	86.3	98.4	45.6	45.0	47.0	45.1	47.3
	Trout Mx3	623	96.3	86.3	100.0	86.0	45.9	46.0	48.1	46.6	47.9
	Trout Mx4	635	86.5	98.4	86.0	100.0	45.8	45.5	47.0	45.2	47.0
	Atlantic Mx1	623	95.3	85.8	97.3	85.4	45.7	45.9	47.9	46.4	47.4
	Atlantic Mx2	638	85.4	94.0	85.4	93.9	45.4	44.8	46.4	44.7	46.7
	Atlantic Mx3	623	96.3	86.5	95.8	86.6	46.5	46.0	48.1	46.7	48.3
	Chinook Mx1	621	98.2	86.8	96.0	86.5	46.2	46.2	47.9	46.7	47.9
	Chinook Mx2	633	83.7	95.0	83.4	95.1	43.6	43.5	45.4	43.5	45.6
	Chinook Mx3	623	95.2	86.0	97.8	86.2	46.2	46.2	48.1	46.3	47.8
	Coho-Mx2	648	84.3	95.2	84.1	95.7	44.7	44.3	45.8	44.2	45.9
	Coho Mx3	623	96.8	85.5	97.8	85.4	46.0	46.2	47.9	46.3	47.8
Charr Mx1	623	96.0	86.6	97.9	86.2	45.7	46.0	48.1	46.1	47.6	
Charr Mx2	638	85.6	94.8	85.6	95.1	45.4	44.9	47.0	45.3	47.3	
SMG2	Trout Mx5	614	46.2	45.6	45.9	45.8	100.0	93.6	61.2	69.4	51.4
	Trout Mx6	606	46.3	45.0	46.0	45.5	93.6	100.0	61.4	69.7	50.8
	Atlantic Mx4	606	45.6	45.4	45.2	45.6	86.0	88.3	62.2	70.1	49.7
	Atlantic Mx5	608	45.4	44.7	45.5	44.8	90.1	88.8	62.7	70.8	50.2
	Atlantic Mx6	627	45.3	44.4	45.0	44.7	83.9	84.4	61.3	67.9	49.5
	Atlantic Mx7	603	46.4	45.0	46.1	45.5	86.6	87.1	61.8	69.9	49.8
	Atlantic Mx8	607	47.0	45.5	46.7	45.6	87.0	90.0	62.5	71.2	50.6
	Coho Mx4	614	46.0	45.2	45.7	45.3	95.1	91.7	61.1	69.9	51.6
	Charr Mx4	607	46.5	45.9	46.2	46.1	90.9	89.8	61.2	69.3	50.2
	Charr Mx5	612	45.1	43.8	45.2	44.2	82.7	85.0	60.6	68.2	49.7
SMG3	Trout Mx7	613	47.9	47.0	48.1	47.0	61.2	61.4	100.0	68.6	49.1
	Trout Mx8	608	46.4	45.1	46.6	45.2	69.4	69.7	68.6	100.0	49.8
	Charr Mx9	549	42.3	41.7	42.4	41.5	53.6	53.4	80.7	57.8	44.5
	Charr Mx10	608	46.1	45.2	46.4	45.4	69.3	69.6	68.7	93.8	49.8
SMG4	Trout Mx9	640	47.7	47.3	47.9	47.0	51.4	50.8	49.1	49.8	100.0
	Atlantic Mx9	642	48.1	47.6	48.2	47.1	51.6	50.8	49.5	49.5	94.9
	Chinook Mx6	638	47.4	47.2	47.8	46.9	51.1	50.5	49.3	49.4	97.0
	Coho Mx6	646	43.2	42.3	43.2	42.1	45.2	44.8	45.4	43.6	80.2
Gar	Spotted gar Mx1	619	73.9	71.3	74.8	71.3	46.6	47.1	50.8	47.0	48.1
	Spotted gar Mx2	684	48.5	47.8	48.5	47.8	38.7	38.6	41.2	39.2	39.4
	Spotted gar Mx3	616	47.6	44.3	47.0	44.4	41.3	40.8	41.1	40.8	37.3
Mammals	Human MxA	670	52.6	52.1	52.4	52.0	40.5	40.4	42.1	41.2	43.1
	Mouse-Mx1	631	51.8	51.2	52.0	51.6	40.4	40.4	42.1	40.9	43.3
	Cow Mx1	648	51.8	52.0	52.1	51.7	39.5	40.7	42.7	41.5	42.8

699

700

701

702 **Figure legend**

703

704 **Fig. 1. Amino acid multiple alignment of rainbow trout Mx family.** The multiple alignment was produced
 705 using ClustalW, and conserved amino acids shaded using BOXSHADE (version 3.21). Human MxA and MxB
 706 were included in the alignment for comparison. The N-terminal GTPase domain and C-terminal GTPase
 707 effector domain are indicated above the alignment. The conserved tripartite GTP-binding motif (GDXXSGKS,
 708 DLPG, and TKPD) and a dynamin signature (LPRXXGXXTR) in the GTPase domain, and leucine residues
 709 that form leucine zipper folds in the GTPase effector domain are in red. The four α -helices and the four loops
 710 connecting them are shown under the alignment as defined by Gao et al. [55]. The potential nuclear
 711 localisation signal (KKRKR) in trout Mx2 and 4, the four lysine residues of human MxA in L4 are in blue.

712 **Fig. 2 Phylogenetic tree (A) and chromosome localisation (B) of salmonid Mx genes.** **A.** The phylogenetic
 713 tree was constructed using a multiple alignment of salmonid and pike Mx aa sequences and the neighbour-
 714 joining method within the MEGA7.0 program. The evolutionary distances were computed using the JTT
 715 matrix-based method with all ambiguous positions removed for each sequence pair. The percentage (>50%) of
 716 replicate trees in which the associated taxa clustered together in the bootstrap test (10,000 replicates) is shown
 717 next to the branches. The accession number for each sequence is given in Table 2. Four salmonid Mx groups
 718 (SMG)1-4 are indicated on the right. **B.** the chromosome localisation of Mx genes in salmonids and pike.

719 **Fig. 3 Phylogenetic tree analysis of bony fish Mx.** The phylogenetic tree was constructed using amino acid
 720 multiple alignments of Mx from selected teleosts and mammals, and the neighbour-joining method within the
 721 MEGA7.0 program. The evolutionary distances were computed using the JTT matrix-based method with all
 722 ambiguous positions removed for each sequence pair. Node values represent percent bootstrap confidence
 723 derived from 10,000 replications. The accession number for each sequence is given after the species and
 724 molecule names. The salmonid clades are highlighted and condensed under the name of SMG (salmonid Mx
 725 group)1-4, which share the same topologies as in **Fig. 2**. The root bootstrap values of mammalian Mx and
 726 teleost Mx group 1-3 are highlighted in red with the tentative groupings indicated on the right of the tree.

727 **Fig. 4. Synteny analysis of Mx loci in bony fish and human.** The synteny was predicted using the
 728 Genomicus program [42] or information extracted from recently released reference genomes at NCBI.

729 **Fig. 5. Comparison of gene organisation of the Mx gene family in rainbow trout, other bony fish and**
 730 **humans.** Boxes represent exons, and lines between exons represent introns. The black and white boxes
 731 represent non-coding and amino acid (aa) coding regions, respectively. The sizes (bp) of each exon are
 732 numbered in the boxes. The gene organization of rainbow trout Mx genes was predicted using the Splign
 733 program based on the sequence information from Table 1 and Figures S1–S6 in Supplementary Material. The
 734 information of other species was extracted from recent released reference genomes at NCBI.

735 **Figure 6. Transcript expression of rainbow trout Mx gene family in tissues and cell lines.** The expression
736 level of Mx1-4 (A), Mx5-6 (B), Mx7-8 (C) and Mx9 (D) was determined by RT-qPCR in 17 tissues from six
737 fish and four replicates of each cell line. The transcript level was calculated using a serial dilution of
738 references that contained equal molar amounts of the probes for each gene and was normalized against the
739 expression level of EF-1 α . The results are presented as the average +SEM.

740 **Fig. 7. The expression ontogeny of rainbow trout Mx gene family.** cDNA samples were prepared from
741 eyed-eggs, immediately post-hatch, pre-first feeding fry or fry 3 weeks after first feeding. Six independent
742 samples for each developmental stage were prepared for real-time quantification of gene expression as
743 described in Fig. 6. The results are presented as the average + SEM. Different letters over bars indicate
744 significant differences ($p \leq 0.05$, one way-ANOVA).

745 **Fig. 8. Modulation of Mx gene expression *in vivo* by poly IC.** Rainbow trout were injected ip with 1 mg
746 poly IC in 0.2 ml PBS or 0.2 ml PBS as control. The spleen, head kidney (HK), gills and intestine were taken
747 at 6 h and 24 h post injection. The quantification of Mx gene expression was as described in Fig. 6. The
748 relative expression is shown, where the average expression level in the control fish at 6 h in each tissue was
749 defined as 1. The results are presented as the mean + SEM of six fish. Different letters over bars in the same
750 tissue indicate significant differences ($p \leq 0.05$, one way-ANOVA).

751 **Fig. 9. Modulation of proinflammatory cytokine gene expression *in vivo* by poly IC.** Rainbow trout were
752 injected ip with 1 mg poly IC in 0.2 ml PBS or 0.2 ml PBS as control. The spleen, head kidney (HK), gills and
753 intestine were taken at 6 h and 24 h post injection. The quantification of gene expression was as described in
754 Fig. 6. The relative expression is shown, where the average expression level in the control fish at 6 h in each
755 tissue was defined as 1. The results are presented as the mean + SEM of six fish. Different letters over bars in
756 the same tissue indicate significant differences ($p \leq 0.05$, one way-ANOVA).

757 **Fig. 10. Modulation of Mx gene expression in RTS-11 cells by poly IC and LPS.** Overnight culture of
758 RTS-11 cells were stimulated with poly IC (50 $\mu\text{g/ml}$), LPS (25 $\mu\text{g/ml}$), or medium as control for 4h, 8 h and
759 24 h, and the expression of trout Mx genes was quantified by RT-qPCR as described in Fig. 6. The data are
760 presented as the mean (+SEM, N=4) fold change calculated as the average expression level of stimulated
761 samples divided by that of time-matched controls. The relative significance of a LSD post hoc test after a
762 significant one-way ANOVA between the stimulated and time-matched controls is shown above the bars as *
763 $p \leq 0.05$; ** $p \leq 0.01$ and *** $p \leq 0.001$.

764 **Fig. 11. Modulation of trout Mx expression in RTS-11 cells by pro-inflammatory cytokines.** Overnight
765 culture of RTS-11 cells were stimulated with IFN γ (20 ng/ml), IFN- α (25 ng/ml), IL-1 β (25 ng/ml), IL-6 (100
766 ng/ml), TNF α (50 ng/ml), or medium as control for 4h, 8 h and 24 h, and the expression of trout Mx genes
767 was quantified by RT-qPCR as described in Fig. 6. The data are presented as the mean (+SEM, N=4) fold
768 change, calculated as the average expression level of stimulated samples divided by that of time-matched

769 controls. The relative significance of a LSD post hoc test after a significant one-way ANOVA between the
770 stimulated and time-matched controls is shown above the bars as * $p \leq 0.05$; ** $p \leq 0.01$ and *** $p \leq 0.001$.

771 **Fig. 12. Modulation of trout Mx expression in RTG-2 and RTGill cell lines by IFNs.** Overnight cultures
772 of cells were stimulated with IFN γ (20 ng/ml), IFN- α (25 ng/ml), or medium as control for 4h, and the
773 expression of trout Mx genes was quantified by RT-qPCR as described in Fig. 6. The data are presented as the
774 mean (+SEM, N=4) fold change, calculated as the average expression level of stimulated samples divided by
775 that of controls. The relative significance of a LSD post hoc test after a significant one-way ANOVA between
776 the stimulated and controls is shown above the bars as * $p \leq 0.05$; ** $p \leq 0.01$ and *** $p \leq 0.001$. The line-
777 connected groups are significantly different.

778 **Fig. 13. Hypothetical evolutionary pathways of teleost Mx gene family.** Three Mx loci (Mx1-3) were
779 present on two chromosomes in ancestral 2R actinopterygians. 3R WGD is expected to have produced 6 Mx
780 loci on four chromosomes that were retained in a lineage specific manner to give rise to the three extant
781 teleost Mx groups. The ancestral vertebrates that evolved into the tetrapod lineage appear to have possessed a
782 cognate Mx locus of gar Ch3. Arrow heads indicate ancestral Mx genes. Representative chromosomal loci are
783 shown.

784

Table 2. Comparison of trout Mx aa sequence identities to Mx from other salmonids, spotted gar and mammals. The amino acid number for each sequence is also shown. Only full-length aa

		No. of aa	SMG1				SMG2		SMG3		SMG4
			Trout Mx1	Trout Mx2	Trout Mx3	Trout Mx4	Trout Mx5	Trout Mx6	Trout Mx7	Trout Mx8	Trout Mx9
SMG1	Trout Mx1	621	100.0	86.5	96.3	86.5	46.2	46.3	47.9	46.4	47.7
	Trout-Mx2	635	86.5	100.0	86.3	98.4	45.6	45.0	47.0	45.1	47.3
	Trout Mx3	623	96.3	86.3	100.0	86.0	45.9	46.0	48.1	46.6	47.9
	Trout Mx4	635	86.5	98.4	86.0	100.0	45.8	45.5	47.0	45.2	47.0
	Atlantic Mx1	623	95.3	85.8	97.3	85.4	45.7	45.9	47.9	46.4	47.4
	Atlantic Mx2	638	85.4	94.0	85.4	93.9	45.4	44.8	46.4	44.7	46.7
	Atlantic Mx3	623	96.3	86.5	95.8	86.6	46.5	46.0	48.1	46.7	48.3
	Chinook Mx1	621	98.2	86.8	96.0	86.5	46.2	46.2	47.9	46.7	47.9
	Chinook Mx2	633	83.7	95.0	83.4	95.1	43.6	43.5	45.4	43.5	45.6
	Chinook Mx3	623	95.2	86.0	97.8	86.2	46.2	46.2	48.1	46.3	47.8
	Coho-Mx2	648	84.3	95.2	84.1	95.7	44.7	44.3	45.8	44.2	45.9
	Coho Mx3	623	96.8	85.5	97.8	85.4	46.0	46.2	47.9	46.3	47.8
	Charr Mx1	623	96.0	86.6	97.9	86.2	45.7	46.0	48.1	46.1	47.6
Charr Mx2	638	85.6	94.8	85.6	95.1	45.4	44.9	47.0	45.3	47.3	
SMG2	Trout Mx5	614	46.2	45.6	45.9	45.8	100.0	93.6	61.2	69.4	51.4
	Trout Mx6	606	46.3	45.0	46.0	45.5	93.6	100.0	61.4	69.7	50.8
	Atlantic Mx4	606	45.6	45.4	45.2	45.6	86.0	88.3	62.2	70.1	49.7
	Atlantic Mx5	608	45.4	44.7	45.5	44.8	90.1	88.8	62.7	70.8	50.2
	Atlantic Mx6	627	45.3	44.4	45.0	44.7	83.9	84.4	61.3	67.9	49.5
	Atlantic Mx7	603	46.4	45.0	46.1	45.5	86.6	87.1	61.8	69.9	49.8
	Atlantic Mx8	607	47.0	45.5	46.7	45.6	87.0	90.0	62.5	71.2	50.6
	Coho Mx4	614	46.0	45.2	45.7	45.3	95.1	91.7	61.1	69.9	51.6
	Charr Mx4	607	46.5	45.9	46.2	46.1	90.9	89.8	61.2	69.3	50.2
Charr Mx5	612	45.1	43.8	45.2	44.2	82.7	85.0	60.6	68.2	49.7	
SMG3	Trout Mx7	613	47.9	47.0	48.1	47.0	61.2	61.4	100.0	68.6	49.1
	Trout Mx8	608	46.4	45.1	46.6	45.2	69.4	69.7	68.6	100.0	49.8
	Charr Mx9	549	42.3	41.7	42.4	41.5	53.6	53.4	80.7	57.8	44.5
	Charr Mx10	608	46.1	45.2	46.4	45.4	69.3	69.6	68.7	93.8	49.8
SMG4	Trout Mx9	640	47.7	47.3	47.9	47.0	51.4	50.8	49.1	49.8	100.0
	Atlantic Mx9	642	48.1	47.6	48.2	47.1	51.6	50.8	49.5	49.5	94.9
	Chinook Mx6	638	47.4	47.2	47.8	46.9	51.1	50.5	49.3	49.4	97.0
	Coho Mx6	646	43.2	42.3	43.2	42.1	45.2	44.8	45.4	43.6	80.2
Gar	Gar Mx1	619	73.9	71.3	74.8	71.3	46.6	47.1	50.8	47.0	48.1
	Gar Mx2	684	48.5	47.8	48.5	47.8	38.7	38.6	41.2	39.2	39.4
	Gar Mx3	616	47.6	44.3	47.0	44.4	41.3	40.8	41.1	40.8	37.3
Mammals	Human MxA	670	52.6	52.1	52.4	52.0	40.5	40.4	42.1	41.2	43.1
	Mouse-Mx1	631	51.8	51.2	52.0	51.6	40.4	40.4	42.1	40.9	43.3
	Cow Mx1	648	51.8	52.0	52.1	51.7	39.5	40.7	42.7	41.5	42.8

sequences were included in the analysis.

Trout Mx1 1 -----MNNYTLNQHYEEKVRPCIDLDLSLRSGLVEKDLALPAIAVIGDQSSGKSSVLEALSGVALPRGSGIVTRCPLELKMRRKKEGEEWHGKISYQDHE
Trout Mx2 1 -----MNYTLNQHYEEKVRPSIDLIDSLRSGLVEKDLALPAIAVIGDQSSGKSSVLEALSGVALPRGSGIVTRCPLELKMRRKKEGEEWHGKISYQDRE
Trout Mx3 1 -----MNNYTLNQHYEEKVRPCIDLDLSLRSGLVEKDLALPAIAVIGDQSSGKSSVLEALSGVALPRGSGIVTRCPLELKMRRKKEGEEWHGKISYQDHE
Trout Mx4 1 -----MNNYTLNQHYEEKVRPSIDLIDSLRSGLVEKDLALPAIAVIGDQSSGKSSVLEALSGVALPRGSGIVTRCPLELKMRRKKEGEEWHGKISYQDRE
Trout Mx5 1 -----MSYD-DGSPMFDQDLAEKVRPFDLIDDMRSTIGDEKELPLPTIAVVGQSSGKSSVLEALSGVALPRGSGIVTRCPLELKLKLCNDR-TVKWDAVISYGGKI
Trout Mx6 1 -----MSHD-DGPRPTFDQDLAEKIRPFDLVDMMRSTIGDEKELPLPTIAVVGQSSGKSSVLEALSGVALPRGSGIVTRCPLELKLKLCNDR-TVKWDAVISYGGKI
Trout Mx7 1 -----MS-EPISSGQFDQMAERVRPFDLIDYLRSTIGIEKELPLPSIAVVGQSSGKSSVLEALSGVALPRGSGIVTRCPLELRLCYVS-GVAMKAVISYRDKR
Trout Mx8 1 -----MSDDEGSPMFDQDLAEKVRPFDLIDYLRSTIGIEKELPLPSIAVVGQSSGKSSVLEALSGVALPRGSGIVTRCPLELRLCYVS-GVAMKAVISYKNTK
Trout Mx9 1 MHRPDAGSEDEERDGIQRGVFYSHLDRHRPFDLIDFLRSTIGIEKDLALPAIAVVGQSSGKSSVLEALSGVALPRGSGIVTRCPLELKLKRCF-GGKWKAVISYQGVV
Human MxA 21 LLNGDATVAQKNPQSVAEENLCSQYEEKVRPCIDLDLSLRSGLVEKDLALPAIAVIGDQSSGKSSVLEALSGVALPRGSGIVTRCPLELKLKLCNDR-TVKWDAVISYRDKR
Human MxB 69 NNQPPGPNRSQPRAMGPENNLYSQYEQKVRPCIDLDLSLRSGLVEKDLALPAIAVIGDQSSGKSSVLEALSGVALPRGSGIVTRCPLELKLKLCNDR-TVKWDAVISYRDKR

Dynamine-like GTPase Domain

Trout Mx1 95 EEIEDPSDVEKKIREAQDEMAGVGVGISDDLISLEIGSPDVPDLTIDLPGIARVAVKQGPENIGEQIKRLIRKFKIMKQETINLVVPCNVDIATTEALKMAQEVDPDGE
Trout Mx2 95 EEIEDPSDVENKIRKAQDEMAGVGVGISDDLISLEIGSPDVPDLTIDLPGIARVAVKQGPENIGEQIKRLIRKFKIMKQETINLVVPCNVDIATTEALKMAQEVDPDGE
Trout Mx3 95 EEIEDPSDVEKKIREAQDEMAGVGVGISDDLISLEIGSPDVPDLTIDLPGIARVAVKQGPENIGEQIKRLIRKFKIMKQETINLVVPCNVDIATTEALKMAQEVDPDGE
Trout Mx4 95 EEIEDPSDVENKIRKAQDEMAGVGVGISDDLISLEIGSPDVPDLTIDLPGIARVAVKQGPENIGEQIKRLIRKFKIMKQETINLVVPCNVDIATTEALKMAQEVDPDGE
Trout Mx5 99 IEFDEPSEVVRHVEQAQNLVLAGKGVGICEDLITLKIPTSSVTCVCLSDLDLPGITRVAVKQGPEDIGVQINNLISEFIKNTITILAVVPCNVDIATTEALKMAQEVDPDGE
Trout Mx6 99 IEFDEPSEVVRHVEQAQNLVLAGKGVGICEDLITLKIPTSSVTCVCLSDLDLPGITRVAVKQGPEDIGVQINNLISKFIKNTITILAVVPCNVDIATTEALKMAQEVDPDGE
Trout Mx7 98 INLIGDPEVAHQVAGNLAEGEGMGIHSLITLKIPTSSVTCVCLSDLDLPGITRVAVKQGPEDIGVQINNLISKFIKNTITILAVVPCNVDIATTEALKMAQEVDPDGE
Trout Mx8 100 FEFDDREEVVRHVEQAQNLVLAGKGVGICEDLITLKIPTSSVTCVCLSDLDLPGITRVAVKQGPEDIGVQINNLISKFIKNTITILAVVPCNVDIATTEALKMAQEVDPDGE
Trout Mx9 110 ETEFEDPSLVEIHVKTQNLVLAGKGVGICEDLITLKIPTSSVTCVCLSDLDLPGITRVAVKQGPEDIGVQINNLISKFIKNTITILAVVPCNVDIATTEALKMAQEVDPDGE
Human MxA 131 IETSDASEVEKEINKAQNALAGEGMGISHELITLKIPTSSVTCVCLSDLDLPGITRVAVKQGPEDIGVQINNLISKFIKNTITILAVVPCNVDIATTEALKMAQEVDPDGE
Human MxB 179 LETLQDPGQVEKEIHKANVMAGNRRGISHELITLKIPTSSVTCVCLSDLDLPGITRVAVKQGPEDIGVQINNLISKFIKNTITILAVVPCNVDIATTEALKMAQEVDPDGE

Trout Mx1 205 RTLGILTKPDLVDKGFTEETVVDIVHNEVIHLTKGYMIVKCRGQKEIMERSVTEATEREKAFKFEHAHLSTLYDEGHATIPKLAEKLTLELVHIEKSLPRLEEIQIEAKL
Trout Mx2 205 RTLGILTKPDLVDKGFTEETVVDIVHNEVIHLTKGYMIVKCRGQKEIMERSVTEATEREKAFKFEHAHLSTLYDEGHATIPKLAEKLTLELVHIEKSLPRLEEIQIEAKL
Trout Mx3 205 RTLGILTKPDLVDKGFTEETVVDIVHNEVIHLTKGYMIVKCRGQKEIMERSVTEATEREKAFKFEHAHLSTLYDEGHATIPKLAEKLTLELVHIEKSLPRLEEIQIEAKL
Trout Mx4 205 RTLGILTKPDLVDKGFTEETVVDIVHNEVIHLTKGYMIVKCRGQKEIMERSVTEATEREKAFKFEHAHLSTLYDEGHATIPKLAEKLTLELVHIEKSLPRLEEIQIEAKL
Trout Mx5 209 RTLAAILTKPDLIDPGAENKVLIEIHNVRVIFLSMGYIVKCRGQKQIDENMSTIRAELEEFQNHHEFRSLVREEKATTKCLAKKLTALVKQIKTYLPQMSEKIKIEQL
Trout Mx6 209 RTLAAILTKPDLIDPGAENKVLIEIHNVRVIFLSMGYIVKCRGQKQIDENMSTIRAELEEFQNHHEFRSLVREEKATTKCLAKKLTALVKQIKTYLPQMSEKIKIEQL
Trout Mx7 208 RTLAAILTKPDLIDRGTEDKVDLIVRNKIIPNLNMGYIVKCRGQKQINDGVTTINDAIEEERDFEFNHDFSSLLDEERVTKCLAAARLTQTLVNHIKQSPMQADQIKQQL
Trout Mx8 210 RTLAAILTKPDLIDRGTEDKVDLIVRNKIIPNLNMGYIVKCRGQKQINDGVTTINDAIEEERDFEFNHDFSSLLDEERVTKCLAAARLTQTLVNHIKQSPMQADQIKQQL
Trout Mx9 220 RTLAAILTKPDLIDRGTEDKVDLIVRNKIIPNLNMGYIVKCRGQKQINDGVTTINDAIEEERDFEFNHDFSSLLDEERVTKCLAAARLTQTLVNHIKQSPMQADQIKQQL
Human MxA 241 RTIGILTKPDLVDKGFTEETVVDIVHNEVIHLTKGYMIVKCRGQKEIMERSVTEATEREKAFKFEHAHLSTLYDEGHATIPKLAEKLTLELVHIEKSLPRLEEIQIEAKL
Human MxB 288 RTIGILTKPDLMDRGTESVMNVNRLTYPLKKGYIVKCRGQKEIMERSVTEATEREKAFKFEHAHLSTLYDEGHATIPKLAEKLTLELVHIEKSLPRLEEIQIEAKL

Trout Mx1 315 SETHAELERYGTGPPEDSAERLYFLIDKVTAFQTDAINLSTGEEMKSGVRLNVFSTLRKEFGKWLHLRSEGEIFNQRIEGEVDDYKTYRGRGRELPGFINYKTFEVMVKD
Trout Mx2 315 EETRTALEKCGTGPPEPKERLYFLIDKVTAFQTDAINLSTGEELKSG-DINVFSTLRTEFGKWKAVVDRSGKNFNKIEKEVADYKRYRGRGRELPGFINYKTFEVIKVD
Trout Mx3 315 SETHAELERYGTGPPEDSAERLYFLIDKVTAFQTDAINLSTGEELKSGVRLNVFSTLRKEFGKWLHLRSGENFNQRIEGEVSNYKTYRGRGRELPGFINYKTFEVMVKD
Trout Mx4 315 EETRTALEKCGTGPPEPKERLYFLIDKVTAFQTDAINLSTGEELKSG-DINVFSTLRTEFGKWKAVVDRSGKNFNKIEKEVADYKRYRGRGRELPGFINYKTFEVIKVD
Trout Mx5 319 GEVKNLSKLEGGPPEPEEKRYLIQVITDFNEQITQLSKGDI---VEENLFLMRKEFTWMECLKNKAKSHYHEVVQVQVDEYDQHRGSELPGFSNRYRVQHVQVQ
Trout Mx6 319 GEVKNLSKLEGGPPEPEEKRYLIQVITDFNEQITQLSKGDI---VEENLFLMRKEFTWMECLKNKAKSHYHEVVQVQVDEYDQHRGSELPGFSNRYRVQHVQVQ
Trout Mx7 318 WVYQTELTKEGPPVDPVGRKRYLIEVYKQFNKIDQLCRGELK---NDENLFINMGNIFAKWFEKLGHSRAGYHKMTQDVVNEFDQKHRGRELPGFNNTYLFESVQVQ
Trout Mx8 320 WNVKALVECEGGPDSLAERKQFELIITFEFNKPTLSTGDNTE---VEENLFLMRKEFTWMECLKNKAKSHYHEVVQVQVDEYDQHRGSELPGFSNRYRVQHVQVQ
Trout Mx9 330 ETVKTELKYSYTPPLERKMGYPYTERLIDFIEKIHLCRIGNS---SEKLNHCLRPVQWDSYLSNFKGSLMKVAAMKKNYDKHRGRELMTSDYCVYEHAVQK
Human MxA 351 QRITTEELKQYGVDPEDENEMFFLIDKVNANFDITALMGQEEVTEGIEDIRLFTLRHPEFKWSTIENNFQEGHILSRKIQFENQYRGRGRELPGFNRYRTEFETIVKQ
Human MxB 398 QKATEELRRCGADIPSQEADKMFFLIEIKMFNQDIEKLVGEEVRENERTLYNKIREDFNKNVWGLATNTQKVKNIIEHEVEYKQYRGRGRELPGFNRYRTEFETIVKQ

Trout Mx1 425 QIKQLEGPVAVKLLKESDAVRKVFLLLAQSSFTGFPNLLSAKTKIEAIKQVNESTAESMLRTQFKMELIVYTQDSTYSHSLCERKREEDD-----QPL
Trout Mx2 424 QIKQLEEPVAVKLLKELSDAARKAFLLLAQNSFTGFPNLLSAKTKIEAIKQVNESTAESMLRTQFKMELIVYTQDSTYSHSLCERKREEDD-----RPLPT
Trout Mx3 425 QIKQLEEPVAVKLLKESDAVRKVFLLLAQSSFTGFPNLLSAKTKIEAIKQVNESTAESMLRTQFKMEMIVYTQDSTYSHSLCERKREEDD-----RPLPT
Trout Mx4 424 QIKQLEEPVAVKLLKELSDAARKAFLLLAQNSFTGFPNLLSAKTKIEAIKQVNESTAESMLRTQFKMELIVYTQDSTYSHSLCERKREEDD-----RPLPT
Trout Mx5 426 LVAELKRPAMSTLQKIRDMVQKQDFHLSSESFKNYPYHLVSKKNITIEQKQSNIVKERVVEQFEMEMQVYVYQDEIFNKVMLEAKSHLLEE-----
Trout Mx6 426 LVAELKRPAMSTLQKIRDMVQKQDFHLSSESFKNYPYHLVSKKNITIEQKQSNIVKERVVEQFEMEMQVYVYQDEIFNKVMLEAKSHLLEE-----
Trout Mx7 425 LVGELKNPAMDTLQKIKDLVQKHFFVVSKSSEFNYPCLQRFSMNTNDDIQKQLTPTVMDRIEEQFEMEM---YTQDEIFARTLTPAQKET-----
Trout Mx8 427 LVAELKRPAMSTLQKIRDMVQKQDFHLSSESFKNYPYHLVSKKNITIEQKQSNIVKERVVEQFEMEMQVYVYQDEIFNKVMLEAKSHLLEE-----
Trout Mx9 437 HILGLQEPALDVLKAIIGMVQAEFRNVECEAFKSPYQLRSMALSKIDEITRQETKVEKRIEYINMERLVYTQDSTIFIKGLKDHKAQFKEAIEEEH-----FYDPEEI
Human MxA 461 QIKALEEPVAVMLHTVTDVIRLAFQDVSINKNFEFFNLHRTAKSLIDRAEQEREKEKLRHLHFQMEQIVYCDQVYRQALQVREKELEEEKQK-----S
Human MxB 508 YIQQLVLEPALSMLQKAMEIQQAFINVAKHFGEFFNLQTVQSTIEDIKVKHTAKAENMIQLQFRMEQMVFCQDQLYSVVLLKVVREIFNPLGTPS-----QNM

alpha 2 helix

alpha 3 helix

GTPase effector domain

Trout Mx1 520 TEIRSTIFSTDNHATLQEMMLHLKSYHIISSQRLADQIPMVIYRVLVQEFASQLQREMLQTLQEKDNIQLLKEDIDIGSKRAALQSKLKRMLKARSYLVEF-----
Trout Mx2 534 VSVRSTVNGLDTHATLREMMLHLKSYHIISSQRLADQIPMVIYRVLVQEFASQLQREMLQTLQEKDNIQLLKEDIDIGSKRAALQSKLKRMLKARSYLVEF-----
Trout Mx3 522 PKIRSTIFSTDNHATLQEMMLHLKSYHIISSQRLADQIPMVIYRVLVQEFASQLQREMLQTLQEKDNIQLLKEDIDIGSKRAALQSKLKRMLKARSYLVEF-----
Trout Mx4 534 VSVRSTVNGLDTHATLREMMLHLKSYHIISSQRLADQIPMVIYRVLVQEFASQLQREMLQTLQEKDNIQLLKEDIDIGSKRAALQSKLKRMLKARSYLVEF-----
Trout Mx5 518 ---GEI-AEDKEQDTRSKYPGLLKAYYEIVVQRLADQVPMIRYFYLKQSAKIVCSEMLDLL-HSDDTNDILQEDSEIGQYRAKLAQAADRLILANDKISILL
Trout Mx6 510 ---EGT-AEGSDHTRSKYPGLLKAYYEIVVQRLADQVPMIRYFYLKQSAKIVCSEMLDLL-HSDDTNDILQEDSEIGQYRAKLAQAADRLILANDKISILL
Trout Mx7 512 ---PGK-TDCSGYDTRSKYPELLNSYFEIVVQRLADQVPMIRYFYLKESARILSSEMLGLL-NREDDLEMLKESEIGRRREALRDKVRLGLANNKISTLWDQSG-
Trout Mx8 512 ---EET-PDCSGYDTRSKYPGLLKAYYEIVVQRLADQVPMIRYFYLKESARILSSEMLGLL-NREDDLEMLKESEIGRRREALRDKVRLGLANNKISTLWDQSG-
Trout Mx9 541 EDLTAT-FNSTPFDSCRKLPKDLGVYIEIVVQRLADQVPMIRYFYLKESARILSSEMLGLL-NREDDLEMLKESEIGRRREALRDKVRLGLANNKISTLWDQSG-
Human MxA 559 WDFGAFQSSASDSSMEEIFQHLMAHYEQEASKRISHIPIIQFVFLQYTGQQQLQKMLQLLQDKDITYSLLKERSDTSDKRRLKLERLARLTQARRLAQFPG-----
Human MxB 608 KLNSHPESSNATSSMEEIFQHLMAHYEQEASKRISHIPIIQFVFLQYTGQQQLQKMLQLLQDKDITYSLLKERSDTSDKRRLKLERLARLTQARRLAQFPG-----

alpha 4 helix

Figure 8







Figure 11












Figure 3







Figure 4


G Domain

GED


Teleost Mx group 1



Teleost Mx group 2



Teleost Mx group 3



Mammalian Mx




Figure 6




Figure 7




Figure 9





Figure 10



Highlights

In addition to Mx1-3, six novel Mx genes (Mx4-9) have been cloned in rainbow trout

Salmonids possess 4 groups of Mx genes residing at four chromosome loci

Trout Mx1-4 are highly expressed in blood but Mx5-9 are highly expressed in intestine

Trout Mx gene expression can be induced by poly IC, type I and type II IFNs, and IL-1 β

The potency of IFN induced Mx expression is gene- and cell line-dependent