

Signature: Med Sci Monit, 2003; 9(5): BR218-222
PMID: 12761446

WWW.MEDSCIMONIT.COM

Basic Research

Received: 2003.03.05
Accepted: 2003.04.15
Published: 2003.05.22

Acetate metabolism in brain mechanisms of adaptation to ethanol

Yuri Kiselevski¹, Nicholas Oganessian¹, Sergey Zimatkin¹, Andrzej Szutowicz², Stefan Angielski², Pavel Niezabitowski³, Wojciech Uracz³, Ryszard Gryglewski³

¹ Department of Clinical Biochemistry, Grodno State Medical University, Grodno, Belarus

² Department of Clinical Biochemistry, Gdansk Medical University, Gdansk, Poland

³ Department of Pharmacology, College of Medicine, Jagiellonian University, Cracow, Poland

Sources of support: KBN (grant No 6 P04A 013 10), Ministry of Education of Poland, Medical University of Gdansk project W-113, Jagellonian Medical Research Center, Cracow branch of 'Wspolnota Polska', 'Okregowa Izba Lekarska' in Cracow, Poland

Summary

Background:	The aim of this study was to estimate the role of acetate-induced metabolic changes in brain mechanisms of resistance to the narcotic effect of ethanol.
Material/Methods:	Wistar rats were treated daily with ethanol (3.5 g/kg i.p. for 7 days). During alcohol treatment, the duration of ethanol-induced sleep was decreased. Levels of acetate, acetyl-CoA, adenosine, and the pool of AMP+ADP, and the activity of acetyl-CoA synthetase and 5'-nucleotidase in the brain were measured. Synaptosomal adenosine and acetylcholine release were measured in the presence of acetate, adenosine and 2-chloroadenosine.
Results:	The concentration of acetate was higher in all investigated brain regions of ethanol rats in comparison to controls. The activity of acetyl-CoA synthetase and 5'-nucleotidase, as well as the levels of adenosine and the pool of AMP+ADP were raised in cerebral cortex of ethanol rats. The synaptosomal level of adenosine was higher in control rats. Acetate caused a 3-fold increase of extrasynaptosomal adenosine level. Synaptosomes from ethanol rats showed higher rates of Ca-dependent releases of acetylcholine. 2-chloroadenosine resulted in inhibition of synaptic acetylcholine release only in control rats.
Conclusions:	Ethanol causes an increase of acetate in brain and adenosine level in cerebral cortex. Acetate causes an increase of the extrasynaptic adenosine level. Prolonged ethanol treatment results in an increase of synaptic Ca-dependent acetylcholine release. Seven-day treatment with ethanol eliminates the inhibitory effect of 2-chloroadenosine – A1 adenosine receptor agonist on synaptic Ca-dependent acetylcholine release.
key words:	resistance to ethanol • acetate metabolism • adenosine • acetylcholine release

Full-text PDF: http://www.MedSciMonit.com/pub/vol_9/no_5/3527.pdf

Word count: 2002

Tables: 1

Figures: 3

References: 29

Author's address: Dr. Yuri Kiselevski, PhD, MD, Gorki Str, 80, Grodno Medical University, Grodno, 230015, Belarus, email: kiselevyuu@yahoo.com

BACKGROUND

Several studies have suggested that acetate may affect the CNS during ethanol consumption via an adenosine-mediated mechanism [1,2]. This has been proven by the correlation between ethanol-induced changes in the characteristics of adenosine receptors, adenosine uptake and release in the brain and certain physiological effects of ethanol and acetate [3].

Acetate is a short-chain fatty acid formed in the liver and released into the general circulation following ethanol injection. The plasma acetate level increases after ethanol injection in humans and animals [3]. Acetate readily crosses the blood-brain barrier and is actively metabolized in the brain [4]. The main entry point for acetate into metabolism in vertebrates is its conversion to acetyl-CoA by acetyl-CoA synthetase (EC 6.2.1.1.) This reaction utilizes adenosine triphosphate (ATP) and yields adenosine 3.5-monophosphate (AMP) [5]. The acetyl-CoA which is formed from acetate can be used for energy generation via TCA; on the other hand, acetyl-CoA is utilized in the brain's cholinergic neurons for acetylcholine (ACh) synthesis. It has been shown that extracellular acetate is accumulated by cholinergic nerve terminals for ACh formation and release [6]. ACh release in the cortex is associated with both behavioral activation and cortical activation, as determined by electroencephalography (EEG) [7]. The rate of efflux of ACh in the cortex is greatest during periods of desynchronized, high-frequency EEG activity, such as that observed during waking and paradoxical sleep, and the lowest during slow-wave sleep. [8]. Factors that modulate the activity of cholinergic neurons could, therefore, alter cortical ACh release and have a major effect on the cortical arousal and behavioral state. Purine nucleoside adenosine is thought to be a modulator of neuronal activity, including cholinergic neurotransmission [9]. In addition, pharmacological experiments suggest that adenosine is a sleep factor, since systemic or intracerebroventricular injections promote sleep and decrease wakefulness [10]. Several studies have shown that some of the central depressant effect of ethanol can be inhibited by adenosine receptor blockers [1]. This confirms the role of adenosine in the central depressant actions of ethanol. Considering the fact that ethanol consumption induces an increase in the adenosine level, it is possible to suppose that the hypnotic effect of ethanol is mediated by adenosine.

The aim of this study was to estimate the role of brain acetate-induced metabolic changes in mechanisms of resistance to the narcotic effect of ethanol.

In order to achieve this aim, we attempted to estimate:

- 1) the effect of chronic ethanol injection on brain acetate and adenosine levels;
- 2) the effect of acetate on synaptic adenosine production and release;
- 3) the effect of adenosine and its receptor agonist on synaptic acetylcholine release in cerebral cortex of alcohol-treated rats.

MATERIAL AND METHODS

Male Wistar rats weighing 100–120 g were used in all studies. The animals were fed standard chow and housed in group cages in an air-conditioned room with a cycle of 12 h light/12 h dark. The rats were divided into 3 groups according to the duration of ethanol-induced sleep time (after an acute test dose of ethanol: 25% solution, 3.5 g/kg i.p.):

- long-sleeping (LS – 178.5±36.6 min);
- medium-sleeping (MS – 113.8±9.3 min);
- short sleeping (SS – 69.2±11 min).

In further experiments we used only MS rats, which made up 70% of the general population. After 14 days of recovery, the MS animals were treated with a daily i. p. injection of 25% ethanol solution, 3.5 g/kg of body weight. The duration of ethanol-induced sleep time was decreased from 108.8±14.3 min. to 13±9 min. by the seventh day of alcohol treatment. The control group received i. p. injections of 0.9% NaCl solution. The rats were sacrificed by decapitation 24 h after the seventh injection of ethanol (for ethanol animals) or NaCl solution (for the control group). Their brains were rapidly removed at the 0–4°C. The brain regions (cerebral cortex, hypothalamus, striatum, medulla oblongata) were dissected, and remained frozen in liquid nitrogen for determination of acetyl-CoA synthetase and 5'nucleotidase activities, as well as the concentration of acetate, acetyl-CoA and adenosine.

The determination of acetyl-CoA synthetase activities was performed in homogenates using the citrate synthetase method [11]. The general activity of 5'nucleotidase was determined in homogenates of brain regions by UV methods with adenosine deaminase and glutamate dehydrogenase [12].

The level of acetyl-CoA was measured by the enzymatic method [13]. The concentration of acetate was determined by gas chromatography (HP 6890) [14]. The concentration of adenosine and the ADP+AMP pools were determined by HPLC (Beckman Gold System) [15].

For metabolic study, nerve terminals were isolated from pooled homogenates of cerebral cortex from 5 animals immediately after dissection. The synaptosomal fraction from cerebral cortex was isolated by differential centrifugation and flotation in Ficoll gradient [16]. The synaptosomes were suspended in 320 mM buffered sucrose. They were incubated for 30 min. at 37°C with shaking at 100 cycles min⁻¹, in depolarizing medium containing a final volume of 2.0 ml: 20 mM Na-HEPES, 1.5 mM Na-phosphate (final pH 7.4), 90 mM NaCl, 30 mM KCl, 2.5 mM Na-pyruvate (or 2.5 mM K-acetate), 2.5 mM Na-L-malate, 0.01 mM choline chloride, 0.01 mM eserine sulfate, 320 mM sucrose and 2.5–3.0 mg of synaptosomal protein. Quantal ACh release was stimulated by addition of 1.0 mM CaCl₂. Adenosine and 2-chloroadenosine were added to the incubation medium (final concentration 10 μM) to study the effects of adenosine on ACh release. Incubation was stopped by centrifugation at 15000 × g for 3 min. in Eppendorf tubes. The adenosine

concentration in the supernatant and pellet was measured using luciferin-luciferase luminometry [17]. The determination of acetylcholine was performed in supernatant by luminometric detection (LKB Pharmacia) [18].

The concentration of protein was measured according to the Bradford method [19].

Data are presented as mean \pm SD. The Mann-Whitney test was used to compare the alcohol-treated group with the controls for metabolic experiments, and the Student test was used for homogenate experiments. The level of significance was set at 0.05.

RESULTS

Table 1 shows that 7-day alcohol treatment resulted in increased acetate concentration in all researched brain regions (cerebral cortex, hypothalamus, striatum and medulla oblongata). The rate of rise of the acetate level was higher in cerebral cortex than in other researched brain regions: about 80% for cerebral cortex, and about 50% for hypothalamus, striatum and medulla oblongata. Simultaneously, there were no significant changes in the brain level of acetyl-CoA in rats treated with alcohol as compared to control animals (Table 1). The activities of acetyl-CoA synthetase and 5'nucleotidase, as well as the levels of adenosine and the pool of AMP+ADP were raised in alcohol-treated rats only in cerebral cortex (Table 1), and showed no significant difference in hypothalamus, striatum and medulla oblongata.

The synaptosomal level of adenosine was higher in control rats than in animals treated with alcohol (Figure 1). In the presence of acetate in the incubation medium there was a 3-fold increase in the release of synaptosomal adenosine (Figure 2).




Figure 1. Effects of 2.5 mM acetate on adenosine concentration in cerebral cortex synaptosomes of control (open column) and alcohol (solid column) rat. Results are means \pm SD from 5 experiments done in duplicate. * Significantly different from respective no-acetate conditions, $p < 0.05$.

7-day alcohol administration caused no significant changes in resting (Ca^{2+} -nondependent) ACh release in synaptosomes (Figure 3). By contrast, in the presence of Ca^{2+} the synaptosomes from ethanol-treated rats showed higher rates of Ca^{2+} -dependent release of ACh (Figure 3). 10 μM of adenosine had no influence on Ca^{2+} stimulated ACh release from the neural endings of either control or experimental rats. However, 10 μM of 2-cloradenosine resulted in inhibition of synaptic ACh release only in control rats, but had no influence on this parameter in animals treated with alcohol (Figure 3).

DISCUSSION

Since acetate is an endogenous metabolite that is considered non-toxic, little attention has been given to the possibility that it may contribute to the intoxicating effects of ethanol. Although earlier studies have suggested that plasma acetate level increased 6 times 30 min. after a single intravenous dose of 0.5 g/kg ethanol in rats [20], there are no data concerning the influence ethanol consumption on the level of acetate in brain [21]. The present data have shown that chronic alcohol treatment resulted in an increased acetate level in brain. As acetate readily crosses the blood-brain barrier, the most probable reason for the rise of brain acetate concentration in alcohol-treated rats is its delivery with blood flow from peripheral tissues, mainly from the liver, which is the main ethanol-metabolizing organ.

The utilization of acetate by tissues [22] and acetate uptake by neurons [6] is dependent on the activities of acetyl-CoA synthetase. It is known that this enzyme is substrate-inducible, but as the acetate level increases both in blood and in brain after ethanol injections, the activation of acetyl-CoA synthetase would be expected. We have shown that 7-day alcohol treatment results in the activa-





Figure 2. Effects of 2.5 mM acetate on extrasynaptosomal adenosine concentration in control (open column) and alcohol (solid column) rat. Results are means \pm SD from 5 experiments done in duplicate. * Significantly different from respective no-acetate conditions, $p < 0.05$.

Table 1. Effect of 7-day ethanol treatment on the concentration of acetate, acetyl-CoA, adenosine, pools of ADP+AMP, activities of acetyl-CoA synthetase and 5'nucleotidase in four brain regions (mean±SD of 6–10 rats).

Parameter (Units)	Groups	Brain regions			
		Cerebral cortex	Hypoth	Striatum	Medulla oblongata
Acetate (mM/g of tissue)	Control	0.31±0.1	0.37±0.07	0.32±0.06	0.32±0.07
	Alcohol	0.62±0.14*	0.56±0.1*	0.48±0.06*	0.51±0.1*
Acetyl-CoA (nM/g of tissue)	Control	82.2±19.1	60.2±7.5	69.3±15.8	91.8±13.6
	Alcohol	88.3±8.8	62.5±11.2	75.3±37.4	97.3±16.4
Acetyl CoA synthetase (nM/mg protein min ⁻¹)	Control	2.56±0.27	2.45±0.27	2.8±0.2	2.97±0.17
	Alcohol	3.48±0.3*	2.96±0.54	3.15±0.3	3.3±0.15
5'nucleotidase (nM/mg protein min ⁻¹)	Control	15.8±4.0	16.3±2.6	13.6±1.2	21±6.4
	Alcohol	33.3±10.0*	15.4±3.9	16±6.2	15±3.8
AMP+ADP (μM/g of tissue)	Control	2.097±0.5	2.7±0.3	2.6±0.35	1.9±0.19
	Alcohol	2.89±0.2*	2.9±1.1	2.5±0.5	1.7±0.5
Adenosine (μM/g of tissue)	Control	0.31±0.03	0.29±0.15	0.42±0.04	0.23±0.03
	Alcohol	0.41±0.05*	0.26±0.09	0.44±0.021	0.23±0.06

* significantly different from respective controls, p<0.05;

Hypoth – Hypothalamus

**Figure 3.** Effects of 10 μM adenosine (ado) and 10 μM 2-chloroadenosine (CLA) on Ca²⁺ stimulated acetylcholine (ACh) release in brain synaptosomes from control (open column) and alcohol (solid column) rat. Results are means ± SD from 5 experiments done in duplicate. * Significantly different from respective controls, p<0.05; + Significantly different from respective no 2-chloradenosine conditions, p<0.05.

tion of acetyl-CoA synthetase in cerebral cortex. Early studies demonstrated that the activities of acetyl-CoA synthetase were also increased in rat liver and heart after chronic ethanol treatment [23]. In one study it was shown that alcohol and acetate produce an increase of the acetyl-CoA level in cultured hepatocytes [5]. On the other hand, there was no such effect of ethanol on brain acetyl-CoA level in the present study, which is consistent with other reports [24]. One possible explanation for this fact is that the stability of the acetyl-CoA level is critically important for the majority of metabolic processes, and significant oscillations of this parameter *in vivo* are inadmissible.

There is much experimental evidence that ethanol and/or acetate cause an increase of the adenosine level,

both in the peripheral and central nervous systems [1]. This is made possible by means of transformation of acetate into acetyl-CoA. The AMP formed from this transformation is subsequently converted to adenosine by 5'nucleotidase. We have found that chronic alcohol treatment results in increasing adenosine concentration in cerebral cortex only. This is in agreement with the rising activities of acetyl-CoA synthetase and 5'nucleotidase also in the same brain region during treatment with alcohol. Simultaneously, several studies have demonstrated that ethanol can increase the extracellular adenosine level by inhibiting its transport [25]. Decreasing adenosine concentration in the cortical nerve endings of alcohol-treated rats is probably the result of impairment of the neuronal mechanism of adenosine uptake by ethanol.

Fredholm et al. have shown that 20 mM ethanol, as well as 5 mM acetate, did cause enhancement of the efflux of adenosine from brain slice [26]. We found a similar effect in cortical synapses. 2.5 mM acetate caused a 3-fold increase of extrasynaptical adenosine concentration. These data justify the supposition that acetate, like ethanol, can block adenosine transport into neuronal cells. Together, these results suggest that ethanol and acetate produce an increased extracellular adenosine level by increasing its production and/or inhibiting adenosine uptake. A prolonged increase of adenosine level in brain during chronic alcohol consumption can lead to decreased sensitivity of the adenosine receptor. Dar et al. have suggested that the adenosine A1 receptor is desensitized by chronic ethanol treatment [3]. Adenosine modulates an 'inhibitory tone' in the CNS and depresses synaptic transmission of various neuromediators, including acetylcholine, by means of activation of the presynaptic A1 receptor [27]. The acceleration of Ca-dependent acetylcholine release from the nerve ending of rats treated with ethanol, according to our findings, is probably caused by elimination of adenosine's 'inhibitory tone' due to desensitization of the presynaptic adenosine A1 receptor. The results of our experiment suggest that the inhibitory effect of selective A1 receptor agonist – 2

chloroadenosine on Ca-stimulated synaptic acetylcholine release is lost during 7-day ethanol injection. It is known that adenosine is a somnogenic factor [28]. According to different experiments, adenosine promotes sleep, at least partly, by activating A1 receptors located on neurons in the basal forebrain and the presynaptic membrane of the nerve endings of these neurons in the cerebral cortex [29]. Accordingly, a decreased duration of ethanol-induced sleep for 7 day treatment with alcohol may result from A1 adenosine receptor desensitization, which is caused by acetate-induced prolonged high extracellular adenosine level.

CONCLUSIONS

1. Ethanol causes an increase of acetate in brain and adenosine level in cerebral cortex.
2. Acetate causes an increase of extrasynaptic adenosine level.
3. Prolonged ethanol treatment results in increased release of synaptic Ca-dependent acetylcholine.
4. Seven-day ethanol treatment eliminates inhibition of the 2-chloroadenosine – A1 adenosine receptor agonist on synaptic Ca-dependent acetylcholine release.

Acknowledgments

This paper was presented at the 10th International Symposium of the Jagellonian Medical Research Center, Krakow, Poland, June 2–4, 2002. This study was made possible thanks to scientific cooperation between Grodno State Medical University (Grodno, Belarus), Collegium Medicum of Jagellonian University (Cracow, Poland) and Gdansk Medical University (Gdansk, Poland), and to financial support from following Polish institutions: the State Research Commission KBN (grant No. 6 P04A 013 10), the Ministry of Education, the Jagellonian Medical Research Center, the Medical University of Gdansk project W-113, the Cracow branch of „Wspolnota Polska“, and the „Okregowa Izba Lekarska“ in Cracow, Poland

REFERENCES:

1. Carmichael FJ, Orrego H, Israel Y: Acetate-induced adenosine mediated effects of ethanol. *Alcohol Alcohol*, 1993; 2(Suppl): 411-418
2. Phillis JW, O'Regan MH, Perkins LM: Actions of ethanol and acetate on rat cortical neurons: ethanol/adenosine interactions. *Alcohol*, 1992; 9: 541-546
3. Dar MS, Clark M: Tolerance to adenosine's accentuation of ethanol-induced motor incoordination in ethanol-tolerant mice. *Alcohol Clin Exp Res*, 1992; 16: 1138-1146
4. Cullen N, Carlen PL: Electrophysiological actions of acetate, a metabolite of ethanol on hippocampal dentate granule neurons: interactions with adenosine. *Brain Research*, 1992; 588: 49-57
5. Smith CM, Israel BC, Iannucci J, Marino KA: Possible role of acetyl-CoA in the inhibition of CoA biosynthesis by ethanol in rats. *Am Inst Nutr*, 1986; 22: 452-460
6. Carroll P: Evidence to suggest that extracellular acetate is accumulated by rat hippocampal cholinergic nerve terminals for acetylcholine formation and release. *Brain Res*, 1997; 753: 47-55
7. Jones BE: The organization of central cholinergic systems and their functional importance in sleep-waking states. *Prog Brain Res*, 1993; 98: 61-71
8. Jasper HH, Tessier J: Acetylcholine liberation from cerebral cortex during paradoxical (REM) sleep. *Science*, 1971; 172: 601-602
9. Kurokawa M, Shiozaki S, Nonaka H et al: In vivo regulation of acetylcholine release via adenosine A1 receptor in rat cerebral cortex. *Neurosci Lett*, 1996; 209: 181-184
10. Ticho SR, Radulovacki M: Role of adenosine in sleep and temperature regulation in the preoptic area of rats. *Pharmac Biochem Behav*, 1991; 40: 33-40
11. Szutowicz A, Stepien M, Piec G: Determination of pyruvate dehydrogenase and acetyl-CoA synthetase activities using citrate synthetase. *Analytical Biochem*, 1981; 115: 81-87
12. 5'-Nucleotidase. In Bergmeyer HU, editor. *Methods of Enzymatic Analysis*. Vol. IV. Florida, Basle: Verlag Chemie Weinheim, 1984; 106-113
13. Kato T: CoA Cycling: An Enzymatic Amplification Method for Determination of CoASH and Acetyl CoA. *Anal Biochem*, 1975; 66: 372-392
14. Giles HG, Meggiorini S, Vidins EI: Semiautomatic analysis of ethanol and acetate in human plasma by head space gas chromatography. *J Fisiol Pharmacol*, 1986; 64: 717-719
15. Dobolyi A, Reichart A, Szikra T et al: Sustained depolarization induced changes in the extracellular concentration of purine and pyrimidine nucleosides in the rat thalamus. *Neurochem Int*, 2000; 37: 71-79
16. Szutowicz A, Lysiak W, Angelski S: The effect of +/- hydroxycitrate on pyruvate metabolism in rat brain synaptosomes. *J Neurochem*, 1977; 29: 375-378
17. Kather H, Wieland E, Waas W: Chemiluminescent determination of adenosine, inosine and hypoxanthine/xanthine. *Anal Biochem*, 1987; 163: 187-206
18. Ricny J, Coupek J, Tucek S: Determination of acetylcholine and choline by flow-injection with immobilized enzymes and fluorimetric or luminometric detection. *Analyt Biochem*, 1986; 176: 221-227
19. Ricny J, Coupek J, Tucek S: Determination of acetylcholine and choline by flow-injection with immobilized enzymes and fluorimetric or luminometric detection. *Analyt Biochem*, 1986; 176: 221-227
20. Carmichael FJ, Israel Y, Crawford M, Minhas K: Central nervous system effects of acetate: contribution to the central effects of ethanol. *J Pharmacol Exp Ther*, 1991; 259: 403-408
21. Wanewski RA, Martin DL: Preferential utilization of acetate by astrocytes is attributable to transport. *J Neurosci*, 1998; 18: 5225-5233
22. Woodnutt G, Parker DS: Acetate metabolism by tissues of the rabbit. *Comp Biochem Physiol*, 1986; 85B: 487-490
23. Lukivskaya OYa, Buko VU: Utilization of ketone bodies by the rat liver, brain and heart in chronic alcohol intoxication. *Alcohol Alcohol*, 1993; 28: 431-6
24. Guynn RW: Effect of ethanol on brain CoA and acetyl-CoA. *J Neurochem*, 1976; 27: 303-304
25. Diamond I, Naggy L, Mochly-Rosen D, Gordon A: The role of adenosine and adenosine transport in ethanol-induced cellular tolerance and dependence. Possible biologic and genetic markers of alcoholism. *Ann NY Acad Sci*, 1991; 625: 473-487
26. Fredholm BB, Wallman-Johansson A: Effects of ethanol and acetate on adenosine production in rat hippocampal slices. *Pharmacol Toxicol*, 1996; 79: 120-123
27. Dunwiddie TV, Fredholm BB: Adenosine neuromodulation. In: Jacobson KA, Jarvis MF, editors. *Purinergic Approaches in Experimental Therapeutics*. New York: Wiley Liss, 1997; 359-82
28. Porkka-Heiskanen T, Strecker RE, McCarley RW: Brain site-specificity of extracellular adenosine concentration changes during sleep deprivation and spontaneous sleep: an in vitro microdialysis study. *Neuroscience*, 2000; 99: 507-517
29. Portas CM, Thakkar M, Rainnie DG: Role of adenosine in behavioral state modulation: a microdialysis study in the freely moving cat. *Neuroscience*, 1997; 79: 225-235