

Human monocytes/macrophages in the antitumour response of the host

ANNA SZAFLARSKA, MACIEJ SIEDLAR, MAREK ZEMBALA

Department of Clinical Immunology, Polish-American Institute of Paediatrics, Jagiellonian University Medical College, Cracow, Poland

Abstract

Monocytes/macrophages play a significant role in the host's response to tumours. This includes: cytotoxic/cytostatic activity, presentation of tumour-associated antigens and induction of specific anticancer response of lymphocytes. Circulating blood monocytes respond to a gradient of chemoattractants produced by the tumour, migrate out from the blood to the tumour bed and form a large part of the cellular infiltrate as tumour infiltrating macrophages (TIM). Monocytes and macrophages produce a large array of factors (cytokines, reactive oxygen and nitrogen intermediates, growth factors, prostacyclins, ect.) with opposing biological activities. Consequently, TIM exhibit both tumour growth promoting and inhibitory activities. Furthermore, tumour-derived molecules also modulate TIM activity. In some circumstances monocytes/macrophages are involved in the metastatic process. This review summarizes the current state of knowledge in this area indicating that in fact macrophage-tumour interactions are quite complicated and a delicate balance exists between antitumour response and protumour effect of TIM and the suppression of TIM activity by the tumour. The clinical implications of these findings are also discussed.

Key words: monocytes/macrophages, tumour cells, cytokines, cytotoxic mediators, tumour infiltrating macrophages, immunotherapy

(*Centr Eur J Immunol* 2003; 28(2): 88–103)

Introduction

Monocytes/macrophages belonging to the mononuclear phagocyte system are involved in the host response against cancer and function not only as cells presenting tumour-associated antigens to (tumour infiltrating) T lymphocytes but also act as cytotoxic/cytostatic effector cells. In addition, they express surface molecules relevant for cell adhesion and cellular interactions and regulate the functions of other cells in the immune system. Monocytes/macrophages are able to distinguish and kill malignant, but not normal, cells and form the major component of the mononuclear cell infiltrate of many tumours, as tumour infiltrating macrophages (TIM). TIM may both inhibit and promote tumour growth and neoangiogenesis. These opposing activities of TIM are best explained by the “macrophage-tumour balance” hypothesis [1]. The exact role of monocytes/macrophages in human malignancy remains not fully understood.

Function of monocytes in malignant diseases

Production of cytokines

Monocytes and macrophages are capable of producing numerous cytokines, e.g. tumour necrosis factor alpha (TNF), interleukins (IL): IL-1, IL-6, IL-10, IL-12, IL-18, colony-stimulating factors (CSF), chemokines and cytotoxic mediators: reactive oxygen (ROI) and nitrogen intermediates (RNI), which appear to play an important role in the regulation of tumour growth.

Tumour necrosis factor

TNF is produced mainly by mononuclear phagocytes and is cytotoxic for some tumour cells. It is proinflammatory cytokine that mediates and induces tumouricidal activity of monocytes and stimulates its own and other cytokine production by monocytes [2], increases their antigen presenting capacity and upregulates HLA-DR and interferon γ receptor

Correspondence: Anna Szaflarska, Medical College, Jagiellonian University, Department of Clinical Immunology, 265 Wielicka st, 30-663 Cracow, Poland. Phone/fax number: +48 12 658 97 30, e-mail: miszafla@cyf-kr.edu.pl

(IFN γ -R) expression on monocytes [3]. TNF downregulates the monocyte CD86 costimulatory molecule expression and mannose receptor-dependent endocytosis [4]. It also enhances HLA-A,B,C and HLA-DR expression on tumour cells [5], but inhibits IFN γ -induced HLA-DR expression on normal differentiated cells (fibroblasts, macrophages) [6]. Tumour cells are able to induce TNF-mRNA expression and TNF production by human monocytes [7]. This capacity is not limited to viable but also metabolically inactive tumour cells and their constituents like membranes or hyaluronan [8, 9]. Several surface molecules (CD44, HLA-DR) [7] and protein kinases: a tyrosine-protein kinases (PTK) and calcium-phospholipid-dependent protein kinases (PKC) are involved in signal transduction for TNF release after stimulation with cancer cells [10]. The antitumour effect of TNF is due to its direct cytotoxic activity on tumour cells and selective damage of endothelial cells of the tumour vasculature leading to apoptosis and necrosis of the tumour [11-13]. The inverse correlation between TNF-mRNA expression and microvessels count was found in non-small lung carcinoma [14]. TNF also induces neutrophil-mediated cytostasis of tumour cells that is mediated by high local concentration of hydrogen peroxide [15].

On the other hand, TNF produced by some tumour cells may enhance tumour spread and metastatic formation by induction of transcription of metalloproteinase (MMP)-9 gene in stromal cells of giant cell tumour of bone [16] and proMMP-9 production in monocytes [17] that cause degradation of extracellular matrix compounds [18]. TNF also facilitates the adherence of tumour cells to vascular endothelium [19, 20]. TNF receptor type I (TNFRI) is involved in the regulation of intercellular adhesion molecule-1 (ICAM-1), E-selectin, vascular adhesion molecule-1 (VCAM-1) and CD44 expression on vascular endothelial cells [21, 22] that regulate monocyte migration from the blood vessels to the tumour site. Hence, TNF has several opposite effects on the tumour growth and influences the TIM-tumour balance.

Production of TNF by lipopolysaccharide (LPS)-stimulated monocytes is reduced in colon cancer patients and this reduction is more pronounced in Dukes' C compared to Dukes' A and B tumour stages. This suppression is not mediated by IL-10 and disappears following surgical resection of the tumour [23]. On the other hand, an increased spontaneous or LPS-stimulated production of TNF by peripheral blood mononuclear cells (monocytes are the major cellular source of TNF) was found in gastric cancer patients [24]. Furthermore, serum levels of TNF are increased in all stages of gastric cancer (our unpublished observations). An increased serum levels of TNF are also found in patients with hepatocellular carcinoma and metastatic liver carcinoma [25]. On the other hand, TNF is undetectable in the serum of patients with metastatic breast carcinoma [26] and patients with gastrointestinal cancer-associated cachexia [27]. These and other observations suggest that TNF has no role in cancer-

related cachexia in man [24, 28] but may play a significant role in the regulation of the inflammatory host response to the growing tumour [29].

Interleukin-1

IL-1 activates tumouricidal activity of monocytes but also trigger the release of IL-1 and TNF by IFN γ -primed monocytes [30, 31]. IL-1 mimics many of the biological actions of TNF [32] and both these cytokines act as stimulators of IFN γ -R synthesis. IL-1 exerts cytotoxic and cytostatic effects *in vitro* on several tumour cell lines [33, 34]. IL-1 and IFN γ have additive growth inhibitory effect on colon cancer cell line [35].

No significant changes in the production of IL-1 by blood monocytes from patients with untreated colorectal [36], lung [37] or head and neck [38] cancers were observed. However, cytoplasmic expression of IL-1 (α and β) in monocytes was reduced in patients with lung and colorectal cancers [36]. The presence of IL-1 and IL-6 was detected in the effusions from ovarian cancer [39]. Only a few patients with metastatic breast cancer had detectable IL-1 β serum levels [26] but in patients with hepatocellular carcinoma, metastatic liver carcinoma and gastrointestinal cancer serum levels of IL-1 α and β were increased [25, 40]. In contrast, no changes in serum levels of these cytokines were found in endometrial and urinary tract cancer [41, 42], while IL-1 release *in vitro* by unstimulated PBMC of patients with urinary tract cancer was decreased [42]. Hence, no consistent results concerning the production of IL-1 are found in different types of cancer. Moreover, IL-1 is not associated with cancer anorexia-cachexia syndrome [28].

Interleukin-6

IL-6 is a pleiotropic cytokine produced by many different cells, mainly monocytes/macrophages, fibroblasts and endothelial cells. This cytokine acts on activated B cells and induces immunoglobulin production. IL-6 is also involved in growth, differentiation and activation of T cells. It synergises with IL-1 in induction of IL-2 production and IL-2 receptor (CD25) expression on T cells. IL-6 is one of the hepatocyte stimulating factors regulating the biosynthesis of acute phase proteins and an important regulator of hematopoiesis [43].

IL-6 plays an important role in the pathogenesis of plasmacytoma [44, 45]. Its increased serum level positively correlates with severity of disease [46]. Monocytes from patients with head and neck cancers produce an increased amount of IL-6 [38]. Raised serum level of IL-6 is observed in patients with pancreatic [47, 48], gastric [25] and liver carcinomas (hepatocellular and metastatic liver carcinomas) [25,49]. In contrast to these observations [25], we have not found any significant changes in the serum levels of IL-6 in patients with different stages of gastric cancer (unpublished). An increased serum level of IL-6 correlates

with tumour progression and poor prognosis in metastatic breast cancer [26]. IL-6 enhances acute phase response and its serum level correlates with poor nutritional status, impaired patient performance status and shorter survival in lung cancer patients [50]. However, other studies revealed no correlation between IL-6 serum levels and the presence of the cancer anorexia-cachexia syndrome [28].

Interleukin-10

IL-10 downregulates antitumour activity of monocytes by suppressing production of IL-1 β and TNF and inhibits HLA-DR expression on antigen presenting cells [51]. Although IL-10 inhibits ROI formation in activated monocytes, it has no inhibitory effect on ROI production by activated macrophages [52]. On the other hand, in experimental systems, it inhibits angiogenesis and suppresses growth and metastasis formation by human melanoma cells [53]. An increased serum level of IL-10 in patients with resectable hepatocellular carcinoma [49] and advanced solid tumours [54] appears to be an independent prognostic factor [49, 54]. Monocytes from breast cancer patients show an increased production of IL-10 and decreased IL-12 [55]. Colon and renal carcinoma cell lines stimulate peripheral blood monocytes and lamina propria mononuclear cells to produce increased levels of IL-10 [4, 56]. Tumour-cell-derived TGF β 1 and PGE₂ are the potent IL-10 synthesis stimulators [56]. In this context, it is of interest that increased serum levels of PGE₂ have been found in some types of cancers and in Hodgkin's disease [4, 57, 58]. However, no changes in the serum levels of IL-10 nor its production by PBMC from gastric cancer were found (our unpublished observations).

Interleukin-12

IL-12 has a powerful antitumour activity and is primarily produced by monocytes/macrophages. It skews the immune response in favour of Th₁ cells that preferentially induce cell-mediated immunity. IL-12 acts as a growth factor for activated NK and T cells [59] and stimulates the production of TNF, IFN- γ , GM-CSF and IL-8 by T lymphocytes and/or NK cells [60, 61]. IL-12 enhances its own production by dendritic cells [61]. On the other hand, IL-12 also stimulates the production of IL-10 by T lymphocytes [61]. Antitumour activity of IL-12 (tumour regression or tumour growth inhibition due to administration of IL-12) was demonstrated *in vivo* using 17 different lines of transplantable murine tumours including carcinomas, sarcomas, melanomas and lymphomas [61, 62]. The antitumour effect of IL-12 may be independent of NK cells, since comparable activity was observed in NK-deficient mice [63]. Tumours undergoing IL-12-mediated regression have large numbers of TIM [63, 64, 65]. There are three main mechanisms of antitumour activity of IL-12: induction of CD8 cytotoxic cells, production of IFN- γ by T and NK cells and inhibition of neoangiogenesis [62].

IL-12 induces cytolytic activity of PBMC from patients with lung cancer against lung cancer cells and Daudi lymphoma cells [66]. However, monocytes from patients with lung cancer show decreased production of IL-12 [66]. Also PBMC from patients with colorectal cancer show decreased IL-12 and increased IL-10 production, the latter known to antagonise IL-12 synthesis. The decrease in IL-12 production is most clearly seen in advanced colorectal cancer [67]. However, in some types of cancer, serum or ascitic fluid levels of IL-12 are increased [68, 69]. There is no correlation between LPS-stimulated IL-12 secretion by blood monocytes and survival of patients with head and neck cancer [70]. The effectiveness of IL-12 therapy was demonstrated in several types of experimental tumours [64, 65, 71-75]. Human IL-12 is undergoing phase I clinical trial in metastatic renal cancer and malignant melanoma [76].

Interleukin-18

IL-18 is monocyte-derived pleiotropic cytokine that synergises with IL-12 and induces IFN- γ , IL-1 β , NO production by T lymphocytes and promotes Th₁-mediated immune response. IL-18 also enhances IL-13 production by T and NK cells [77]. Antitumour effects of IL-18 may also involve FasL-mediated apoptosis of tumour cells by cytokine enhanced Fas-ligand (Fas-L) expression on NK cells [78] and inhibition of neoangiogenesis in experimental tumours [79]. The decreased production of this cytokine by colon adenocarcinoma cells in comparison to normal epithelial cells of the colon mucosa is associated with immunosuppression observed in colon cancer [80]. However, our unpublished observations indicate an increased IL-18 serum level in patients with gastric cancer.

Colony stimulating factors

G-CSF is produced primarily by activated monocytes/macrophages and enhances the expression of complement receptor (CR) type 1, 3, Fc γ RI (CD64) and Fc γ RIII (CD16) on monocytes and upregulates their tumouricidal capacity [81, 82]. M-CSF is also produced by monocytes/macrophages and stimulates the differentiation of monocytes and macrophages from their progenitor cells. Its high serum levels were found in patients with breast, endometrial and ovarian cancers and correlated with poor prognosis [83, 84]. M-CSF is also produced by tumour cells. Expression of M-CSF and its receptor by breast cancer cells is associated with high macrophage infiltration and poor prognosis [85, 86]. Monocytes from patients undergoing GM-CSF therapy showed a significant increase in MHC class I and II expression, production of TNF and monocyte-mediated cytotoxicity against U937 tumour cells [81]. GM-CSF from transfected human colon cancer cells stimulates monocytes to secrete monocyte chemotactic protein (MCP)-1 and induces expression of the CD11b adhesion molecule [87].

Production of reactive nitrogen intermediates

NO is involved in tumouricidal activity of monocytes/macrophages [88, 89]. Cytotoxicity of NO is mainly due to peroxynitrite (ONOO⁻) or nitrosothiols (RSNO) production [90, 91]. Peroxynitrite causes the inhibition of mitochondrial respiration and damage of variety of mitochondrial components, nitrosothiols inhibit respiratory complex I, while NO inhibits cytochrome oxidase [91]. Furthermore, NO selectively inhibits IL-12 synthesis by activated monocytes [92, 93] and suppresses T lymphocyte proliferation [94].

Inducible nitric oxide synthase (iNOS) is responsible for biosynthesis of NO by monocytes. Some cytokines (IL-1 β , IFN- γ , IL-2, TNF) and LPS induce iNOS-mRNA synthesis but not NO release by monocytes [95-98]. However, some cancer cells may stimulate monocytes for de novo production of NO [95]. Both iNOS-mRNA and iNOS protein were observed in monocytes stimulated with colon carcinoma cell line, but not with human pancreatic cancer cell line [96]. We have recently found that following stimulation with cancer cells, CD14⁺/CD16⁺ subpopulation of monocytes show an increased expression of iNOS protein and release of NO in comparison to "classical" (CD14⁺) monocytes [97]. Human urothelial carcinoma cell line also fails to induce NO production by monocytes, while tumour cells display iNOS expression and NO production in cocultures with monocytes [98]. The above findings may cast some doubts about the ability of monocytes to produce NO. However, our other observations indicate the expression of iNOS in monocytes stimulated by the supernatants from the culture of cancer cell lines. This phenomenon is probably due to microparticles released by tumour cells. The CD29, CD44, CD58, HLA-DR, and MHC class I of monocytes are engaged in tumour cell-induced production of NO [99]. The signal transduction pathways for NO and TNF production seem to be different as at least three protein kinases: PKC, PTK and cAMP-dependent kinase (PKA) are involved in the induction of NO by monocytes stimulated with tumour cells [10].

L-arginine is the substrate molecule for NO synthesis, but some tumours stimulate monocytes for biosynthesis of ornithine, a precursor for polyamine growth factors: putrescine, spermidine and spermine [100]. Polyamines may promote experimental tumour growth not only by increasing proliferation of tumour cells, but also by induction of neoangiogenesis [101]. NO is involved in apoptosis of some cells (it promotes or inhibits apoptosis) and its effect is dose-dependent and cell-type specific [102]. In cholangiocarcinoma cells, NO inhibits apoptosis directly by blocking caspase 9 activation [103]. On the other hand, the presence of iNOS in pancreatic cancer cells positively correlates with their apoptosis [104]. Also mononuclear cells, including macrophages infiltrating colorectal cancer, show both the increased apoptosis and expression of iNOS [105].

Production of reactive oxygen intermediates

In man, there is no evidence for the local production of ROI in the tumour bed. The role of ROI in the antitumour response in human has been indirectly implicated by observation that myeloperoxidase-deficient individuals show an increased incidence of malignancy [106]. Patients with renal cancer show an increased production of ROI by peripheral blood monocytes [107]. *In vitro* stimulation of monocytes with tumour cells, but not with untransformed cells, induces the production of ROI [108]. O₂⁻, hydrogen peroxide, OH[•] and probably hypohalites are involved in the spontaneous cytotoxic activity of monocytes towards tumour cells [108]. On the other hand, the significant inhibition of ROI formation in *in vitro* co-cultures of macrophages and tumour spheroids of colon carcinoma cell lines or supernatants from cultures of tumour cells is observed [52]. TGF- β 1, IL-10 and IL-4 are not involved in this tumour-induced suppression of ROI production [52]. It is interesting that ROI may suppress lymphocyte and NK cell function [109]. CD18, CD29 and CD44 adhesion molecules are engaged in the induction of ROI production by monocytes stimulated with cancer cells [110]. Hyaluronan, the major ligand for CD44, which is overexpressed on many cancer cells, triggers ROI generation by monocytes via ligation of CD44 and allows them to distinguish cancer from non-malignant cells. On the other hand, blocking of CD44 on monocytes by free hyaluronan inhibits their response to tumour cells [110].

Production of eicosanoids

Cancer cells may affect monocytes function through alteration of arachidonic acid (AA) metabolism and production of eicosanoids: prostaglandins (PGs), thromboxane, leukotriens (LTs) and hydroxyeicosatetraenoic acid [111]. Serum level of PGE₂ is increased in cancer patients [112]. Stimulation by cancer cells induces monocyte PGE₂ production [4]. Also the level of PGs in tumour tissue is increased [113, 114]. PGs may promote tumour cell proliferation [115]. Blood monocytes and peritoneal macrophages from ovarian cancer patients show enhanced tumouricidal activity following inhibition of cyclooxygenase. However, this has no effect on tumouricidal activity of alveolar macrophages from lung cancer patients [116]. Metabolism of AA occurs preferentially via lipoxygenase pathway [117], which is not altered in circulating monocytes or TIM in cancer patients [116]. Antitumour activity of peritoneal macrophages is correlated with the production of PGE₂ and positively associated with synthesis of LTC₄ and LTD₄ [118].

Chemotactic response of monocytes to the tumour

Migration of blood monocytes to the tumour bed involves a response to a positive gradient of chemoattractants and induces the adherence to vascular

endothelium, emigration from the blood vessels and directed movement within the extracellular matrix (ECM). Tumour cells secrete chemotactic factors, such as IL-8 and monocyte chemoattractant protein-1, 2, 3 (MCP-1, 2, 3), the members of CC chemokine family: CCL-2, CCL-8, CCL-7. These molecules selectively attract monocytes, but not neutrophils [119]. MCP-1 gene expression within the tumour, predominantly in stromal cells, is correlated with the degree of invasiveness of breast carcinomas [120] and MCP-1 expression by both tumour cells and macrophages is positively associated with macrophages infiltration [121]. MCP-1 regulates the cell surface expression of adhesion molecules, especially β_2 integrins (CD11b/CD18 and CD11c/CD18) on monocytes, thus facilitating their adherence to vascular endothelium. MCP-1 also induces production of IL-1 and IL-6 by monocytes [122]. In ovarian cancer, MCP-1 serum level correlates with the histological grade of the tumour and the age of patients [123]. The production of MCP-1 by human tumours engrafted into mice enables early recruitment of monocytes and tumour growth inhibition [124].

Defective monocyte chemotaxis is observed in patients with head and neck, lung, gastric, breast and genitourinary cancers [125-129] and melanoma [130]. This defect is more apparent in advanced disease and reversed by tumour removal [125, 126, 130]. Chemokine receptor expression is an important factor for monocyte chemotaxis. TIM isolated from ovarian carcinoma and, to lesser extent, blood monocytes display defective mRNA and surface expression of chemokine receptor CCR2 (MCP-1 receptor). Downregulation of CCR2 is largely dependent on the local production of TNF [131]. However, monocyte chemotactic activity and the levels of CC chemokines are higher in non-small-cell lung cancer than in normal lung tissue [132]. Furthermore, blood monocytes from patients with breast cancer display a higher transendothelial migration than those from patients with benign diseases of the breast. It is not concerned with the differences in monocytes phenotype (HLA-DR, CD64, CD11a and CD11b expression) [133].

Cytotoxic activity of monocytes

Human monocytes possess high spontaneous cytotoxic activity against malignant, but not normal, cells [134-136]. ROI, TNF and NO may act synergistically in the cytotoxic damage of neoplastic cells. In most instances, an increased cytotoxic or cytostatic activity of monocytes in patients with different neoplasms, e.g. lung, breast, and gastrointestinal cancers [137], primary and metastatic brain tumours [138], squamous cell carcinoma [139] is observed. However, no changes in cytotoxic activity of monocytes are observed in patients with renal cancer [140] and non-Hodgkin's lymphoma [141]. A variety of agents are capable to induce tumouricidal activity of monocytes, e.g.: LPS, IL-1, IL-2, GM-CSF, laminin, LPS or IFN γ [142-148]. IFN γ prevents the loss of cytotoxic activity, which occurs during monocyte

maturation to macrophages [143]. Cytotoxic potential of monocytes is age dependent. Monocytes from aged healthy subjects show decreased *in vitro* cytotoxicity against tumour cells which is associated with compromised IL-1, ROI and RNI production [149].

Tumour cells are heterogeneous in their susceptibility to cytotoxic activity of monocytes. Human tumours of the same histological origin are affected to different degrees by monocytes [150] or their cytotoxic mediators [151]. The maximal tumouricidal activity usually requires a direct contact between monocytes and target cells [152]. It is suggested that outer membrane phosphatidyloserine [153] or hyaluronan [110] may be involved in the "recognition" of tumour cells by activated monocytes [153].

Phenotypic characteristics of monocytes in malignancy

Changes in monocyte function in malignant diseases are often correlated with changes in the expression of functionally important cell surface molecules.

Fc γ R1

The receptor for Fc part of IgG (Fc γ R1) of monocytes is involved in antibody-dependent cellular cytotoxicity (ADCC) against tumour cells [154-156]. Its expression is increased in patients with lung, colon [157], kidney [158] and gastric [159] cancers but decreased in patients with squamous cell carcinoma [139]. However, monocytes from metastatic squamous cell carcinoma show an increased expression of Fc γ R1 [160]. In patients with breast cancer the expression of Fc γ R1 on monocytes is unchanged [161].

MHC class II

MHC class II expression is critical for antigen presentation [161]. Furthermore, HLA-DR determinants of monocytes play a role in signal transduction for TNF gene activation [7] and NO production [99]. Monocytes from the *in vitro* co-culture with tumour cells show significant enhancement of HLA-DR expression [162]. Macrophages from the cellular infiltrate surrounding tumour express an abundant quantity of HLA class II determinants, which suggests that they are activated in the tumour bed [163]. There is also an opposite observation that cancer cells may induce a down-regulation of HLA-DR expression on monocytes [4]. In patients with squamous cell and breast carcinoma, HLA-DR expression on monocytes remains unchanged [139, 161].

Subpopulations of monocytes

On the basis of CD14 expression, two main monocyte subpopulations are distinguished. The major population that shows an enhanced expression of CD14 antigen (CD14⁺⁺ monocytes) and the minor one with a weak expression of

CD14 and the presence of CD16 (CD14⁺/CD16⁺ monocytes). In healthy donors CD14⁺/CD16⁺ subpopulation account for 5 +/- 3% of total monocytes. This subpopulation appears to represent more mature monocytes [164], is the main producer of TNF and show low or no production of IL-10 following stimulation with LPS. Therefore, CD14⁺/CD16⁺ subpopulation has been defined as "pro-inflammatory" monocytes [165, 166]. CD14⁺/CD16⁺ monocytes are also defined as the subpopulation containing dendritic cell precursors [167, 168]. The absolute number of CD14⁺/CD16⁺ monocytes is increased in various inflammatory diseases, like bacterial sepsis, viral infections, major trauma [166, 169, 170] and in patients with metastatic gastrointestinal cancers and other solid tumours [171]. We have also observed an elevated absolute number of these cells in the blood of gastric cancer patients (unpublished observations). However, no changes in the expression of CD16 on monocytes are found in patients with kidney cancer [158]. Patients with gastrointestinal carcinoma treated with M-CSF show a significant increase in the percentage of CD16⁺ monocytes in comparison to healthy subjects [172]. Our unpublished observations indicate that among all monocytes, CD14⁺/CD16⁺ cells possess the highest antitumour activity as they are the main producers of TNF, IL-12, NO and ROI (O₂⁻) and produce low levels of immunosuppressive IL-10.

Adhesion molecules

The CD11b (CR3 α chain) is an important receptor for phagocytosis and subsequent activation of respiratory burst in mononuclear phagocytes [173]. The interaction of CD11b with ICAM-1 promotes attachment to endothelium and extravasation of leukocytes [174]. CD11b molecule is lost upon their migration to the tissues [175]. Anti-CD11b monoclonal antibodies inhibit monocytes recruitment to both MCP-1 producing and nonproducing human tumours [176]. CD11b is involved in adhesion of MCP-1-stimulated monocytes to laminin of ECM. [177]. MCP-1 induces the expression of CD11b and CD11c and IL-1 and IL-6 production by monocytes [178]. The expression of CR3 on monocytes from kidney, but not breast, cancer patients is increased [179, 161]. Monocytes also express β 1 integrin - very late antigen-4 (VLA-4) and use this molecule in interactions with activated endothelial cells [180] or with tumour cells [181]. The decreased surface expression of VLA-4 on monocytes is observed during tumour growth, which suggests a reduced monocytes ability to bind ECM.

Immunoregulatory activity

The progression of malignant diseases is associated with immune dysfunction. Different monocyte populations may act as suppressor cells. The elevated suppressor activity of monocytes in cancer patients is related to tumour burden and the stage of disease. The presence of activated monocytes,

which also showed an increased suppressor activity for T cells, is associated with favourable prognosis in some patients with gastric cancer [182]. An increased monocyte-mediated cytostasis of lymphoid cell lines has been observed also in breast and lung cancer patients [183]. Monocytes of some patients with gastrointestinal cancer possess suppressor activity as well as increased cytostatic capacity against L1210 lymphoma cell line [184]. The question arises whether monocyte cytostatic and suppressor activities are interrelated and both indicate an activated state of the cell.

Tumours produce a number of factors (vascular endothelial growth factor, VEGF, M-CSF, IL-6) that block the differentiation of CD34⁺ stem cells into dendritic cells. However, tumours may promote the altered maturation and early apoptosis of human monocyte-derived dendritic cells. Upregulation of surface markers (CD80, CD86, HLA-DR), nuclear translocation of RelB and allostimulatory activity is associated with the lack of capacity to produce IL-12 and rapid apoptosis of monocytes [185]. Apoptosis of monocytes is also induced during their direct contact with cancer cells *in vitro* [162]. It may be one of the mechanisms by which tumours evade the immune response of the host. The inhibition of the cellular immune response of the host is also due to the gangliosides shedding by the tumour cells and its binding to leukocytes in the tumour microenvironment [186, 187].

Tumour infiltrating macrophages (TIM)

Function of TIM in the tumour growth

Macrophages represent a major component of the mononuclear cell infiltrate of tumours [163, 188, 189] and may consist up to 80% of the total tumour mass [190]. They are located within the tumour mass (intratumourally) or at the periphery of the tumour (peritumourally).

The process of leukocyte migration from the circulation into the tumour involves their interactions with vascular endothelium, i.e. leukocyte rolling and adhesion. This may be reduced in tumour microvessels due to decreased expression of adhesion molecules caused by tumour-derived angiogenic factors [191]. TIM have been demonstrated in the stroma of numerous malignant tumours including colon [192], breast [192, 193], skin [194], lung, ovary or thyroid gland cancers and melanoma [195]. The number of TIM is more increased in advanced (Duke's C) than in early colon cancer and in tumours expressing MHC class II molecules [196]. The composition of the cellular infiltrate depends on the properties of invaded tissue. The extent of TIM infiltration in tumours of the same histological origin varies, but the average number of TIM in particular tumour during its growth is relatively stable. Having divergent functional properties, TIM may modulate tumour growth by affecting cell proliferation, vascularization (angiogenesis), stroma formation, killing and dissolution of neoplastic cells.

The role of TIM in human malignancy is complex. Experimental evidence indicates that the intratumoural as well as peritumoural TIM limit tumour size in early stages of tumour development [174, 176]. In colon cancer TIM accumulate along the invasive edge, are in a direct contact with lymphocytes and express costimulatory molecules: CD80 (B7-1) as well as CD86 (B7-2). In contrast, the expression of CD80 and CD86 is usually inconspicuous in the tumour stroma [197]. These observations are in accordance with clinicopathological observations suggesting that peritumoural lymphocytic infiltration is a favourable prognostic factor in colorectal cancer [198]. The large number of CD16⁺ macrophages was found in renal cancer, melanoma and colon carcinoma [188]. As these tumours are susceptible to immunotherapy with lymphokine activated killer cells, it may indicate that CD16⁺ macrophages are involved in antitumour cytotoxic response [188]. In contrast, high TIM content within breast cancer stroma is associated with poor prognosis. This is due to a significant number of suppressor macrophages producing IL-10 that decreases the expression of MHC class II determinants and IL-2R on T cells [192]. However, no significant difference in the spontaneous and LPS-stimulated IL-10 production by alveolar macrophages is observed in patients with metastatic lung cancer [199]. The presence of iNOS was observed in TIM, especially intratumoural, infiltrating gastric [200] and breast cancers [201, 202]. The expression of iNOS in infiltrating cells positively correlates with the metastasis formation in breast cancer [202]. Although TIM from ovarian and colorectal cancers show the presence of TNF-mRNA [203, 204], the production of proinflammatory cytokines like IL-1 and IL-6 by TIM from ovarian cancer, upon stimulation with endotoxin, is decreased in comparison to monocytes from the same patients [205]. TIM from non-small-cell lung cancer exhibit decreased tumouricidal potential after stimulation with different stimuli in comparison to peripheral blood monocytes and normal alveolar macrophages [137].

Antitumour activity of TIM is considerably decreased in comparison to blood monocytes [206]. This indicates the suppressive role of tumour microenvironment on TIM *in situ*. In advanced stages of breast cancer, TIM may be ineffective or even promote tumour growth [189]. It is known that activated macrophages are able to produce several growth factors, including: TGF- α and - β , fibroblast growth factor (FGF), IL-1 and endothelial growth factor (EGF). There is a positive association between the degree of TIM infiltration and progression of breast tumours [207]. In breast cancer TIM are involved in stroma formation by transformation into fibroblast-like cells, which produce collagen type I [207]. Tumour may influence the activity of TIM by modulating the binding of TIM to ECM proteins. TIM can secrete proteases, which degrade the surrounding tissue and could facilitate tumour cell expansion and infiltration of the tissues. The activities of two families of

proteases: MMPs and urokinases are associated with tumour invasiveness and are important in angiogenesis. MMPs facilitate tumour invasion and metastasis through degradation of ECM compounds like collagens, laminins, proteoglycans and modulation of cell adhesion. MMPs may paradoxically stimulate the creation of biologic active proteins including chemotactic molecules derived from laminin-5 and angiotatin from plasminogen [208, 209, 210]. Colorectal cancer cells are able to stimulate monocytes production of MMP-2 and MMP-9, and this is dependent on metastatic potential of tumour cells. Also soluble products of metastatic colorectal cancer cells induce the expression of MMP-9 in monocytes [211]. The role of tissue inhibitors of MMPs (TIMPs) in cancer is complex. They are produced both by tumour cells and stromal fibroblasts. In experimental tumours TIMPs reduce tumour growth, metastasis and angiogenesis. On the other hand they may promote tumourigenesis and cancer progression through the influence on cell proliferation, apoptosis and MMP activity [212, 213].

Cell adhesion molecules are required for the cellular interactions and development of effective immune response. Contact between macrophages and cancer cells induces changes in the expression of adhesion molecules on both types of interacting cells. TIM from gastrointestinal cancers, especially localised along the invasive edge, show the expression of ICAM-1 (CD54) and lymphocytes from the invasive margin express LFA-1 (receptor of ICAM-1). In diffuse-type gastric cancer, majority of TIM are ICAM-negative [214]. ICAM-1 expression has also been observed on malignant cells including lymphomas [215], melanomas [216] and carcinomas [217-220]. Its expression may be upregulated by macrophage products: IFN- γ , TNF, IL-1 α , β , IL-6 and ROI [218, 219, 221, 222] and by interaction of hyaluronan (present on or shed from cancer cells) with CD44 [223]. Expression of ICAM-1 on melanoma cells is associated with their susceptibility to monocyte cytotoxicity [224]. LFA-1 is also present on cancer cells. The enhancement of this molecule on cancer cells and concomitant upregulation of ICAM-1 on monocytes occur after coculture of these cells [162].

The role of TIM in neoangiogenesis

In breast cancer a significant correlation exists between the vascular grade of the tumour, shortened patient survival and number of TIM [190]. Several soluble products of TIM are responsible for neoangiogenesis. These include EGF, VEGF, FGF, platelets derived growth factor (PDGF), GM-CSF, TGF- α and β , IL-1, IL-6 and PGs. PDGF expression in TIM from breast cancer positively correlates with the tumour size and microvessel count [225]. The role of VEGF in the tumour growth is not limited to promotion of angiogenesis as it also stimulates extravasation of plasma fibrinogen that leads to fibrin deposition and increase of ECM within the tumour. This in turn promotes the ingrowth of TIM, fibroblasts and endothelial cells [226]. The release

of VEGF is stimulated by hypoxia [227]. Thus, TIM from avascular and necrotic areas of breast cancer display an increased production of VEGF [228]. VEGF has also a crucial role in carcinoma-dependent ascites formation [229]. Not only TIM but also tumour cells are able to produce VEGF, which in turn acts as chemoattractant for macrophages [230]. Angiogenesis is also regulated by chemokines. ECR-CXC chemokines, including CXCL8 (IL-8), CXCL7 (NAP-2), CXCL5 (ENA-78) and CXCL1 (GRO α), are a potent angiogenic factors, whereas non-ELR-CXC chemokines such as CXCL10 (IP-10) and CXCL9 (Mig) are angiostatic [231, 232]. Also the member of CC chemokine family, CCL2 (MCP-1), is the angiogenic factor that acts by augmenting both TIM accumulation and angiogenesis [233].

Both tumour cells and monocytes release urokinase plasminogen activator (uPA). It promotes angiogenesis through plasminogen activation and degradation of stroma components and vessel walls (probably the first step of neoangiogenesis) [234]. Some ECM compounds after degradation by plasmin induce angiogenesis [235]. In breast cancer uPA is mainly localized in peripheral parts of the tumour [236]. A specific cell surface uPA receptor (uPAR) has been identified on human monocytes and a variety of cancer cells. In colon adenocarcinoma, uPAR is expressed by tumour cells and by TIM localized at the invasive edge [237], which may facilitate tumour invasion and metastasis [238].

The role of TIM in metastasis formation

There is evidence that TIM are involved in the metastatic process. The association between the presence of large numbers of TIM and lymph node metastases in human breast cancer has been observed [239]. Highly invasive and metastatic tumours can secrete glycoproteins that act as tumour-associated antigens evoking production of antibodies that promote tumour cells invasion and growth. It is due to activation of tumour infiltrating immune cells and proteases secretion followed by ECM degradation and angiogenesis [240].

Tumour invasion and metastasis formation begins from blood vessels basement membranes degradation due to release of matrix metalloproteinases by tumour cells and tumour cells activated monocytes/macrophages [241]. The next step is the formation of emboli in the microvasculature of different organs. PBMC form aggregates with renal cancer cells through the Siglec7 - the receptor for disialogangliosides, which is expressed by monocytes and NK cells. Metastatic potential of renal cell carcinoma is associated with the expression of gangliosides on tumour cells [242]. Coagulation associated with metastasis formation may also be the result of inappropriate expression of tissue factor in monocytes. Tissue factor (CD142), the main initiator of blood clotting, is produced by activated monocytes [243]. Tissue factor induces thrombin and in turn fibrin formation. Fibrin is known to stimulate the

migration of endothelial cells and thus potentiate angiogenesis [244]. Cancer patients have higher monocyte CD142 expression [245] that correlates with tumour progression [246]. uPA produced by macrophages and tumour cells, play also an important role in tissue invasion and metastases formation [247]. High uPA levels are correlated with high vessel density in tumour and higher vascular invasion of tumour cells. uPA and its receptor is localised mainly in the periphery of tumour [236, 237]. uPA content in peripheral parts of the tumour is increased in patients with metastases [236]. It is connected with the proteolytic activity of plasmin on ECM, basement membrane and vessel walls [234, 248]. Cathepsins (lysosomal proteinases) are also involved in metastasis formation. Their expression have been found in TIM from bladder tumours [249] and breast cancers [250].

Some studies suggested the fusion of monocytes with certain haematopoietic tumour cells as an important mechanism of metastases formation [251]. Highly metastatic variant of T cell lymphoma cell line is derived *in vitro* from the spontaneous fusion of the lymphoma cells with the host macrophages [252]. However, the mechanisms of fusion *in vivo* remain unknown. TIM also participate in the osteolysis associated with bone metastases. They are the major cellular component of the inflammatory infiltrates in the bones and can release local mediators that stimulate osteoclast activity. Moreover, they can also resorb the bone on their own and differentiate into osteoclast-like cells [253]. Although no data on the role of TIM in the peritoneal dissemination of human malignancy are available, in the murine model milky spots, which are aggregates of macrophages, are considered to be the locus for early peritoneal metastases formation [254, 255].

Monocytes in cancer immunotherapy

In local and systemic adoptive immunotherapy, the autologous effector cells harvested from the blood are activated *in vitro* and reinfused into the host [256-258]. Macrophages can be activated for tumour cell killing by some immunopotentiators [259]. Antitumour activity of monocytes may be enhanced by their incubation in the presence of IFN- γ , LPS, GM-CSF and (OH)₂ VitD3 [147, 260-262]. In 1987, Stevenson et al. reported the first clinical trial with adoptive transfer of activated macrophages. They used IFN γ -activated macrophages for intraperitoneal infusions in patients with colorectal cancer [263]. Adoptive transfer of macrophages has undergone phase I clinical trials for patients with metastatic cancer (colon, ovarian, lung, renal, pancreatic cancer and melanoma) infused systemically or intraperitoneally [261, 264-266]. Monocytes differentiated in the presence of IFN γ were used for phase II adoptive therapy of advanced colorectal cancer. Commonly, continuous flow centrifugal leukapheresis and counterflow centrifugal elutriation are used for monocyte

isolation. However, the clinical results of adoptive immunotherapy are still controversial [264]. No significant partial or complete responses have been reported, though prolonged disease free intervals were observed [261, 266].

Monocyte cytotoxic activity can be enhanced by chemotherapeutic drugs, e.g. cisplatin [268], IFN γ and murapeptides [269]. There is some evidence that activated monocytes are cytotoxic to drug resistant tumour cells [270, 271]. They can also carry cytotoxic drugs and immunomodulators [272, 273]. Immunostimulation with IFN (may reverse monocyte deactivation in patients with chemotherapy-induced neutropenia and the serious infections and cause clinical improvement and increase the level of CD14+ DR+ circulating monocytes [274]. The novel approach is to utilize the ability of macrophages to migrate into hypoxic areas of the tumour and use them for delivering gene therapy [275]. However, currently adoptive therapy with the use of monocytes/macrophages is still at infancy.

Abbreviations

AA – arachidonic acid
 CCL – CC chemokine family
 CSF – colony stimulating factor
 CCR – CC receptors
 CR – chemokine receptor
 ECM – extracellular matrix
 EGF – endothelial growth factor
 FasL – Fas-ligand
 FcR – the receptor of Fc part of immunoglobulin
 FGF – fibroblast growth factor
 GM-CSF - ggranulocyte-macrophage colony stimulating factor
 ICAM-1 – intercellular adhesion molecule-1
 IFN – interferon
 IL – interleukin
 iNOS – inducible nitric oxide synthase
 LFA-3 - leucocyte functional antigen-3
 LPS – lipopolysaccharide
 LT – leukotrien
 MCP – monocyte chemotactic protein
 M-CSF – macrophage colony stimulating factor
 MMPs – matrix metalloproteinases
 NO – nitric oxide
 PAI – inhibitor of plasminogen activation
 PBMC – peripheral blood mononuclear cells
 PDGF – plated derived growth factor
 PG – prostaglandin
 PKA – camp-dependent kinase
 PKC – calcium-phospholipid-dependent protein kinase
 PTK – tyrosine protein kinase
 RNI– reactive nitrogen intermediates
 ROI – reactive oxygen intermediates
 TGF – transforming growth factor
 TIM – tumour infiltrating macrophages
 TIMP – tissue inhibitor of matrix metalloproteinases
 TNF – tumour necrosis factor

TX – tromboxane
 uPA – urokinase plasminogen activator
 VCAM-1 - vascular adhesion molecule-1
 VEGF – vascular endothelial growth factor
 VLA-4 - very late antigen-4

References

- Mantovani A, Bottazzi B, Colotta F, Sozzani S, Ruco L (1992): The origin and function of tumor-associated macrophages. *Immunol Today* 13: 265-270.
- Grace H, Wong W, Goeddel D: Tumour necrosis factor. In: Human monocytes. Ed. Zembala M, Asherson G: Academic Press, London 1989.
- Krakauer T, Oppenheim J (1993): IL-1 and tumor necrosis factor- α each up-regulate both the expression of IFN- γ receptors and enhance IFN- γ -induced HLA-DR expression on human monocytes and a human monocytic cell line (THP-1). *J Immunol* 150: 1205-1211.
- Ishii N, Chiba M, Iizuka M, Horie Y, Masamune O (1994): Induction of HLA-DR antigen on human colonic epithelium by tumor necrosis factor- α and interferon- γ . *Scan J Gastroenterol* 29: 903-907.
- Menetrier-Caux C, Bain C, Favrot M, Duc A, Blay J (1999): Renal cell carcinoma induces interleukin 10 and prostaglandin E2 production by monocytes. *Br J Cancer* 79: 119-130.
- Pfizenmaier K, Scheurich P, Schluter C, Kronke M (1987): Tumor necrosis factor enhances HLA-A,B,C and HLA-DR gene expression in human tumor cells. *J Immunol* 138: 975-980.
- Watanabe Y, Jacob C (1991): Regulation of MHC class II antigen expression. Opposing effects of tumor necrosis factor- α on IFN- γ -induced HLA-DR and Ia expression depends on the maturation and differentiation stage of the cell. *J Immunol* 146: 899-905.
- Zembala M, Siedlar M, Ruggiero I, et al. (1994): The MHC class-II and CD44 molecules are involved in the induction of tumour necrosis factor (TNF) gene expression by human monocytes stimulated with tumour cells. *Int J Cancer* 56: 269-274.
- Janicke R, Mannel D (1990): Distinct tumor cell membrane constituents activate human monocytes for tumor necrosis factor synthesis. *J Immunol* 144: 1144-1150.
- Mytar B, Woloszyn M, Baj-Krzyworzeka M, et al.: Tumour cell-induced deactivation of human monocytes. (submitted).
- Siedlar M, Mytar B, Hyszko M, Zembala M (1999): Involvement of protein kinases in signalling for nitric oxide (NO) and tumour necrosis factor alpha (TNF) production by monocytes stimulated with colorectal DeTa cancer cells: the lack of evidence for the role of TNF in the regulation of NO production. *Immunol Letters* 65: 189-195.
- Cohen M, Bereta M, Bereta J (1997): Effect of cytokines on tumour cell-endothelial interactions. *Indian J Biochem Biophys* 34: 199-204.
- Vassali P (1992): The pathophysiology of tumor necrosis factors. *Annu Rev Immunol* 10: 411-452.
- Lejeune F, Ruegg C, Lienard D (1998): Clinical applications of TNF- α in cancer. *Curr Opin Immunol* 10: 573-580.
- Boldrini L, Calciani A, Samaritani E, et al. (2000): Tumour necrosis factor- α and transforming growth factor- β are significantly associated with better prognosis in non-small cell lung carcinoma: putative relation with BCL-2-mediated neovascularization. *British J Cancer* 83: 480-486.
- Shau H (1988): Characteristic and mechanisms of neutrophil-mediated cytostasis induced by tumor necrosis factor. *J Immunol* 141: 234-240.

16. Rao V, Singh R, Delimont D, et al. (1999): Transcriptional regulation of MMP-9 expression in stromal cells of human giant cell tumor of bone by tumor necrosis factor- α . *Int J Oncol* 14: 291-300.
17. Leber T, Balkwill F (1998): Regulation of monocyte MMP-9 production by TNF- α and a tumour-derived soluble factor (MMPSF). *Br J Cancer* 78: 724-732.
18. Macura-Biegun A (1998): The role of extracellular matrix in tumor progression. *Central-European J Immunol* 23: 8-14.
19. Rice G, Gimbrone M, Bevilacqua M (1988): Tumor cell-endothelial interactions: increased adhesion of human melanoma cells to activated vascular endothelium. *Am J Pathol* 133: 204-206.
20. Bereta M, Bereta J, Cohen S, Zaifert K, Cohen M (1991): Effect of inflammatory cytokines on the adherence of tumor cells to endothelium in a murine model. *Cell Immunol* 136: 263-267.
21. Mackay F, Loetscher H, Stueber D, Gehr G, Lesslauer W (1993): Tumor necrosis factor α (TNF- α)-induced cell adhesion to human endothelial cells is under dominant control of one TNF receptor type, TNF-R55. *J Exp Med* 177: 1277-86
22. Mackay F, Loetscher H, Stueber D, Gehr G, Lesslauer W (1993): Tumor necrosis factor α (TNF- α)-induced cell adhesion to human endothelial cells is under dominant control of one TNF receptor type, TNF-R55. *J Exp Med* 177:1277-1286.
23. Heriot A, Marriott J, Cookson S, Kumar D, Dalglish A (2000): Reduction in cytokine production in colorectal cancer patients: association with stage and reversal resection. *Br J Cancer* 82: 1009-1012.
24. Zembala M, Mytar B, Wołoszyn M, et al. (1988): Monocyte TNF production in gastrointestinal cancer [letter]. *Lancet* 26: 1262.
25. Nakazaki H (1992): Preoperative and postoperative cytokines in patients with cancer. *Cancer* 70: 709-713.
26. Zhang G, Adachi I (1999): Serum interleukin-6 levels correlate to tumour progression and prognosis in metastatic breast carcinoma. *Anticancer Res* 19: 1427-1232.
27. Socher S, Martinez D, Craig J, Kuhn J, Oliff A (1988): Tumor necrosis factor not detectable in patients with clinical cancer cachexia. *J Natl Cancer Inst* 80: 595-598.
28. Maltoni M, Fabbri L, Nanni O, et al. (1997): Serum levels of tumour necrosis factor α and other cytokines do not correlate with weight loss and anorexia in cancer patients. *Support Care Cancer* 5: 130-135
29. Blakwill F, Mantovani A (2001): Inflammation and cancer: back to Virchow? *Lancet* 357: 539-542.
30. Sodhi A, Singh R, Singh S (1992) Effect of interferon- γ priming on the activation of murine peritoneal macrophages to tumouricidal state by cisplatin, IL-1 and tumour necrosis factor (TNF): production of IL-1 and TNF. *Clin Exp Immunol* 88: 350-355.
31. Roberts A, O'Connell S, Ebert E (1993) tinal intraepithelial lymphocytes bind to colon cancer cells by HML-1 and CD11a. *Cancer Res* 53: 1608-1611.
32. Le J, Vilcek J(1987): Biology of disease. Tumor necrosis factor and interleukin 1: cytokines with multiple overlapping biological activities. *Lab Invest* 56: 234-248.
33. Onozaki K, Matsushima K, AggrawalB, Oppenheim J (1985): Human interleukin 1 is a cytotoxic factor for several tumor cell lines. *J Immunol*135: 3962-3968.
34. Ichinose Y, Tsao J, Fidler I (1988): Destruction of tumor cells by monokines released from activated human blood monocytes: evidence for parallel and additive effects of IL-1 and TNF. *Cancer Immunol Immunother* 27: 7-12.
35. Raitano B, Korc M (1993): Growth inhibition of a human colorectal carcinoma cell line by interleukin 1 is associated with enhanced expression of gamma-interferon receptors. *Cancer Res* 53:636-640.
36. Chasseing N, Trejo Y, Bordenave H, Zanoni L, Rumi L (1997): Intracytoplasmatic and extracellular interleukin-1 production by monocytes of lung and colorectal cancer patients. *Acta Physiol Pharmacol Ther Latinoam* 47: 147-156.
37. Sone S, Utsugi T, Tandon P, Yanagawa H, Okubo A, Ogura T (1990): Tumor cytotoxicity and interleukin 1 production of blood monocytes of lung cancer patients. *Cancer Immunol Immunother* 30: 357-362.
38. Aggarwal B, Kohr W, Hass P, et al. (1985): Human tumor necrosis factor. Production, purification, and characterization. *J Biol Chem* 260: 2345-2354.
39. Kutteh W, Kutteh C (1992): Quantitation of tumor necrosis factor- α , interleukin-1 β and interleukin-6 in the effusions of ovarian epithelial neoplasms. *Am J Obstetrics Gynecology* 167: 1864-1869.
40. Kabir S, Daar A (1995): Serum levels of interleukin-1, interleukin-6 and tumour necrosis factor- α in patients with gastric carcinoma. *Cancer Lett* 95: 207-212
41. Chopra V, Dinh T, Hannigan E. (1997): Serum levels of interleukins, growth factors and angiogenin in patients with endometrial cancer. *J Cancer Res Clin Oncol* 123: 167-172.
42. Stefanovic V, Bogicevic M, Stamenic T, et al. (1994): Cytokine levels in patients with urinary tract cancer. *Pathol Biol (Paris)* 42: 842-846.
43. Matsuda T, Hirano T: IL-6. In: *Cytokine References*, vol. 1, Academic Press. London. 2001.
44. Hirano T (1991): Interleukin 6 (IL-6) and its receptor: their role in plasma cell neoplasias. *Int J Cell Cloning*, 9: 166-184.
45. Kawano M, Hirano T, Matsuda T, et al. (1988): Autocrine generation and requirement of BSF-2/IL-6 for human multiple myelomas. *Nature* 332: 83-85.
46. Bataille R, Jourdan M, Zhang X, Klein B (1989): Serum levels of interleukin 6, a potent myeloma cell growth factor, as a reflect of disease severity in plasma cell dyscrasias. *J Clin Invest* 84: 2008-2011.
47. Barber M, Fearon K, Ross J (1999): Relationship of serum levels of interleukin-6, soluble interleukin-6 receptor and tumour necrosis factor receptors to the acute-phase protein response in advanced pancreatic cancer. *Clin Sci (Lond)* 96: 83-87.
48. Okada S, Okusaka T, Ishii H, et al. (1998): Elevated serum interleukin-6 levels in patients with pancreatic cancer. *Jpn J Clin Oncol* 28: 12-15.
49. Chau G, Wu C, Lui W, et al. (2000): Serum interleukin-10 but not interleukin-6 is related to clinical outcome in patients with resectable hepatocellular carcinoma. *Ann Surg* 231: 552-558.
50. Martin F, Santolaria F, Batista N, et al. (1999): Cytokine levels (IL-6 and IFN- γ), acute phase response and nutritional status as prognostic factors in lung cancer. *Cytokine* 11: 80-86.
51. Yano S, Sone S, Nishioka Y, et al. (1995): Differential effect of anti-inflammatory cytokines (IL-4, IL-10 and IL-13) on tumoricidal and chemotactic properties of human monocytes induced by monocyte chemotactic and activating factor. *J Leukoc Biol* 57: 303-309.
52. Siegert A, Denkert C, Leclere A, Hauptmann S (1999): Suppression of the reactive oxygen intermediates production of human macrophages by colorectal adenocarcinoma cell lines. *Immunology* 98: 551-556.
53. Huang S, Xie K, Bucana C, Ullrich S, Bar-Eli M (1996): Interleukin 10 suppresses tumor growth and metastasis of human melanoma cells: potential inhibition of angiogenesis. *Clin Cancer Res* 2: 1969-1979.
54. DeVita F, Orditura M, Galizia G, et al. (2000): Serum interleukin-10 is an independent prognostic factor in advanced solid tumors. *Oncol Rep* 7: 357-361.

55. Merendino R, Gangemi S, Misefari A, et al. (1999): Interleukin-12 and interleukin-10 production by mononuclear phagocytic cells from breast cancer patients. *Immunol Lett* 68: 355-358.
56. Kucharzik T, Luger N, Winde G, Domschke W, Stoll R (1997): Colon carcinoma cell lines stimulate monocytes and lamina propria mononuclear cells to produce IL-10. *Clin Exp Immunol* 110: 296-302.
57. Klimberg V, Kornbluth J, Cao Y, et al. (1996): Glutamine suppresses PGE2 synthesis and breast cancer growth. *J Surg Res* 63: 293-297.
58. Godwin J, Ceuppens J (1983): Regulation of the immune response by prostaglandins. *J Clin Immunol* 3: 295-315.
59. Shmitt E, Hoehn P, Huels C, et al. (1994): T helper type I development of naive CD4+ T cells requires the co-ordinate action of IL-12 and IFN- γ and is inhibited by TGF- β . *Eur J Immunol* 24: 793-798.
60. Germann T, Rude E (1995): Interleukin 12. *Int Arch Allergy Immunol* 108: 103-112.
61. Esche C, Shurin M, Lotze M: IL-12. In: *Cytokine Reference*, vol. 1. Academic Press. London. 2001.
62. Trinchieri G, Scott P (1994): The role of interleukin 12 in the immune response, disease and therapy. *Immunology Today* 15: 460-463.
63. Di Carlo E, Comes A, Basso S, et al. (1995): The combined action of IL-15 and IL-12 gene transfer can induce tumor cell rejection without T and NK cell involvement. *J Immunol* 165: 3111-3118.
64. Ha S, Lee C, Lee S, et al. (1999): A novel function of IL-12p40 as a chemotactic molecule for macrophages. *J Immunol* 163: 2902-2908.
65. Ha S, Lee S, Kim C, et al. (1998): Rapid recruitment of macrophages in interleukin-12-mediated tumour regression. *Immunology* 95: 156-163.
66. Haku T, Yanagawa H, Nabioullin R, et al. (1997): Interleukin-12-mediated killer activity in lung cancer patients. *Cytokine* 9: 846-852.
67. O'Hara R, Greenman J, MacDonald W, et al. (1998): Advanced colorectal cancer is associated with impaired interleukin 12 and enhanced interleukin 10 production. *Clin Cancer Res* 4: 1943-1948.
68. Kovacs E (2000): Serum level of IL-12 and the production of IFN- γ , IL-2, IL-4 by peripheral blood mononuclear cells (PBMC) in cancer patients treated with *Viscum album* extract. *Biomed Pharmacother* 54: 305-310.
69. Zeimet A, Widschwendter M, Knabbe C, et al. (1998): Ascitic interleukin-12 is an independent prognostic factor in ovarian cancer. *J Clin Oncol* 16: 1861-1868.
70. Heimdal J, Aarstad H, Olofsson J (2000): Peripheral blood T-lymphocyte and monocyte function and survival in patients with head and neck carcinoma. *Laryngoscope* 110: 402-407.
71. Fujiwara H, Hamaoka T (1996): Antitumor and antimetastatic effects of interleukin 12. *Cancer Chemother Pharmacol* 38, Suppl: 22-26.
72. Brunda M, Luistro L, Rumennik L, et al. (1996): Antitumor activity of interleukin 12 in preclinical models. *Cancer Chemother Pharmacol* 38, Suppl: 16-21.
73. Mu J, Zou J, Yamamoto N, et al. (1995): Administration of recombinant interleukin 12 prevents outgrowth of tumor cells metastasizing spontaneously to lung and lymph nodes. *Cancer Res* 55: 4404-4408.
74. Caminschi I, Venetsanakos E, Leong C, et al. (1998): Interleukin-12 induces an effective antitumor response in malignant mesothelioma. *Am J Respir Cell Mol Biol* 19: 738-746.
75. Mazzolini G, Qian C, Xie X, et al. (1999): Regression of colon cancer and induction of antitumor immunity by intratumoral injection of adenovirus expressing interleukin-12. *Cancer Gene Ther* 6: 514-522.
76. Gollob J, Mier J, Veenestra K, et al. (2000): Phase I trial of twice-weekly intravenous interleukin 12 in patients with metastatic renal cell cancer or malignant melanoma: ability to maintain IFN- γ induction is associated with clinical response. *Clin Cancer Res* 6: 1678-1692.
77. Lebel-Binay S, Berger A, Zinzindohoue F, et al. (2000): Interleukin-18: biological properties and clinical implications. *Eur Cytokine Netw* 11: 15-26.
78. Hashimoto W, Osaki T, Okamura H, et al. (1999): Differential antitumor effects of administration of recombinant IL-18 or recombinant IL-12 are mediated primarily by Fas-Fas ligand- and perforin-induced tumor apoptosis, respectively. *J Immunol* 163: 583-589.
79. Coughlin C, Salhany K, Wysocka M, et al (1998): W: Interleukin-12 and interleukin-18 synergistically induce murine tumor regression which involves inhibition of angiogenesis. *J Clin Invest* 101: 1441-1449.
80. Pages F, Berger A, Henglein B, et al. (1999): Modulation of interleukin-18 expression in human colon carcinoma: consequences for tumor immune surveillance. *Int J Cancer* 84: 326-330.
81. Wiltshcke C, Krainer M, Wagner A, et al. (1995): Influence of in vivo administration of GM-CSF and G-CSF on monocyte cytotoxicity. *Exp Hematol* 23: 402-406.
82. Ohsaka A, Saionji K, Kuwaki T, et al. (1995): Granulocyte colony-stimulating factor administration modulates the surface expression of effector cell molecules on human monocytes. *Br J Haematol* 89: 465-472. Comment in: *Br J Haematol* 91: 513-515.
83. Kaciński B (1995): CSF-1 and its receptor in ovarian, endometrial and breast cancer. *Ann Med* 27: 79-85.
84. Chambers S, Kaciński B, Ivins C, Carcangiu M (1997): Overexpression of epithelial macrophage colony-stimulating factor (CSF-1) and CSF-1 receptor: a poor prognostic factor in epithelial ovarian cancer, contrasted with a protective effect of stromal CSF-1. *Clin Cancer Res* 3: 999-1007.
85. Tang R, Beuvon F, Ojeda M, et al. (1992): M-CSF (monocyte colony stimulating factor) and M-CSF receptor expression by breast tumour cells: M-CSF mediated recruitment of tumour infiltrating monocytes? *J Cell Biochem* 50: 350-356.
86. Scholl S, Pallud C, Beuvon F, et al. (1994): Anti-colony-stimulating factor-1 antibody staining in primary breast adenocarcinomas correlates with marked inflammatory cell infiltrates and prognosis. *J Natl Cancer Inst* 86: 120-126.
87. Shinohara H, Yano S, Bucana C, Fidler I (2000): Induction of chemokine secretion and enhancement of contact-dependent macrophage cytotoxicity by engineered expression of granulocyte-macrophage colony-stimulating factor in human colon cancer cells. *J Immunol* 164: 2728-2737.
88. Stuehr D, Nathan C (1989): Nitric oxide. A macrophage product responsible for cytostasis and respiratory inhibition in tumor target cells. *J Exp Med* 169: 1543-1555.
89. Hibbs J, Taintor R, Vavrin Z, Rachlin E (1988): Nitric oxide: a cytotoxic activated macrophage effector molecule. *Biochem Biophys Res Commun* 157: 87-94.
90. Patel R, McAndrews J, Sellak H, et al. (1999): Biological aspects of reactive nitrogen species. *Biochim Biophys Acta* 1411: 385-400.
91. Brown G (2001): Regulation of mitochondrial respiration by nitric oxide inhibition of cytochrome c oxidase. *Biochem Biophys Acta* 1504: 46-57
92. Kurzawa H, Wysocka M, Aruga E, et al. (1998): Interleukin 12 enhances cellular immune responses to vaccination only after a period of suppression. *Cancer Res* 58: 491-499.

93. Koblish H, Hunter C, Wysocka M, et al. (1998): Immune suppression by recombinant interleukin (rIL)-12 involves interferon γ induction of nitric oxide synthase 2 (iNOS) activity: inhibitors of NO generation reveal the extent of sIL-12 vaccine adjuvant effect. *J Exp Med* 188: 1603-1610.
94. Allione A, Bernabei P, Bosticardo M, et al. (1999): Nitric oxide suppress human T lymphocyte proliferation through IFN- γ -dependent and IFN- γ -independent induction of apoptosis. *J Immunol* 163: 4182-4191.
95. Zembala M, Siedlar M, Marcinkiewicz J, Pryjma J (1994): Human monocytes are stimulated for nitric oxide release *in vitro* by some tumor cells but not by cytokines and lipopolisaccharide. *Eur J Immunol* 24: 435-439.
96. Albina J (1995): On the expression of nitric oxide synthase by human macrophages. Why no NO? *J Leukoc Biol* 58: 643-649.
97. Schneemann M, Schoedon G, Hofer S, et al. (1993): Nitric oxide synthase is not constituent of the antimicrobial armature of human mononuclear phagocytes. *J Infect Dis* 167: 1358-1363.
98. Denis M (1994): Human monocytes/macrophages: NO or no NO? *J Leukoc Biol* 55: 682-684.
99. Siedlar M, Marcinkiewicz J, Zembala M (1995): MHC class I and class II determinants and some adhesion molecules are engaged in the regulation of nitric oxide production *in vitro* by human monocytes stimulated with colon carcinoma cells. *Clin Immunol Immunopathol* 77: 380-384.
100. Mills C, Shearer J, Evans R, Caldwell M (1992): Macrophage arginine metabolism and the inhibition or stimulation of cancer. *J Immunol* 149: 2709-2714.
101. Takigawa M, Enomoto M, Nishide Y, et al. (1990): Tumor angiogenesis and polyamines: α -difluoromethylornithine, an irreversible inhibitor of ornithine decarboxylase, inhibits B16 melanoma-induced angiogenesis *in ovo* and the proliferation of vascular endothelial cells *in vitro*. *Cancer Res* 50: 4131-4138.
102. Kim P, Zamora R, Petrosko P, Billiar T (2001): The regulatory role of nitric oxide in apoptosis. *Int Immunopharmacol* 1: 1421-1441.
103. Torok N, Higuchi H, Bronk S, Gores G (2002): Nitric oxide inhibits apoptosis downstream of cytochrome C release by nitrosylating caspase 9. *Cancer Res* 62: 1648-1653.
104. Kong G, Kim E, Kim W, et al. (2001): Inducible nitric oxide synthase (iNOS) immunoreactivity and its relationship to cell proliferation, apoptosis, angiogenesis, clinicopathologic characteristic, and patient survival in pancreatic cancer. *Int J Pancreatol* 29: 133-140.
105. Chen G, Lee J, Chan U, et al. (2002): Increased apoptosis in infiltrating mononuclear cells of colorectal cancer. *Arch Pathol Lab Med* 126: 686-691.
106. Lanza F, Fietta A, Spisani S, et al. (1987): Does a relationship exist between neutrophil myeloperoxidase deficiency and the occurrence of neoplasms? *J Clin Lab Immunol* 22: 175-180.
107. Trulsson A, Nilsson S, Brekkan E, Venge P (1994): Patients with renal cancer have a larger proportion of high-density blood monocytes with increased lucigenin-enhanced chemiluminescence. *Inflammation* 18: 99-105.
108. Mytar B, Siedlar M, Wołoszyn M, et al. (1999): Induction of reactive oxygen intermediates in human monocytes by tumour cells and their role in spontaneous monocyte cytotoxicity. *Br J Cancer* 79: 737-743.
109. Kono K, Salazar-Onfray F, Petersson M, et al. (1996): Hydrogen peroxide secreted by tumor-derived macrophages downmodulates signal-transducing zeta molecules and inhibits tumor-specific T-cell- and natural killer cell-mediated cytotoxicity. *Eur J Immunol* 26: 1308-1313.
110. Mytar B, Siedlar M, Wołoszyn M, Colizzi V, Zembala M (2001): Cross-talk between human monocytes and cancer cells during reactive oxygen intermediates generation: the essential role of hyaluronan. *Int J Cancer* 94: 727-732.
111. Kurland J, Bockman R, Broxmeyer H, Moore M (1978): Limitation of excessive myelopoiesis by the intrinsic modulation of macrophage-derived prostaglandin E. *Science* 199: 552-555.
112. Nara K, Odagiri H, Fujii M, et al. (1987): Increased production of tumor necrosis factor and prostaglandin E2 by monocytes in cancer patients and its unique modulation by their plasma. *Cancer Immunol Immunother* 25: 126-132.
113. Bennet A, Carroll M, Stamford I, et al. (1992): Prostaglandins and human lung carcinomas. *Br J Cancer* 46: 888-893.
114. Karmali R, Wustrow T, Thaler H, Strong E (1984): Prostaglandins in carcinomas of the head and neck. *Cancer Lett* 22: 333-336.
115. Yang C, Meng C (1994): Regulation of PG synthase by EGF and PDGF in human oral, breast, stomach and fibrosarcoma cancer cell lines. *J Dent Res* 73: 1407-1415.
116. Braun D, Ahn M, Harris J, et al. (1993): Sensitivity of tumoricidal function of macrophages from different anatomical sites of cancer patients to modulation of arachidonic acid metabolism. *Cancer Res* 53: 3362-3368.
117. Tripp C, Needleman P (1987): Regulation of macrophage arachidonic acid metabolism during the immune response. *Adv Prostaglandin Thromboxane Leukot Res* 17B: 1085-1090.
118. Ben-Efraim S, Bonta I (1994): Modulation of antitumour activity of macrophages by regulation of eicosanoids and cytokine production. *Int J Immunopharmacol* 16: 397-399.
119. Van Damme J, Proost P, Lenaerts J, Opdenakker G (1992): Structural and functional identification of two human, tumor-derived monocyte chemotactic proteins (MCP-2 and MCP-3) belonging to the chemokine family. *J Exp Med* 176: 59-65.
120. Valkovic T, Lucin K, Krstulja M, et al. (1998): Expression of monocyte chemotactic protein-1 in human invasive ductal breast cancer. *Pathol Res Pract* 194: 335-340.
121. Ueno T, Masakazu T, Saji H, et al. (2000): Significance of macrophage chemo-attractant protein-1 on macrophage recruitment, angiogenesis and survival in human breast cancer. *Clin Cancer Res* 6: 3282-3289.
122. Jiang J, Beller D, Frenzl G, Graves D (1992): Monocyte chemoattractant protein-1 regulates adhesion molecule expression and cytokine production in human monocytes. *J Immunol* 148: 2423-2428.
123. Hefler L, Tempfer C, Heinze G, et al. (1999): Monocyte chemoattractant protein-1 serum levels in ovarian cancer patients. *Br J Cancer* 81: 855-859.
124. Zhang L, Khayat A, Cheng H, Graves D (1997): The pattern of monocyte recruitment in tumors is modulated by MCP-1 expression and influences the rate of tumor growth. *Lab Invest* 76: 579-590.
125. Tan I, Drexhage H, Scheper R, et al. (1986): Defective monocyte chemotaxis in patients with head and neck cancer. Restoration after treatment. *Arch Otolaryngol Head Neck Surg* 112: 541-544.
126. Nielsen H, Bennedson J, Dombernowsky P (1982): Normalization of defective monocyte chemotaxis during chemotherapy in patients with small cell anaplastic carcinoma of the lung. *Cancer Immunol Immunother* 14: 13-15.
127. Yamane T, Sakita M, Kasuga M, et al. (1981): Monocyte count, monocyte chemotaxis and chemotactic factor inactivator in gastric cancer patient. *Jpn J Surg* 11: 422-427.
128. Snyderman R, Meadows L, Holder W, Wells S (1978): Abnormal monocyte chemotaxis in patients with breast

- cancer: evidence for a tumor-mediated effect. *J Natl Cancer Inst* 60: 737-740.
129. Hausman M, Brosman S, Snyderman R, et al. (1975): Defective monocyte function in patients with genitourinary carcinoma. *J Natl Cancer Inst* 55: 1047-1054.
 130. Snyderman R, Seigler H, Meadows L (1977): Abnormalities of monocyte chemotaxis in patients with melanoma: effects of immunotherapy and tumor removal. *J Natl Cancer Inst*, 58: 37-41.
 131. Sica A, Saccani A, Bottazzi B, et al. (2000): Defective expression of the monocyte chemotactic protein-1 receptor CCR2 in macrophages associated with human ovarian carcinoma. *J Immunol* 164: 733-738.
 132. Arenberg D, Keane M, DiGiovine B, et al. (2000): Macrophage infiltration in human non-small-cell lung cancer: the role of CC chemokines. *Cancer Immunol Immunother* 49: 63-70.
 133. Gebhard B, Gnant M, Schutz G, et al. (2000): Different transendothelial migration behaviour pattern of blood monocytes derived from patients with benign and malignant diseases of the breast. *Anticancer Res* 20: 4599-4604.
 134. Mantovani A, Jerrells T, Dean J, Herberman R (1979): Cytolytic and cytostatic activity on tumor cells of circulating human monocytes. *Int J Cancer* 23: 18-27.
 135. Davies B, Edwards S (1992): Interactions between human monocytes and tumour cells. *Br J Cancer* 66: 463-469.
 136. Gerrard T, Terz J, Kaplan A (1980): Cytotoxicity to tumour cells of monocytes from normal individuals and cancer patients. *Int J Cancer* 26: 585-593.
 137. Siziopikou K, Harris J, Casey L, et al. (1991): Impaired tumoricidal function of alveolar macrophages from patients with non-small cell lung cancer. *Cancer* 68: 1035-1044.
 138. Barna B, Rogers L, Thomassen M, et al. (1991): Monocyte tumoricidal activity and tumor necrosis factor production in patients with malignant brain tumors. *Cancer Immunol Immunother* 33: 314-318.
 139. Murali P, Somasundaram R, Chiplunkar S, et al. (1989): Monocyte/macrophage functions in patients with squamous cell carcinoma of the oral cavity. *J Oral Pathol Med* 18: 539-543.
 140. Galligioni E, Quaia M, Spada A, et al. (1993): Activation of cytolytic activity in peripheral blood monocytes of renal cancer patients against non-cultured autologous tumor cells. *Int J Cancer* 55: 380-385.
 141. Jhaver K, De A, Advani S, Nadkarni J (1991): Production of interleukin-1 and tumour necrosis factor in non-Hodgkin' lymphoma patients. *Cancer Immunol Immunother* 34: 123-127.
 142. Mace K, Ehrke M, Hori K, et al. (1988): Role of tumor necrosis factor in macrophage activation and tumoricidal activity. *Cancer Res* 48: 5427-5432.
 143. Fisher D, Rubinstein M (1986): Human monocytes tumoricidal activity: the role of interferon-gamma and bacterial lipopolysaccharide in its stimulation, preservation and decay. *Immunobiology* 172: 110-119.
 144. Nishimura Y, Higashi N, Tsuji T, et al. (1992): Activation of human monocytes by interleukin-2 and various cytokines. *J Immunother* 12: 90-97.
 145. Bosco M, Pulkki K, Rowe T, et al. (1995): IL-4 inhibits IL-2 induced tumoricidal activity and secretory functions of human monocytes. Modulation of IL-2 receptor beta gamma chain expression. *J Immunol* 155: 1411-1419.
 146. Grabstein K, Urdal D, Tushinski R, et al. (1986): Induction of macrophage tumoricidal activity by granulocyte-macrophage colony-stimulating factor. *Science* 232: 506-508.
 147. Howard A, Erickson K (1995): The induction and augmentation of macrophage tumoricidal responses by plated-activating factor. *Cell Immunol* 164: 105-112.
 148. Perri R, Vercellotti G, McCarthy J, et al. (1985): Laminin selectively enhances monocyte-macrophage-mediated tumoricidal activity. *J Lab Clin Med* 105: 30-35.
 149. McLachlan J, Serkin C, Morrey K, Bakouche O (1995): Antitumoral properties of aged human monocytes. *J Immunol* 154: 832-843.
 150. Peri G, Polentarutti N, Sessa C, et al. (1981): Tumoricidal activity of macrophages isolated from human ascitic and solid ovarian carcinomas: augmentation by interferon, lymphokines and endotoxin. *Int J Cancer* 28: 143-152
 151. Nio Y, Zigelboim J, Berek J, Bonavida B (1990): Cycliheximide-induced modulation of TNF-mediated cytotoxicity in sensitive and resistant ovarian tumor cells. *Cancer Chemother Pharmacol* 26: 1-8.
 152. Mantovani A: Cytotoxic killing of tumour cells by monocytes. In: *Human monocytes*. Ed. Zembala M, Asherson G: Academic Press, London, 1989, 303.
 153. Utsugi T, Schroit A, Connor J, et al. (1991): Elevated expression of phosphatidylserine in outer membrane leaflet of human tumour cells and recognition by activated human blood monocytes. *Cancer Res* 61: 241-248.
 154. Tripathi A, Taplits M, Puri J, Hoffman T (1991): Down-regulation of surface FcRI and decrease in antibody-dependent cellular cytotoxicity of cultured monocytes. *J Immunol* 146: 1309-1315.
 155. Fanger M, Shen L, Graziano R, Guyre M (1989): Cytotoxicity mediated by human Fc receptors for IgG. *Immunol Today* 10: 92-99.
 156. Hellstrom I, Garrigues U, Lavie E, Hellstrom K (1988): Antibody-mediated killing of human tumor cells by attached effector cells. *Cancer Res* 48: 624-627.
 157. Moutsatsos I, Davis J, Wang J (1986): Endogenous lectins from cultured cells: subcellular localisation of carbohydrate-binding protein 35 in 3T3 fibroblasts. *J Cell Biol* 102: 477-483.
 158. Salminen E, Kankuri M, Nuutila J, et al. (2001): Modulation of IgG and complement receptor expression of phagocytes in kidney cancer patients during treatment with interferon-alpha. *Anticancer Res* 21: 2049-2055
 159. Uracz W, Pituch-Noworolska A, Zembala M, et al. (1982): "Activated" monocytes in gastric cancer patients. An increased Fc receptor expression, antibody dependent cellular cytotoxicity and NBT reduction. *J Cancer Res Clin Oncol* 104: 181-184.
 160. Vena G, Angelini G, D'Ovidio R, et al. (1983): Monocyte Fc-IgG receptors expression and soluble suppressor factor in skin squamous cell carcinomas. *Acta Derm Venereol* 63: 507-512.
 161. Gebhard B, Gant M, Schutz G, et al. (2000): Different transendothelial migration behaviour pattern of blood monocytes derived from patients with benign and malignant diseases of the breast. *Anticancer Res* 20: 4599-4604.
 162. Mytar B, Baran J, Gawlicka M, et al. (2002): Immunophenotypic changes and induction of apoptosis of monocytes and tumour cells during their interactions *in vitro*. *Anticancer Research* 22: 2789-2796.
 163. Zembala M, Buckle A: Monocytes in malignant disease. In: *Human monocytes*. Ed. Zembala M, Asherson G: Academic Press, London 1989, 514-528.
 164. Ziegler-Heitbrock L, Fingerle G, Strobel M, et al. (1993) The novel subset of CD14⁺/CD16⁺ blood monocytes exhibits features of tissue macrophages. *Eur J Immunol* 23: 2053-2056.
 165. Frankenberg M, Sternsdorf T, Pechumer H, et al. (1996): Differential cytokine expression in human blood monocyte subpopulation: a polymerase chain reaction analysis. *Blood* 87: 373-379.
 166. Fingerle G, Pforte A, Passlick B, et al. (1993): The novel subset of CD14⁺/CD16⁺ blood monocytes is expanded in sepsis patients. *Blood* 82: 3170-3175.

167. Thomas R, Lipsky P (1994): Human peripheral blood dendritic cell subsets. Isolation and characterisation of precursor and mature antigen presenting cells. *J Immunol* 153: 4016-4028.
168. Siedlar M, Frankenberger M, Ziegler-Heitbrock HWL, Belge KU (2000): The M-DC8-positive leukocytes are a subpopulation of the CD14⁺/CD16⁺ monocytes. *Immunobiol* 202: 11-17.
169. Locher C, Vanham G, Kestens L, et al. (1994): Expression patterns of Fc(receptors, HLA-DR and selected adhesion molecules on monocytes from normal and HIV-infected individuals. *Cli Exp Immunol* 98: 115-121.
170. Fingerle-Rowson G, Auers J, Kreuzer P, et al. (1998): Expansion of CD14⁺/CD16⁺ monocytes in critically ill cardiac surgery patients. *Inflammation* 22: 367-372.
171. Saleh M, Goldman S, LoBugilo A, et al. (1995): CD16⁺ monocytes in patients with cancer: spontaneous elevation and pharmacologic induction by recombinant human macrophage colony-stimulating factor. *Blood* 85: 2910-2917.
172. Schmid I, Baldwin G, Jacobs E, et al. (1995): Alternations in phenotype and cell-surface antigen expression levels of human monocytes: differential response to in vivo administration of rhM-CSF or rhGM-CSF. *Cytometry* 22: 103-110.
173. Beller D, Springer T, Schreiber R (1982): Anti-Mac-1 selectively inhibits the mouse and human type three complement receptor. *J Exp Med* 56: 1000-1009.
174. Shang X, Issekutz A (1998): Contribution of CD11a/CD18, CD11b/CD18, ICAM-1 (CD54) -2 (CD102) to human monocyte migration through endothelium and connective tissue barriers. *Eur J Immunol* 28: 1970-1979.
175. Allen C, Hogg N (1987): Elevation of infiltrating mononuclear phagocytes in human colorectal tumors. *J Natl Cancer Inst* 78: 465-470.
176. Zhang L, Yoshimura T, Graves D (1997): Antibody to Mac-1 or monocyte chemoattractant protein-1 inhibits monocyte recruitment and promotes tumor growth. *J Immunol* 158: 4855-4861.
177. Jiang Y, Zhu J, Lusinskas F, Graves D (1994): MCP-1-stimulated monocyte attachment to laminin is mediated by beta 2-integrins. *Am J Physiol* 267: 1112-1118.
178. Jiang Y, Beller D, Frenzl G, Graves D (1992): Monocyte chemoattractant protein-1 regulates adhesion molecule expression and cytokine production in human monocytes. *J Immunol* 148: 2423-2428.
179. Salminen E, Kankuri M, Nuutila J, et al. (2001): Modulation of IgG and complement receptor expression of phagocytes in kidney cancer patients during treatment with interferon-alpha. *Anticancer Res* 21: 2049-2055.
180. Carlos T, Harlan J (1990): Membrane proteins involved in phagocyte adherence to endothelium. *Immunol-Rev* 114: 5-28.
181. Chuluyan H, Lang B, Issekutz A (2000): Differential mechanisms of neutrophil and monocytes adhesion on neuroblastoma cells: CD18 and VLA-4 integrins mediate adhesion to SK-N-SH, but not to SK-N-MC cell line. *J Neurosci Res* 60: 649-655.
182. Zembala M, Mytar B, Ruggiero I, et al. (1982): Suppressor cells and survival of patients with advanced gastric patients. *J Natl Cancer Inst* 70: 223-228.
183. Jerrells T, Dean J, Richardson G, et al. (1979): Increased monocyte-mediated cytostasis of lymphoid cell lines in breast and lung cancer patients. *Int J Cancer* 23: 768-772.
184. Mytar B, Zembala M, Uracz W, Czupryna A (1982): Cytostatic activity on tumour cells of monocytes from patients with gastrointestinal cancer. *Cancer Immunol Immunother* 13: 190-193.
185. Kiertscher S, Luo J, Dubinett S, Roth M (2000): Tumors promote altered maturation and early apoptosis of monocyte-derived dendritic cells. *J Immunol* 164: 1269-1276.
186. Ladisch S, Li R, Olson E (1994): Ceramide structure predicts tumor ganglioside immunosuppressive activity. *Proc Natl Acad Sci USA*, 91: 1974-1978.
187. Olshefski R, Ladisch S (1996): Intercellular transfer of shed tumor cell gangliosides. *FEBS Letters* 386: 11-14.
188. van Ravenswaay Claasen H, Kluin P, Fleuren G (1992): Tumor infiltrating cells in human cancer. On the possible role of CD16⁺ macrophages in antitumor cytotoxicity. *Lab Invest* 67: 166-174.
189. Leek RD, Lewis CE, Whitehouse R, et al. (1996): Association of macrophage infiltration with angiogenesis and prognosis in invasive breast carcinoma. *Cancer Res* 56: 4625-4629.
190. Lewis C, Leek R, Harris A, McGee J (1995): Cytokine regulation of angiogenesis in breast cancer: the role of tumor-associated macrophages. *J Leukoc Biol* 57: 747-751.
191. Tromp S, oude Egbrink M, Dings R, et al. (1999): Tumor angiogenesis factors reduce leukocyte adhesion in vivo. *Int Immunol* 12: 671-676.
192. Toomey D, Harmeij J, Condron C, et al. (1999): Phenotyping of immune cell infiltrates in breast and colorectal tumours. *Immunol Invest* 28: 29-41.
193. Gottlinger H, Rieber P, Gokel J, et al. (1985): Infiltrating mononuclear cells in human breast carcinoma: predominance of T4⁺ monocytic cells in the tumor stroma. *Int J Cancer* 35: 199-280.
194. Broker E, Zwadlo G, Holzmann B, et al. (1988) *Int J Cancer* 41: 562-567.
195. Loercher A, Nash M, Kavanagh J, et al. (1999): Identification of an IL-10-producing HLA-DR-negative monocyte subset in the malignant ascites of patients with ovarian carcinoma that inhibits cytokine protein expression and proliferation of autologous T cells. *J Immunol* 163: 6251-6260.
196. Allen A, Hogg N (1987): Association of colorectal tumor epithelium expressing HLA-D/DR with CD8 positive T-cells and mononuclear phagocytes. *Cancer Res* 47: 2919-2923.
197. Chau P, Moutet M, Faivre J, et al. (1996): Inflammatory cells infiltrating human colorectal carcinomas express HLA class II but not B7-1 and B7-2 costimulatory molecules of the T-cell activation. *Lab Invest* 74: 975-983.
198. Ropponen K, Eskelinen M, Lipponen P, et al. (1997): Prognostic value of tumour-infiltrating lymphocytes (TILs) in colorectal cancer. *J Pathol* 182: 318-324.
199. Yanagawa H, Takeuchi E, Suzuki Y, et al. (1999): Production of interleukin-10 by alveolar macrophages from lung cancer patients. *Respir Med* 93: 666-671.
200. Doi C, Noguchi Y, Marat D, et al. (1999): Expression of nitric oxide synthase in gastric cancer. *Cancer Lett* 144: 161-167.
201. Thomsen L, Miles D, Happerfield L, et al. (1995): Nitric oxide synthase activity in human breast cancer. *Br J Cancer* 72: 41-44.
202. Vakkala M, Kahlos K, Lakari E, et al. (2000): Inducible nitric oxide synthase expression, apoptosis, and angiogenesis in in situ and invasive breast carcinomas. *Clin Cancer Res* 6: 2408-2416.
203. Naylor M, Malik S, Stamp G, et al. (1990): *In situ* detection of tumour necrosis factor in human ovarian cancer specimens. *Eur J Cancer* 26: 1027-1030.
204. Naylor M, Stamp G, Balkwill F (1990): Demonstration of mRNA for tumor necrosis factor and other cytokines in human colorectal cancer. *Cancer Res* 50: 4436-4440.
205. Erroi A, Simoni M, Chiaffarino F, et al. (1989): IL-1 and IL-6 release by tumor-associated macrophages from human ovarian carcinoma. *Int J Cancer* 44: 795-801.
206. Mantovani A, Dean J, Jerrells T, Herberman R (1980): Augmentation of tumoricidal activity of human monocytes and macrophages by lymphokines. *Int J Cancer* 25: 691-699.

207. Van Netten J, Ashamead B, Cavers D (1992): Macrophages and their putative significance in breast cancer. *Br J Cancer* 66: 220-221.
208. Kleiner D, Stetler-Stevenson W (1999): Matrix metalloproteinases and metastasis. *Cancer Chemother Pharmacol* 43, Suppl: 42-51.
209. Gorrin-Rivas MJ, Arie S, Mori A, et al (2000): Implications of human macrophage metalloelastase and vascular endothelial growth factor gene expression in angiogenesis of hepatocellular carcinoma. *Ann Surg* 231: 67-73.
210. Patterson BC, Sang QA (1997): Angiostatin-converting enzyme activities of human matrilysin (MMP-7) and gelatinase B/typ IV collagenase (MMP-9). *J Biol Chem* 272: 28823-28825.
211. Swallow C, Murray M, Guillem J (1996): Metastatic colorectal cancer cells induce matrix metalloproteinase release by human monocytes. *Clin Exp Metastasis* 14: 3-11.
212. Jiang Y, Goldberg I, Shi Y (2002): Complex roles of tissue inhibitors of metalloproteinases in cancer. *Oncogene* 28: 2245-2252.
213. Chang C, Werb Z (2001): The many faces of metalloproteases: cell growth, invasion, angiogenesis and metastasis. *Trends Cell Biol* 11: 37-43.
214. Mizoi T, Ohtani H, Suzuki Y, et al. (1995): Intercellular adhesion molecule-1 expression by macrophages in human gastrointestinal carcinoma: possible roles as host immune/inflammatory reaction. *Pathol Int* 45: 565-572.
215. Stauder R, Greil R, Schulz T, et al. (1989): Expression of leukocyte function-associated antigen-1 and 7F7 antigen, an adhesion molecule related to intracellular adhesion molecule-1 (ICAM-1) in non-Hodkin lymphomas and leukaemias: possible influence on growth pattern and leukaemic behaviour. *Clin Exp Immunol* 77: 234-238.
216. Johnson J, Stade B, Holzmann B, et al. (1989): *De novo* expression of intracellular-adhesion molecule 1 in melanoma correlates with increased risk of metastasis. *Proc Natl Acad Sci USA* 86: 641-644.
217. Vogetseder W, Feichtinger H, Schulz T, et al. (1989): Expression of 7F7 antigen, a human adhesion molecule identical to intercellular adhesion molecule-1 (ICAM-1) in human carcinomas and their stromal fibroblasts. *Int J Cancer* 43: 768-773.
218. Tomita Y, Nishiyama T, Watanabe H, Fujiwara M, Sato S (1990): Expression of intercellular adhesion molecule-1 (ICAM-1) on renal cell cancer: possible significance in host immune responses. *Int J Cancer* 46: 1001-1006.
219. Schardt C, Heymanns J, Schardt C, et al. (1993): Differential expression of the intercellular adhesion molecule-1 (ICAM-1) in lung cancer cell lines of various histological types. *Eur J Cancer* 29: 2250-2255.
220. Hutchins D, Steel CM (1994): Regulation of ICAM-1 (CD54) expression in human breast cancer cell lines by interleukin 6 and fibroblast-derived factors. *Int J Cancer*, 58: 80-84.
221. Morandin R, Boeynaems J, Duhant X, et al. (1999): SODs are involved in the regulation of ICAM-1 expression in human melanoma and endothelial cells. *Cell Biol Mol* 45: 1053-1063.
222. Hutchins D, Steel CM (1994): Regulation of ICAM-1 (CD54) expression in human breast cancer cell lines by interleukin 6 and fibroblast-derived factors. *Int J Cancer* 58: 808-814.
223. McBride W, Bard J (1979): Hyaluronidase-sensitive halos around adherent cells. Their role in blocking lymphocyte mediated cytotoxicity. *J Exp Med* 149: 507-515.
224. Jonjic N, Alberti S, Bernasconi S, et al. (1992): Heterogeneous susceptibility of human melanoma clones to monocyte cytotoxicity: role of ICAM-1 defined by antibody blocking and gene transfer. *Eur J Immunol* 22: 2255-2260.
225. Nagaoka H, Iino Y, Takei H, Morishita Y (1998): Platelet-derived endothelial cell growth factor/thymidine phosphorylase expression in macrophages correlates with tumor angiogenesis and prognosis in invasive breast cancer. *Int J Oncol* 13: 449-454.
226. Senger D, Van de Water L, Brown L, et al. (1993): Vascular permeability factor (VPF, VEGF) in tumor biology. *Cancer Metastasis Rev* 12: 303-324.
227. Harmey J, Dimitriadis E, Kay E, et al. (1998): Regulation of macrophage production of vascular endothelial growth factor (VEGF) by hypoxia and transformigf growth factor β -1. *Ann Surg Oncol* 5: 271-278.
228. Leek R, Hunt N, Landres R, et al. (2000): Macrophage infiltration is associated with VEGF and EGFR expression in breast cancer. *J Pathol* 190: 430-436.
229. Shibuya M, Luo J, Toyoda M, Yamaguchi S (1999): Involvement of VEGF and its receptors in ascites tumor formation. *Cancer Chemother Pharmacol* 43 Suppl: 72-77.
230. Clauss M, Gerlach M, Gerlach H, et al. (1990): Vascular permeability factor: a tumor-derived polypeptide that induces endothelial cell and monocyte procoagulant activity, and promotes monocyte migration. *J Exp Med* 172: 1535-1545.
231. Strieter R, Polverini P, Arenberg D, Kunkel S (1995): The role of CXC chemokines as regulators of angiogenesis. *Shock* 4: 155-160.
232. Strieter R, Kunkel S, Arenberg D, et al. (1995): Interferon gamma-inducible protein 10 (IP-10), a member of the C-X-C chemokine family, is an inhibitor of angiogenesis. *Biochem Biophys Res Commun* 210: 51-57.
233. Salcedo R, Ponce M, Young H, et al. (2000): Human endothelial cells express CCR2 and respond to MCP-1: direct role of MCP-1 in angiogenesis and tumor progression. *Blood* 96: 34-40.
234. Mahadevan V, Hart J (1990): Metastases and angiogenesis. *Acta Oncol* 29: 319-331.
235. West D, Hapson I, Arnold F, Kumar S (1995): Angiogenesis induced by degradation products of hyaluronic acid. *Science* 228: 1324-1326.
236. Hildenbrand R, Dilger I, Stutte H (1995): Urokinase and macrophages in tumour angiogenesis. *Br J Cancer* 72: 818-823.
237. Pyke C, Kristensen P, Ralfkiaer E, et al. (1991): Urokinase-type plasminogen activator is expressed in stromal cells and its receptor in cancer cells at invasive foci in human colon adenocarcinomas. *Am J Pathol* 138: 1059-1067.
238. Mahadevan V, Hart J (1990): Metastases and angiogenesis. *Acta Oncol* 29: 97-103.
239. Zuk J, Walker R (1987): Immunohistochemical analysis of HLA antigens and mononuclear infiltrates of benign and malignant breast. *J Pathol* 152: 275-285.
240. Barbera-Guillem E, May K, Nyhus J, Nelson M (1999): Promotion of tumor invasion by cooperation of granulocytes and macrophages activated by anti-tumor antibodies. *Neoplasia* 1: 453-460.
241. Swallow C, Murray M, Guillem J (1996): Metastatic colorectal cancer cells induce matrix metalloproteinase release by human monocytes. *Clin Exp Metastasis* 14: 3-11.
242. Ito A, Handa K, Withers D, Satoh M, Hakomori S (2001): Binding specificity of siglec7 to disialogangliosides of renal cell carcinoma: possible role of disialogangliosides in tumor progression. *FEBS Lett* 504: 82-6.
243. Lwaleed BA, Bass PS, Cooper AJ (2001): The biology and tumour-related properties of monocyte tissue factor. *J Pathol* 193: 3-12.

244. Bingle L, Brown N, Lewis C (2002): The role of tumour-associated macrophages in tumour progression: implications for new anticancer therapies. *J Pathol* 196: 254-265.
245. Edwards R, Rickles F, Cronlund M (1981): Abnormalities of blood coagulation in patients with cancer. Mononuclear cell tissue factor generation. *J Lab Clin Med* 98: 917-928.
246. Lwaleed B, Chisholm M, Francis J (1999): The significance of measuring monocyte tissue factor in patients with breast and colorectal cancer. *Br J Cancer* 80: 279-285.
247. Dano K, Andreassen P, Grondhal-Hansen J, et al. (1985): Plasminogen activators. Tissue degradation and cancer. *Adv Cancer Res* 44: 139-266.
248. Dvorak H (1986): Tumours: wounds that do not heal. *N Engl J Med* 315: 161-165.
249. Lipponen P (1996): Expression of cathepsin D in transitional cell bladder tumours. *J Pathol* 178: 59-64.
250. Lah T, Kalman E, Najjar D, et al. (2000): Cells producing cathepsins B, D and L in human breast carcinoma and their association with prognosis. *Hum Pathol* 31: 149-160.
251. Spötl L, Sarti A, Dierich M, Most J (1995): Cell membrane labeling with fluorescent dyes for the demonstration of cytokine-induced fusion between monocytes and tumor cells. *Cytometry* 21:160-169.
252. Larizza L, Schirmacher V, Graf L, et al. (1984): Suggestive evidence that the highly metastatic variant Esb of the T-cell lymphoma Eb is derived from spontaneous fusion with a host macrophage. *Int J Cancer* 34: 699-707.
253. Teitelbaum S, Stewart C, Kahn A (1979): Rodent peritoneal macrophages as bone resorbing cells. *Calcif Tissue Int* 27: 255-261.
254. Yonemura Y, Endo Y, Yamaguchi T, et al. (1996): Mechanisms of the formation of the peritoneal dissemination in gastric cancer. *Int J Oncology* 8: 795-802.
255. Hagiwara A, Takahashi T, Sawai K, et al. (1993): Milky spots as the implantation site for malignant cells in peritoneal dissemination in mice. *Cancer Res* 53: 687-692.
256. Rosenberg S, Lotze M, Muul L, Chang A, Avis F (1987): Progress report on treatment of 157 patients with advanced cancer using LAK cells and IL2 or IL2 alone. *New Engl Med* 316: 889-897.
257. Stevenson HC: Adoptive cellular immunotherapy of cancer. Marcel Dekker, New York, 1989.
258. Bartholeyns J, Lopez M, Andreessen R (1991): Adoptive immunotherapy of solid tumors with activated macrophages; experimental and clinical results. *Anticancer Res* 11: 1201-1204
259. Fidler I, Jessup J, Fogler W, et al. (1986): Activation of tumoricidal properties in peripheral blood monocytes of patients with colorectal carcinoma. *Cancer Res* 46: 994-998.
260. Chokri M, Lopez M, Oleron C, et al. (1992): Production of human macrophages with potent antitumor properties (MAK) by culture of monocytes in the presence of GM-CSF and 1,25-dihydroxy vitamin D3. *Anticancer Res* 12: 2257-2260.
261. Hennemann B, Beckman G, Eichelmann A, et al. (1998): Phase I trial of adoptive immunotherapy of cancer patients using monocyte-derived macrophages activated with interferon γ and lipopolisaccharide. *Cancer Immunol Immunother* 45: 250-256.
262. Chokri M, Girard A, Borrelly M, et al. (1992): Adoptive immunotherapy with bispecific antibodies: targeting through macrophages. *Res Immunol* 143: 95-99.
263. Stevenson H, Keenan A, Woodhouse C, et al. (1987): Fate of IFN-gamma activated killer blood monocytes adoptively transferred into the abdominal cavity of patients with peritoneal carcinoma. *Cancer Res* 47: 6100-6103.
264. Andreessen R, Scheibenbogen C, Brugger W, et al. (1990): Adoptive transfer of tumor cytotoxic macrophages generated *in vitro* from circulating monocytes: a new approach to cancer immunotherapy. *Cancer Res* 50: 7450-7456.
265. Faradji A, Bohbot A, Schmitt M, et al. (1991): Phase I trial of IV infusion of ex vivo activated blood-derived macrophages in patients with non-small cell lung cancer: toxicity and immunomodulatory effects. *Cancer Immunol Immunother* 33: 319-326.
266. Lopez M, Fechtenbaum J, David B, et al. (1992): Adoptive immunotherapy with activated macrophages grown *in vitro* from blood monocytes in cancer patients: a pilot study. *J Immunother* 11: 209-217.
267. Anderson C (1992): Gene therapy researcher under fire over controversial cancer trials. *Nature* 360: 399-403.
268. Walder S, Haynes H, Beitler J, et al. (1996): Phase II clinical trial with 5-fluorouracil, recombinant interferon-alpha-2b, and cisplatin for patients with metastatic or regionally advanced carcinoma of the esophagus. *Cancer* 78: 30-34.
269. Wiesel M, Faradji A, Grunebaum L, et al. (1992): Hemostatic changes in human adoptive immunotherapy with activated blood monocytes or derived macrophages. *Ann Hematol* 65: 778-782.
270. Allavena P, Grandi M, D'Incalci M, et al. (1987): Human tumor cell lines with pleiotropic drug resistance are efficiently killed by interleukin-2 activated killer cells and by activated monocytes. *Int J Cancer* 40: 104-107.
271. Allavena P, Damia G, Colombo T, et al. (1989): Lymphokine-activated killer (LAK) and monocyte-mediated cytotoxicity on tumor cell lines resistant to antitumor agent. *Cell Immunol* 120: 250-258.
272. Bartholeyns J, Lopez M (1994): Immune control of neoplasia by adoptive transfer of macrophages: potentiality for antigen presentation and gene transfer. *Anticancer Res* 14: 2673-2676.
273. Lei H, Ju D, Yu Y, et al. (2000): Induction of potent antitumor response by vaccination with tumor lysate-pulsed macrophages engineered to secrete macrophage colony-stimulating factor and interferon-gamma. *Gene Ther* 7: 707-713.
274. Böni H, Körholz D, Lex C, et al. (2000): Monocyte deactivation and its reversal in a patient with chemotherapy-induced leukopenia and severe systemic infection. *Med. Pediatr Oncol* 34: 39-42.
275. Griffiths L, Binley K, Iqbal S, et al. (2000): The macrophage - a novel system to deliver gene therapy to pathological hypoxia. *Gene Ther* 7: 255-262.