

## Existence of positive solutions for a singular fractional boundary value problem

Johnny Henderson<sup>a</sup>, Rodica Luca<sup>b</sup>

<sup>a</sup>Department of Mathematics, Baylor University,  
Waco, Texas, 76798-7328 USA  
johnny\_henderson@baylor.edu

<sup>b</sup>Department of Mathematics, Gh. Asachi Technical University,  
Iasi 700506, Romania  
rluca@math.tuiasi.ro

**Received:** March 15, 2016 / **Revised:** May 27, 2016 / **Published online:** December 2, 2016

**Abstract.** We study the existence of positive solutions for a nonlinear Riemann–Liouville fractional differential equation with a sign-changing nonlinearity, subject to multi-point fractional boundary conditions.

**Keywords:** Riemann–Liouville fractional differential equation, fractional boundary conditions, positive solutions, sign-changing nonlinearity.

### 1 Introduction

We consider the nonlinear fractional differential equation

$$D_{0+}^{\alpha} u(t) + \lambda f(t, u(t)) = 0, \quad t \in (0, 1), \quad (\text{E})$$

with the multi-point fractional boundary conditions

$$\begin{aligned} u(0) = u'(0) = \dots = u^{(n-2)}(0) = 0, \\ D_{0+}^p u(t)|_{t=1} = \sum_{i=1}^m a_i D_{0+}^q u(t)|_{t=\xi_i}, \end{aligned} \quad (\text{BC})$$

where  $\lambda$  is a positive parameter,  $\alpha \in \mathbb{R}$ ,  $\alpha \in (n-1, n]$ ,  $n \in \mathbb{N}$ ,  $n \geq 3$ ,  $\xi_i \in \mathbb{R}$  for all  $i = 1, \dots, m$  ( $m \in \mathbb{N}$ ),  $0 < \xi_1 < \dots < \xi_m < 1$ ,  $p, q \in \mathbb{R}$ ,  $p \in [1, n-2]$ ,  $q \in [0, p]$ ,  $D_{0+}^{\alpha}$  denotes the Riemann–Liouville derivative of order  $\alpha$ , and the nonlinearity  $f$  may change sign and may be singular at  $t = 0$  or  $t = 1$ .

We present intervals for parameter  $\lambda$  such that problem (E)–(BC) has at least one positive solution. By a positive solution of (E)–(BC) we mean a function  $u \in C([0, 1])$  satisfying (E) and (BC) with  $u(t) > 0$  for all  $t \in (0, 1]$ . This problem is a generalization

of the problem studied in [7], where  $p \in \mathbb{N}$  and  $q = 1$ . Other particular cases of problem (E)–(BC) and of problem from [7] were investigated in [13] (where  $n = 3$ ,  $p = 0$  and  $a_i = 0$  for all  $i = 1, \dots, m$ ) and in [14] (where  $p = 0$  and  $a_i = 0$  for all  $i = 1, \dots, m$  and  $n$  is an arbitrary natural number,  $n \geq 3$ ). In [11], the author studied the existence of pseudo solutions of problem (E)–(BC) in a reflexive Banach space with  $\lambda = 1$ ,  $f(t, u) = g(t)f(t, u)$ ,  $p = q = 0$  and instead of derivative  $D_{0+}^\alpha$ , he considered the pseudo fractional derivative  $D^\alpha$ . For some recent results on the existence and multiplicity of positive solutions for systems of fractional differential equations with various boundary conditions, see the monograph [5].

Fractional differential equations describe many phenomena in various fields of engineering and scientific disciplines such as physics, biophysics, chemistry, biology, economics, control theory, signal and image processing, aerodynamics, viscoelasticity, electromagnetics, and so on (see [1–3, 8–10, 12, 15]).

The paper is organized as follows. Section 2 contains some auxiliary results, which investigate a nonlocal boundary value problem for fractional differential equations. In Section 3, we give the existence theorems for the positive solutions with respect to a cone for our problem (E)–(BC). Finally, in Section 4, two examples are given to illustrate our main results.

## 2 Auxiliary results

We present here the definitions of the Riemann–Liouville fractional integral and the Riemann–Liouville fractional derivative and some auxiliary results that will be used to prove our main results.

**Definition 1.** The (left-sided) fractional integral of order  $\alpha > 0$  of a function  $f : (0, \infty) \rightarrow \mathbb{R}$  is given by

$$(I_{0+}^\alpha f)(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} f(s) \, ds, \quad t > 0,$$

provided the right-hand side is pointwise defined on  $(0, \infty)$ , where  $\Gamma(\alpha)$  is the Euler gamma function defined by  $\Gamma(\alpha) = \int_0^\infty t^{\alpha-1} e^{-t} \, dt$ ,  $\alpha > 0$ .

**Definition 2.** The Riemann–Liouville fractional derivative of order  $\alpha \geq 0$  for a function  $f : (0, \infty) \rightarrow \mathbb{R}$  is given by

$$\begin{aligned} (D_{0+}^\alpha f)(t) &= \left( \frac{d}{dt} \right)^n (I_{0+}^{n-\alpha} f)(t) \\ &= \frac{1}{\Gamma(n-\alpha)} \left( \frac{d}{dt} \right)^n \int_0^t \frac{f(s)}{(t-s)^{\alpha-n+1}} \, ds, \quad t > 0, \end{aligned}$$

where  $n = \lfloor \alpha \rfloor + 1$ , provided that the right-hand side is pointwise defined on  $(0, \infty)$ .

The notation  $[\alpha]$  stands for the largest integer not greater than  $\alpha$ . If  $\alpha = m \in \mathbb{N}$ , then  $D_{0+}^m f(t) = f^{(m)}(t)$  for  $t > 0$ , and if  $\alpha = 0$ , then  $D_{0+}^0 f(t) = f(t)$  for  $t > 0$ .

We consider now the fractional differential equation

$$D_{0+}^\alpha u(t) + \tilde{x}(t) = 0, \quad 0 < t < 1, \tag{1}$$

with the boundary conditions

$$\begin{aligned} u(0) = u'(0) = \dots = u^{(n-2)}(0) = 0, \\ D_{0+}^p u(t)|_{t=1} = \sum_{i=1}^m a_i D_{0+}^q u(t)|_{t=\xi_i}, \end{aligned} \tag{2}$$

where  $\alpha \in (n - 1, n]$ ,  $n \geq 3$ ,  $0 < \xi_1 < \dots < \xi_m < 1$ ,  $p, q \in \mathbb{R}$ ,  $p \in [1, n - 2]$ ,  $q \in [0, p]$ , and  $\tilde{x} \in C(0, 1) \cap L^1(0, 1)$ . We denote by  $\Delta = \Gamma(\alpha)/\Gamma(\alpha - p) - (\Gamma(\alpha)/\Gamma(\alpha - q)) \sum_{i=1}^m a_i \xi_i^{\alpha-q-1}$ .

**Lemma 1.** *If  $\Delta \neq 0$ , then the function  $u \in C([0, 1])$  given by*

$$\begin{aligned} u(t) = & -\frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} \tilde{x}(s) \, ds + \frac{t^{\alpha-1}}{\Delta \Gamma(\alpha-p)} \int_0^1 (1-s)^{\alpha-p-1} \tilde{x}(s) \, ds \\ & - \frac{t^{\alpha-1}}{\Delta \Gamma(\alpha-q)} \sum_{i=1}^m a_i \int_0^{\xi_i} (\xi_i-s)^{\alpha-q-1} \tilde{x}(s) \, ds, \quad t \in [0, 1], \end{aligned} \tag{3}$$

is solution of problem (1)–(2).

*Proof.* We denote by

$$\begin{aligned} c_1 = & \frac{1}{\Delta \Gamma(\alpha-p)} \int_0^1 (1-s)^{\alpha-p-1} \tilde{x}(s) \, ds \\ & - \frac{1}{\Delta \Gamma(\alpha-q)} \sum_{i=1}^m a_i \int_0^{\xi_i} (\xi_i-s)^{\alpha-q-1} \tilde{x}(s) \, ds. \end{aligned}$$

Then the continuous function  $u$  from (3) can be written as

$$\begin{aligned} u(t) = & c_1 t^{\alpha-1} - I_{0+}^\alpha \tilde{x}(t) \\ = & c_1 t^{\alpha-1} - \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} \tilde{x}(s) \, ds, \quad t \in [0, 1]. \end{aligned} \tag{4}$$

Because  $D_{0+}^\alpha u(t) = c_1 D_{0+}^\alpha (t^{\alpha-1}) - D_{0+}^\alpha I_{0+}^\alpha \tilde{x}(t) = -\tilde{x}(t)$  for all  $t \in (0, 1)$ , we deduce that  $u$  satisfies equation (1). In addition, we have  $u(0) = u'(0) = \dots = u^{(n-2)}(0) = 0$ .

Because

$$D_{0+}^p u(t) = c_1 \frac{\Gamma(\alpha)}{\Gamma(\alpha-p)} t^{\alpha-p-1} - I_{0+}^{\alpha-p} \tilde{x}(t),$$

$$D_{0+}^q u(t) = c_1 \frac{\Gamma(\alpha)}{\Gamma(\alpha-q)} t^{\alpha-q-1} - I_{0+}^{\alpha-q} \tilde{x}(t),$$

by a simple computation we conclude that  $D_{0+}^p u(t)|_{t=1} = \sum_{i=1}^m a_i D_{0+}^q u(t)|_{t=\xi_i}$ , that is

$$\begin{aligned} & c_1 \frac{\Gamma(\alpha)}{\Gamma(\alpha-p)} - \frac{1}{\Gamma(\alpha-p)} \int_0^1 (1-s)^{\alpha-p-1} \tilde{x}(s) \, ds \\ &= \sum_{i=1}^m a_i \left( c_1 \frac{\Gamma(\alpha)}{\Gamma(\alpha-q)} \xi_i^{\alpha-q-1} - \frac{1}{\Gamma(\alpha-q)} \int_0^{\xi_i} (\xi_i-s)^{\alpha-q-1} \tilde{x}(s) \, ds \right). \end{aligned}$$

Therefore, we deduce that  $u$  is solution of problem (1)–(2).  $\square$

**Lemma 2.** *If  $\Delta \neq 0$ , then the solution  $u$  of problem (1)–(2) given by (3) can be written as*

$$u(t) = \int_0^1 G(t,s) \tilde{x}(s) \, ds, \quad t \in [0,1], \quad (5)$$

where

$$G(t,s) = g_1(t,s) + \frac{t^{\alpha-1}}{\Delta} \sum_{i=1}^m a_i g_2(\xi_i,s), \quad (6)$$

and

$$\begin{aligned} g_1(t,s) &= \frac{1}{\Gamma(\alpha)} \begin{cases} t^{\alpha-1}(1-s)^{\alpha-p-1} - (t-s)^{\alpha-1}, & 0 \leq s \leq t \leq 1, \\ t^{\alpha-1}(1-s)^{\alpha-p-1}, & 0 \leq t \leq s \leq 1, \end{cases} \\ g_2(t,s) &= \frac{1}{\Gamma(\alpha-q)} \begin{cases} t^{\alpha-q-1}(1-s)^{\alpha-p-1} - (t-s)^{\alpha-q-1}, & 0 \leq s \leq t \leq 1, \\ t^{\alpha-q-1}(1-s)^{\alpha-p-1}, & 0 \leq t \leq s \leq 1, \end{cases} \end{aligned} \quad (7)$$

for all  $(t,s) \in [0,1] \times [0,1]$ .

*Proof.* By Lemma 1 and relation (3), we deduce

$$\begin{aligned} u(t) &= \frac{1}{\Gamma(\alpha)} \left\{ \int_0^t [t^{\alpha-1}(1-s)^{\alpha-p-1} - (t-s)^{\alpha-1}] \tilde{x}(s) \, ds \right. \\ &\quad \left. + \int_t^1 t^{\alpha-1}(1-s)^{\alpha-p-1} \tilde{x}(s) \, ds \right\} - \frac{1}{\Gamma(\alpha)} \int_0^1 t^{\alpha-1}(1-s)^{\alpha-p-1} \tilde{x}(s) \, ds \end{aligned}$$

$$\begin{aligned}
& + \frac{t^{\alpha-1}}{\Delta\Gamma(\alpha-p)} \int_0^1 (1-s)^{\alpha-p-1} \tilde{x}(s) \, ds - \frac{t^{\alpha-1}}{\Delta\Gamma(\alpha-q)} \sum_{i=1}^m a_i \int_0^{\xi_i} (\xi_i-s)^{\alpha-q-1} \tilde{x}(s) \, ds \\
& = \frac{1}{\Gamma(\alpha)} \left\{ \int_0^t [t^{\alpha-1}(1-s)^{\alpha-p-1} - (t-s)^{\alpha-1}] \tilde{x}(s) \, ds \right. \\
& \quad \left. + \int_t^1 t^{\alpha-1}(1-s)^{\alpha-p-1} \tilde{x}(s) \, ds \right\} - \frac{1}{\Delta\Gamma(\alpha-p)} \int_0^1 t^{\alpha-1}(1-s)^{\alpha-p-1} \tilde{x}(s) \, ds \\
& \quad + \frac{1}{\Delta\Gamma(\alpha-q)} \sum_{i=1}^m a_i \xi_i^{\alpha-q-1} \int_0^1 t^{\alpha-1}(1-s)^{\alpha-p-1} \tilde{x}(s) \, ds \\
& \quad + \frac{t^{\alpha-1}}{\Delta\Gamma(\alpha-p)} \int_0^1 (1-s)^{\alpha-p-1} \tilde{x}(s) \, ds - \frac{t^{\alpha-1}}{\Delta\Gamma(\alpha-q)} \sum_{i=1}^m a_i \int_0^{\xi_i} (\xi_i-s)^{\alpha-q-1} \tilde{x}(s) \, ds \\
& = \frac{1}{\Gamma(\alpha)} \left\{ \int_0^t [t^{\alpha-1}(1-s)^{\alpha-p-1} - (t-s)^{\alpha-1}] \tilde{x}(s) \, ds \right. \\
& \quad \left. + \int_t^1 t^{\alpha-1}(1-s)^{\alpha-p-1} \tilde{x}(s) \, ds \right\} \\
& \quad + \frac{t^{\alpha-1}}{\Delta\Gamma(\alpha-q)} \sum_{i=1}^m a_i \left[ \int_0^1 \xi_i^{\alpha-q-1} (1-s)^{\alpha-p-1} \tilde{x}(s) \, ds - \int_0^{\xi_i} (\xi_i-s)^{\alpha-q-1} \tilde{x}(s) \, ds \right] \\
& = \frac{1}{\Gamma(\alpha)} \left\{ \int_0^t [t^{\alpha-1}(1-s)^{\alpha-p-1} - (t-s)^{\alpha-1}] \tilde{x}(s) \, ds \right. \\
& \quad \left. + \int_t^1 t^{\alpha-1}(1-s)^{\alpha-p-1} \tilde{x}(s) \, ds \right\} \\
& \quad + \frac{t^{\alpha-1}}{\Delta\Gamma(\alpha-q)} \sum_{i=1}^m a_i \left\{ \int_0^{\xi_i} [\xi_i^{\alpha-q-1} (1-s)^{\alpha-p-1} - (\xi_i-s)^{\alpha-q-1}] \tilde{x}(s) \, ds \right. \\
& \quad \left. + \int_{\xi_i}^1 \xi_i^{\alpha-q-1} (1-s)^{\alpha-p-1} \tilde{x}(s) \, ds \right\}
\end{aligned}$$

$$= \int_0^1 g_1(t, s) \tilde{x}(s) \, ds + \frac{t^{\alpha-1}}{\Delta} \sum_{i=1}^m a_i \int_0^1 g_2(\xi_i, s) \tilde{x}(s) \, ds = \int_0^1 G(t, s) \tilde{x}(s) \, ds,$$

where  $G$  and  $g_1, g_2$  are given in (6) and (7).

Therefore, we obtain expression (5) for the solution  $u$  of problem (1)–(2) given by (3).  $\square$

**Lemma 3.** *The functions  $g_1$  and  $g_2$  given by (7) have the properties:*

- (a)  $g_1(t, s) \leq h_1(s)$  for all  $ts \in [0, 1]$ , where  $h_1(s) = (1-s)^{\alpha-p-1}(1-(1-s)^p)/\Gamma(\alpha)$ ,  $s \in [0, 1]$ ;
- (b)  $g_1(t, s) \geq t^{\alpha-1}h_1(s)$  for all  $t, s \in [0, 1]$ ;
- (c)  $g_1(t, s) \leq t^{\alpha-1}/\Gamma(\alpha)$  for all  $t, s \in [0, 1]$ ;
- (d)  $g_2(t, s) \geq t^{\alpha-q-1}h_2(s)$  for all  $t, s \in [0, 1]$ , where  $h_2(s) = (1-s)^{\alpha-p-1} \times (1-(1-s)^{p-q})/\Gamma(\alpha-q)$ ,  $s \in [0, 1]$ ;
- (e)  $g_2(t, s) \leq (1/\Gamma(\alpha-q))t^{\alpha-q-1}$  for all  $t, s \in [0, 1]$ ;
- (f) *The functions  $g_1$  and  $g_2$  are continuous on  $[0, 1] \times [0, 1]$ ;  $g_1(t, s) \geq 0$ ,  $g_2(t, s) \geq 0$  for all  $t, s \in [0, 1]$ ;  $g_1(t, s) > 0$ ,  $g_2(t, s) > 0$  for all  $t, s \in (0, 1)$ .*

*Proof.* For the proof of properties (a)–(c), see [6].

(d) From (7), if  $s \leq t$ , we obtain

$$\begin{aligned} g_2(t, s) &= \frac{1}{\Gamma(\alpha-q)} [t^{\alpha-q-1}(1-s)^{\alpha-p-1} - (t-s)^{\alpha-q-1}] \\ &\geq \frac{1}{\Gamma(\alpha-q)} [t^{\alpha-q-1}(1-s)^{\alpha-p-1} - (t-ts)^{\alpha-q-1}] \\ &= \frac{1}{\Gamma(\alpha-q)} t^{\alpha-q-1} [(1-s)^{\alpha-p-1} - (1-s)^{\alpha-q-1}] \\ &= \frac{1}{\Gamma(\alpha-q)} t^{\alpha-q-1} (1-s)^{\alpha-p-1} (1-(1-s)^{p-q}) \\ &= t^{\alpha-q-1} h_2(s), \end{aligned}$$

where  $h_2(s) = (1/\Gamma(\alpha-q))(1-s)^{\alpha-p-1}(1-(1-s)^{p-q})$ ,  $s \in [0, 1]$ .

If  $t \leq s$ , we deduce

$$g_2(t, s) = \frac{1}{\Gamma(\alpha-q)} t^{\alpha-q-1} (1-s)^{\alpha-p-1} \geq t^{\alpha-q-1} h_2(s).$$

Hence, we conclude  $g_2(t, s) \geq t^{\alpha-q-1} h_2(s)$  for all  $t, s \in [0, 1]$ .

(e) We have

$$g_2(t, s) \leq \frac{1}{\Gamma(\alpha-q)} t^{\alpha-q-1} (1-s)^{\alpha-p-1} \leq \frac{t^{\alpha-q-1}}{\Gamma(\alpha-q)} \quad \forall t, s \in [0, 1].$$

(f) This property follows from the definitions of  $g_1$  and  $g_2$  and from properties (b) and (d) above.  $\square$

**Lemma 4.** Assume that  $a_i \geq 0$  for all  $i = 1, \dots, m$  and  $\Delta > 0$ . Then the function  $G$  given by (6) is a continuous function on  $[0, 1] \times [0, 1]$  and satisfies the inequalities:

- (a)  $G(t, s) \leq J(s)$  for all  $t, s \in [0, 1]$ , where  $J(s) = h_1(s) + (1/\Delta) \sum_{i=1}^m a_i g_2(\xi_i, s)$ ,  $s \in [0, 1]$ ;
- (b)  $G(t, s) \geq t^{\alpha-1} J(s)$  for all  $t, s \in [0, 1]$ ;
- (c)  $G(t, s) \leq \sigma t^{\alpha-1}$  for all  $t, s \in [0, 1]$ , where  $\sigma = 1/\Gamma(\alpha) + \sum_{i=1}^m a_i \xi_i^{\alpha-q-1} / (\Delta \Gamma(\alpha - q))$ .

*Proof.* By definition of the function  $G$  we deduce that  $G$  is a continuous function. In addition, by using Lemma 3, we obtain for all  $t, s \in [0, 1]$

- (a)  $G(t, s) \leq h_1(s) + \frac{1}{\Delta} \sum_{i=1}^m a_i g_2(\xi_i, s) = J(s)$ ;
- (b)  $G(t, s) \geq t^{\alpha-1} h_1(s) + \frac{t^{\alpha-1}}{\Delta} \sum_{i=1}^m a_i g_2(\xi_i, s) = t^{\alpha-1} J(s)$ ;
- (c)  $G(t, s) \leq \frac{t^{\alpha-1}}{\Gamma(\alpha)} + \frac{t^{\alpha-1}}{\Delta \Gamma(\alpha - q)} \sum_{i=1}^m a_i \xi_i^{\alpha-q-1} = \sigma t^{\alpha-1}$ . □

**Lemma 5.** Assume that  $a_i \geq 0$  for all  $i = 1, \dots, m$ ,  $\Delta > 0$ ,  $\tilde{x} \in C(0, 1) \cap L^1(0, 1)$  and  $\tilde{x}(t) > 0$  for all  $t \in (0, 1)$ . Then the solution  $u$  of problem (1)–(2) given by (3) satisfies the inequality  $u(t) \geq t^{\alpha-1} u(t')$  for all  $t, t' \in [0, 1]$ .

*Proof.* By using Lemma 4, we obtain for all  $t, t' \in [0, 1]$

$$\begin{aligned}
 u(t) &= \int_0^1 G(t, s) \tilde{x}(s) \, ds \geq \int_0^1 t^{\alpha-1} J(s) \tilde{x}(s) \, ds \\
 &\geq t^{\alpha-1} \int_0^1 G(t', s) \tilde{x}(s) \, ds = t^{\alpha-1} u(t').
 \end{aligned}$$
□

In the proof of our main results, we shall use the Guo–Krasnosel’skii fixed point theorem presented below (see [4]).

**Theorem 1.** Let  $X$  be a Banach space and let  $C \subset X$  be a cone in  $X$ . Assume  $\Omega_1$  and  $\Omega_2$  are bounded open subsets of  $X$  with  $0 \in \Omega_1 \subset \overline{\Omega_1} \subset \Omega_2$  and let  $\mathcal{A} : C \cap (\overline{\Omega_2} \setminus \Omega_1) \rightarrow C$  be a completely continuous operator such that, either

- (i)  $\|\mathcal{A}u\| \leq \|u\|$ ,  $u \in C \cap \partial\Omega_1$ , and  $\|\mathcal{A}u\| \geq \|u\|$ ,  $u \in C \cap \partial\Omega_2$ , or
- (ii)  $\|\mathcal{A}u\| \geq \|u\|$ ,  $u \in C \cap \partial\Omega_1$ , and  $\|\mathcal{A}u\| \leq \|u\|$ ,  $u \in C \cap \partial\Omega_2$ .

Then  $\mathcal{A}$  has a fixed point in  $C \cap (\overline{\Omega_2} \setminus \Omega_1)$ .

### 3 Main results

In this section, we investigate the existence of positive solutions for our problem (E)–(BC). First, we present the assumptions that we shall use in the sequel.

- (H1)  $\alpha \in \mathbb{R}$ ,  $\alpha \in (n-1, n]$ ,  $n \in \mathbb{N}$ ,  $n \geq 3$ ,  $\xi_i \in \mathbb{R}$  for all  $i = 1, \dots, m$  ( $m \in \mathbb{N}$ ),  $0 < \xi_1 < \dots < \xi_m < 1$ ,  $p, q \in \mathbb{R}$ ,  $p \in [1, n-2]$ ,  $q \in [0, p]$ ,  $a_i \geq 0$  for all  $i = 1, \dots, m$ ,  $\lambda > 0$ ,  $\Delta = \Gamma(\alpha)/\Gamma(\alpha-p) - (\Gamma(\alpha)/\Gamma(\alpha-q)) \sum_{i=1}^m a_i \xi_i^{\alpha-q-1} > 0$ .
- (H2) The function  $f \in C((0, 1) \times [0, \infty), \mathbb{R})$  may be singular at  $t = 0$  and/or  $t = 1$ , and there exist the functions  $r, z \in C((0, 1), [0, \infty))$ ,  $g \in C([0, 1] \times [0, \infty), [0, \infty))$  such that  $-r(t) \leq f(t, x) \leq z(t)g(t, x)$  for all  $t \in (0, 1)$  and  $x \in [0, \infty)$  with  $0 < \int_0^1 r(t) dt < \infty$ ,  $0 < \int_0^1 z(t) dt < \infty$ .
- (H3) There exists  $c \in (0, 1/2)$  such that  $f_\infty = \lim_{u \rightarrow \infty} \min_{t \in [c, 1-c]} f(t, u)/u = \infty$ .
- (H4) There exists  $c \in (0, 1/2)$  such that  $\liminf_{u \rightarrow \infty} \min_{t \in [c, 1-c]} f(t, u) > L_0$ , with  $L_0 = (2\sigma \int_0^1 r(s) ds) / (c^{\alpha-1} \int_c^{1-c} J(s) ds)$ , and  $g_\infty = \lim_{u \rightarrow \infty} \max_{t \in [0, 1]} g(t, u)/u = 0$ , where  $J$  and  $\sigma$  are given in Section 2 (Lemma 4).

We consider the fractional differential equation

$$D_{0+}^\alpha x(t) + \lambda(f(t, [x(t) - \lambda w(t)]^*) + r(t)) = 0, \quad 0 < t < 1, \quad (8)$$

with the multi-point fractional boundary conditions

$$\begin{aligned} x(0) = x'(0) = \dots = x^{(n-2)}(0) = 0, \\ D_{0+}^p x(t)|_{t=1} = \sum_{i=1}^m a_i D_{0+}^q x(t)|_{t=\xi_i}, \end{aligned} \quad (9)$$

where  $\lambda > 0$  and  $\zeta(t)^* = \zeta(t)$  if  $\zeta(t) \geq 0$ , and  $\zeta(t)^* = 0$  if  $\zeta(t) < 0$ . Here  $w(t) = \int_0^1 G(t, s)r(s) ds$ ,  $t \in [0, 1]$ , is solution of problem

$$\begin{aligned} D_{0+}^\alpha w(t) + r(t) = 0, \quad 0 < t < 1, \\ w(0) = w'(0) = \dots = w^{(n-2)}(0) = 0, \\ D_{0+}^p w(t)|_{t=1} = \sum_{i=1}^m a_i D_{0+}^q w(t)|_{t=\xi_i}. \end{aligned}$$

Under assumptions (H1)–(H2), we have  $w(t) \geq 0$  for all  $t \in [0, 1]$ . We shall prove that there exists a solution  $x$  of problem (8)–(9) with  $x(t) \geq \lambda w(t)$  on  $[0, 1]$  and  $x(t) > \lambda w(t)$  on  $(0, 1)$ . In this case,  $u = x - \lambda w$  represents a positive solution of problem (E)–(BC). Therefore, in what follows, we shall investigate problem (8)–(9).

By using Lemma 2, a solution of equation

$$x(t) = \lambda \int_0^1 G(t, s)(f(s, [x(s) - \lambda w(s)]^*) + r(s)) ds, \quad t \in [0, 1],$$

is a solution for problem (8)–(9).



We consider the Banach space  $X = C([0, 1])$  with the supremum norm  $\|\cdot\|$ , and we define the cone

$$P = \{x \in X: x(t) \geq t^{\alpha-1}\|x\| \ \forall t \in [0, 1]\}.$$

For  $\lambda > 0$ , we introduce the operator  $T : X \rightarrow X$  defined by

$$Tx(t) = \lambda \int_0^1 G(t, s)(f(s, [x(s) - \lambda w(s)]^*) + r(s)) \, ds, \quad t \in [0, 1], x \in X.$$

It is clear that if  $x$  is a fixed point of operator  $T$ , then  $x$  is a solution of problem (8)–(9).

**Lemma 6.** *If (H1) and (H2) hold, then operator  $T : P \rightarrow P$  is a completely continuous operator.*

*Proof.* Let  $x \in P$  be fixed. By using (H1) and (H2), we deduce that  $Tx(t) < \infty$  for all  $t \in [0, 1]$ . Besides, by Lemma 4, we obtain for all  $t, t' \in [0, 1]$

$$\begin{aligned} Tx(t) &\leq \lambda \int_0^1 J(s)(f(s, [x(s) - \lambda w(s)]^*) + r(s)) \, ds, \\ Tx(t) &\geq \lambda \int_0^1 t^{\alpha-1} J(s)(f(s, [x(s) - \lambda w(s)]^*) + r(s)) \, ds \\ &\geq t^{\alpha-1}Tx(t'). \end{aligned}$$

Therefore,  $Tx(t) \geq t^{\alpha-1}\|Tx\|$  for all  $t \in [0, 1]$ . We deduce that  $Tx \in P$ , and hence,  $T(P) \subset P$ .

By using standard arguments, we conclude that operator  $T : P \rightarrow P$  is a completely continuous operator. □

**Theorem 2.** *Assume that (H1), (H2) and (H3) hold. Then there exists  $\lambda^* > 0$  such that, for any  $\lambda \in (0, \lambda^*]$ , the boundary value problem (E)–(BC) has at least one positive solution.*

*Proof.* We choose a positive number  $R_1 > \sigma \int_0^1 r(s) \, ds > 0$ , and we define the set  $\Omega_1 = \{x \in P: \|x\| < R_1\}$ .

We introduce

$$\lambda^* = \min \left\{ 1, R_1 \left( M_1 \int_0^1 J(s)(z(s) + r(s)) \, ds \right)^{-1} \right\}$$

with  $M_1 = \max\{\max_{t \in [0,1], u \in [0, R_1]} g(t, u), 1\}$ .

Let  $\lambda \in (0, \lambda^*]$ . Because  $w(t) \leq \sigma t^{\alpha-1} \int_0^1 r(s) \, ds$  for all  $t \in [0, 1]$ , we deduce for any  $x \in P \cap \partial\Omega_1$  and  $t \in [0, 1]$

$$[x(t) - \lambda w(t)]^* \leq x(t) \leq \|x\| \leq R_1$$

and

$$\begin{aligned} x(t) - \lambda w(t) &\geq t^{\alpha-1} \|x\| - \lambda \sigma t^{\alpha-1} \int_0^1 r(s) \, ds = t^{\alpha-1} \left( R_1 - \lambda \sigma \int_0^1 r(s) \, ds \right) \\ &\geq t^{\alpha-1} \left( R_1 - \lambda^* \sigma \int_0^1 r(s) \, ds \right) \geq t^{\alpha-1} \left( R_1 - \sigma \int_0^1 r(s) \, ds \right) \geq 0. \end{aligned}$$

Then for any  $x \in P \cap \partial\Omega_1$  and  $t \in [0, 1]$ , we obtain

$$\begin{aligned} Tx(t) &\leq \lambda \int_0^1 J(s) (z(s)g(s, [x(s) - \lambda w(s)]^*) + r(s)) \, ds \\ &\leq \lambda^* M_1 \int_0^1 J(s) (z(s) + r(s)) \, ds \leq R_1 = \|x\|. \end{aligned}$$

Therefore, we conclude

$$\|Tx\| \leq \|x\| \quad \forall x \in P \cap \partial\Omega_1. \quad (10)$$

On the other hand, for  $c$  from (H3), we choose a positive constant  $L > 0$  such that

$$L \geq 2 \left( \lambda c^{2(\alpha-1)} \int_c^{1-c} J(s) \, ds \right)^{-1}.$$

By (H3), we deduce that there exists a constant  $M_0 > 0$  such that

$$f(t, u) \geq Lu \quad \forall t \in [c, 1-c], u \geq M_0. \quad (11)$$

Now we define  $R_2 = \max\{2R_1, 2M_0/c^{\alpha-1}\}$  and let  $\Omega_2 = \{x \in P: \|x\| < R_2\}$ .

Then for any  $x \in P \cap \partial\Omega_2$ , we obtain

$$\begin{aligned} x(t) - \lambda w(t) &\geq t^{\alpha-1} \|x\| - \lambda \sigma t^{\alpha-1} \int_0^1 r(s) \, ds \geq t^{\alpha-1} \left( R_2 - \sigma \int_0^1 r(s) \, ds \right) \\ &\geq t^{\alpha-1} \left( R_1 - \sigma \int_0^1 r(s) \, ds \right) \geq 0 \quad \forall t \in [0, 1]. \end{aligned}$$

Therefore, we conclude

$$\begin{aligned} [x(t) - \lambda w(t)]^* &= x(t) - \lambda w(t) \geq t^{\alpha-1} \left( R_2 - \sigma \int_0^1 r(s) \, ds \right) \\ &\geq \frac{c^{\alpha-1} R_2}{2} \geq M_0 \quad \forall t \in [c, 1-c]. \end{aligned} \quad (12)$$

Then for any  $x \in P \cap \partial\Omega_2$  and  $t \in [c, 1 - c]$ , by (11) and (12) we deduce

$$\begin{aligned} Tx(t) &\geq \lambda \int_c^{1-c} G(t, s)(f(s, [x(s) - \lambda w(s)]^*) + r(s)) \, ds \\ &\geq \lambda \int_c^{1-c} G(t, s)L[x(s) - \lambda w(s)] \, ds \geq \lambda L \int_c^{1-c} t^{\alpha-1} J(s) \frac{c^{\alpha-1} R_2}{2} \, ds \\ &\geq \frac{\lambda L c^{2(\alpha-1)} R_2}{2} \int_c^{1-c} J(s) \, ds \geq R_2 = \|x\|. \end{aligned}$$

Then

$$\|Tx\| \geq \|x\| \quad \forall x \in P \cap \partial\Omega_2. \tag{13}$$

By (10), (13) and Theorem 1(i) we conclude that  $T$  has a fixed point  $x_1 \in P \cap (\bar{\Omega}_2 \setminus \Omega_1)$ , that is  $R_1 \leq \|x_1\| \leq R_2$ . Since  $\|x_1\| \geq R_1$ , we deduce

$$\begin{aligned} x_1(t) - \lambda w(t) &\geq t^{\alpha-1} \left( \|x_1\| - \sigma \lambda \int_0^1 r(s) \, ds \right) \geq t^{\alpha-1} \left( R_1 - \sigma \int_0^1 r(s) \, ds \right) \\ &= A_1 t^{\alpha-1}, \end{aligned}$$

and so  $x_1(t) \geq \lambda w(t) + A_1 t^{\alpha-1}$  for all  $t \in [0, 1]$ , where  $A_1 = R_1 - \sigma \int_0^1 r(s) \, ds > 0$ .

Let  $u_1(t) = x_1(t) - \lambda w(t)$  for all  $t \in [0, 1]$ . Then  $u_1$  is a positive solution of problem (E)–(BC) with  $u_1(t) \geq A_1 t^{\alpha-1}$  for all  $t \in [0, 1]$ . This completes the proof of Theorem 2.  $\square$

**Theorem 3.** Assume that (H1), (H2) and (H4) hold. Then there exists  $\lambda_* > 0$  such that, for any  $\lambda \geq \lambda_*$ , the boundary value problem (E)–(BC) has at least one positive solution.

*Proof.* By (H4) there exists  $M_2 > 0$  such that

$$f(t, u) \geq 2\sigma \int_0^1 r(s) \, ds \left( c^{\alpha-1} \int_c^{1-c} J(s) \, ds \right)^{-1} \quad \forall t \in [c, 1 - c], u \geq M_2.$$

We define

$$\lambda_* = M_2 \left( c^{\alpha-1} \sigma \int_0^1 r(s) \, ds \right)^{-1}.$$

We assume now  $\lambda \geq \lambda_*$ . Let  $R_3 = 2\lambda\sigma \int_0^1 r(s) ds$  and  $\Omega_3 = \{x \in P: \|x\| < R_3\}$ . Then for any  $x \in P \cap \partial\Omega_3$ , we deduce

$$\begin{aligned} x(t) - \lambda w(t) &\geq t^{\alpha-1}\|x\| - \lambda\sigma t^{\alpha-1} \int_0^1 r(s) ds = t^{\alpha-1} \left( R_3 - \lambda\sigma \int_0^1 r(s) ds \right) \\ &= t^{\alpha-1} \lambda\sigma \int_0^1 r(s) ds \geq t^{\alpha-1} \lambda_* \sigma \int_0^1 r(s) ds \\ &= \frac{M_2 t^{\alpha-1}}{c^{\alpha-1}} \geq 0 \quad \forall t \in [0, 1]. \end{aligned}$$

Therefore, for any  $x \in P \cap \partial\Omega_3$  and  $t \in [c, 1-c]$ , we have

$$[x(t) - \lambda w(t)]^* = x(t) - \lambda w(t) \geq \frac{M_2 t^{\alpha-1}}{c^{\alpha-1}} \geq M_2.$$

Hence, for any  $x \in P \cap \partial\Omega_3$  and  $t \in [c, 1-c]$ , we conclude

$$\begin{aligned} Tx(t) &\geq \lambda \int_c^{1-c} G(t,s) f(s, [x(s) - \lambda w(s)]^*) ds \\ &\geq \lambda \int_c^{1-c} t^{\alpha-1} J(s) f(s, u(s) - \lambda w(s)) ds \\ &\geq \lambda \int_c^{1-c} t^{\alpha-1} J(s) \left( 2\sigma \int_0^1 r(s) ds \right) \left( c^{\alpha-1} \int_c^{1-c} J(s) ds \right)^{-1} ds \\ &= \frac{R_3 t^{\alpha-1}}{c^{\alpha-1}} \geq R_3. \end{aligned}$$

Therefore, we obtain

$$\|Tx\| \geq \|x\| \quad \forall x \in P \cap \partial\Omega_3. \quad (14)$$

On the other hand, we consider the positive number  $\varepsilon = (2\lambda \int_0^1 J(s)z(s) ds)^{-1}$ . Then by (H4) we deduce that there exists  $M_3 > 0$  such that  $g(t, u) \leq \varepsilon u$  for all  $t \in [0, 1]$ ,  $u \geq M_3$ . Therefore, we obtain  $g(t, u) \leq M_4 + \varepsilon u$  for all  $t \in [0, 1]$ ,  $u \geq 0$ , where  $M_4 = \max_{t \in [0, 1], u \in [0, M_3]} g(t, u)$ .

We define now

$$R_4 > \max \left\{ R_3, 2\lambda \max\{M_4, 1\} \int_0^1 J(s)(z(s) + r(s)) ds \right\}$$

and  $\Omega_4 = \{x \in P: \|x\| < R_4\}$ .

Then for any  $x \in P \cap \partial\Omega_4$ , we have

$$\begin{aligned} x(t) - \lambda w(t) &\geq t^{\alpha-1} \|x\| - \lambda \sigma t^{\alpha-1} \int_0^1 r(s) \, ds = t^{\alpha-1} \left( R_4 - \lambda \sigma \int_0^1 r(s) \, ds \right) \\ &\geq t^{\alpha-1} \left( R_3 - \lambda \sigma \int_0^1 r(s) \, ds \right) = t^{\alpha-1} \lambda \sigma \int_0^1 r(s) \, ds \\ &\geq t^{\alpha-1} \lambda_* \sigma \int_0^1 r(s) \, ds = \frac{M_2 t^{\alpha-1}}{c^{\alpha-1}} \geq 0 \quad \forall t \in [0, 1]. \end{aligned}$$

Then for any  $x \in P \cap \partial\Omega_4$ , we obtain

$$\begin{aligned} Tx(t) &\leq \lambda \int_0^1 J(s) [z(s)g(s, [x(s) - \lambda w(s)]^*) + r(s)] \, ds \\ &\leq \lambda \int_0^1 J(s) [z(s)(M_4 + \varepsilon(x(s) - \lambda w(s))) + r(s)] \, ds \\ &\leq \lambda \max\{M_4, 1\} \int_0^1 J(s) (z(s) + r(s)) \, ds + \lambda \varepsilon R_4 \int_0^1 J(s) z(s) \, ds \\ &\leq \frac{R_4}{2} + \frac{R_4}{2} = R_4 = \|x\| \quad \forall t \in [0, 1]. \end{aligned}$$

Therefore,

$$\|Tx\| \leq \|x\| \quad \forall x \in P \cap \partial\Omega_4. \quad (15)$$

By (14), (15) and Theorem 1(ii) we conclude that  $T$  has a fixed point  $x_1 \in P \cap (\bar{\Omega}_4 \setminus \Omega_3)$ , so  $R_3 \leq \|x_1\| \leq R_4$ .

In addition, we deduce that for all  $t \in [0, 1]$ ,

$$\begin{aligned} x_1(t) - \lambda w(t) &\geq x_1(t) - \lambda \sigma t^{\alpha-1} \int_0^1 r(s) \, ds \geq t^{\alpha-1} \|x_1\| - \lambda \sigma t^{\alpha-1} \int_0^1 r(s) \, ds \\ &\geq t^{\alpha-1} R_3 - \lambda \sigma t^{\alpha-1} \int_0^1 r(s) \, ds = \lambda \sigma t^{\alpha-1} \int_0^1 r(s) \, ds \\ &\geq \lambda_* \sigma t^{\alpha-1} \int_0^1 r(s) \, ds = \frac{M_2 t^{\alpha-1}}{c^{\alpha-1}}. \end{aligned}$$

Let  $u_1(t) = x_1(t) - \lambda w(t)$  for all  $t \in [0, 1]$ . Then  $u_1(t) \geq \tilde{\Lambda}_1 t^{\alpha-1}$  for all  $t \in [0, 1]$ , where  $\tilde{\Lambda}_1 = M_2/c^{\alpha-1}$ . Hence, we conclude that  $u_1$  is a positive solution of problem (E)–(BC), which completes the proof of Theorem 3.  $\square$

In a similar manner to that used in the proof of Theorem 3, we deduce the following result.

**Theorem 4.** Assume that (H1), (H2) and

(H4) There exists  $c \in (0, 1/2)$  such that  $\hat{f}_\infty = \lim_{u \rightarrow \infty} \min_{t \in [c, 1-c]} f(t, u) = \infty$  and  $g_\infty = \lim_{u \rightarrow \infty} \max_{t \in [0, 1]} g(t, u)/u = 0$

hold. Then there exists  $\tilde{\lambda}_* > 0$  such that, for any  $\lambda \geq \tilde{\lambda}_*$ , the boundary value problem (E)–(BC) has at least one positive solution.

## 4 Examples

Let  $\alpha = 10/3$  ( $n = 4$ ),  $p = 3/2$ ,  $q = 4/3$ ,  $m = 3$ ,  $\xi_1 = 1/4$ ,  $\xi_2 = 1/2$ ,  $\xi_3 = 3/4$ ,  $a_1 = 1$ ,  $a_2 = 1/2$ ,  $a_3 = 1/3$ .

We consider the fractional differential equation

$$D_{0+}^{10/3} u(t) + \lambda f(t, u(t)) = 0, \quad t \in (0, 1), \quad (\text{E}^*)$$

with the boundary conditions

$$\begin{aligned} u(0) = u'(0) = u''(0) = 0, \\ D_{0+}^{3/2} u(t)|_{t=1} = D_{0+}^{4/3} u(t)|_{t=1/4} + \frac{1}{2} D_{0+}^{4/3} u(t)|_{t=1/2} + \frac{1}{3} D_{0+}^{4/3} u(t)|_{t=3/4}. \end{aligned} \quad (\text{BC}^*)$$

Then we obtain  $\Delta = \Gamma(10/3)(1/\Gamma(11/6) - 3/4) \approx 0.86980822$ . So assumption (H1) is satisfied. Besides, we deduce

$$g_1(t, s) = \frac{1}{\Gamma(10/3)} \begin{cases} t^{7/3}(1-s)^{5/6} - (t-s)^{7/3}, & 0 \leq s \leq t \leq 1, \\ t^{7/3}(1-s)^{5/6}, & 0 \leq t \leq s \leq 1, \end{cases}$$

$$g_2(t, s) = \begin{cases} t(1-s)^{5/6} - t + s, & 0 \leq s \leq t \leq 1, \\ t(1-s)^{5/6}, & 0 \leq t \leq s \leq 1, \end{cases}$$

$$G(t, s) = g_1(t, s) + \frac{t^{7/3}}{\Delta} \left( g_2\left(\frac{1}{4}, s\right) + \frac{1}{2} g_2\left(\frac{1}{2}, s\right) + \frac{1}{3} g_2\left(\frac{3}{4}, s\right) \right)$$

for all  $t, s \in [0, 1]$ .

We also obtain  $h_1(s) = (1/\Gamma(10/3))(1-s)^{5/6}(1-(1-s)^{3/2})$  for all  $s \in [0, 1]$ ,  $\sigma = 1/\Gamma(10/3) + 3/(4\Delta) \approx 1.22220971$  and

$$J(s) = \begin{cases} h_1(s) + \frac{1}{\Delta} \left[ \frac{3}{4}(1-s)^{5/6} + \frac{22s-9}{12} \right], & 0 \leq s < \frac{1}{4}, \\ h_1(s) + \frac{1}{\Delta} \left[ \frac{3}{4}(1-s)^{5/6} + \frac{5s-3}{6} \right], & \frac{1}{4} \leq s < \frac{1}{2}, \\ h_1(s) + \frac{1}{\Delta} \left[ \frac{3}{4}(1-s)^{5/6} + \frac{4s-3}{12} \right], & \frac{1}{2} \leq s < \frac{3}{4}, \\ h_1(s) + \frac{3}{4\Delta}(1-s)^{5/6}, & \frac{3}{4} \leq s \leq 1. \end{cases}$$

*Example 1.* We consider the function

$$f(t, u) = \frac{u^2 + u + 1}{\sqrt[3]{t(1-t)^2}} + \ln t, \quad t \in (0, 1), \quad u \geq 0.$$

We have  $r(t) = -\ln t$  and  $z(t) = 1/\sqrt[3]{t(1-t)^2}$  for all  $t \in (0, 1)$ ,  $g(t, u) = u^2 + u + 1$  for all  $t \in [0, 1]$  and  $u \geq 0$ ,  $\int_0^1 r(t) dt = 1$ ,  $\int_0^1 z(t) dt = \Gamma(2/3)\Gamma(1/3) \approx 3.63$ . Therefore, assumption (H2) is satisfied. In addition, for  $c \in (0, 1/2)$  fixed, assumption (H3) is also satisfied ( $f_\infty = \infty$ ).

After some computations, we deduce that  $\int_0^1 J(s)(z(s) + r(s)) ds \approx 1.12036124$ . We choose  $R_1 = 2$  ( $R_1 > \sigma \int_0^1 r(s) ds$ ), and then we obtain  $M_1 = 7$  and  $\lambda^* \approx 0.255$ . By Theorem 2 we conclude that problem (E\*)–(BC\*) has at least one positive solution for any  $\lambda \in (0, \lambda^*]$ .

*Example 2.* We consider the function

$$f(t, u) = \frac{\sqrt{u+1}}{\sqrt[4]{t^3(1-t)}} - \frac{1}{\sqrt{t}}, \quad t \in (0, 1), \quad u \geq 0.$$

Here we have  $r(t) = 1/\sqrt{t}$  and  $z(t) = 1/\sqrt[4]{t^3(1-t)}$  for all  $t \in (0, 1)$ ,  $g(t, u) = \sqrt{u+1}$  for all  $t \in [0, 1]$  and  $u \geq 0$ . For  $c \in (0, 1/2)$  fixed, assumptions (H2) and (H4) are satisfied ( $\int_0^1 r(t) dt = 2$ ,  $\int_0^1 z(t) dt \approx 4.44$ ,  $\lim_{u \rightarrow \infty} \min_{t \in [c, 1-c]} f(t, u) = \infty$  and  $g_\infty = 0$ ).

For  $c = 1/4$ , we obtain  $\int_{1/4}^{3/4} J(s) ds \approx 0.23472325$  and  $L_0 \approx 529.001$ . From the proof of Theorem 3 we deduce that  $M_2 \approx 91277.5$  and  $\lambda_* \approx 948407$ . Then by Theorem 3 we conclude that, for any  $\lambda \geq \lambda_*$ , our problem (E\*)–(BC\*) has at least one positive solution.

## References

1. J. Caballero, I. Cabrera, K. Sadarangani, Positive solutions of nonlinear fractional differential equations with integral boundary value conditions, *Abstr. Appl. Anal.*, **2012**:303545, 2012.
2. S. Das, *Functional Fractional Calculus for System Identification and Controls*, Springer, New York, 2008.

3. J.R. Graef, L. Kong, Q. Kong, M. Wang, Uniqueness of positive solutions of fractional boundary value problems with non-homogeneous integral boundary conditions, *Fract. Calc. Appl. Anal.*, **15**(3):509–528, 2012.
4. D. Guo, V. Lakshmikantham, *Nonlinear Problems in Abstract Cones*, Academic Press, New York, 1988.
5. J. Henderson, R. Luca, *Boundary Value Problems for Systems of Differential, Difference and Fractional Equations. Positive solutions*, Elsevier, Amsterdam, 2016.
6. J. Henderson, R. Luca, On a system of Riemann–Liouville fractional boundary value problems, *Commun. Appl. Nonlinear Anal.*, **23**(2):1–19, 2016.
7. Y. Jia, X. Zhang, Positive solutions for a class of fractional differential equation multi-point boundary value problems with changing sign nonlinearity, *J. Appl. Math. Comput.*, **47**:15–31, 2015.
8. A.A. Kilbas, H.M. Srivastava, J.J. Trujillo, *Theory and Applications of Fractional Differential Equations*, North-Holland Math. Stud., Vol. 204, Elsevier, Amsterdam, 2006.
9. I. Podlubny, *Fractional Differential Equations*, Academic Press, San Diego, CA, 1999.
10. J. Sabatier, O.P. Agrawal, J.A.T. Machado (Eds.), *Advances in Fractional Calculus: Theoretical Developments and Applications in Physics and Engineering*, Springer, Dordrecht, 2007.
11. H. Salen, On the fractional order  $m$ -point boundary value problem in reflexive Banach spaces and weak topologies, *J. Comput. Appl. Math.*, **224**:565–572, 2009.
12. S.G. Samko, A.A. Kilbas, O.I. Marichev, *Fractional Integrals and Derivatives. Theory and Applications*, Gordon and Breach, Yverdon, 1993.
13. Y. Wang, L. Liu, Y. Wu, Positive solutions for a nonlocal fractional differential equation, *Nonlinear Anal., Theory Methods Appl., Ser. A, Theory Methods*, **74**(11):3599–3605, 2011.
14. C. Yuan, Multiple positive solutions for  $(n - 1, 1)$ -type semipositone conjugate boundary value problems of nonlinear fractional differential equations, *Electron. J. Qual. Theory Differ. Equ.*, **2010**:36, 2010.
15. C. Yuan, Two positive solutions for  $(n - 1, 1)$ -type semipositone integral boundary value problems for coupled systems of nonlinear fractional differential equations, *Commun. Nonlinear Sci. Numer. Simul.*, **17**(2):930–942, 2012.