

IL NUOVO CIMENTO  
DOI 10.1393/ncc/i2014-11718-2

VOL. 37 C, N. 1

Gennaio-Febbraio 2014

COLLOQUIA: IFAE 2013

## Measurement of the $\tau$ lepton electric dipole moment at BaBar

S. MARTELOTTI

*Dipartimento di Fisica "E. Amaldi", Università di Roma Tre  
Via della Vasca Navale 84, 00146 Roma, Italy and  
INFN, Laboratori Nazionali di Frascati  
Via Enrico Fermi 40, 00044 Frascati (RM), Italy*

ricevuto l'1 Ottobre 2013

**Summary.** — The search for a  $CP$  violation signature arising from an electric dipole moment of the  $\tau$  lepton in the  $e^+e^- \rightarrow \tau^+\tau^-$  reaction is currently in progress using  $470 \text{ fb}^{-1}$  of data collected with the BaBar detector at the PEP II collider from 1999 to 2008. In this paper the EDM search analysis method will be illustrated.

PACS 14.60.Fg – Taus.

PACS 31.30.jn – Electric dipole moments.

PACS 13.40.Gp – Form factors electromagnetic.

The SM prediction for  $CP$  violation in the lepton sector is negligibly small and has never been observed so far. However, the presence of physics beyond the SM could introduce these effects at experimentally accessible levels [1], making any observation of  $CP$  violation in this sector a clear evidence of new physics. In some models such effects are expected to be enhanced for the  $\tau$  leptons due to its very large mass compared to other leptons, since new bosons and Higgs strongly couple with heavy particles through quantum loop effects. The contribution from  $CP$  violating interactions in the  $\tau$  pair production process can be parametrized at the leading order, in a model-independent way, using an electric dipole moment (EDM) of the  $\tau$  lepton,  $d_\tau$ . Some theoretical SM predictions are given in [2, 3]. Presently, the best bound on  $d_\tau$  is from the Belle collaboration [4]. The Belle EDM analysis was performed using  $29.5 \text{ fb}^{-1}$  of data, finding 95% confidence level limits of  $-2.2 < \text{Re}(d_\tau) < 4.5(10^{-17} \text{ e cm})$  and  $-2.5 < \text{Im}(d_\tau) < 0.8(10^{-17} \text{ e cm})$ . The analysis discussed in the present paper will be based on all the data recorded by the BaBar detector [5] at the PEP-II asymmetric-energy  $e^+e^-$  storage rings operated at the SLAC National Accelerator Laboratory. An integrated luminosity of about  $470 \text{ fb}^{-1}$  was collected.

Taking into account the EDM contribution in the Lagrangian, the cross section of the process:  $e^+(\vec{p})e^-(-\vec{p}) \rightarrow \tau^+(\vec{k}, \vec{S}_+)\tau^-(-\vec{k}, \vec{S}_-)$ , where  $\vec{p}$  is the momentum vector of  $e^+$ ,  $\vec{k}$  is the momentum vector of the  $\tau^+$  and  $\vec{S}_\pm$  are the spin vectors for  $\tau^\pm$ , all expressed in the center-of-mass frame, is proportional to the squared spin density matrix [6]:  $\mathcal{M}_{\text{prod}}^2 =$




Fig. 1. – Real observable distribution for  $\tau^+\tau^- \rightarrow \pi^+\pi^-\nu_\tau\bar{\nu}_\tau$  events. Dotted line is the background contribution.




Fig. 2. – Correlations between  $\langle \mathcal{O}_{\text{Re}} \rangle$  and  $\text{Re}(d_\tau)$  for the analysed MC sample.

$\mathcal{M}_{\text{SM}}^2(\vec{k}, \vec{S}_\pm) + \text{Re}(d_\tau)\mathcal{M}_{\text{Re}}^2(\vec{k}, \vec{S}_\pm) + \text{Im}(d_\tau)\mathcal{M}_{\text{Im}}^2(\vec{k}, \vec{S}_\pm) + |d_\tau|^2\mathcal{M}_{d^2}^2(\vec{k}, \vec{S}_\pm)$ . In addition to the standard term, the interference terms between the SM and the  $CP$  violating amplitudes  $\mathcal{M}_{\text{Re}}^2$  and  $\mathcal{M}_{\text{Im}}^2$ , respectively related to the real and imaginary part of the  $d_\tau$ , appear. For the EDM measurement we adopt the same method of Belle, the so-called optimal observable method [7], which maximizes the sensitivity to  $d_\tau$ . The optimal observables are defined as

$$\mathcal{O}_{\text{Re}} = \frac{\mathcal{M}_{\text{Re}}^2}{\mathcal{M}_{\text{SM}}^2}, \quad \mathcal{O}_{\text{Im}} = \frac{\mathcal{M}_{\text{Im}}^2}{\mathcal{M}_{\text{SM}}^2}.$$

The mean values of these observables  $\langle \mathcal{O}_{\text{Re}} \rangle$  and  $\langle \mathcal{O}_{\text{Im}} \rangle$  are linear functions of  $d_\tau$ :  $\langle \mathcal{O}_{\text{Re}} \rangle = a_{\text{Re}} \cdot \text{Re}(d_\tau) + b_{\text{Re}}$ ,  $\langle \mathcal{O}_{\text{Im}} \rangle = a_{\text{Im}} \cdot \text{Im}(d_\tau) + b_{\text{Im}}$ .

In order to extract the value of  $d_\tau$  from the mean values of the observables measured on the data, we have to know the coefficients  $a_j$  and the offsets  $b_j$ . The parameters  $a_j$  and  $b_j$  are estimated from the correlations between  $\langle \mathcal{O}_{\text{Re}} \rangle$  ( $\langle \mathcal{O}_{\text{Im}} \rangle$ ) and  $\text{Re}(d_\tau)$  ( $\text{Im}(d_\tau)$ ) extracted by a full Monte Carlo (MC) simulation including the detector simulation with acceptance effects and event selection efficiency. In the MC different EDM values are introduced. With a  $30\text{fb}^{-1}$  MC sample the final state in which both  $\tau$ 's decay hadronically into

$\tau^\pm \rightarrow \pi^\pm \nu$  has been analysed so far. Distribution of the real optimal observable for this channel is shown in fig. 1. Dotted line is the background contribution, with a preliminary event selection the purity of the sample is of about 50%. In fig. 2 the correlations between the real observable means  $\langle O_{\text{Re}} \rangle$  and  $\text{Re}(d_\tau)$  of this MC sample is shown. By fitting the linear correlation plot of fig. 2, the parameters  $a_{\text{Re}}$  and  $b_{\text{Re}}$  are obtained. The slope  $a_{\text{Re}}$  represents the real EDM sensitivity, the offset  $b_{\text{Re}}$  represent the difference from zero of the observable mean when  $d_\tau = 0$ . Sample purity is under improvement with a BDT multivariate analysis and the EDM sensitivity evaluation with the full MC sample is in progress.

## REFERENCES

- [1] BERNREUTHER W. *et al.*, *Phys. Lett. B*, **391** (1997) 413. *ibid.*, **412** (1997) 425.
- [2] HUANG T., LU W. and TAO Z., *Phys. Rev. D*, **55** (1997) 1643.
- [3] BERNREUTHER W. and NACHTMANN O., *Phys. Rev. Lett.*, **63** (1989) 2787.
- [4] INAMI K. *et al.* (BELLE COLLABORATION), *Phys. Lett. B*, **551** (2003) 16.
- [5] AUBERT B. *et al.* (BABAR COLLABORATION), *Nucl. Instrum. Methods A*, **479** (2002) 1.
- [6] BERNREUTHER W., NACHTMANN O. and OVERMANN P., *Phys. Rev. D*, **48** (1993) 78.
- [7] ATWOOD D. and SONI A., *Phys. Rev. D*, **45** (1992) 2405.