

Measurement of ZZ production cross section and limits on anomalous triple gauge couplings with the ATLAS detector

A. MENGARELLI(*)

Università di Bologna e INFN, Sezione di Bologna - Bologna, Italy

ricevuto l'1 Ottobre 2013

Summary. — The measurement of the ZZ production cross section performed by the ATLAS detector in LHC proton-proton collisions at $\sqrt{s} = 7$ TeV is discussed. The results are based on an integrated luminosity of 4.6 fb^{-1} collected by ATLAS in 2011 with a fully operational detector and stable beam conditions. The normalized differential cross sections in bins of various kinematic variables together with limits on ZZZ and $ZZ\gamma$ anomalous triple gauge couplings derived using the transverse momentum of the leading Z boson are also presented.

PACS 13.38.Dg – Decays of Z bosons.

1. – Introduction

In several studies, processes involving the electroweak sector of the SM have received much attention as a strong test of the model by the fact that these interactions are intimately related to the gauge group of the model. Among them, Z -boson pair production cross section measurement has a particular importance since it plays a crucial role also in the Higgs boson study. In fact, ZZ decaying in the four charged leptons channel ($ZZ \rightarrow l^+l^-l^+l^-$) is an irreducible background that surrounds the emergence of the Higgs boson events.

Another fundamental aspect carried by interactions involving pair of Z bosons, is the sensitivity to anomalous triple gauge couplings (nTGC). In particular, since a direct coupling of three neutrally charged gauge bosons is forbidden in the SM, a deviation of sensitive parameters from the SM prediction would provide important information about new physics beyond it [1].

(*) E-mail: Alberto.Mengarelli@bo.infn.it




Fig. 1. – Anomalous nTGC 95% confidence intervals from LHC, LEP [1] and Tevatron [5] experiments. Integrated luminosities, centre-of-mass energy and cut-off Λ for each experiment are shown.

2. – Analysis description and results

Events are selected in two channels [2] $ZZ \rightarrow l^+l^-l^+l^-$ and $ZZ \rightarrow l^+l^-\nu\nu$. First, a fiducial cross section is measured to reduce systematic uncertainties then the total cross section is calculated using a fiducial acceptance for total phase space extrapolation. Correction factors and fiducial acceptances to measure fiducial and total cross sections are estimated using NLO POWHEGBOX [3] generator and the $gg \rightarrow ZZ$ contribution is modelled with GG2ZZ [4] generator.

The total ZZ production cross section combining the $ZZ \rightarrow l^+l^-l^+l^-$ and $ZZ \rightarrow l^+l^-\nu\nu$ channels is determined to be

$$(1) \quad \sigma_{ZZ}^{\text{tot}} = 6.7 \pm 0.7(\text{stat.})_{-0.3}^{+0.4}(\text{syst.}) \pm 0.3(\text{lumi.}) \text{ (pb)}.$$

The result in eq. (1) is consistent with the NLO Standard Model prediction of $5.89_{-0.18}^{+0.22}$ pb, calculated with Z bosons with a mass between 66 and 116 GeV. Unfolded distributions of the fiducial cross sections are derived for the transverse momentum of the leading Z (p_T^Z), the angular difference between the two leptons forming the leading Z ($\Delta\phi(l, l)$), and the mass of the four leptons system (M_{4l}) are in agreement with SM predictions. The event yields as a function of the p_T^Z for the $ZZ \rightarrow l^+l^-l^+l^-$ and $ZZ \rightarrow l^+l^-\nu\nu$ selections are used to derive 95% confidence intervals for anomalous nTGC as shown in fig. 1.

REFERENCES

- [1] THE LEP COLLABORATIONS ALEPH, DELPHI, L3, OPAL and the LEP ELECTROWEAK WORKING GROUP, *A Combination of preliminary electroweak measurements and constraints on the standard model*, hep-ex/0612034.
- [2] THE ATLAS COLLABORATION, *JHEP*, **03** (2013) 128.
- [3] ALIOLI S., NASON P., OLEARI C. and RE E., *JHEP*, **1006** (2010) 043.
- [4] BINOTH T., KAUER N. and MERTSCH P., *Gluon-induced QCD corrections to $pp \rightarrow ZZ \rightarrow l^+l^-l^+l^-$* , arXiv:0807.0024.
- [5] D0 COLLABORATION, *Phys. Rev. Lett.*, **100** (2008) 131801.