


# Using Surface Enhanced Raman Scattering to Analyze the Interactions of Protein Receptors with Bacterial Quorum Sensing Modulators

Celina Costas,<sup>†,‡</sup> Vanesa López-Puente,<sup>†,‡</sup> Gustavo Bodelón,<sup>\*,†</sup> Concepción González-Bello,<sup>‡</sup> Jorge Pérez-Juste,<sup>†</sup> Isabel Pastoriza-Santos,<sup>†</sup> and Luis M. Liz-Marzán<sup>\*,†,§,||</sup>

<sup>†</sup>Departamento de Química Física, Universidade de Vigo, 36301 Vigo, Spain, <sup>‡</sup>Centro Singular de Investigación en Química Biológica y Materiales Moleculares (CIQUS), Universidad de Santiago de Compostela, 15782 Santiago de Compostela, Spain, <sup>§</sup>Bionanoplasmonics Laboratory, CIC biomaGUNE, Paseo de Miramón 182, 20009 Donostia-San Sebastian, Spain, and <sup>||</sup>Ikerbasque, Basque Foundation for Science, 48013 Bilbao, Spain. <sup>#</sup>These authors contributed equally to this work.

**ABSTRACT** Many members of the LuxR family of quorum sensing (QS) transcriptional activators, including LasR of *Pseudomonas aeruginosa*, are believed to require appropriate acyl-homoserine lactone (acyl-HSL) ligands to fold into an active conformation. The failure to purify ligand-free LuxR homologues in nonaggregated form at the high concentrations required for their structural characterization has limited the understanding of the mechanisms by which QS receptors are activated. Surface-enhanced Raman scattering (SERS) is a vibrational



spectroscopy technique that can be applied to study proteins at extremely low concentrations in their active state. The high sensitivity of SERS has allowed us to detect molecular interactions between the ligand-binding domain of LasR (LasR<sub>LBD</sub>) as a soluble apoprotein and modulators of *P. aeruginosa* QS. We found that QS activators and inhibitors produce differential SERS fingerprints in LasR<sub>LBD</sub>, and in combination with molecular docking analysis provide insight into the relevant interaction mechanism. This study reveals signal-specific structural changes in LasR upon ligand binding, thereby confirming the applicability of SERS to analyze ligand-induced conformational changes in proteins.

**KEYWORDS:** nanoplasmonics · SERS · quorum sensing · protein receptors · *Pseudomonas aeruginosa*

Surface-enhanced Raman scattering (SERS) spectroscopy is an ultra-sensitive and ultra-rapid analytical technique that can achieve detection limits down to the single molecule level thanks to nanoplasmonic effects.<sup>1,2</sup> As a vibrational spectroscopy technique, SERS can provide detailed information about the structure and conformation of a molecular target, which has been used to identify molecules, proteins and their mutual interactions.<sup>3</sup> A major advantage of SERS is that it can be performed under physiological conditions, thereby preserving the active state of proteins and revealing protein kinetics and dynamics. SERS does not require high protein concentrations as is the case for NMR or X-ray crystallography and it can be applied without prior knowledge of the structure of the protein.<sup>4</sup> Hence, SERS has emerged as an attractive analytical tool to investigate protein–protein and protein–ligand interactions.<sup>5,6</sup>


In Gram-negative bacteria, quorum sensing (QS) is mostly achieved through the biosynthesis and sensing of *N*-acyl  $\ell$ -homoserine lactones (acyl-HSLs) by LuxI/R-type QS protein receptor systems.<sup>7,8</sup> In such QS circuits, LuxI-type synthases catalyze the production of acyl-HSLs that freely diffuse into and out of cells, where they bind LuxR-type proteins that serve as signal receptors. When acyl-HSLs reach a threshold concentration, the acyl-HSL-LuxR complex activates the expression of QS-dependent genes. Biochemical and structural studies have shown that LuxR homologues are homodimers, and each monomer folds into two functional domains: an amino-terminal ligand-binding domain (LBD) and a carboxyl-terminal DNA-binding domain (DBD), joined by a short linker region. The cognate ligand binds to a narrow cavity in the LBD, formed by a cluster of hydrophobic and aromatic residues, and it is believed that

\* Address correspondence to gbodelon@uvigo.es, lizmarzan@cicbiomagune.es.

Received for review March 25, 2015 and accepted April 30, 2015.

Published online April 30, 2015  
10.1021/acsnano.5b01800

© 2015 American Chemical Society



**Figure 1.** (A) Co-crystal structure of LasR<sub>LBD</sub> (green) bound to C12-HSL (red) (PDB identity code 2UV0). The amino (N-) and carboxy (C-)terminal regions of the polypeptide are indicated. (B) Schematic representation of full-length LasR and the corresponding LBD bearing a hexahistidine-tag (6×His) and a C-terminal cysteine (Cys). (C) SDS-PAGE and Coomassie blue staining of LasR<sub>LBD</sub> expression and purification. Lane 1, uninduced total fraction; lane 2, IPTG-induced total fraction; lane 3, IPTG-induced soluble fraction; lane 4, affinity chromatography flow-through; lane 5, eluted protein. (D) Cross-linking assay of LasR<sub>LBD</sub>. The polypeptide expressed in bacteria in the absence (lanes 1–3) or in the presence (lanes 4–6) of C12-HSL was solubilized (S), affinity purified and subjected to crosslinking (+) with 0.5 mM DSS (lanes 3,6) or not (lanes 2,5). Bands corresponding to dimeric (\*\*\*) and monomeric (\*) forms are indicated on the right.

this triggers a conformational change in the polypeptide, presumably enabling the binding of the protein to specific promoters, ultimately leading to transcriptional activation of QS genes.<sup>9,10</sup> Despite of the availability of a wealth of information, the precise molecular mechanisms of LuxR receptor activation still remain mostly unknown. This is largely due to the failure to purify the relevant regulators in non-aggregated form outside the cellular environment, in the absence of an appropriate acyl-HSL.

Much of the information regarding acyl-HSL-regulated QS systems has been obtained through the study of *Pseudomonas aeruginosa* bacteria. *P. aeruginosa* contains two pairs of LuxI/LuxR-type systems, termed LasI/LasR and RhII/RhIR. LasI and RhII are synthases that catalyze production of *N*-(3-oxododecanoyl)-homoserine lactone (C12-HSL) and *N*-(butyryl)-homoserine lactone (C4-HSL), which are sensed by the LasR and RhIR transcriptional receptors, respectively.<sup>11,12</sup> LasR is known to initiate the QS regulatory system in this bacterial species,<sup>13</sup> and therefore, it has attracted significant interest as a target for QS modulation.<sup>14–16</sup> The X-ray structure of the LBD of LasR forming a complex with its natural ligand C12-HSL<sup>17</sup> (Figure 1A) revealed a symmetric homodimer with the molecule buried in a pocket enclosed by  $\alpha$ -helices ( $\alpha$ 3,  $\alpha$ 4 and  $\alpha$ 5) and the concave side of the central  $\beta$ -sheet of each LBD.<sup>17,18</sup>

We demonstrate in this paper the application of SERS spectroscopy, combined with structural analysis, molecular docking studies and assays of biological activity, to detect molecular interactions between LasR and diverse activators and inhibitors of *P. aeruginosa* QS.

To this end the LBD of LasR (LasR<sub>LBD</sub>) bearing a hexahistidine (6×His) tag and a cysteine in its carboxy-terminus was expressed and purified in a soluble, ligand-free (*i.e.*, apoprotein) active form. Oriented attachment of the polypeptide onto plasmonic gold nanoparticle thin films and label-free SERS allowed us to detect structural changes of LasR<sub>LBD</sub> as a result of its interaction with cognate C12-HSL ligands, C4-HSL agonists and Furanone C30 or acetylsalicylic acid (SA) antagonists. Remarkably, highly sensitive and reproducible SERS spectra allowed us to discriminate between activators and inhibitors of QS, through distinctive vibrational features. Our study thus provides insight into the molecular interactions between LasR and modulators of *P. aeruginosa* QS, while featuring SERS as a powerful tool for the dynamic study and screening of protein–ligand interactions.

## RESULTS AND DISCUSSION

### Expression of Apo LasR<sub>LBD</sub> and Analysis of Functional Activity.

We schematically show in Figure 1 the lactone-binding domain (LBD) and the DNA-binding domain (DBD) of LasR. To carry out the SERS investigation of conformational changes in LasR upon ligand interaction, we focused on the LBD region of the protein as the DBD is not required for ligand binding activity.<sup>17,19</sup> Therefore, the LBD of *lasR* gene, coding from amino-acids encompassing Met-1 to Lys-173,<sup>17</sup> was PCR amplified from the pMHLAS plasmid<sup>20</sup> and cloned into the pET21a(+) vector in frame with a 6×His tag for affinity purification and a C-terminal cysteine that is known to strongly bind onto gold surfaces through the

formation of Au–S bonds (Figure 1B). The resulting construct, termed pET21-LasR<sub>LB</sub>, was transformed in *Escherichia coli* BL21(DE3)pLysS cells for protein expression. *E. coli* bacteria have been widely used as suitable hosts to study the influence of different acyl-HSLs in the expression of LuxR homologues as they do not produce acyl-HSLs endogenously.<sup>9</sup>


The SERS study of the interactions of apo LasR<sub>LB</sub> with QS modulators required the expression and purification of the polypeptide in the absence of its cognate C12-HSL ligand. However, it has been shown that overexpression of recombinant LuxR homologues in *E. coli* without the appropriate signaling molecules in the bacterial growth medium renders these proteins insoluble and prone to degradation by cellular proteases.<sup>9,17</sup> Therefore, we first assessed the stability and solubility of the LasR<sub>LB</sub> polypeptide produced in *E. coli* BL21(DE3)pLysS bacteria grown without C12-HSL ligand (see Materials and Methods). Briefly, bacterial cultures in log phase (OD<sub>600</sub>–0.6) were induced with 0.5 mM isopropyl β-D-1-thiogalactopyranoside (IPTG) at 28 °C for 4 h and then total proteins and soluble protein fractions were separated on a 12% sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) under reducing conditions and stained with Coomassie Blue. As shown in Figure 1C, a protein band with the expected molecular weight corresponding to LasR<sub>LB</sub> (~19 kDa) can be seen in the total protein extract from IPTG-induced cells (Figure 1C, lane 2) but not in the total protein extract from uninduced cells (Figure 1C, lane 1). The expression of LasR<sub>LB</sub> was also confirmed by Western blot and immunostaining with anti-His tag antibodies (data not shown). Importantly, apo LasR<sub>LB</sub> was found in the soluble fraction (Figure 1C, lane 3), which allowed us to purify it upon elution from a nickel resin (Figure 1C, lane 5). Similar levels of apo LasR<sub>LB</sub> relative to the rest of proteins in the soluble fraction (Figure 1C, lane 3) and in the total protein extract (Figure 1C, lane 2), suggested that the polypeptide was highly soluble. The amount of soluble apo LasR<sub>LB</sub> (Figure 1D, lane 1) was similar to that obtained from bacteria grown in the presence of the C12-HSL ligand (Figure 1D, lane 4), again confirming that the polypeptide is stable and soluble in its apoprotein form. Importantly, increasing either the expression of apo LasR<sub>LB</sub> or its concentration resulted in decreased solubility (data not shown).

Dimerization of LasR, as well as other LuxR-type receptors, is a requisite for transcriptional activation of target QS genes.<sup>21,22</sup> The oligomeric state of apo LasR<sub>LB</sub> was assessed by chemical cross-linking with disuccinimidyl suberate (DSS), which is a commonly used strategy for the characterization of dimers of LuxR homologues.<sup>23–25</sup> To this end, affinity purified LasR<sub>LB</sub> obtained from IPTG-induced BL21(DE3)pLysS bacteria grown in the presence or in the absence of C12-HSL was cross-linked with DSS and analyzed by SDS-PAGE.


As shown in Figure 1D, cross-linked dimeric complexes of LasR<sub>LB</sub> were detected at similar levels regardless of the presence (lane 6) or absence (lane 3) of the ligand, indicating that LasR<sub>LB</sub> does exist in the form of homodimers even when no C12-HSL is present. We then investigated the activity of purified apo LasR<sub>LB</sub> as a quorum-quencher of C12-HSL molecules, by employing an *E. coli* MT102 biosensor strain harboring the pMHLAS reporter system that expresses green fluorescent protein (GFP) upon activation of LasR by exogenous addition of C12-HSL<sup>20</sup> (Supporting Information Figure S1). We hypothesized that functional LasR<sub>LB</sub> would sequester extracellular C12-HSLs, reducing the levels of free ligand that would be available for binding and activating intracellular LasR, resulting in decreased GFP expression (Supporting Information Figure S1). To test this hypothesis, C12-HSL was preincubated with either affinity purified LasR<sub>LB</sub> or BSA for 1 h, and the mixtures were subsequently added to liquid cultures of *E. coli* MT102 bacteria bearing pMHLAS and further incubated for 90 min to allow GFP expression. The expression of GFP was then evaluated by epifluorescence microscopy and quantified by fluorescence spectroscopy. As shown in Figure 2, the preincubation of C12-HSL (0.5 nmol) with apo LasR<sub>LB</sub> significantly reduced the levels of GFP whereas preincubation with BSA (0.5 nmol) as a control did not affect GFP expression. This indicated that apo LasR<sub>LB</sub> did indeed bind C12-HSLs, preventing full induction of the acyl-HSL reporter system.

A first conclusion of this study is thus related to the general belief that LasR and other LuxR family members require the presence of appropriate acyl-HSL molecules to properly fold into their active conformations, which is mostly based on the production of large amounts of soluble and stable protein upon supplementing the bacterial growth medium with cognate acyl-HSLs. This strategy has been widely applied toward the biochemical and structural characterization of several LuxR homologues including LasR,<sup>22</sup> TraR<sup>21,26</sup> or QscR.<sup>27</sup> This idea has been however challenged by several groups,<sup>28,29</sup> who provided experimental evidence demonstrating that LasR folds into an active conformation in the absence of C12-HSLs and can remain in a properly folded signal-free state. Our results indeed confirm these findings as we demonstrate that LasR<sub>LB</sub> can be produced as a soluble, dimeric and functional protein in extracts of bacteria that were grown in the absence of cognate C12-HSL ligands.

**LasR Modulation by QS Activators and Inhibitors.** C4-HSL is the natural signaling ligand of *P. aeruginosa* RhlR QS receptor, and it has been shown to slightly activate the expression of a LasR-dependent *lasB-lacZ* translational fusion in *E. coli*.<sup>30</sup> On the other hand, Furanone C30 and SA are synthetic compounds that inhibit LasR signaling,<sup>31,32</sup> presumably by interacting with the



**Figure 2.** (A) Fluorescence microscopy images of MT102/pMHLAS bacteria treated with C12-HSL previously incubated with either LasR<sub>LBD</sub> or BSA, as labeled. Upper and lower panels show DAPI staining of bacteria and GFP expression, respectively. (B) Induction of GFP expression from pMHLAS in *E. coli* MT102 after incubation of bacteria with C12-HSL or C12-HSL and LasR<sub>LBD</sub> polypeptide. Error bars represent standard deviation calculated for three independent experiments.




**Figure 3.** (A) Compounds used in LasR modulation studies: *N*-3-oxododecanoyl-homoserine lactone (C12-HSL), *N*-butyryl-homoserine lactone (C4-HSL, C4), acetylsalicylic acid (SA) and 4-bromo-5-(bromomethylene)-2(5*H*)-furanone (C30). (B) Evaluation of GFP expression from pMHLAS in *E. coli* MT102 bacteria upon incubation with the indicated amounts of C4, SA or C30 and subsequently treated without (white bars) or with (gray bars) C12-HSL. The mean from at least two independent experiments was normalized to control inductions (control) lacking C4, SA and C30.

acyl-HSL binding pocket of LasR as shown by molecular modeling studies.<sup>11,32,33</sup> We analyzed the activity of these LasR modulating compounds by means of the *E. coli* MT102 biosensor bearing pMHLAS. A liquid culture of *E. coli* containing pMHLAS plasmid was divided into several subcultures, and incubated with C4-HSL, Furanone C30 or SA at different concentrations for 20 min. Subsequently, they were either treated or untreated with C12-HSL for 90 min, as explained in the Materials and Methods section, and GFP expression was determined by fluorescence spectroscopy. As shown in Figure 3B, C4-HSL was capable to induce GFP expression, though to a lesser extent than C12-HSL, demonstrating an agonistic activity over LasR. On the other hand, SA and Furanone C30 displayed strong

inhibitory effects on GFP expression, while not affecting bacterial growth (not shown), which confirms their role as LasR antagonists.

**SERS Analysis of Apo LasR<sub>LBD</sub>.** The SERS substrate was a gold nanoparticle thin film fabricated using polyelectrolyte layer-by-layer (LbL) assembly, by means of alternate deposition of a positively charged polyelectrolyte, poly(diallyldimethylammonium chloride) (PDDA), and negatively charged, citrate-stabilized Au NPs (60 nm diameter), on a glass slide. The resulting LbL film comprising three bilayers (PDDA-Au NPs) displayed a long-range uniform Au deposition (Supporting Information Figure S2), displaying a broad extinction band centered at 850 nm due to significant plasmon coupling and a less intense band at around 550 nm from single particles.<sup>34</sup> Since nonspecific binding of LasR<sub>LBD</sub> to the Au nanoparticle surface would lead to nonhomogenous SERS signals, a cysteine amino acid was genetically introduced at the C-terminus of the polypeptide, immediately after the hexa-histidine tag. The high affinity toward Au-thiol binding has been widely employed for site-directed attachment of functional antibody fragments, enzymes or proteins on Au surfaces.<sup>35,36</sup> In addition, the hexa-histidine tag has been reported to bind the polypeptide to gold surfaces through metal-histidine coordination.<sup>37</sup> Thus, the attachment of LasR<sub>LBD</sub> onto the Au nanoparticle-based substrate should take place through its C-terminus, placing the polypeptide in the conformation illustrated in Figures 1A and 5. As shown in Figure 4, highly reproducible SERS spectra were obtained, most likely due to the targeted binding of LasR<sub>LBD</sub> to the gold surface. The main features of the SERS spectra were assigned according to the existing literature (Table 1). The SERS spectra (Figure 4) display the characteristic feature of Au–S vibrational mode at 386 cm<sup>-1</sup> confirming the specific interaction between the C-terminal cysteine and the Au surface.<sup>38</sup> In addition, the spectra show vibrational bands that could also be assigned to this amino acid (448, 690, and 788 cm<sup>-1</sup>) and to the histidine




**Figure 4.** SERS spectra of apo LasR<sub>LBD</sub> measured in three different regions of a gold nanoparticle-based substrate. Excitation with a 785 nm laser line was used to avoid protein damage.


**TABLE 1.** Experimental Vibrational Frequencies (cm<sup>-1</sup>) for LasR<sub>LBD</sub> with Tentative Assignments

LasR/cm <sup>-1</sup>	band assignment
386	Au–S
448	Cys
690	Cys
788	Cys
830	Tyr
944	Glu, Asp
1000	Phe
1028	Phe
1099	His, CαN
1142	CCαN
1178	Tyr, Phe
1260	Tyr
1325	Amide III
1439	His, Glu, Lys
1537	Tyr
1578	NH and NH <sub>3</sub> <sup>+</sup> def.

amino acid (1099, 1439, and 1578 cm<sup>-1</sup>) abundantly present in the hexa-histidine tag.<sup>39</sup> The targeted attachment of LasR<sub>LBD</sub> orients the L3 loop toward the Au surface (Figure 5). Therefore, the following vibrational bands can be ascribed to amino acids present in the L3 loop and in its vicinity, such as Tyr-47 (peaks at 830, 1178, 1260, and 1537 cm<sup>-1</sup>), Phe-167 (peaks at 1000, 1028, and 1178 cm<sup>-1</sup>), Asp-43, Asp-46 and Glu-48, 124, and 168 (peaks at 944 and 1439 cm<sup>-1</sup>).<sup>39–43</sup> Furthermore, the enhancement of modes corresponding to the asymmetric stretching of CCαN (1142 cm<sup>-1</sup>) suggests interactions of nitrogen atoms in histidine, arginine, lysine or glutamine amino-acids with the gold surface.<sup>39</sup> The intense peak at 1325 cm<sup>-1</sup> can be assigned to an amide III (in phase combination of the N–H bending and C–H stretching vibrations).<sup>39</sup> We confirmed that the SERS spectral peaks shown in Figure 4 were characteristic of LasR<sub>LBD</sub>, as neither the buffer nor the gold nanoparticle substrate produced significant Raman signals (data not shown). Importantly, SERS spectra recorded at 1, 3, and 6 h after binding the polypeptide to the gold substrate did not show any significant changes, thus proving that the protein is



**Figure 5.** Schematic representation of possible contact points between LasR<sub>LBD</sub> and a gold surface. Side (A) and front (B) view of LasR<sub>LBD</sub> (PDB identification code 2UV0). Highlighted are relevant structural features (α3, α4, α5, 3<sub>10</sub>-helix, L3 loop) and interacting amino acids: Lys-42 (light blue), Asp-43 (orange), Gln 45 (red), Asp-46 (orange), Tyr-47 (blue), Glu-48 (yellow), Phe-167 (magenta), and Glu-168 (yellow).



**Figure 6.** SERS spectra (after baseline correction) of LasR<sub>LBD</sub> polypeptide incubated with C12-HSL, C4-HSL, SA or Furanone C30. The green, red, and blue areas indicate common, inhibitor-specific, and activator-specific SERS fingerprints, respectively. Illumination with a 785 nm laser line was used to avoid protein damage.

stable under the assay conditions (Supporting Information Figure S3).

**Detection of Interactions between LasR<sub>LBD</sub> and QS Modulators by SERS.** SERS was then used to analyze whether incubation of apo LasR<sub>LBD</sub> with either its cognate C12-HSL or with agonist C4-HSL ligands can induce conformational changes in the polypeptide. We attached apo LasR<sub>LBD</sub> (ca. 0.5 nmol in 10 mM sodium phosphate buffer at pH 8.0, PB buffer) to the Au nanoparticle-based substrate, followed by incubation with C12-HSL or C4-HSL (100 μM). As shown in Figure 6,

the SERS spectra of LasR<sub>LBD</sub> forming a complex with either C12-HSL or C4-HSL, resemble that of ligand-free LasR<sub>LBD</sub> but display some remarkable differences: reduction in SERS intensity at 448, 690, and 788 cm<sup>-1</sup> corresponding to the C-terminal cysteine and appearance of additional peaks at 408, 1405 and 1630 cm<sup>-1</sup>. The peak at 408 cm<sup>-1</sup> was assigned to the deformation of the C–N vibrational mode,<sup>40</sup> whereas the 1630 cm<sup>-1</sup> band was assigned to an Amide I peak characteristic of a relatively uncommon secondary structure called the 3<sub>10</sub>-helix.<sup>44,45</sup> Interestingly, LasR<sub>LBD</sub> possesses a 3<sub>10</sub>-helix conformed by the amino acids from Gly-68 to Arg-71, which follows the L3 loop (Figure 5) and has been shown to participate in the lactone-binding pocket.<sup>17</sup> Finally, the band at 1405 cm<sup>-1</sup> was assigned to the COO<sup>-</sup> symmetric vibration of aspartic and glutamic amino acids, both present in the L3 loop and 3<sub>10</sub>-helix (Figure 5).<sup>43</sup> As neither C12-HSL nor C4-HSL or the PB buffer produced any significant SERS signals at the concentration used in the binding assay when tested separately (data not shown), we can state that, the observed peaks are protein-specific. Altogether, these results show that incubation of apo LasR<sub>LBD</sub> with both QS activators gives rise to changes in the vibrational SERS spectra of the polypeptide, thereby indicating conformational changes linked to ligand interaction.

Analysis of the interactions of LasR<sub>LBD</sub> with SA (100 μM) and Furanone C30 (10 μM) antagonists (Figure 6) showed changes in the SERS spectra, which were similar to those obtained with the acyl-HSLs, such as the appearance of the C–N stretching at 408 cm<sup>-1</sup> and the decrease of the cysteine peaks at 448, 690, and 788 cm<sup>-1</sup>. Additionally, the LasR antagonists induced a new peak at 1267 cm<sup>-1</sup>, assigned to C=O stretching vibrations of an α-helix,<sup>46</sup> which was not observed for C12-HSL or C4-HSL. This specific SERS signal could be attributed to movements of the α-helices α3, α4 and/or α5, which constitute the ligand-binding pocket (Figure 5). Importantly, the antagonists did not induce the 1405 and 1630 cm<sup>-1</sup> peaks associated with structural movements involving the L3 loop and the 3<sub>10</sub>-helix, thus confirming these SERS signals as signatures of the QS activators. SA was not detected by SERS on bare gold nanoparticle substrates at the concentration used in the binding assay (data not shown). On the other hand, Furanone C30 did yield strong SERS signals when analyzed separately (Supporting Information Figure S6). Interestingly, the SERS peaks corresponding to Furanone C30 itself were not detected upon incubation with LasR<sub>LBD</sub>, indicating that it did not reach the gold surface in the presence of the polypeptide, likely due to binding to LasR<sub>LBD</sub>. Finally, SERS measurements performed at 1, 3, and 6 h yielded similar spectra for each group of QS modulators (Supporting Information Figures S4 and S5) demonstrating the reliability of the data. These experiments also revealed that

**TABLE 2. SERS Fingerprints of LasR<sub>LBD</sub> upon Binding to C12-HSL or C4-HSL (QS Activators) and SA or Furanone C30 (QS Inhibitors)**


QS activators	QS inhibitors	assignments
408	408	δ(CN)
448	448	Cys
690	690	Cys
788	788	Cys
	1267	v(C=O) (α-helix)
1405		COO <sup>-</sup> sym. stretch.
1630		Amide I (3 <sub>10</sub> -helix)

conformational changes in LasR<sub>LBD</sub> were evident within the first hour of incubation. The results, summarized in Table 2, clearly show SERS detection of structural changes in LasR<sub>LBD</sub>, associated with interactions of both activators and inhibitors of QS, and that each group of QS modulators produced specific SERS fingerprints.

#### Molecular Docking analysis of LasR<sub>LBD</sub>–Ligand Interactions.

Molecular docking analysis was used to study the binding mode of C4-HSL, SA and Furanone C30 to LasR<sub>LBD</sub>, so as to obtain an insight into their functional roles as QS modulators. These studies were carried out using the program GOLD (version 5.2) and the protein geometries found in the crystal structure of the LasR<sub>LBD</sub>/C12-HSL binary complex (PDB identification code 2UV0). C12-HSL was initially docked as a control and the result compared to the PDB entry 2UV0. The efficacy of a compound to activate LasR was directly correlated with its ability to maintain the binding interactions observed in the LasR<sub>LBD</sub>/C12-HSL binary complex (Figure 7A). As illustrated in Figure 7B, the binding mode of the agonist C4-HSL is similar to that of the natural ligand as it retains most of the binding interactions of the homoserine lactone and amide moieties of C12-HSL, including four intermolecular hydrogen bonds involving Tyr-56, Trp-60, Asp-73 and Ser-129, CH–II interactions with Trp-88 and a lipophilic contact with Ala-105. Interestingly, despite of the different lengths of their acyl chains, the two QS activators did not show any significant distinct SERS signals. This suggests that the molecular interactions of C12-HSL and C4-HSL with the recognition center of LasR<sub>LBD</sub> detected by SERS mainly depend on the homoserine lactone moieties. On the other hand, it has been proposed that exposure of buried hydrophobic residues in the lactone-binding pocket to the bulk solvent could be responsible for protein instability and degradation within the cell.<sup>18</sup> Therefore, as C4-HSL lacks a full-length acyl chain it cannot meet most of the hydrophobic interactions established by C12-HSL (*cf.* Figures 7A and 7B), thus explaining why the noncognate C4-HSL molecule is not able to sustain full activation of LasR (Figure 3B).

Regarding LasR antagonists, molecular docking analysis revealed that both SA (Figure 7C) and Furanone



**Figure 7.** Modeling of QS ligand interactions with LasR<sub>LBD</sub>. (A) Binding mode of C12-HSL (yellow) in the recognition center of LasR receptor (gray). Hydrogen bonding (red) and lipophilic interactions (black) are shown. (B) Comparison between the predicted binding mode of C4-HSL (purple) and the natural autoinducer C12-HSL (yellow). (C) Plausible binding mode of SA (light orange). (D) Predicted binding mode of Furanone C30 (green).

C30 (Figure 7D) would establish similar polar contacts with the lactone-binding pocket of LasR<sub>LBD</sub>, involving Tyr-56, Trp-60, Thr-75 and Arg-61 residues. Small differences were however observed in the lipophilic contacts of both ligands, in particular involving Leu-110, Val-76 and Ala-127 residues. The shared topology may be responsible for inducing similar SERS spectra as experimentally observed (Figure 6). Although each compound (Figure 7C,D) seemed to establish some of the strongest contacts predicted for the homoserine lactone moieties (Figure 7B), the type of chemical interactions and their respective strengths were different (Supporting Information Table S1), which may explain their inhibitory effect on LasR signaling (Figure 3B). We speculate that SA and Furanone C30 do not establish contacts with key amino-acid residues in LasR<sub>LBD</sub>, thus failing to properly stabilize the polypeptide, which would result in an accelerated proteolytic degradation. Indeed, it has been reported that the half-life of LuxR protein of *Vibrio fischeri* is reduced up to 100-fold in the presence of Furanone C30, suggesting that furanones control expression of acyl-HSL-dependent QS by destabilizing LuxR.<sup>31</sup> Muh and co-workers used ultrahigh-throughput screening to search for inhibitors in a library of 200 000 compounds. The screening identified triphenyl (TP) compounds that are structurally different from acyl-HSLs, bearing either agonist or antagonist activity of the LasR-dependent fluorescent reporter. *In silico* docking studies suggested

that hydrogen bond interactions of these molecules with Trp-60 and Asp-73 influenced whether the TP compounds act as an agonist or an antagonist.<sup>9</sup> The authors indicated that only those molecules establishing contacts with both Trp-60 and Asp-73 function as agonists. Interestingly, our modeling study showed that whereas C12-HSL and C4-HSL were predicted to maintain interactions with Trp-60 and Asp-73, SA and Furanone C30 were predicted to interact with Trp-60 only (Figure 7). In a follow-up study, Zou and Nair obtained crystal structures of LasR<sub>LBD</sub> complexing TP compounds of agonistic effect in LasR, demonstrating that these molecules do establish hydrogen bond interactions with Trp-60 and Asp-73 amino acids. Unfortunately, the authors could not demonstrate the importance of these residues in the molecular mechanism for an agonist to antagonist switching as they could not determine the X-ray crystallographic structure of LasR<sub>LBD</sub> bound to the antagonistic triphenyl compound TP-5, due to instability of this complex.<sup>19</sup>

## CONCLUSIONS

Upon demonstration by biochemical and biological activity assays that the LBD of LasR receptor proteins can fold into a functional conformation in the absence of its cognate acyl-HSL ligand, SERS spectroscopy allowed us to obtain direct experimental evidence of conformational changes in LasR<sub>LBD</sub> upon interaction with activators and inhibitors of *P. aeruginosa* quorum





chains. Ligand geometries were minimized using the AM1 Hamiltonian as implemented in Gaussian 09<sup>49</sup> and used as MOL2 files. Each ligand was docked in 25 independent genetic algorithm (GA) runs, and for each of these a maximum number of 100 000 GA operations were performed on a single population of 50 individuals. Operator weights for crossover, mutation and migration in the entry box were used as default parameters (95, 95, and 10, respectively), as well as the hydrogen bonding (4.0 Å) and van der Waals (2.5 Å) parameters. The position of C12-HSL in the crystal structure was used to define the active-site and the radius was set to 10 Å. The “flip ring corners” flag was switched on, while all the other flags were off. The GOLD scoring function was used to rank the ligands in order of fitness.

**Conflict of Interest:** The authors declare no competing financial interest.

**Acknowledgment.** This work has been funded by the European Research Council (ERC Advanced Grant #267867 Plasmaquo). We are grateful to Michael Givskov for kindly providing the pMHLAS vector. V.L.-P. acknowledges an FPI scholarship from the Spanish MINECO. C.G.-B. acknowledges funding from the Spanish MINECO (SAF2013-42899-R) and Xunta de Galicia (GRC2013-041).

**Supporting Information Available:** Various figures, schemes and tables including additional SERS spectra showing stability and reproducibility of substrates. The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acsnano.5b01800.

## REFERENCES AND NOTES

- Moskovits, M. Surface-Enhanced Raman Spectroscopy: A Brief Retrospective. *J. Raman Spectrosc.* **2005**, *36*, 485–496.
- Willems, K. A.; Van Duyn, R. P. Localized Surface Plasmon Resonance Spectroscopy and Sensing. *Annu. Rev. Phys. Chem.* **2007**, *58*, 267–297.
- Schlucker, S. Surface-Enhanced Raman Spectroscopy: Concepts and Chemical Applications. *Angew. Chem., Int. Ed.* **2014**, *53*, 4756–4795.
- Siddhanta, S.; Narayana, C. Surface Enhanced Raman Spectroscopy of Proteins: Implications for Drug Designing Invited Review Article. *Nanomater. Nanotechnol.* **2012**, *2*, 1–10.
- Ahijado-Guzman, R.; Gomez-Puertas, P.; Alvarez-Puebla, R. A.; Rivas, G.; Liz-Marzan, L. M. Surface-Enhanced Raman Scattering-Based Detection of the Interactions between the Essential Cell Division FtsZ Protein and Bacterial Membrane Elements. *ACS Nano* **2012**, *6*, 7514–7520.
- Karthigeyan, D.; Siddhanta, S.; Kishore, A. H.; Perumal, S. S. R. R.; Agren, H.; Sudevan, S.; Bhat, A. V.; Balasubramanyam, K.; Subbegowda, R. K.; Kundu, T. K.; et al. SERS and MD Simulation Studies of a Kinase Inhibitor Demonstrate the Emergence of a Potential Drug Discovery Tool. *Proc. Natl. Acad. Sci. U.S.A.* **2014**, *111*, 10416–10421.
- Case, R. J.; Labbate, M.; Kjelleberg, S. AHI-Driven Quorum-Sensing Circuits: Their Frequency and Function among the Proteobacteria. *ISME J.* **2008**, *2*, 345–349.
- Galloway, W. R. J. D.; Hodgkinson, J. T.; Bowden, S. D.; Welch, M.; Spring, D. R. Quorum Sensing in Gram-Negative Bacteria: Small-Molecule Modulation of AHI and AI-2 Quorum Sensing Pathways. *Chem. Rev.* **2011**, *111*, 28–67.
- Churchill, M. E. A.; Chen, L. L. Structural Basis of Acyl-Homoserine Lactone-Dependent Signaling. *Chem. Rev.* **2011**, *111*, 68–85.
- Li, Z.; Nair, S. K. Quorum Sensing: How Bacteria Can Coordinate Activity and Synchronize Their Response to External Signals? *Protein Sci.* **2012**, *21*, 1403–1417.
- Fuqua, C.; Greenberg, E. P. Listening in on Bacteria: Acyl-Homoserine Lactone Signaling. *Nat. Rev. Mol. Cell Biol.* **2002**, *3*, 685–695.
- Schuster, M.; Sexton, D. J.; Diggle, S. P.; Greenberg, E. P. Acyl-Homoserine Lactone Quorum Sensing: From Evolution to Application. *Annu. Rev. Microbiol.* **2013**, *67*, 43–63.
- Pesci, E. C.; Pearson, J. P.; Seed, P. C.; Iglewski, B. H. Regulation of Las and Rhl Quorum Sensing in *Pseudomonas aeruginosa*. *J. Bacteriol.* **1997**, *179*, 3127–3132.
- Pearson, J. P.; Gray, K. M.; Passador, L.; Tucker, K. D.; Eberhard, A.; Iglewski, B. H.; Greenberg, E. P. Structure of the Autoinducer Required for Expression of *Pseudomonas aeruginosa* Virulence Genes. *Proc. Natl. Acad. Sci. U.S.A.* **1994**, *91*, 197–201.
- Muh, U.; Hare, B. J.; Duerkop, B. A.; Schuster, M.; Hanzelka, B. L.; Heim, R.; Olson, E. R.; Greenberg, E. P. A Structurally Unrelated Mimic of a *Pseudomonas aeruginosa* Acyl-Homoserine Lactone Quorum-Sensing Signal. *Proc. Natl. Acad. Sci. U.S.A.* **2006**, *103*, 16948–16952.
- Borlee, B. R.; Geske, G. D.; Blackwell, H. E.; Handelsman, J. Identification of Synthetic Inducers and Inhibitors of the Quorum-Sensing Regulator LasR in *Pseudomonas aeruginosa* by High-Throughput Screening. *Appl. Environ. Microbiol.* **2010**, *76*, 8255–8258.
- Bottomley, M. J.; Muraglia, E.; Bazzo, R.; Carfi, A. Molecular Insights into Quorum Sensing in the Human Pathogen *Pseudomonas aeruginosa* from the Structure of the Virulence Regulator LasR Bound to Its Autoinducer. *J. Biol. Chem.* **2007**, *282*, 13592–13600.
- Zou, Y. Z.; Nair, S. K. Molecular Basis for the Recognition of Structurally Distinct Autoinducer Mimics by the *Pseudomonas aeruginosa* LasR Quorum-Sensing Signaling Receptor. *Chem. Biol.* **2009**, *16*, 961–970.
- Kiratisin, P.; Tucker, K. D.; Passador, L. LasR, a Transcriptional Activator of *Pseudomonas aeruginosa* Virulence Genes, Functions as a Multimer. *J. Bacteriol.* **2002**, *184*, 4912–4919.
- Hentzer, M.; Riedel, K.; Rasmussen, T. B.; Heydorn, A.; Andersen, J. B.; Parsek, M. R.; Rice, S. A.; Eberl, L.; Molin, S.; Hoiby, N.; et al. Inhibition of Quorum Sensing in *Pseudomonas aeruginosa* Biofilm Bacteria by a Halogenated Furanone Compound. *Microbiology* **2002**, *148*, 87–102.
- Zhu, J.; Winans, S. C. The Quorum-Sensing Transcriptional Regulator Trar Requires Its Cognate Signaling Ligand for Protein Folding, Protease Resistance, and Dimerization. *Proc. Natl. Acad. Sci. U.S.A.* **2001**, *98*, 1507–1512.
- Schuster, M.; Urbanowski, M. L.; Greenberg, E. P. Promoter Specificity in *Pseudomonas aeruginosa* Quorum Sensing Revealed by DNA Binding of Purified LasR. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, *101*, 15833–15839.
- Qin, Y. P.; Luo, Z. Q.; Smyth, A. J.; Gao, P.; von Bodman, S. B.; Farrand, S. K. Quorum-Sensing Signal Binding Results in Dimerization of Trar and Its Release from Membranes into the Cytoplasm. *EMBO J.* **2000**, *19*, 5212–5221.
- Welch, M.; Todd, D. E.; Whitehead, N. A.; McGowan, S. J.; Bycroft, B. W.; Salmond, G. P. C. *N*-Acyl Homoserine Lactone Binding to the Carr Receptor Determines Quorum-Sensing Specificity in *Erwinia*. *EMBO J.* **2000**, *19*, 631–641.
- Schu, D. J.; Ramachandran, R.; Geissinger, J. S.; Stevens, A. M. Probing the Impact of Ligand Binding on the Acyl-Homoserine Lactone-Hindered Transcription Factor Esar of *Pantoea stewartii* Subsp. *Stewartii*. *J. Bacteriol.* **2011**, *193*, 6315–6322.
- Zhang, R. G.; Pappas, T.; Brace, J. L.; Miller, P. C.; Oulmassov, T.; Molyneaux, J. M.; Anderson, J. C.; Bashkin, J. K.; Winans, S. C.; Joachimiak, A. Structure of a Bacterial Quorum-Sensing Transcription Factor Complexed with Pheromone and DNA. *Nature* **2002**, *417*, 971–974.
- Lintz, M. J.; Oinuma, K. I.; Wysoczynski, C. L.; Greenberg, E. P.; Churchill, M. E. A. Crystal Structure of Qsqr, a *Pseudomonas aeruginosa* Quorum Sensing Signal Receptor. *Proc. Natl. Acad. Sci. U.S.A.* **2011**, *108*, 15763–15768.
- Sappington, K. J.; Dandekar, A. A.; Oinuma, K. I.; Greenberg, E. P. Reversible Signal Binding by the *Pseudomonas aeruginosa* Quorum-Sensing Signal Receptor LasR. *MBio* **2011**, *2*, e00011.
- Claussen, A.; Jakobsen, T. H.; Bjarnsholt, T.; Givskov, M.; Welch, M.; Ferkinghoff-Borg, J.; Sams, T. Kinetic Model for Signal Binding to the Quorum Sensing Regulator LasR. *Int. J. Mol. Sci.* **2013**, *14*, 13360–13376.
- Pearson, J. P.; Pesci, E. C.; Iglewski, B. H. Roles of *Pseudomonas aeruginosa* Las and Rhl Quorum-Sensing Systems in Control of Elastase and Rhamnolipid Biosynthesis Genes. *J. Bacteriol.* **1997**, *179*, 5756–5767.

31. Manefield, M.; Rasmussen, T. B.; Hentzer, M.; Andersen, J. B.; Steinberg, P.; Kjelleberg, S.; Givskov, M. Halogenated Furanones Inhibit Quorum Sensing through Accelerated Luxr Turnover. *Microbiology* **2002**, *148*, 1119–1127.
32. El-Mowafy, S. A.; Abd El Galil, K. H.; El-Messery, S. M.; Shaaban, M. I. Aspirin Is an Efficient Inhibitor of Quorum Sensing, Virulence and Toxins in *Pseudomonas aeruginosa*. *Microb. Pathog.* **2014**, *74*, 25–32.
33. Yang, L.; Rybtke, M. T.; Jakobsen, T. H.; Hentzer, M.; Bjarnsholt, T.; Givskov, M.; Tolker-Nielsen, T. Computer-Aided Identification of Recognized Drugs as *Pseudomonas aeruginosa* Quorum-Sensing Inhibitors. *Antimicrob. Agents Chemother.* **2009**, *53*, 2432–2443.
34. Vial, S.; Pastoriza-Santos, I.; Perez-Juste, J.; Liz-Marzan, L. M. Plasmon Coupling in Layer-by-Layer Assembled Gold Nanorod Films. *Langmuir* **2007**, *23*, 4606–4611.
35. Backmann, N.; Zahnd, C.; Huber, F.; Bietsch, A.; Pluckthun, A.; Lang, H. P.; Guntherodt, H. J.; Hegner, M.; Gerber, C. A Label-Free Immunosensor Array Using Single-Chain Antibody Fragments. *Proc. Natl. Acad. Sci. U.S.A.* **2005**, *102*, 14587–14592.
36. Park, K.; Lee, J. M.; Jung, Y.; Habtemariam, T.; Salah, A. W.; Fermin, C. D.; Kim, M. Combination of Cysteine- and Oligomerization Domain-Mediated Protein Immobilization on a Surface Plasmon Resonance (Spr) Gold Chip Surface. *Analyst* **2011**, *136*, 2506–2511.
37. Kogot, J. M.; England, H. J.; Strouse, G. F.; Logan, T. M. Single Peptide Assembly onto a 1.5 nm Au Surface via a Histidine Tag. *J. Am. Chem. Soc.* **2008**, *130*, 16156–16157.
38. Wang, H.; Levin, C. S.; Halas, N. J. Nanosphere Arrays with Controlled Sub-10-Nm Gaps as Surface-Enhanced Raman Spectroscopy Substrates. *J. Am. Chem. Soc.* **2005**, *127*, 14992–14993.
39. Arif, M.; Karthigeyan, D.; Siddhanta, S.; Kumar, G. V.; Narayana, C.; Kundu, T. K. Analysis of Protein Acetyltransferase Structure-Function Relation by Surface-Enhanced Raman Scattering (Sers): A Tool to Screen and Characterize Small Molecule Modulators. *Methods Mol. Biol.* **2013**, *981*, 239–261.
40. Aliaga, A. E.; Aguayo, T.; Garrido, C.; Clavijo, E.; Hevia, E.; Gomez-Jeria, J. S.; Leyton, P.; Campos-Vallette, M. M.; Sanchez-Cortes, S. Surface-Enhanced Raman Scattering and Theoretical Studies of the C-Terminal Peptide of the Beta-Subunit Human Chorionic Gonadotropin without Linked Carbohydrates. *Biopolymers* **2011**, *95*, 135–143.
41. Wei, F.; Zhang, D. M.; Halas, N. J.; Hartgerink, J. D. Aromatic Amino Acids Providing Characteristic Motifs in the Raman and Sers Spectroscopy of Peptides. *J. Phys. Chem. B* **2008**, *112*, 9158–9164.
42. Podstawka, E.; Ozaki, Y.; Proniewicz, L. M. Part I: Surface-Enhanced Raman Spectroscopy Investigation of Amino Acids and Their Homodipeptides Adsorbed on Colloidal Silver. *Appl. Spectrosc.* **2004**, *58*, 570–580.
43. Barth, A.; Zscherp, C. What Vibrations Tell Us About Proteins. *Q. Rev. Biophys.* **2002**, *35*, 369–430.
44. Pande, J.; Pande, C.; Gilg, D.; Vasak, M.; Callender, R.; Kagi, J. H. R. Raman, Infrared, and Circular-Dichroism Spectroscopic Studies on Metallothionein—A Predominantly Turn-Containing Protein. *Biochemistry* **1986**, *25*, 5526–5532.
45. Gullekson, C.; Lucas, L.; Hewitt, K.; Kreplak, L. Surface-Sensitive Raman Spectroscopy of Collagen I Fibrils. *Biophys. J.* **2011**, *100*, 1837–1845.
46. Kumar, G. V. P.; Reddy, B. A. A.; Arif, M.; Kundu, T. K.; Narayana, C. Surface-Enhanced Raman Scattering Studies of Human Transcriptional Coactivator P300. *J. Phys. Chem. B* **2006**, *110*, 16787–16792.
47. Bastus, N. G.; Comenge, J.; Puntès, V. Kinetically Controlled Seeded Growth Synthesis of Citrate-Stabilized Gold Nanoparticles of up to 200 nm: Size Focusing versus Ostwald Ripening. *Langmuir* **2011**, *27*, 11098–11105.
48. CDC Web page. [http://www.cdc.cam.ac.uk/products/life\\_sciences/gold/](http://www.cdc.cam.ac.uk/products/life_sciences/gold/).
49. Frisch, M. J. T., G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G. A.; Nakatsuji, H. et al. *Gaussian 09*, Revision A.2 ed.; Gaussian, Inc.: Wallingford, CT, 2009.