

Research Article

Asymptotic Behavior of Solutions to Abstract Stochastic Fractional Partial Integrodifferential Equations

Zhi-Han Zhao,¹ Yong-Kui Chang,² and Juan J. Nieto^{3,4}

¹ Department of Mathematics and Information Engineering, Sanming University, Sanming, Fujian 365004, China

² Department of Mathematics, Lanzhou Jiaotong University, Lanzhou 730070, China

³ Departamento de Análisis Matemático, Facultad de Matemáticas, Universidad de Santiago de Compostela, 15782 Santiago de Compostela, Spain

⁴ Department of Mathematics, Faculty of Science, King Abdulaziz University, P.O. Box 80203, Jeddah 21589, Saudi Arabia

Correspondence should be addressed to Zhi-Han Zhao; zhaozhihan841110@126.com

Received 28 June 2013; Accepted 1 December 2013

Academic Editor: Dumitru Baleanu

Copyright © 2013 Zhi-Han Zhao et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The existence of asymptotically almost automorphic mild solutions to an abstract stochastic fractional partial integrodifferential equation is considered. The main tools are some suitable composition results for asymptotically almost automorphic processes, the theory of sectorial linear operators, and classical fixed point theorems. An example is also given to illustrate the main theorems.

1. Introduction

This paper is mainly concerned with the existence and uniqueness of square-mean asymptotically almost automorphic mild solutions to the following stochastic fractional partial integrodifferential equation in the form

$$\begin{aligned} & d[x(t) - f(t, x(t))] \\ &= \int_0^t \frac{(t-s)^{\alpha-2}}{\Gamma(\alpha-1)} A [x(s) - f(s, x(s))] ds dt \\ &+ g(t, x(t)) dW(t), \quad t \geq 0, \\ &x(0) = u_0, \end{aligned} \quad (1)$$

where $1 < \alpha < 2$, $A : D(A) \subset L^2(\mathbb{P}, \mathbb{H}) \rightarrow L^2(\mathbb{P}, \mathbb{H})$ is a linear densely defined operator of sectorial type on a Hilbert space $L^2(\mathbb{P}, \mathbb{H})$, $W(t)$ is a two-sided standard one-dimensional Brownian motion defined on the filtered probability space $(\Omega, \mathcal{F}, \mathbb{P}, \mathcal{F}_t)$, where $\mathcal{F}_t = \sigma\{W(u) - W(v); u, v \leq t\}$, and u_0 is an \mathcal{F}_0 -adapted, \mathbb{H} -valued random variable independent of the Wiener process W . Here f and g are appropriate functions to be specified later. The convolution integral in

(1) is understood in the Riemann-Liouville fractional integral (see, e.g., [1, 2]). We notice that fractional order can be complex in viewpoint of pure mathematics and there is much interest in developing the theoretical analysis and numerical methods to fractional equations, because they have recently proven to be valuable in various fields of science and engineering (see, e.g., [3–11] and references therein).

The concept of asymptotically almost automorphic functions was firstly introduced by N'Guérékata in [12]. Since then these functions have become of great interest to several mathematicians and gained lots of developments and applications, we refer the reader to [13–16] and the references listed therein.

Recently, the existence of almost automorphic and pseudo almost automorphic solutions to some stochastic differential equations has been considered in many publications such as [17–27] and the references therein. In a very recent paper [28], the authors introduced a new notation of square-mean asymptotically almost automorphic stochastic processes including a composition theorem. However, to the best of our knowledge, the existence of square-mean asymptotically almost automorphic mild solutions to the problem (1) is an untreated topic. Therefore, motivated by the works [16, 28], the main purpose of this paper is to investigate

the existence and uniqueness of square-mean asymptotically almost automorphic mild solutions to the problem (1). Then, we present an example as an application of our main results.

The rest of this paper is organized as follows. In Section 2, we recall some basic definitions and facts which will be used throughout this paper. In Section 3, we prove some existence results of square-mean asymptotically almost automorphic mild solutions to the problem (1). Finally, we give an example as an application of our abstract results.

2. Preliminaries

In this section, we introduce some basic definitions, notations, and preliminary facts which will be used in the sequel. For more details on this section, we refer the reader to [28–30].

Throughout the paper, $(\mathbb{H}, \|\cdot\|, \langle \cdot, \cdot \rangle)$ stands for a real separable Hilbert space. $(\Omega, \mathcal{F}, \mathbb{P})$ denotes a complete probability space, and $L^2(\mathbb{P}, \mathbb{H})$ stands for the space of all \mathbb{H} -valued random variables x such that

$$E\|x\|^2 = \int_{\Omega} \|x\|^2 d\mathbb{P} < \infty. \tag{2}$$

Note that $L^2(\mathbb{P}, \mathbb{H})$ is a Hilbert space equipped with the norm

$$\|x\|_2 := \left(\int_{\Omega} \|x\|^2 d\mathbb{P} \right)^{1/2}, \quad \text{for each } x \in L^2(\mathbb{P}, \mathbb{H}). \tag{3}$$

We denote by $C_0(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$ the collection of all bounded continuous stochastic processes φ from \mathbb{R}^+ into $L^2(\mathbb{P}, \mathbb{H})$ such that $\lim_{t \rightarrow +\infty} E\|\varphi(t)\|^2 = 0$. It is then easy to check that $C_0(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$ is a Banach space when it is endowed with the norm $\|\varphi\|_{C_0} := \sup_{t \in \mathbb{R}^+} \|\varphi(t)\|_2$. Similarly, $C_0(\mathbb{R}^+ \times L^2(\mathbb{P}, \mathbb{H}); L^2(\mathbb{P}, \mathbb{H}))$ stands for the space of the continuous stochastic processes $f : \mathbb{R}^+ \times L^2(\mathbb{P}, \mathbb{H}) \rightarrow L^2(\mathbb{P}, \mathbb{H})$ such that

$$\lim_{t \rightarrow +\infty} E\|f(t, x)\|^2 = 0 \tag{4}$$

uniformly for $x \in K$, where $K \subset L^2(\mathbb{P}, \mathbb{H})$ is any bounded subset. Additionally, $W(t)$ will be a two-sided standard one-dimensional Brownian motion defined on the filtered probability space $(\Omega, \mathcal{F}, \mathbb{P}, \mathcal{F}_t)$, where $\mathcal{F}_t = \sigma\{W(u) - W(v); u, v \leq t\}$.

2.1. Sectorial Linear Operators. A closed and linear operator A is said to be sectorial of type ϖ and angle θ if there exist $0 < \theta < \pi/2, M > 0$, and $\varpi \in \mathbb{R}$ such that its resolvent exists outside the sector $\varpi + S_{\theta} := \{\varpi + \lambda : \lambda \in \mathbb{C}, |\arg(-\lambda)| < \theta\}$ and $\|(\lambda - A)^{-1}\| \leq M/|\lambda - \varpi|, \lambda \notin \varpi + S_{\theta}$.

Definition 1 (see [2]). Let A be a closed and linear operator with domain $D(A)$ defined on a Banach space \mathbb{X} . We call A the generator of a solution operator if there exist $\varpi \in \mathbb{R}$ and a strongly continuous function $S_{\alpha} : \mathbb{R}^+ \rightarrow \mathcal{L}(\mathbb{X})$ such that $\{\lambda^{\alpha} : \operatorname{Re}(\lambda) > \varpi\} \subset \rho(A)$ and $\lambda^{\alpha-1}(\lambda^{\alpha} - A)^{-1}x = \int_0^{\infty} e^{-\lambda t} S_{\alpha}(t)x dt, \operatorname{Re}(\lambda) > \varpi, x \in \mathbb{X}$. In this case, $S_{\alpha}(\cdot)$ is called the solution operator generated by A .

We note that if A is sectorial of type ϖ with $0 \leq \theta < \pi(1 - \alpha/2)$, then A is the generator of a solution operator given by $S_{\alpha}(t) := (1/2\pi i) \int_{\gamma} e^{\lambda t} \lambda^{\alpha-1}(\lambda^{\alpha} - A)^{-1} d\lambda, t \geq 0$, where γ is a suitable path lying outside the sector $\varpi + S_{\theta}$. Recently, Cuesta in [1] proved that if A is a sectorial operator of type $\varpi < 0$ for some $M > 0$ and $0 \leq \theta < \pi(1 - \alpha/2)$, then there exists a constant $C > 0$ such that

$$\|S_{\alpha}(t)\| \leq \frac{CM}{1 + |\varpi|t^{\alpha}}, \quad t \geq 0. \tag{5}$$

Remark 2. Note that $S_{\alpha}(t)$ is, in fact, integrable. For more details on the solution family $S_{\alpha}(t)$ and related issues, we refer the reader to [31–33].

2.2. Square-Mean Asymptotically Almost Automorphic Processes. We recall some basic facts for a symptomatically almost automorphic processes which will be used in the sequel.

Definition 3 (see [22]). A stochastic process $x : \mathbb{R} \rightarrow L^2(\mathbb{P}, \mathbb{H})$ is said to be stochastically continuous if

$$\lim_{t \rightarrow s} E\|x(t) - x(s)\|^2 = 0. \tag{6}$$

Definition 4 (see [17]). A stochastically continuous stochastic process $x : \mathbb{R} \rightarrow L^2(\mathbb{P}, \mathbb{H})$ is said to be square-mean almost automorphic if, for every sequence of real numbers $\{s'_n\}_{n \in \mathbb{N}}$, there exist a subsequence $\{s_n\}_{n \in \mathbb{N}}$ and a stochastic process $y : \mathbb{R} \rightarrow L^2(\mathbb{P}, \mathbb{H})$ such that

$$\begin{aligned} \lim_{n \rightarrow \infty} E\|x(t + s_n) - y(t)\|^2 &= 0, \\ \lim_{n \rightarrow \infty} E\|y(t - s_n) - x(t)\|^2 &= 0 \end{aligned} \tag{7}$$

hold for each $t \in \mathbb{R}$. The collection of all square-mean almost automorphic stochastic processes $x : \mathbb{R} \rightarrow L^2(\mathbb{P}, \mathbb{H})$ is denoted by $AA(\mathbb{R}; L^2(\mathbb{P}, \mathbb{H}))$.

Definition 5 (see [17]). A function $f : \mathbb{R} \times L^2(\mathbb{P}, \mathbb{H}) \rightarrow L^2(\mathbb{P}, \mathbb{H}), (t, x) \rightarrow f(t, x)$, which is jointly continuous, is said to be square-mean almost automorphic if $f(t, x)$ is square-mean almost automorphic in $t \in \mathbb{R}$ uniformly for all $x \in \mathbb{K}$, where \mathbb{K} is any bounded subset of $L^2(\mathbb{P}, \mathbb{H})$. That is to say, for every sequence of real numbers $\{s'_n\}_{n \in \mathbb{N}}$, there exists a subsequence $\{s_n\}_{n \in \mathbb{N}}$ and a function $\tilde{f} : \mathbb{R} \times L^2(\mathbb{P}, \mathbb{H}) \rightarrow L^2(\mathbb{P}, \mathbb{H})$ such that

$$\begin{aligned} \lim_{n \rightarrow \infty} E\|f(t + s_n, x) - \tilde{f}(t, x)\|^2 &= 0, \\ \lim_{n \rightarrow \infty} E\|\tilde{f}(t - s_n, x) - f(t, x)\|^2 &= 0 \end{aligned} \tag{8}$$

for each $t \in \mathbb{R}$ and each $x \in \mathbb{K}$. Denote by $AA(\mathbb{R} \times L^2(\mathbb{P}, \mathbb{H}); L^2(\mathbb{P}, \mathbb{H}))$ the set of all such functions.

Lemma 6 (see [22]). $(AA(\mathbb{R}; L^2(\mathbb{P}, \mathbb{H})), \|\cdot\|_{\infty})$ is a Banach space equipped with the norm

$$\|x\|_{\infty} := \sup_{t \in \mathbb{R}} \|x(t)\|_2 = \sup_{t \in \mathbb{R}} (E\|x(t)\|^2)^{1/2}, \tag{9}$$

for $x \in AA(\mathbb{R}; L^2(\mathbb{P}, \mathbb{H}))$.

Lemma 7 (see [17]). *Let $f : \mathbb{R} \times L^2(\mathbb{P}, \mathbb{H}) \rightarrow L^2(\mathbb{P}, \mathbb{H})$, $(t, x) \rightarrow f(t, x)$ be square-mean almost automorphic, and assume that $f(t, \cdot)$ is uniformly continuous on each bounded subset $\mathbb{K} \subset L^2(\mathbb{P}, \mathbb{H})$ uniformly for $t \in \mathbb{R}$; that is, for all $\varepsilon > 0$, there exists $\delta > 0$ such that $x, y \in \mathbb{K}$ and $E\|x - y\|^2 < \delta$ imply that $E\|f(t, x) - f(t, y)\|^2 < \varepsilon$ for all $t \in \mathbb{R}$. Then for any square-mean almost automorphic process $x : \mathbb{R} \rightarrow L^2(\mathbb{P}, \mathbb{H})$, the stochastic process $F : \mathbb{R} \rightarrow L^2(\mathbb{P}, \mathbb{H})$ given by $F(\cdot) := f(\cdot, x(\cdot))$ is square-mean almost automorphic.*

Definition 8 (see [25]). A stochastically continuous process $f : \mathbb{R}^+ \rightarrow L^2(\mathbb{P}, \mathbb{H})$ is said to be square-mean asymptotically almost automorphic if it can be decomposed as $f = g + h$, where $g \in AA(\mathbb{R}; L^2(\mathbb{P}, \mathbb{H}))$ and $h \in C_0(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$. Denote by $AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$ the collection of all the square-mean asymptotically almost automorphic processes $f : \mathbb{R}^+ \rightarrow L^2(\mathbb{P}, \mathbb{H})$.

Definition 9 (see [28]). A function $f : \mathbb{R}^+ \times L^2(\mathbb{P}, \mathbb{H}) \rightarrow L^2(\mathbb{P}, \mathbb{H})$, $(t, x) \rightarrow f(t, x)$, which is jointly continuous, is said to be square-mean asymptotically almost automorphic if it can be decomposed as $f = g + h$, where $g \in AA(\mathbb{R} \times L^2(\mathbb{P}, \mathbb{H}); L^2(\mathbb{P}, \mathbb{H}))$ and $h \in C_0(\mathbb{R}^+ \times L^2(\mathbb{P}, \mathbb{H}); L^2(\mathbb{P}, \mathbb{H}))$. Denote by $AAA(\mathbb{R}^+ \times L^2(\mathbb{P}, \mathbb{H}); L^2(\mathbb{P}, \mathbb{H}))$ the set of all such functions.

Lemma 10 (see [28]). *If f, f_1 , and f_2 are all square-mean asymptotically almost automorphic stochastic processes, then the following hold true:*

- (I) $f_1 + f_2$ is square-mean asymptotically almost automorphic;
- (II) λf is square-mean asymptotically almost automorphic for any scalar λ ;
- (III) there exists a constant $M > 0$ such that $\sup_{t \in \mathbb{R}^+} E\|f(t)\|^2 \leq M$.

Lemma 11 (see [28]). *Suppose that $f \in AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$ admits a decomposition $f = g + h$, where $g \in AA(\mathbb{R}; L^2(\mathbb{P}, \mathbb{H}))$ and $h \in C_0(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$. Then $\{g(t) : t \in \mathbb{R}\} \subset \{f(t) : t \in \mathbb{R}^+\}$.*

Corollary 12 (see [28]). *The decomposition of a square-mean asymptotically almost automorphic process is unique.*

Lemma 13 (see [28]). *$AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$ is a Banach space when it is equipped with the norm*

$$\|f\|_{AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))} := \sup_{t \in \mathbb{R}} \|g(t)\|_2 + \sup_{t \in \mathbb{R}^+} \|h(t)\|_2, \quad (10)$$

where $f = g + h \in AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$ with $g \in AA(\mathbb{R}; L^2(\mathbb{P}, \mathbb{H}))$, $h \in C_0(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$.

Lemma 14 (see [28]). *$AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$ is a Banach space with the norm*

$$\|f\|_{\infty} := \sup_{t \in \mathbb{R}^+} \|f(t)\|_2 = \sup_{t \in \mathbb{R}^+} (E\|f(t)\|^2)^{1/2}. \quad (11)$$

Remark 15 (see [28]). In view of the previous lemmas it is clear that the two norms are equivalent in $AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$.

Lemma 16 (see [28]). *Let $f \in AA(\mathbb{R} \times L^2(\mathbb{P}, \mathbb{H}); L^2(\mathbb{P}, \mathbb{H}))$ and let $f(t, x)$ be uniformly continuous in any bounded subset $\mathbb{K} \subset L^2(\mathbb{P}, \mathbb{H})$ uniformly for $t \in \mathbb{R}^+$. Then $f(t, x)$ is uniformly continuous in any bounded subset $\mathbb{K} \subset L^2(\mathbb{P}, \mathbb{H})$ uniformly for $t \in \mathbb{R}$.*

Lemma 17 (see [28]). *Let $f \in AAA(\mathbb{R}^+ \times L^2(\mathbb{P}, \mathbb{H}); L^2(\mathbb{P}, \mathbb{H}))$ and suppose that $f(t, x)$ is uniformly continuous in any bounded subset $\mathbb{K} \subset L^2(\mathbb{P}, \mathbb{H})$ uniformly for $t \in \mathbb{R}^+$. If $u(t) \in AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$, then $f(\cdot, u(\cdot)) \in AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$.*

We now give the following concept of mild solution of (1).

Definition 18. Let $S_\alpha(t)$ be an integrable solution operator on $L^2(\mathbb{P}, \mathbb{H})$ with generator A . An \mathcal{F}_t -adapted stochastic process $x : [0, +\infty) \rightarrow L^2(\mathbb{P}, \mathbb{H})$ is called a mild solution of the problem (1) if $x(0) = u_0$ is \mathcal{F}_0 -measurable and $x(t)$ satisfies the corresponding stochastic integral equation:

$$x(t) = S_\alpha(t) [u_0 - f(0, u_0)] + f(t, x(t)) + \int_0^t S_\alpha(t-s) g(s, x(s)) dW(s), \quad t \geq 0. \quad (12)$$

3. Main Results

In this section, we establish the existence of square-mean asymptotically almost automorphic mild solutions to the problem (1). For that, we need the following technical results.

First, we list the following basic assumptions.

- (H1) The operator A is a sectorial operator of type $\varpi < 0$ for some $M > 0$ and $0 \leq \theta < \pi(1 - \alpha/2)$, and then there exists $C > 0$ such that

$$\|S_\alpha(t)\| \leq \frac{CM}{1 + |\varpi|t^\alpha}, \quad t \geq 0, \quad (13)$$

where $S_\alpha(t)$ is the solution operator generated by A .

- (H2) The function $f \in AAA(\mathbb{R}^+ \times L^2(\mathbb{P}, \mathbb{H}); L^2(\mathbb{P}, \mathbb{H}))$ and there exists a continuous and nondecreasing function $L_f : [0, +\infty) \rightarrow [0, +\infty)$ such that for each $r \geq 0$ and for all $E\|x\|^2 \leq r, E\|y\|^2 \leq r$,

$$E\|f(t, x) - f(t, y)\|^2 \leq L_f(r) E\|x - y\|^2 \quad (14)$$

for all $t \in \mathbb{R}^+$.

- (H3) The function $g \in AAA(\mathbb{R}^+ \times L^2(\mathbb{P}, \mathbb{H}); L^2(\mathbb{P}, \mathbb{H}))$ and there exists a continuous and nondecreasing function $L_g : [0, +\infty) \rightarrow [0, +\infty)$ such that for each $r \geq 0$ and for all $E\|x\|^2 \leq r, E\|y\|^2 \leq r$,

$$E\|g(t, x) - g(t, y)\|^2 \leq L_g(r) E\|x - y\|^2 \quad (15)$$

for all $t \in \mathbb{R}^+$.

(H4) We have

$$\sup_{r>0} \left[\frac{r}{6(CM)^2} - \frac{L_f(r)r}{(CM)^2} - \frac{|\omega|^{-1/\alpha} (1-1/\alpha) \pi L_g(r)r}{\alpha \sin(\pi/\alpha)} - \lambda r \right] > \left(1 + \frac{1}{(CM)^2} \right) M_f + \frac{|\omega|^{-1/\alpha} (1-1/\alpha) \pi}{\alpha \sin(\pi/\alpha)} M_g, \quad (16)$$

where $M_f = \sup_{t \in \mathbb{R}^+} E \|f(t, x(t))\|^2$ and $M_g = \sup_{t \in \mathbb{R}^+} E \|g(t, x(t))\|^2$.

(H5) The operator A is a sectorial operator of type ω with $0 \leq \theta < \pi(1 - \alpha/2)$, and there exists $\phi(\cdot) \in L^1(\mathbb{R}^+)$ such that

$$\|S_\alpha(t)\|^2 \leq \phi(t) \quad \forall t \geq 0, \quad \lim_{t \rightarrow +\infty} \phi(t) = 0, \quad (17)$$

where $S_\alpha(t)$ is the solution operator generated by A .

Lemma 19. Suppose that assumption (H1) holds and let $f \in AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$. If F is the function defined by

$$F(t) := \int_0^t S_\alpha(t-s) f(s) dW(s), \quad t \geq 0, \quad (18)$$

then $F \in AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$.

Proof. Since $f \in AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$, we have by definition that $f = g + h$, where $g \in AA(\mathbb{R}; L^2(\mathbb{P}, \mathbb{H}))$ and $h \in C_0(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$. Then

$$\begin{aligned} F(t) &= \int_0^t S_\alpha(t-s) g(s) dW(s) \\ &\quad + \int_0^t S_\alpha(t-s) h(s) dW(s) \\ &= \int_{-\infty}^t S_\alpha(t-s) g(s) dW(s) \\ &\quad - \int_{-\infty}^0 S_\alpha(t-s) g(s) dW(s) \\ &\quad + \int_0^t S_\alpha(t-s) h(s) dW(s) \\ &= G(t) + H(t), \end{aligned} \quad (19)$$

where $G(t) = \int_{-\infty}^t S_\alpha(t-s) g(s) dW(s)$ and $H(t) = - \int_{-\infty}^0 S_\alpha(t-s) g(s) dW(s) + \int_0^t S_\alpha(t-s) h(s) dW(s)$.

First we prove that $G(t) \in AA(\mathbb{R}; L^2(\mathbb{P}, \mathbb{H}))$. Let $\{s'_n\}_{n \in \mathbb{N}}$ be an arbitrary sequence of real numbers. Since $g \in AA(\mathbb{R}; L^2(\mathbb{P}, \mathbb{H}))$, there exists a subsequence $\{s_n\}_{n \in \mathbb{N}}$ of $\{s'_n\}_{n \in \mathbb{N}}$ such that for a certain stochastic process \tilde{g}

$$\begin{aligned} \lim_{n \rightarrow \infty} E \|g(t + s_n) - \tilde{g}(t)\|^2 &= 0, \\ \lim_{n \rightarrow \infty} E \|\tilde{g}(t - s_n) - g(t)\|^2 &= 0 \end{aligned} \quad (20)$$

hold for each $t \in \mathbb{R}$. Now, let $\tilde{W}(\sigma) := W(\sigma + s_n) - W(s_n)$ for each $\sigma \in \mathbb{R}$. Note that \tilde{W} is also a Brownian motion and has the same distribution as W . Moreover, if we let $\tilde{G}(t) = \int_{-\infty}^t S_\alpha(t-s) \tilde{g}(s) dW(s)$, then by making a change of variables $\sigma = s - s_n$ to get (see the equation (10.6.6) in [34])

$$\begin{aligned} E \|G(t + s_n) - \tilde{G}(t)\|^2 &= E \left\| \int_{-\infty}^{t+s_n} S_\alpha(t + s_n - s) g(s) dW(s) - \int_{-\infty}^t S_\alpha(t-s) \tilde{g}(s) dW(s) \right\|^2 \\ &= E \left\| \int_{-\infty}^t S_\alpha(t-\sigma) [g(\sigma + s_n) - \tilde{g}(\sigma)] d\tilde{W}(\sigma) \right\|^2 \end{aligned} \quad (21)$$

and, hence, using the Ito's isometry property of stochastic integral, we have the following estimations

$$\begin{aligned} E \|G(t + s_n) - \tilde{G}(t)\|^2 &\leq E \left(\int_{-\infty}^t \|S_\alpha(t-\sigma) [g(\sigma + s_n) - \tilde{g}(\sigma)]\|^2 d\sigma \right) \\ &\leq \int_{-\infty}^t \|S_\alpha(t-\sigma)\|^2 E \|g(\sigma + s_n) - \tilde{g}(\sigma)\|^2 d\sigma \\ &\leq \sup_{t \in \mathbb{R}} E \|g(t + s_n) - \tilde{g}(t)\|^2 \int_{-\infty}^t \left(\frac{CM}{1 + |\omega|(t-\sigma)^\alpha} \right)^2 d\sigma \\ &\leq (CM)^2 \sup_{t \in \mathbb{R}} E \|g(t + s_n) - \tilde{g}(t)\|^2 \int_0^\infty \frac{1}{(1 + |\omega|s^\alpha)^2} ds \\ &\leq \frac{(CM)^2 |\omega|^{-1/\alpha} (1-1/\alpha) \pi}{\alpha \sin(\pi/\alpha)} \sup_{t \in \mathbb{R}} E \|g(t + s_n) - \tilde{g}(t)\|^2. \end{aligned} \quad (22)$$

Then by (20), we obtain that $\lim_{n \rightarrow \infty} E \|G(t + s_n) - \tilde{G}(t)\|^2 = 0$ for each $t \in \mathbb{R}$. In a similar way, we can show that $\lim_{n \rightarrow \infty} E \|\tilde{G}(t - s_n) - G(t)\|^2 = 0$ for each $t \in \mathbb{R}$. Thus we conclude that $G(\cdot) \in AA(\mathbb{R}; L^2(\mathbb{P}, \mathbb{H}))$.

Next, let us show that $H(\cdot) \in C_0(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$. Since $h \in C_0(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$ and $1/(1 + |\omega|s^\alpha)^2$ is integrable in $[0, +\infty)$, for any sufficiently small $\varepsilon > 0$, there exists a constant $T > 0$ such that $E \|h(s)\|^2 \leq \varepsilon$ and $\int_T^\infty (1/(1 + |\omega|s^\alpha)^2) ds \leq \varepsilon$ for all $s \geq T$. Then, for all $t \geq 2T$, we obtain

$$\begin{aligned} E \|H(t)\|^2 &= E \left\| \int_0^{t/2} S_\alpha(t-s) h(s) dW(s) + \int_{t/2}^t S_\alpha(t-s) h(s) dW(s) - \int_{-\infty}^0 S_\alpha(t-s) g(s) dW(s) \right\|^2 \end{aligned}$$

$$\begin{aligned}
 &\leq 3E \left(\int_0^{t/2} \|S_\alpha(t-s)h(s)\|^2 ds \right) \\
 &\quad + 3E \left(\int_{t/2}^t \|S_\alpha(t-s)h(s)\|^2 ds \right) \\
 &\quad + 3E \left(\int_{-\infty}^0 \|S_\alpha(t-s)g(s)\|^2 ds \right) \\
 &\leq 3 \int_0^{t/2} \left(\frac{CM}{1+|\omega|(t-s)^\alpha} \right)^2 E \|h(s)\|^2 ds \\
 &\quad + 3 \int_{t/2}^t \left(\frac{CM}{1+|\omega|(t-s)^\alpha} \right)^2 E \|h(s)\|^2 ds \\
 &\quad + 3 \int_{-\infty}^0 \left(\frac{CM}{1+|\omega|(t-s)^\alpha} \right)^2 E \|g(s)\|^2 ds \\
 &\leq 3(CM)^2 \sup_{t \in \mathbb{R}^+} E \|h(t)\|^2 \int_{t/2}^t \frac{1}{(1+|\omega|s^\alpha)^2} ds \\
 &\quad + 3(CM)^2 \varepsilon \int_0^{t/2} \frac{1}{(1+|\omega|s^\alpha)^2} ds \\
 &\quad + 3(CM)^2 \sup_{t \in \mathbb{R}} E \|g(t)\|^2 \int_t^\infty \frac{1}{(1+|\omega|s^\alpha)^2} ds \\
 &\leq 3(CM)^2 \sup_{t \in \mathbb{R}^+} E \|h(t)\|^2 \int_T^\infty \frac{1}{(1+|\omega|s^\alpha)^2} ds \\
 &\quad + 3(CM)^2 \varepsilon \int_0^\infty \frac{1}{(1+|\omega|s^\alpha)^2} ds \\
 &\quad + 3(CM)^2 \sup_{t \in \mathbb{R}} E \|g(t)\|^2 \int_T^\infty \frac{1}{(1+|\omega|s^\alpha)^2} ds \\
 &\leq 3(CM)^2 \varepsilon \left(\sup_{t \in \mathbb{R}^+} E \|h(t)\|^2 + \frac{|\omega|^{-1/\alpha} (1-1/\alpha) \pi}{\alpha \sin(\pi/\alpha)} \right. \\
 &\quad \left. + \sup_{t \in \mathbb{R}} E \|g(t)\|^2 \right). \tag{23}
 \end{aligned}$$

This inequality proves the assertion since ε is arbitrary. Recalling that $F(t) = G(t) + H(t)$ for all $t \geq 0$, we get $F(t) \in AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$. The proof is completed. \square

It is easy to see that, by arguments similar to those in the proof of Lemma 19, we have the following result.

Lemma 20. *Suppose that assumption (H5) holds and let $f \in AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$. If F is the function defined by*

$$F(t) := \int_0^t S_\alpha(t-s) f(s) dW(s), \quad t \geq 0, \tag{24}$$

then $F \in AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$.

Now, we are ready to establish our main results.

Theorem 21. *Assume that (H1)–(H4) hold. Then there exists $\varepsilon > 0$ such that for each $u_0 \in B_\varepsilon(0, L^2(\mathbb{P}, \mathbb{H}))$ there exists a unique square-mean asymptotically almost automorphic mild solution $x(\cdot, u_0)$ of the problem (1) on $[0, \infty)$ such that $x(0, u_0) = u_0$.*

Proof. We define a nonlinear operator Y by

$$\begin{aligned}
 (Yx)(t) &= S_\alpha(t) [u_0 - f(0, u_0)] + f(t, x(t)) \\
 &\quad + \int_0^t S_\alpha(t-s) g(s, x(s)) dW(s) \quad t \geq 0. \tag{25}
 \end{aligned}$$

First we prove that $Y(AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))) \subseteq AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$. Given $x \in AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$, from the properties of $\{S_\alpha(t)\}_{t \geq 0}$, f , and g , we infer that Yx is well defined and continuous. Since $x(t)$ is bounded, we can choose a bounded subset \mathbb{K} of $L^2(\mathbb{P}, \mathbb{H})$ such that $x(t) \in \mathbb{K}$ for all $t \in \mathbb{R}^+$. It follows from (H2) and (H3) that both $f(t, x)$ and $g(t, x)$ are uniformly continuous on the bounded subset \mathbb{K} uniformly for $t \in \mathbb{R}^+$. Moreover, from Lemmas 17 and 19 and taking into account (H1), it follows that $Yx \in AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$.

Now, by (H4), there exists a constant $r > 0$ such that

$$\begin{aligned}
 &\frac{r}{6(CM)^2} - \frac{L_f(r)r}{(CM)^2} - \frac{|\omega|^{-1/\alpha} (1-1/\alpha) \pi L_g(r)r}{\alpha \sin(\pi/\alpha)} - \lambda r \\
 &> \left(1 + \frac{1}{(CM)^2} \right) M_f + \frac{|\omega|^{-1/\alpha} (1-1/\alpha) \pi}{\alpha \sin(\pi/\alpha)} M_g. \tag{26}
 \end{aligned}$$

Let $0 < \lambda < 1$. We affirm that the assertion holds for $\varepsilon = \lambda r$. In fact, let $u_0 \in B_\varepsilon(0, L^2(\mathbb{P}, \mathbb{H}))$. Define the space $\mathbb{D} = \{x \in AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H})) : x(0) = u_0, E \|x(t)\|^2 \leq r, t \geq 0\}$. Then \mathbb{D} is a closed subspace of $AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$. We claim that $Y\mathbb{D} \subseteq \mathbb{D}$. If $x \in \mathbb{D}$ and $t \in \mathbb{R}^+$, we get

$$\begin{aligned}
 &E \|(Yx)(t)\|^2 \\
 &\leq 3E \|S_\alpha(t) [u_0 - f(0, u_0)]\|^2 + 3E \|f(t, x(t))\|^2 \\
 &\quad + 3E \left\| \int_0^t S_\alpha(t-s) g(s, x(s)) dW(s) \right\|^2 \\
 &\leq 6(CM)^2 [E \|u_0\|^2 + E \|f(0, u_0)\|^2] \\
 &\quad + 3E \|f(t, x(t)) - f(t, 0) + f(t, 0)\|^2 \\
 &\quad + 3E \left(\int_0^t \|S_\alpha(t-s) g(s, x(s))\|^2 ds \right) \\
 &\leq 6(CM)^2 \lambda r + 6(CM)^2 E \|f(0, u_0)\|^2 \\
 &\quad + 6L_f(r)r + 6 \sup_{t \in \mathbb{R}^+} E \|f(t, 0)\|^2 \\
 &\quad + 6(CM)^2 \left[\int_0^t \left(\frac{1}{1+|\omega|(t-s)^\alpha} \right)^2 L_g(r) r ds \right. \\
 &\quad \left. + \int_0^t \left(\frac{1}{1+|\omega|(t-s)^\alpha} \right)^2 E \|g(s, 0)\|^2 ds \right]
 \end{aligned}$$

$$\begin{aligned} &\leq 6(CM)^2\lambda r + 6(CM)^2M_f + 6L_f(r)r + 6M_f \\ &\quad + \frac{6(CM)^2|\bar{\omega}|^{-1/\alpha}(1-1/\alpha)\pi}{\alpha\sin(\pi/\alpha)} [L_g(r)r + M_g], \end{aligned} \quad (27)$$

which from (26) implies that $E\|(\Upsilon x)(t)\|^2 \leq r$ for all $t \geq 0$, and so that $\Upsilon\mathbb{D} \subseteq \mathbb{D}$.

Next, to complete the proof, we need to show that $\Upsilon(\cdot)$ is a contraction from \mathbb{D} into \mathbb{D} . By (26), we know that

$$\frac{r}{6(CM)^2} - \frac{L_f(r)r}{(CM)^2} - \frac{|\bar{\omega}|^{-1/\alpha}(1-1/\alpha)\pi L_g(r)r}{\alpha\sin(\pi/\alpha)} - \lambda r > 0. \quad (28)$$

That is,

$$\frac{r}{6(CM)^2} > \frac{L_f(r)r}{(CM)^2} + \frac{|\bar{\omega}|^{-1/\alpha}(1-1/\alpha)\pi L_g(r)r}{\alpha\sin(\pi/\alpha)} + \lambda r. \quad (29)$$

Then, one has

$$6L_f(r) + \frac{6(CM)^2|\bar{\omega}|^{-1/\alpha}(1-1/\alpha)\pi}{\alpha\sin(\pi/\alpha)}L_g(r) + 6\lambda(CM)^2 < 1. \quad (30)$$

For any $x, y \in \mathbb{D}$ and $t \geq 0$, we have

$$\begin{aligned} &E\|(\Upsilon x)(t) - (\Upsilon y)(t)\|^2 \\ &\leq 2E\|f(t, x(t)) - f(t, y(t))\|^2 \\ &\quad + 2E\left\|\int_0^t S_\alpha(t-s)[g(s, x(s)) - g(s, y(s))]dW(s)\right\|^2 \\ &\leq 2L_f(r)\sup_{t \in \mathbb{R}^+}E\|x(t) - y(t)\|^2 \\ &\quad + 2(CM)^2\int_0^t\left(\frac{1}{1+|\bar{\omega}|(t-s)^\alpha}\right)^2 \\ &\quad \quad \times E\|g(s, x(s)) - g(s, y(s))\|^2 ds \\ &\leq 2L_f(r)\sup_{t \in \mathbb{R}^+}E\|x(t) - y(t)\|^2 \\ &\quad + \frac{2(CM)^2|\bar{\omega}|^{-1/\alpha}(1-1/\alpha)\pi}{\alpha\sin(\pi/\alpha)}L_g(r) \\ &\quad \times \sup_{t \in \mathbb{R}^+}E\|x(t) - y(t)\|^2 \\ &\leq \left[2L_f(r) + \frac{2(CM)^2|\bar{\omega}|^{-1/\alpha}(1-1/\alpha)\pi}{\alpha\sin(\pi/\alpha)}L_g(r)\right] \\ &\quad \times \sup_{t \in \mathbb{R}^+}E\|x(t) - y(t)\|^2. \end{aligned} \quad (31)$$

Thus, we get

$$\begin{aligned} &\|Yx - Yy\|_\infty \\ &= \sup_{t \in \mathbb{R}^+} \left(E\|(\Upsilon x)(t) - (\Upsilon y)(t)\|^2 \right)^{1/2} \\ &\leq \sqrt{2L_f(r) + \frac{2(CM)^2|\bar{\omega}|^{-1/\alpha}(1-1/\alpha)\pi}{\alpha\sin(\pi/\alpha)}L_g(r)} \|x - y\|_\infty. \end{aligned} \quad (32)$$

It follows from (30) that Υ is a contraction mapping on \mathbb{D} . So by the Banach contraction mapping principle, we draw a conclusion that there exists a unique fixed point $x(\cdot)$ for Υ in \mathbb{D} . It is clear that x is a square-mean asymptotically almost automorphic mild solution of (1). The proof is complete. \square

The next result is proved using the similar steps as in the proof of the previous result, so we omit the details.

Theorem 22. Assume that (H1)–(H3) hold. If $L_f(r) \equiv L_f$ and $L_g(r) \equiv L_g$ for all $r \geq 0$ and $L_f + (CM)^2|\bar{\omega}|^{-1/\alpha}(1-1/\alpha)\pi L_g/\alpha\sin(\pi/\alpha) < 1/2$, then for every $u_0 \in L^2(\mathbb{P}, \mathbb{H})$, there exists a unique square-mean asymptotically almost automorphic mild solution $x(\cdot, u_0)$ of the problem (1) on $[0, \infty)$ such that $x(0, u_0) = u_0$.

Theorem 23. Suppose that assumptions (H2), (H3), and (H5) hold. If $L_f(r) \equiv L_f$ and $L_g(r) \equiv L_g$ for all $r \geq 0$ and $L_f + L_g\|\phi\|_1 < 1/2$, then for every $u_0 \in L^2(\mathbb{P}, \mathbb{H})$, there exists a unique square-mean asymptotically almost automorphic mild solution $x(\cdot, u_0)$ of the problem (1) on $[0, \infty)$ such that $x(0, u_0) = u_0$.

Proof. Consider the nonlinear operator Υ given by

$$\begin{aligned} (\Upsilon x)(t) &= S_\alpha(t)[u_0 - f(0, u_0)] + f(t, x(t)), \\ &\quad + \int_0^t S_\alpha(t-s)g(s, x(s))dW(s), \quad t \geq 0. \end{aligned} \quad (33)$$

First we prove that Υ maps $AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$ into itself. Given $x \in AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$, from the properties of $\{S_\alpha(t)\}_{t \geq 0}$, f and g , we infer that Υx is well defined and continuous. Since $x(t)$ is bounded, we can choose a bounded subset \mathbb{K} of $L^2(\mathbb{P}, \mathbb{H})$ such that $x(t) \in \mathbb{K}$ for all $t \in \mathbb{R}^+$. It follows from conditions (H2) and (H3) that both $f(t, x)$ and $g(t, x)$ are uniformly continuous on the bounded subset \mathbb{K} uniformly for $t \in \mathbb{R}^+$. Moreover, from Lemmas 17 and 20 and taking into account (H5), it follows that $\Upsilon x \in AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$.

Next we prove that Υ is a contraction mapping from $AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$ into itself. Note that we have already proved $\Upsilon(AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))) \subseteq AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$. Moreover, for any $x, y \in AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$ and $t \geq 0$, we have

$$\begin{aligned} &E\|(\Upsilon x)(t) - (\Upsilon y)(t)\|^2 \\ &\leq 2E\|f(t, x(t)) - f(t, y(t))\|^2 \end{aligned}$$

$$\begin{aligned}
 &+ 2E \left\| \int_0^t S_\alpha(t-s) [g(s, x(s)) - g(s, y(s))] dW(s) \right\|^2 \\
 \leq &2L_f \sup_{t \in \mathbb{R}^+} E \|x(t) - y(t)\|^2 \\
 &+ 2 \int_0^t \phi(t-s) E \|g(s, x(s)) - g(s, y(s))\|^2 ds \\
 \leq &2L_f \sup_{t \in \mathbb{R}^+} E \|x(t) - y(t)\|^2 \\
 &+ 2L_g \sup_{t \in \mathbb{R}^+} E \|x(t) - y(t)\|^2 \int_0^\infty \phi(s) ds \\
 \leq &[2L_f + 2L_g \|\phi\|_1] \sup_{t \in \mathbb{R}^+} E \|x(t) - y(t)\|^2.
 \end{aligned} \tag{34}$$

Therefore

$$\begin{aligned}
 \|\Upsilon x - \Upsilon y\|_\infty &= \sup_{t \in \mathbb{R}^+} (E \|\Upsilon x(t) - \Upsilon y(t)\|^2)^{1/2} \\
 &\leq \sqrt{2L_f + 2L_g \|\phi\|_1} \|x - y\|_\infty.
 \end{aligned} \tag{35}$$

That is, Υ is a contraction mapping on $AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$. By the Banach contraction mapping principle, Υ has a unique fixed point $x(t) \in AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$. It is clear that x is a square-mean asymptotically almost automorphic mild solution of (1). The proof is then complete. \square

4. An Example

In this section, we apply the results obtained previously to investigate the existence of square-mean asymptotically almost automorphic mild solutions for the following partial stochastic fractional differential system:

$$\begin{aligned}
 &\frac{\partial}{\partial t} \left[u(t, \xi) - \int_{-\infty}^t a(t) a_1(t-s) u(s, \xi) ds \right] \\
 &= J_t^{\alpha-1} \left(\frac{\partial^2}{\partial \xi^2} - \nu \right) \left[u(t, \xi) - \int_{-\infty}^t a(t) a_1(t-s) u(s, \xi) ds \right] \\
 &+ \int_{-\infty}^t a(t) a_2(t-s) u(s, \xi) ds dW(t), \\
 &u(t, 0) = u(t, \pi) = 0, \quad t \geq 0, \\
 &u(0, \xi) = u_0(\xi), \quad \xi \in I = [0, \pi],
 \end{aligned} \tag{36}$$

where $a \in AAA(\mathbb{R}^+; \mathbb{R})$, $a_1, a_2 : \mathbb{R}^+ \rightarrow \mathbb{R}$ are continuous functions, and $\nu > 0$ is a fixed constant.

Let $\mathbb{H} = L^2([0, \pi])$ with the norm $\|\cdot\|$ and $A : D(A) \subseteq \mathbb{H} \rightarrow \mathbb{H}$ be the operator defined by $Ax = x'' - \nu x$ domain $D(A) = \{x \in \mathbb{H} : x'' \in \mathbb{H}, x(0) = x(\pi) = 0\}$. It is well known that $\Delta x = x''$ is the infinitesimal generator of an analytic semigroup $\{S(t)\}_{t \geq 0}$ on \mathbb{H} . Furthermore, A is sectorial of type $\omega = -\nu < 0$.

In the sequel, we assume

$$\begin{aligned}
 L_f &= \|a\|_\infty \left(\int_{-\infty}^0 |a_1(-s)|^2 ds \right)^{1/2} < \infty, \\
 L_g &= \|a\|_\infty \left(\int_{-\infty}^0 |a_2(-s)|^2 ds \right)^{1/2} < \infty.
 \end{aligned} \tag{37}$$

Now, we can define the functions $f, g : \mathbb{R}^+ \times L^2(\mathbb{P}, \mathbb{H}) \rightarrow L^2(\mathbb{P}, \mathbb{H})$ by

$$\begin{aligned}
 f(t, x)(\xi) &= a(t) \int_{-\infty}^0 a_1(-s) x(s, \xi) ds, \\
 g(t, x)(\xi) &= a(t) \int_{-\infty}^0 a_2(-s) x(s, \xi) ds, \\
 J_t^{\alpha-1} h(t) &= \int_0^t \frac{(t-s)^{\alpha-2}}{\Gamma(\alpha-1)} h(s) ds,
 \end{aligned} \tag{38}$$

which permits to transform the system (36) into the abstract system (1). Moreover, it is not difficult to see that f, g are continuous and Lipschitz in the second variable with Lipschitz constants L_f and L_g , respectively.

The next result is a consequence of Theorem 22.

Theorem 24. *Under the previous assumptions, (36) has a unique mild solution $x \in AAA(\mathbb{R}^+; L^2(\mathbb{P}, \mathbb{H}))$ whenever L_f and L_g are small enough.*

Acknowledgments

The first author was supported by Research Fund for Young Teachers of Sanming University (B2011071Q). The second author was supported by NSF of China (11361032), Program for New Century Excellent Talents in University (NCET-10-0022), and NSF of Gansu Province of China (1107RJZA091). And the third author was partially supported by Ministerio de Economía y Competitividad (Spain), project MTM2010-15314, and cofinanced by the European Community fund FEDER.

References

- [1] E. Cuesta, "Asymptotic behaviour of the solutions of fractional integro-differential equations and some time discretizations," *Discrete and Continuous Dynamical Systems Series A*, supplement, pp. 277-285, 2007.
- [2] C. Cuevas and J. C. de Souza, "S-asymptotically ω -periodic solutions of semilinear fractional integro-differential equations," *Applied Mathematics Letters*, vol. 22, no. 6, pp. 865-870, 2009.
- [3] R. P. Agarwal, V. Lakshmikantham, and J. J. Nieto, "On the concept of solution for fractional differential equations with uncertainty," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 72, no. 6, pp. 2859-2862, 2010.
- [4] V. V. Anh and R. McVinish, "Fractional differential equations driven by Lévy noise," *Journal of Applied Mathematics and Stochastic Analysis*, vol. 16, no. 2, pp. 97-119, 2003.
- [5] K. Diethelm and N. J. Ford, "Analysis of fractional differential equations," *Journal of Mathematical Analysis and Applications*, vol. 265, no. 2, pp. 229-248, 2002.

- [6] A. M. A. El-Sayed, "Nonlinear functional-differential equations of arbitrary orders," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 33, no. 2, pp. 181–186, 1998.
- [7] R. Gorenflo and F. Mainardi, "Fractional calculus: integral and differential equations of fractional order," in *Fractals and Fractional Calculus in Continuum Mechanics*, vol. 378, pp. 223–276, Springer, New York, NY, USA, 1997.
- [8] B. I. Henry and S. L. Wearne, "Existence of Turing instabilities in a two-species fractional reaction-diffusion system," *SIAM Journal on Applied Mathematics*, vol. 62, no. 3, pp. 870–887, 2002.
- [9] H. Hilfer, *Applications of Fractional Calculus in Physics*, World Scientific, River Edge, NJ, USA, 2000.
- [10] V. Lakshmikantham and A. S. Vatsala, "Basic theory of fractional differential equations," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 69, no. 8, pp. 2677–2682, 2008.
- [11] K. S. Miller and B. Ross, *An Introduction to the Fractional Calculus and Fractional Differential Equations*, A Wiley-Interscience Publication, John Wiley & Sons, New York, NY, USA, 1993.
- [12] G. M. N'Guérékata, "Sur les solutions presque-automorphes d'équations différentielles abstraites," *Les Annales des Sciences Mathématiques du Québec*, vol. 5, no. 1, pp. 69–79, 1981.
- [13] D. Bugajewski and G. M. N'Guérékata, "On the topological structure of almost automorphic and asymptotically almost automorphic solutions of differential and integral equations in abstract spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 59, no. 8, pp. 1333–1345, 2004.
- [14] T. Diagana, E. M. Hernández, and J. P. C. dos Santos, "Existence of asymptotically almost automorphic solutions to some abstract partial neutral integro-differential equations," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 71, no. 1-2, pp. 248–257, 2009.
- [15] H.-S. Ding, T.-J. Xiao, and J. Liang, "Asymptotically almost automorphic solutions for some integrodifferential equations with nonlocal initial conditions," *Journal of Mathematical Analysis and Applications*, vol. 338, no. 1, pp. 141–151, 2008.
- [16] J. P. C. dos Santos and C. Cuevas, "Asymptotically almost automorphic solutions of abstract fractional integro-differential neutral equations," *Applied Mathematics Letters*, vol. 23, no. 9, pp. 960–965, 2010.
- [17] Y.-K. Chang, Z.-H. Zhao, and G. M. N'Guérékata, "A new composition theorem for square-mean almost automorphic functions and applications to stochastic differential equations," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 74, no. 6, pp. 2210–2219, 2011.
- [18] Y.-K. Chang, Z.-H. Zhao, and G. M. N'Guérékata, "Square-mean almost automorphic mild solutions to non-autonomous stochastic differential equations in Hilbert spaces," *Computers & Mathematics with Applications*, vol. 61, no. 2, pp. 384–391, 2011.
- [19] Y.-K. Chang, Z.-H. Zhao, and G. M. N'Guérékata, "Square-mean almost automorphic mild solutions to some stochastic differential equations in a Hilbert space," *Advances in Difference Equations*, vol. 2011, article 9, 2011.
- [20] Y.-K. Chang, Z.-H. Zhao, G. M. N'Guérékata, and R. Ma, "Stepanov-like almost automorphy for stochastic processes and applications to stochastic differential equations," *Nonlinear Analysis: Real World Applications*, vol. 12, no. 2, pp. 1130–1139, 2011.
- [21] J. Cao, Q. Yang, and Z. Huang, "Existence and exponential stability of almost automorphic mild solutions for stochastic functional differential equations," *Stochastics*, vol. 83, no. 3, pp. 259–275, 2011.
- [22] M. Fu and Z. Liu, "Square-mean almost automorphic solutions for some stochastic differential equations," *Proceedings of the American Mathematical Society*, vol. 138, no. 10, pp. 3689–3701, 2010.
- [23] M. Fu, "Almost automorphic solutions for nonautonomous stochastic differential equations," *Journal of Mathematical Analysis and Applications*, vol. 393, no. 1, pp. 231–238, 2012.
- [24] M. Fu and F. Chen, "Almost automorphic solutions for some stochastic differential equations," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 80, pp. 66–75, 2013.
- [25] Z. Chen and W. Lin, "Square-mean pseudo almost automorphic process and its application to stochastic evolution equations," *Journal of Functional Analysis*, vol. 261, no. 1, pp. 69–89, 2011.
- [26] G. M. N'Guérékata, *Almost Automorphic and almost Periodic Functions in Abstract Spaces*, Kluwer Academic/Plenum Publishers, New York, NY, USA, 2001.
- [27] G. M. N'Guérékata, *Topics in Almost Automorphy*, Springer, New York, NY, USA, 2005.
- [28] Z.-H. Zhao, Y.-K. Chang, and J. J. Nieto, "Square-mean asymptotically almost automorphic process and its application to stochastic integro-differential equations," *Dynamic Systems and Applications*, vol. 22, no. 2-3, pp. 269–284, 2013.
- [29] R. P. Agarwal, B. de Andrade, and C. Cuevas, "Weighted pseudo-almost periodic solutions of a class of semilinear fractional differential equations," *Nonlinear Analysis: Real World Applications*, vol. 11, no. 5, pp. 3532–3554, 2010.
- [30] C. Cuevas and C. Lizama, "Almost automorphic solutions to a class of semilinear fractional differential equations," *Applied Mathematics Letters*, vol. 21, no. 12, pp. 1315–1319, 2008.
- [31] O. Fattorini, *Second Order Differential Equations in Banach Spaces*, North-Holland, Amsterdam, The Netherlands, 1985.
- [32] M. Haase, *The Functional Calculus for Sectorial Operators*, vol. 169, Birkhäuser, Basel, Switzerland, 2006.
- [33] J. Prüss, *Evolutionary Integral Equations and Applications*, vol. 87, Birkhäuser, Basel, Switzerland, 1993.
- [34] H.-H. Kuo, *Introduction to Stochastic Integration*, Springer, New York, NY, USA, 2006.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

