

Screening of *Ipomoea tuba* Leaf Extract for Identification of Bioactive Compounds and Evaluation of Its *in vitro* Antiproliferative Activity Against MCF-7 and HeLa Cells

Thirupati Chinna Venkateswarulu¹, Gaddam Eswaraiah¹, Srirama Krupanidhi¹, Karlapudi Abraham Peele¹, Indira Mikkili¹, Alugunulla Venkata Narayana¹, Bharath Kumar Ravuru², John Babu Dulla¹ and Ranga Rao Ambati^{1*}

¹Department of Bio-Technology, Vignans Foundation for Science, Technology & Research (Deemed to be University), Vadlamudi-522213, Guntur, Andhra Pradesh, India

²School of Life Sciences, Rayalaseema University, Kurnool-518002, Andhra Pradesh, India

Received: 2 May 2019

Accepted: 17 March 2020



SUMMARY

Mangroves contain a wide range of bioactive compounds with pharmacological activities. In the present study, we analysed the separation and detection of phytoconstituents with the methanol extract of *Ipomoea tuba* leaf using gas chromatography-mass spectrometry (GC-MS) and tested its *in vitro* cytotoxicity effect against MCF-7 and HeLa cells. Phytochemical compounds such as docosanoic, octadecatrienoic and *cis*-9-octadecanoic acids, triterpenoid γ -sitosterol, and terpene alcohol in methanol extract of *I. tuba* leaf were identified. Furthermore, *in vitro* antiproliferative activity of the extract of *I. tuba* leaf was evaluated using MCF-7 and HeLa cells. The results indicated a reduction of cell viability of 37.43 and 41.89 % of MCF-7 and HeLa cells respectively. The methanol extract of *I. tuba* leaf proved to be effective in protecting the cells against oxidative stress. This is the first report on the *in vitro* cytotoxicity effect of *I. tuba* leaf extract on MCF-7 and HeLa cells.

Key words: mangrove species, *Ipomoea tuba*, bioactive compounds, antiproliferative activity

INTRODUCTION

Mangrove ecosystem plays a major role in the human life to protect us from natural disasters like tsunami, floods, high tides and soil erosion. Mangrove plants grow in water logging region and offer a shelter for wide ranges of endemic fauna and flora (1,2). They absorb and remove five times more carbon dioxide than normal terrestrial plants (3). The mangrove species, namely *Suaeda maritima*, commonly known as seablite is used in homemade foods such as salad, curry, soy sauce, and spicy soup in Thailand (4). In recent years, bioactive compounds produced from the plants have attracted the interest of pharmaceutical industries for formulation of drugs because the effectiveness of synthetic antibiotics against several pathogenic strains is slowly decreasing (5). The natural compounds and related drugs are used to treat different human diseases (6). The crude methanol extract of different medicinal plants contains many bioactive compounds having anticancer activity against several cancers like gastric, colon and breast cancer cell lines (7). Phytoconstituents and cytotoxicity of *I. tuba* have not been studied so far. Hence, the present study aims to evaluate the phytoconstituents of methanol extract of *I. tuba* leaf and their cytotoxicity effects on MCF-7 and HeLa cells.

MATERIALS AND METHODS

Sample preparation

Ipomoea tuba sample was collected from Nizampatnam mangroves, Guntur, Andhra Pradesh, India. The sample was prepared from leaves of *I. tuba* by soaking 50 g of powdered sample in 50 mL of absolute methanol for 72 h. The sample was filtered through Whatman No. 42 filter paper and then methanol was evaporated from the test sample by rotary vacuum evaporator (EV11; Equitron Medica Pvt Ltd, Mumbai, India). The final crude extract was dissolved in 100 % dimethyl sulfoxide (DMSO; Sigma-Aldrich Chemicals Pvt Ltd, Merck, Bangalore, India), made to final concentration of 100 mg/mL and used for antiproliferative studies. The concentration of DMSO maintained in the wells was less than 1 %, which is not toxic to the cell lines (8,9).

*Corresponding author:

E-mail: arangarao99@gmail.com

Identification of compounds by GC-MS analysis

Bioactive compounds in leaf extract of *I. tuba* were identified by GC-MS (6890 series; Agilent, Santa Clara, CA, USA). The following chromatographic conditions were maintained: initial column temperature 30 °C, heated up to 300 °C at 10 °C/5 min, flow rate 1.0 mL/min and helium was used as carrier gas in split mode. The bioactive compounds were identified based on retention times and quantified by integration of peak area. Similarity of compounds was compared with known compounds using NIST-based AMDIS software (10).

3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay for cell viability

Ipomoea tuba leaf extract was screened for *in vitro* cytotoxicity activity on MCF-7 and HeLa cells ($5 \cdot 10^3$ cell/well) using MTT (Sigma-Aldrich, Merck, St. Louis, MO, USA) assay. The sample (100 μ L diluted plant extract) was added to 100 μ L of Dulbecco's Modified Eagle's medium (DMEM), then the cell lines were added to the 96-well microtiter plate, and incubated for 48 h at 37 °C. The MTT was added and allowed to incubate for 2 h until the purple precipitate was formed. Then, absorbance values were measured at 520 nm using UV-Vis spectrophotometer (Cary 60; Agilent Technologies, Selangor Darul Ehsan, Malaysia). The dose-response curve was plotted for evaluation of IC_{50} values (11).

Statistical analysis

The experimental data of both cell lines were statistically analyzed using ANOVA method. The value $p < 0.05$ is considered statistically significant for the analysis of the percentage of inhibition of cell viability.

RESULTS AND DISCUSSION

GC-MS analysis for compound identification

The chromatogram confirmed the presence of compounds such as fatty acids: docosanoic, octadecatrienoic and *cis*-9-octadecanoic acids, triterpenoid γ -sitosterol, and terpene alcohol in the leaf extract of mangrove plant *Ipomoea tuba*. The compounds were identified based on retention times. **Table 1** gives the molecular mass and retention times of each compound. Angaye *et al.* (12) reported various bioactive compounds in the extracts of mangroves *Rhizophora mangle*, *Rhizophora racemosa*, *Avicennia germinans* and *Laguncularia racemosa*. Phytochemical compounds such as

β -sitosterol, eicosanol and taraxerol are found in *Bruguiera cylindrica* extract and showed their cytotoxicity against neuro2A cancer cell lines (13).

The bioactive compounds from *Avicennia marina*, *Salvadora persica* and *Avicennia officinalis* contain several types of secondary metabolites like flavonoids, tannins, alkaloids and saponins (14,15). In a previous study, GC-MS analysis of the extracts of *Acrostichum aureum* confirmed the presence of bioactive compounds such as stigmaterol, γ -sitosterol, campesterol and 24-methylene cycloartenol with potential activity against adenocarcinoma, carcinoma and other human cancer cell lines (16). Ganesh and Vennila (17) reported the presence of different bioactive compounds, namely terpenoids, steroids, saponins, catechol and phenols in the methanol extracts of *Acanthus ilicifolius* and *Avicennia officinalis*. Report from Basyuni *et al.* (18) showed that the mangrove species *Acanthus ilicifolius*, *Rhizophora apiculata* and *Sonneratia caseolaris* contain a wide range of phytoconstituents like triterpenoids, taraxerol, germanicol and tannins, with anti-inflammatory, anticarcinogenic, antimicrobial and antiprotozoan activities. The mangrove plant *Rhizophora mucronata* leaf extract contains vindoline, catharanthine and serpentine, the major alkaloids with free radical scavenging and cytotoxic activity (19). The findings of Grozav *et al.* (20) proved that the derivative of thiazole synthesized from mangrove plant leaves has potential anticancer effect against ovarian cancer cell lines A2780 and HeLa cell lines.

In vitro antiproliferative activity of *I. tuba* leaf extract on MCF-7 cells

Extract of *I. tuba* leaves showed significant *in vitro* antiproliferative effect on MCF-7, and the viability of MCF-7 was reduced with the increase in the concentration of the sample. High reduction of MCF-7 cells was observed at the concentration of *I. tuba* leaf extract of 100 μ g/mL (**Table 2**) and the IC_{50} value against MCF-7 cells was found to be (40.4 ± 0.1) μ g/mL. During experiments, it was observed that the increase in sample concentration altered the morphology of MCF-7 cells, leading to cell death (**Fig. 1**). Similar findings are reported for *in vitro* cytotoxicity effect of *Avicennia marina* extracts on different cancerous cells (21-23). Patra and Thatoi (24) reported the antiproliferative activity of the methanol extract of *Heritiera fomes* leaves against melanoma cell lines and achieved 40% inhibition. The methanol extract of *Avicennia marina* leaf showed anticancer activity against MDA-MB 231 and MCF-7 cell (25).

Table 1. Bioactive compounds identified in the extract of *Ipomoea tuba* leaf by GC-MS

| Peak no. | t_r /min | Compound name | Formula | M /(g/mol) | CAS number |
|----------|------------|---|--------------------|--------------|------------|
| 1 | 19.49 | docosanoic acid | $C_{21}H_{44}COOH$ | 340.59 | 112-85-6 |
| 2 | 21.24 | 3,7,11,15-tetramethyl-2-hexadecene-1-ol | $C_{20}H_{40}O$ | 296.00 | 7541-49-3 |
| 3 | 27.86 | octadecatrienoic acid-ethyl ester | $C_{19}H_{32}O$ | 292.46 | 1191-41-9 |
| 4 | 31.07 | <i>cis</i> -9-octadecanoic acid | $C_{18}H_{34}O_2$ | 282.00 | 112-80-1 |
| 5 | 32.80 | γ -sitosterol | $C_{29}H_{50}O$ | 414.71 | 83-47-6 |

Table 2. Inhibition of MCF-7 and HeLa cells using *Ipomoea tuba* leaf extract

| γ (extract)/($\mu\text{g}/\text{mL}$) | Viability of MCF-7 cells/% | Viability of HeLa cells/% |
|--|-------------------------------|-------------------------------|
| 100 | (37.4 \pm 0.1) ^f | (41.4 \pm 0.1) ^f |
| 75 | (42.0 \pm 0.2) ^e | (44.7 \pm 0.2) ^e |
| 50 | (45.4 \pm 0.6) ^d | (48.3 \pm 1.0) ^d |
| 25 | (53.5 \pm 0.8) ^c | (51.4 \pm 0.4) ^c |
| 10 | (55.8 \pm 0.5) ^b | (53.9 \pm 0.3) ^b |
| 5 | (60.7 \pm 0.3) ^a | (55.6 \pm 0.8) ^a |

Values are expressed as mean \pm S.D. Values with different letters in superscript in the same column are significantly different ($p < 0.05$) determined by ANOVA

In vitro antiproliferative activity of *I. tuba* leaf extracts on HeLa cells

HeLa cell viability was decreased with the increased concentration of leaf extract and the maximum reduction in HeLa cells was observed at 100 $\mu\text{g}/\text{mL}$ (Table 2), with the IC_{50} value of (37.4 \pm 0.1) $\mu\text{g}/\text{mL}$. It was observed that after the treatment with *I. tuba* extract, the HeLa cells slowly detached from one another. Fig. 2 shows the change in morphology of HeLa cells. Khajure and Rathod (26) reported that the extract of *A. ilicifolius* had cytotoxic activity against HeLa and KB cells. Rajeswari *et al.* (27) also reported that the flavone molecule from *Excoecaria agallocha* has the cytotoxic activity against HeLa cells.




Fig. 1. Morphology of MCF-7 cells after the treatment with the extract of *Ipomoea tuba* leaf: a) untreated MCF-7 cell lines, and b-g) treated with different concentrations (5, 10, 25, 50, 75 and 100 $\mu\text{g}/\text{mL}$ respectively) of the leaf extract




Fig. 2. Morphology of HeLa cells after the treatment with the extract of *Ipomoea tuba* leaf: a) untreated HeLa cell lines, and b-g) treated with different concentrations (5, 10, 25, 50, 75, and 100 $\mu\text{g}/\text{mL}$ respectively) of the leaf extract

CONCLUSIONS

In conclusion, GC-MS analysis confirmed the presence of different phytoconstituents and the *Ipomoea tuba* extracts were proved to have antiproliferative effect on MCF-7 and HeLa cells. This is the first report of high antiproliferative activity of the extract of *I. tuba* leaf on MCF-7 and HeLa cells. Furthermore, these bioactive compounds could be used in functional food applications for health benefits.









ACKNOWLEDGEMENTS

Authors acknowledge the research facilities supported by Fund for Improvement of Science & Technology Infrastructure in Higher Educational Institutions (FIST Project No: LSI-576/2013), Department of Science and Technology, Government of India and Vignan's Foundation for Science, Technology and Research University.

CONFLICT OF INTEREST

Authors declare that there is no conflict of interest.

ORCID ID

T.C. Venkateswarulu  <https://orcid.org/0000-0002-7822-8860>
 G. Eswaraiyah  <https://orcid.org/0000-0002-7480-3610>
 S. Krupanidhi  <https://orcid.org/0000-0001-7702-8974>
 K. Abraham Peele  <https://orcid.org/0000-0001-8173-9672>
 I. Mikkili  <https://orcid.org/0000-0002-7141-6682>
 A. Venkata Narayana  <https://orcid.org/0000-0002-5349-0491>
 B. Kumar Ravuru  <https://orcid.org/0000-0002-1385-0798>
 J. Babu Dulla  <https://orcid.org/0000-0003-4641-915X>
 A. Ranga Rao  <https://orcid.org/0000-0002-5735-4489>

REFERENCES

1. Eswaraiyah G, Venkateswarulu TC, Krupanidhi S, Abraham Peele K, Indira M, Venkata Narayana A, *et al.* GC-MS analysis for leaf extract of *Suaeda nudiflora* and screening of their *in vitro* anti-proliferative effect against MCF 7 and HeLa cells. *Agri Res & Tech: Open Access J.* 2019;22(1):556187. <https://doi.org/10.19080/ARTOAJ.2019.22.556187>
2. Eswaraiyah G, Abraham Peele K, Krupanidhi S, Kumar RB, Venkateswarulu TC. Identification of bioactive compounds in leaf extract of *Avicennia alba* by GC-MS analysis and evaluation of its *in-vitro* anticancer potential against MCF7 and HeLa cell lines. *J King Saud Univ Sci.* 2020;32(1):740-4. <https://doi.org/10.1016/j.jksus.2018.12.010>
3. Record S, Charney ND, Zakaria RM, Ellison AM. Projecting global mangrove species and community distributions under climate change. *Ecosphere.* 2013;4(3):1-23. <https://doi.org/10.1890/ES12-00296.1>
4. Pornpitakdamrong A, Sudjaroen Y. Seablite (*Suaeda maritima*) product for cooking, Samut Songkram province, Thailand. *Food Nutr Sci.* 2014;5(9):850-6. <https://doi.org/10.4236/fns.2014.59094>
5. Teiten MH, Gaascht F, Dicato M, Diederich M. Anticancer bioactivity of compounds from medicinal plants used in European medieval traditions. *Biochem Pharmacol.* 2013;86(9):1239-47. <https://doi.org/10.1016/j.bcp.2013.08.007>
6. Newman DJ, Cragg GM. Natural products as sources of new drugs over the last 25 years. *J Nat Prod.* 2007;70(3):461-77. <https://doi.org/10.1021/np068054v>
7. Akter R, Uddin SJ, Grice ID, Tiralongo E. Cytotoxic activity screening of Bangladeshi medicinal plant extracts. *J Nat Med.* 2014;68:246-52. <https://doi.org/10.1007/s11418-013-0789-5>
8. Eswaraiyah G, Abraham Peele K, Krupanidhi S, Indira M, Kumar RB, Venkateswarulu TC. GC-MS analysis for compound identification in leaf extract of *Lumnitzera racemosa* and evaluation of its *in vitro* anticancer effect against MCF7 and HeLa cell lines. *J King Saud Univ Sci.* 2020;32(1):780-3. <https://doi.org/10.1016/j.jksus.2019.01.014>
9. Eswaraiyah G, Abraham Peele K, Krupanidhi S, Kumar RB, Venkateswarulu TC. Studies on phytochemical, antioxidant, antimicrobial analysis and separation of bioactive leads of leaf extract from the selected mangroves. *J King Saud Univ Sci.* 2020;32(1):842-7. <https://doi.org/10.1016/j.jksus.2019.03.002>
10. Kulkarni A, Govindappa M, Ramachandra YL, Koka P. GC-MS analysis of methanol extract of *Cassia fistula* and its *in vitro* anticancer activity on human prostate cancer cell line. *Indo Am J Pharm Res.* 2015;5(2):937-44.
11. Bhat RP. Anticancer activities of plant extracts of *Gymnocranthera farquhariana* (Hook. f. & Thomson) Warb., *Myristica fatua* Houtt. var. *magnifica* (Beddome) Sinclair and *Samadera indica* Gaertner. *Adv Obes Weight Manag Control.* 2017;6(5):167-71. <https://doi.org/10.15406/aowmc.2017.06.00173>
12. Angaye TCN, Ohimain EI, Siasia EP, Asaigbe PI, Finomo OA. Larvicidal activities of the leaves of Niger Delta mangrove plants against *Anopheles gambiae*. *Sky J Microbiol Res.* 2014;2(7):045-050.
13. Nithyamol Kalappurakkal V, Bhattacharya D, Chakravarty S, Venkata Uppuluri M. Isolation, synthesis and AChE inhibitory potential of some novel cinnamyl esters of taraxerol, the major metabolite of the Mangrove *Bruguiera cylindrica*. *Chem Biodivers.* 2018;15(4):e1800008. <https://doi.org/10.1002/cbdv.201800008>
14. Bholay AD, Ingale MH, Gaur A. Therapeutic potential of mangrove and its associate plant extracts from thane creek, against human respiratory tract MDR pathogens. *J Biol Env Sci.* 2015;7(4):118-26.
15. Thatoi H, Samantaray D, Das SK. The genus *Avicennia*, a pioneer group of dominant mangrove plant species with potential medicinal values: A review. *Front Life Sci.* 2016;9(4):267-91. <https://doi.org/10.1080/21553769.2016.1235619>

16. Thomas A, Prashob PKJ, Chandramohanakumar N. A profiling of anti-tumour potential of sterols in the mangrove fern *Acrostichum aureum*. *Int J Pharmacogn Phytochem Res.* 2016;8(11):1828-32.
17. Ganesh S, Vennila J. Phytochemical analysis of *Acanthus ilicifolius* and *Avicennia officinalis* by GC-MS. *Res J Phytochem.* 2011;5(1):60-5.
<https://doi.org/10.3923/rjphyto.2011.60.65>
18. Basyuni M, Putri LAP, Oku H. Phytomedicinal investigation from six mangrove tree species, North Sumatra, Indonesia (Investigasi Fitomedisinal Enam Spesies Mangrove, Sumatera Utara, Indonesia). *Ilmu Kelautan.* 2013;18(3):157-64.
<https://doi.org/10.14710/ik.ijms.18.3.157-164>
19. Gurudeeban S, Ramanathan T, Satyavani K. Antimicrobial and radical scavenging effects of alkaloid extracts from *Rhizophora mucronata*. *Pharm Chem J.* 2013;47:50-3.
<https://doi.org/10.1007/s11094-013-0895-4>
20. Grozav A, Porumb ID, Găină LI, Filip L, Hanganu D. Cytotoxicity and antioxidant potential of novel 2-(2-((1*H*-indol-5yl)methylene)-hydrazinyl)-thiazole derivatives. *Molecules.* 2017;22(2):260.
<https://doi.org/10.3390/molecules22020260>
21. Karami L, Majd A, Mehrabian S, Nabiuni M, Salehi M, Irian S. Antimutagenic and anticancer effects of *Avicennia marina* leaf extract on *Salmonella typhimurium* TA100 bacterium and human promyelocytic leukaemia HL-60 cells. *Sci Asia.* 2012;38(3):349-55.
<https://doi.org/10.2306/scienceasia1513-1874.2012.38.349>
22. Sukhramani PS, Patel PM. Biological screening of *Avicennia marina* for anticancer activity. *Pelagia Res Lib.* 2013;4(2):125-30.
23. Shanthi N, Murugesan S, Janetta Nithia SM, Kotteswari M. *In vitro* anticancer activity of methanol extracts of *Avicennia marina* (Forssk) Vireh against HepG2 liver hepatocellular carcinoma cancer cell line. *Int J Recent Sci Res.* 2018;9(10):29122-6.
<https://doi.org/10.24327/ijrsr.2018.0910.2796>
24. Patra JK, Thatoi H. Anticancer activity and chromatography characterization of methanol extract of *Heritiera fomes* Buch. Ham., a mangrove plant from Bhitarkanika, India. *Orient Pharm Exp Med.* 2013;13:133-42.
<https://doi.org/10.1007/s13596-013-0113-7>
25. Momtazi-borojeni AA, Behbahani M, Sadeghi-aliabadi H. Antiproliferative activity and apoptosis induction of crude extract and fractions of *Avicennia marina*. *Iran J Basic Med Sci.* 2013;16(11):1203-8.
26. Khajure PV, Rathod JL. Potential anticancer activity of *Acanthus ilicifolius* extracted from the mangroves forest of Karwar, West coast of India. *World J Sci Technol.* 2011;1:1-6.
27. Rajeswari K, Rao TB, Sharma GVR, Muralikrishna R. Anticancer activity of flavone of *Excoecaria agallocha* in cytotoxicity studies using Hela Cells. *Asian J Chem.* 2017;29(11): 2406-08.
<https://doi.org/10.14233/ajchem.2017.20724>