



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

23rd International Grassland Congress

---

## Conservation Prioritization of Semi-Arid Savannas in India: A Multi-Scale Landscape Approach

Abi Tamim Vanak

*Ashoka Trust for Research in Ecology and the Environment, India*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/23/3-6-1/5>

The 23rd International Grassland Congress (Sustainable use of Grassland Resources for Forage Production, Biodiversity and Environmental Protection) took place in New Delhi, India from November 20 through November 24, 2015.

Proceedings Editors: M. M. Roy, D. R. Malaviya, V. K. Yadav, Tejveer Singh, R. P. Sah, D. Vijay, and A. Radhakrishna

Published by Range Management Society of India

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## **Conservation prioritization of semi-arid savannas in India: a multi-scale landscape approach**

**Abi Tamim Vanak**

ATREE, Bangalore, India

Corresponding author e-mail: [avanak@atree.org](mailto:avanak@atree.org)

**Keywords:** Biodiversity, Endangered, Grasslands, Indian subcontinent

### **Introduction**

The semi-arid savanna grasslands of peninsular India are unique habitats that support a vast proportion of India's agro-pastoralist community. They are also home to critically endangered species such as the Great Indian bustard and other endangered and endemic species such as the Indian wolf (*Canis lupus pallipes*) and blackbuck antelope (*Antelope cervicapra*). Unfortunately, savanna grasslands have not received the attention of conservationists or policy makers, resulting in a lack of protection for both the endangered and endemic wildlife as well as the habitat. Given the high human-dependence on these biomes, traditional conservation measures such as large protected areas are not feasible. The management and conservation of these fragmented and human-dominated regions requires delineation of high-priority areas which are most likely to persist in the long-term.

### **Materials and Methods**

Delineation of semi-arid savannas with standard remote-sensing analysis techniques does not yield desirable results due to spectral mismatching with rain-fed agriculture. Therefore, to determine the extent of savanna grasslands in India, we used a multi-scale multi-method landscape level analysis. We first used multi-date NDVI derived from coarse resolution MODIS imageries to create a map of grasslands. We then used a rule-set based on a regression tree to predict areas which are bioclimatically suitable for grasslands. We then focussed on four states in central-south India (Madhya Pradesh, Maharashtra, Karnataka and Andhra Pradesh), and derived medium resolution (30m) maps using a Supervised classification of Landsat ETM+ imageries and extensive ground truthing. We used a filter approach to determine the largest contiguous patches in these four states (>100ha). Finally, we derived high-resolution (5.6 m LISS IV) maps of the largest contiguous patches in these four states. We combined these maps in the software package ZONATION with information on species presence and conservation compatible and incompatible human-use to generate areas of conservation prioritisation.

### **Results and Discussion**

Based on our exercise, we found that Karnataka had only 2.1% of its geographic area under grassland >100 ha; Andhra Pradesh (unified) had 4.2%, Madhya Pradesh had 7.9% and Maharashtra had 12.8%. Within these states, we identified nine areas ranging in size from 34 sq. km to 922 sq. km. for further survey and conservation action. These areas not only have the potential to conserve the inherent biological values of this biome, but at the same time support traditional livelihood practices.

### **Conclusion**

We propose these areas be declared as conservation reserves that allows for mixed use of the landscape, by engaging all stakeholders in developing a joint landscape conservation program.

### **Acknowledgement**

Ministry of Environment and Forests, Maharashtra Forest Department, Abhijeet Kulkarni, Ameya Gode, Chintan Sheth, Krithi Karanth and Enrico DiMinin are all gratefully acknowledged for their contributions to this study.