



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

23rd International Grassland Congress

Photosynthesis and Biomass Studies in *Lasiurus indicus* of Chandan Grassland in Thar Desert

P. Raja

Central Arid Zone Research Institute, India

Maharaj Singh

Central Arid Zone Research Institute, India

Nilendu Singh

Wadia Institute of Himalayan Geology, India

Nawalesh Kumar Sinha

Indian Institute of Agricultural Biotechnology, India

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/23/4-1-3/14>

The 23rd International Grassland Congress (Sustainable use of Grassland Resources for Forage Production, Biodiversity and Environmental Protection) took place in New Delhi, India from November 20 through November 24, 2015.

Proceedings Editors: M. M. Roy, D. R. Malaviya, V. K. Yadav, Tejveer Singh, R. P. Sah, D. Vijay, and A. Radhakrishna

Published by Range Management Society of India

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Photosynthesis and biomass studies in *Lasiurus indicus* of Chandan grassland in Thar Desert

P. Raja^{1*}, Maharaj Singh², Nilendu Singh³, N.K. Sinha⁴

¹ICAR-Central Arid Zone Research Institute, Jodhpur, India

²ICAR-Central Arid Zone Research Institute, Regional Research Station, Jaisalmer, India

³Wadia Institute of Himalayan Geology, Dehradun, India

⁴ICAR-Indian Institute of Agricultural Biotechnology, Ranchi, India

*Corresponding author e-mail: prajanbss@gmail.com

Keywords: Biomass, Ecophysiology, *Lasiurus indicus*, Light use efficiency, Water use efficiency

Introduction

Grasslands comprise about one third of the world's area of natural vegetation (Adams *et al.*, 1990). Uncertainties prevail in the arid ecosystem with reference to carbon balance and fluxes which are primarily attributed to the sensitivity of grasslands to variation in annual precipitation, temperature and other regulating mechanisms of eco-physiological processes (Flanagan *et al.*, 2002). It is therefore necessary to study radiation and water use efficiency of *Lasiurus indicus* which is a predominant grass species in Chandan grassland of Thar desert to understand and evaluate its growth in relation to variation in climate. Also, it is pertinent to resolve whether grassland ecosystem function as a source or sink to atmospheric CO₂. We therefore, measured diurnal variability of net photosynthesis, diurnal trend in light and water use efficiency and root and shoot biomass studies (10 days interval) of *L. indicus* (*Sewan*) grass species in relation to ambient micrometeorological conditions during growth period along three points *viz.*, initial, peak and declining phenophase to have basic understanding of eco-physiological responses to prevailing micrometeorology to evaluate its carbon use efficiency.

Materials and Methods

Present study was conducted at Central Arid Zone Research Institute (CAZRI) experimental area at Chandan (26°50.83'N, 71°18.083'E) near Jaisalmer, covering ≈96 hectares of grassland ecosystem predominantly consisting of *Lasiurus indicus* locally known as *Sewan* grass (≈60%) with associated grass species *viz.*, *Cenchrus ciliaris* (≈15%) and *Panicum antidotale* (≈5%). The climatic conditions are characterized by highly variable precipitation (≈100-300 mm), extremes of temperatures (48°C in summer and -3°C in winter), high wind speed (≈33kmh⁻¹) high insolation and thus leading to very high evaporative demand. The soil texture is sandy with low organic carbon content and low fertility status. The phenology of the *Lasiurus indicus* varies (Plate 1) as the beginning of the growing season during south-west monsoon (Mid of July-August end), followed by good green grass growth in post-monsoon season (September - Mid of October) and senescence phase in winter (3rd week of October to - January end).



Plate. 1: Pre - (a) and post-monsoon (b) phenology of *L. indicus* grass species in the study area

The leaf area index (LAI) was measured at 10-day interval from the growth initiation (greening) phase from the first dekad of July (Jul 1d). The LAI measurements lasted up to senescence (browning) phase. The LAI measurements were taken using plant canopy analyzer (LAI-2000; Li-Cor, Inc., Lincoln, NE). In each quadrat, the mean LAI was obtained by taking average of eight readings. Mean LAI of all the quadrats were computed to represent the temporal dynamics of LAI. Leaf level photosynthesis (L_{nar}) data was acquired using Li-Cor's LI-6400XT portable photosynthesis system was upscaled to canopy level according to Campbell and Norman (1998) assuming single light assimilation response relation for all the leaves in the canopy. Biomass of *L. indicus* was collected in 3 replicates in 1x1m quadrats at every 15 days interval through destructive method from 2nd Week of April to 3rd week of August from 1m² area using permanent iron quadrats. Fresh weights of both shoots and roots were taken using digital balance and the dry weights after keeping them in oven at 55° C for 24 hours.

Results and Discussion

During initial phenophase of new leaf initialization, leaf level photosynthesis (L_{nar}) of *L. indicus* was measured in daytime with an average of 6.4 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$. LAI of *L. indicus* species was also minimum at 0.65. Canopy level photosynthesis (C_{nar}) noted approximately half to that of L_{nar} as LAI was minimum. At peak growth phenophase, L_{nar} of *L. indicus* was at life cycle maximum with an average of 8.3 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$. LAI at this phase of growth was 1.75. At this time C_{nar} was about 20-35 per cent higher than L_{nar} . At senescence phase, with decline in soil moisture regime and unfavorable winter conditions, *L. indicus* reached physiologically dormant state. This was manifested in declining LAI (0.8) and degreening of leaves in grass. During dormant stage an average L_{nar} was found 3.2 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ and C_{nar} was 20-25 per cent lower than L_{nar} , as LAI decline. *L. indicus* exhibited bimodal distribution in its diurnal photosynthetic rates with its first peak (13 to 15 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$) at about 10 Indian Standard Time (IST) and second but higher peak (18 to 20 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$) at around 16 hrs. The maximum light use efficiency (2.73 gMJ^{-1}) was around 10 and 16IST with occasional dip (0.47 gMJ^{-1}) at around 14IST (Fig. 1).




Fig. 1: Eco-physiological responses (Diurnal) of *L. indicus* during peak growth stage

Stomatal closure was affected by extreme temperature and atmospheric forcings could be the cause for the dip in photosynthesis rate during noon hours. Daytime peak canopy CO_2 exchange rate in the present study is comparable to that obtained by eddy covariance system in semiarid environment of south India (Lalrammawia and Paliwal, 2010). The diffuse fraction was also found to have sharp bimodal trend (15-51%) with peaks in the morning and late afternoon which might have contributed to observed peaks in photosynthesis (Zhang *et al.*, 2009). During evening hours after setting of declining limb of winter condition, with decrease in daylight hours, L_{nar} turned negative, *i.e.*, respiration exceeded photosynthesis rate. The *L. indicus* showed maximum water use efficiency (WUE) during morning and evening hours. Consequently, light use efficiency (LUE) *L. indicus*, showed similar trend *viz.*, minimum (0.43 gMJ^{-1}) during noon hours and peaks ($2.96, 2.03 \text{ gMJ}^{-1}$) at morning and afternoon hours (Fig 1). Combination of higher diffuse fraction and lower ambient temperatures in the morning and afternoon hours in contrast to noon hours might have resulted into generalized bimodal trend in the *LUE* of *L. indicus*. Biomass studies of *L. indicus* (*Sewan*) grass species indicates that average shoot and root weight is 296 and 414 gm/m^2 respectively. Biomass studies and related analyses showed that *L. indicus* acted as a carbon sink to atmospheric CO_2 and has fixed carbon to about 3.20 tons per hectare.

Conclusion

The study concludes that at the initial and final phenophase in the life cycle of *L. indicus*, leaf level assimilation rate is greater than that of canopy level. Diurnal trend of photosynthesis, LUE and WUE exemplifies the adaptation strategy of this grass species. Stomatal closure in *L. indicus* during peak day time hours controls net carbon assimilation and transpiration. *L. indicus* grass species in Chandan grassland acted as a carbon sink to the atmospheric CO₂ and has the ability to ameliorate the extreme climate in Thar desert region through fixation of carbon in vegetation and soil.

References

- Adams, J. M., H. Faure, L. Fauredenard, J. M. Mcglade and F. I. Woodward. 1990. Increases in terrestrial carbon storage from the last glacial maximum to the present. *Nature* 348: 711–714.
- Campbell, G. S. and J. M. Norman. 1998. *Introduction to Environmental Biophysics*. Springer Science + Business Media Inc., New York, USA.
- Flanagan, L. B., L. A. Wever and P. J. Carlson. 2002. Seasonal and inter annual variation in carbon dioxide exchange and carbon balance in a northern temperate grassland. *Global Change Biology* 8: 599–615.
- Lalrammawia, C. and K. Paliwal. 2010. Seasonal changes in net ecosystem exchange of CO₂ and respiration of *Cenchrus ciliaris* L. grassland ecosystem in semi-arid tropics: an eddy covariance measurement. *Current Science* 98:1211-1218.
- Zhang, M., G. R. Yu, L. M. Zhang, X. M. Sun, X. F. Wen, S. J. Han and J. H. Yan. 2009. Impact of cloudiness on net ecosystem exchange of carbon dioxide in different types of forest ecosystems in China. *Biogeosciences Discussions* 6: 8215–8245.

Acknowledgement

This work has been carried out at ICAR- Central Arid Zone Research Institute (CAZRI) under the project titled “Energy and Mass Exchange in Arid Grassland System” which forms a part of SAC national programme “Energy and Mass Exchange in Vegetative Systems (EME-VS)”. Authors are thankful to Space Applications Centre (ISRO) for funding this project and Directors, SAC (ISRO), Ahmedabad and ICAR-CAZRI, Jodhpur for providing logistics support.