

Old Dominion University
ODU Digital Commons

Community & Environmental Health Faculty
Publications

Community & Environmental Health

2020

Strategies for Controlling Violence Against Healthcare Workers: Application of Fuzzy Analytical Hierarchy Process and Fuzzy Additive Ratio Assessment

Fazel Rajabi

Mehdi Jahangiri


Farnaz Bagherifard

Sean Banaee

Old Dominion University, sbanaee@odu.edu

Payam Fahadi

Follow this and additional works at: https://digitalcommons.odu.edu/commhealth_fac_pubs

 Part of the [Bioethics and Medical Ethics Commons](#), [Health and Medical Administration Commons](#),
and the [Health Services Administration Commons](#)

Original Publication Citation

Rajabi, F., Jahangiri, M., Bagherifard, F., Banaee, S., & Fahadi, P. (2020). Strategies for controlling violence against healthcare workers: Application of fuzzy analytical hierarchy process and fuzzy additive ratio assessment. *Journal of Nursing Management*. doi:10.1111/jonm.12989

This Article is brought to you for free and open access by the Community & Environmental Health at ODU Digital Commons. It has been accepted for inclusion in Community & Environmental Health Faculty Publications by an authorized administrator of ODU Digital Commons. For more information, please contact digitalcommons@odu.edu.



Title: Strategies for Controlling Violence Against Healthcare Workers: Application of Fuzzy Analytical Hierarchy Process and Fuzzy Additive Ratio Assessment

Short title: Ranking of control measures for violence

Fazel Rajabi¹, Mehdi Jahangiri^{2*}, Farnaz Bagherifard³, Sean Banaee⁴, Payam Fahadi⁵

¹ PhD student of occupational health engineering, School of Health, Shiraz University of Medical Sciences, Shiraz, Iran

Email: fazel.rajabi70@gmail.com

ORCID: <https://orcid.org/0000-0003-3239-6110>

² Research Center for Health Science, Institute of Health, Associate Professor, Department of Occupational Health, Shiraz University of Medical Sciences

* Address correspondence to; Occupational Health Department, School of Health, Shiraz University of Medical Sciences; POBox: 71645-111, Shiraz, I.R.Iran, Tel: +98 711 7251020, Fax: +98 711 7260225

ORCID : <https://orcid.org/0000-0002-4703-2523>

Email: jahangiri_m@sums.ac.ir

³ MSc of occupational health engineering, Jahrom University of Medical Sciences, Jahrom , Iran

Email: lord.bagherifard@gmail.com

⁴ Assistant Professor, College of Health Sciences, Old Dominion University, Norfolk, Virginia, USA

Email: sbanaee@odu.edu

⁵ Department of Management, Zand Higher Education Institute, Shiraz, Iran.

Email: payamfarhadi88@gmail.com

ORCID: <https://orcid.org/0000-0003-2314-566X>

- **Conflict of Interest statement**

No conflict of interest has been declared by the authors.

- **Funding Statement**

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/JONM.12989](https://doi.org/10.1111/JONM.12989)

This article is protected by copyright. All rights reserved

Strategies for Controlling Violence Against Healthcare Workers: Application of Fuzzy Analytical Hierarchy Process and Fuzzy Additive Ratio Assessment

Abstract

Objective: The present study aimed to identify and prioritize control measures of violence against Healthcare Workers (HWs) using the Fuzzy Analytical Hierarchy Process (FAHP) and Fuzzy Additive Ratio Assessment (F-ARAS).

Background: Occupational violence is a pervasive problem in healthcare centers. Reducing violence against staff is one of the challenges for healthcare managers.

Method: At the first stage, the most common criteria and control options for violence against HWs were identified and extracted using a review of previous studies. At the next stage, criteria for selection of control measures were prioritized using the FAHP. Finally, control measures of workplace violence were prioritized using the F-ARAS method.

Results: Results of the FAHP indicated that safety and efficiency were the most important criteria. Results of the F-ARAS also revealed that “increasing number of security personnel” and “training staff” were the best recommendations for controlling violence against HWs.

Conclusion: Based on expert’s opinions, administrative measures are the optimal ways to control violence at health centers; therefore, it is suggested that violence control programs should be more focused on administrative measures.

Implications for Nursing Management: These results could assist nursing management to take best strategies for controlling occupational violence based on multi-criteria decision-making methods.

Keywords: Occupational violence; Healthcare workers; Ranking; Control strategies; Fuzzy Analytical Hierarchy Process (FAHP); Fuzzy Additive Ratio Assessment (F-ARAS)

Accepted Article

1- Introduction

Violence is a widespread and worrisome problem in workplaces (CDC, 2006; Henry & Ginn, 2002). The National Institute for Occupational Safety and Health (NIOSH) introduced workplace violence as a major occupational hazard, including aggressive behavior such as physical attacks against employees or the threat of physical invasion while at work (CDC, 2006; OSHA, 2015). Workplace violence can be classified into four categories: physical violence (punching, kicking, slapping, pushing, biting, pinching, and wounding by sharp objects), verbal violence (insulting, humiliating, scaring, mocking and abusing), racial violence (threatening due to color and language, nationality, religion, place of birth, or any other situation), and sexual violence (any violent behavior related to gender which is considered as an offense by a person and threatens, insults, or embarrasses people) (Moraveji, Soleymannejad, & Bazargan, 2012).

According to the Bureau of Labor Statistics, during the 7-year period from 2011 to 2017, an average of 447 homicides and 712 intentional injuries by employees occurred annually in the United States (BLS, 2017). Among all working groups and professionals, HWs are the major victims of occupational violence, with their risk of exposure to occupational violence being 16 times higher than other employees (Moeini, Fallahi, Hossaini, & Dalvandi, 2017). Occupational Safety and Health Administration (OSHA) also reported rates of violence in the workplace in healthcare centers as much higher than with social service providers and industries (OSHA, 2015). Results of most studies indicate that more than half of nurses experience violence in the workplace during a year, despite the fact that the incidence rate of violence is not the same in different communities or sectors of healthcare centers (Cezar & Marziale, 2006; Gacki-Smith et al., 2009; Lanza, Zeiss, & Rierdan, 2006; Lin & Liu, 2005; O'Connell, Young, Brooks, Hutchings, & Lofthouse, 2000; Rafati Rahimzadeh, Zabihi, & Hosseini, 2011; Talas, Kocaöz, & Akgüç, 2011).

Violence in healthcare centers has broad consequences (Lanctôt & Guay, 2014; Nikathil, Olaussen, Gocentas, Symons, & Mitra, 2017). Lanctot and Guay (Lanctôt & Guay, 2014) in a comprehensive review study, categorized the consequences of violence in previous studies into seven categories: (1) physical (such as physical injuries), (2) psychological (such as posttraumatic stress disorder, Irritability, sleep disturbances, depression, Burnout, and anxiety), (3) emotional (e.g. anger, sadness, fear, disgust, and surprise), (4) workplace

satisfaction functioning (such as quitting the job, changing employers, reducing performance and productivity and negative effects on job satisfaction) 5) patients' dissatisfaction affecting quality of care, (6) social or personal (e.g. negative effects on victim's relationships with colleagues and family) and (7) financial.

Due to high rates of workplace violence and its negative consequences for victims, it is important to implement strategies that prevent workplace violence or mitigate its negative effects. There is no possibility of performing or testing all control strategies due to various financial and technical limitations. Due to the wide variety of criteria for choosing control strategies, it is difficult to identify optimal strategies. There are many ways to classify control strategies. The use of multiple criteria decision-making (MCDM) method is the most efficient way to achieve this goal. The MCDM is a decision-making tool for complex multi-criteria issues that considers qualitative and quantitative aspects of problems in the decision-making process. These methods scientifically and systematically determine decision-maker's priorities to help them choose between several options that usually have different outcomes (Garfi, Tondelli, & Bonoli, 2009).

Despite the fact that a great number of studies have been conducted on occupational violence against HWs, most of them have not reported statistics for the incidence of violence and its contributing factors (Cezar & Marziale, 2006; Cheraghi et al., 2014; Dehnadi-Moghaddam et al., 2013; Gillespie, Gates, Miller, & Howard, 2010; Moraveji et al., 2012; Mozafari & Tavan, 2013; Soheili, Mohammadpour, Jafarizadeh, Habibzadeh, & Mehryar, 2014; Talas et al., 2011); moreover few studies have investigated strategies for preventing or controlling violence (Havaei, Macphee, & Lee, 2019; Morphet, Griffiths, Beattie, Reyes, & Innes, 2018; Spelten et al., 2017; Vladutiu, Casteel, Nocera, Harrison, & Peek-Asa, 2016; Zamanzadeh & Abdollahzadeh, 2007). The aim of this study is to prioritize violence control strategies using scientific and valid methods such as the MCDM and to cover the existing limitations.

2- Method

The present study employed a cross-sectional, qualitative and applied research design to identify and prioritize methods for preventing or reducing violence against HWs. The research was conducted at three dependent stages as shown in Fig. 1. The total sample size of the study consisted of 40 staff and managers (experts) of hospitals (Blinded for peer review)(15 in the second phase and 25 in the third phase of the study). The selection of effective criteria for choosing control strategies was conducted after consulting 15 experts,

including occupational health experts working in health centers and hospital managers. The second statistical sample consisted of 25 physicians, nurses, security forces, and managers of hospitals who had higher knowledge about the risk of occupational violence and violence prevention methods or who were more exposed to violence. Both panels of experts had more than 5 years of experience. To avoid bias, two samples were selected independently. Like other studies conducted by the MCDM, a purposive sampling method was done in the present study; and the sample size was calculated according to similar studies (Cancela, Fico, & Waldmeyer, 2015; Hsu, Hsieh, Yang, & Lu, 2015; Mühlbacher, Juhnke, & Kaczynski, 2016; Olivieri et al., 2012; Rajabi, Jahangiri, Molaeifar, Honarbakhsh, & Farhadi, 2018). While there is no absolute law on the number and method of choosing members of the expert group in MCDM methods, sampling methods in most studies conducted in this field are purposive (non-random). The sample size is less than 30 (Mühlbacher et al., 2016; Rajabi et al., 2018).

Figure 1: Steps of study [about here]

Stage 1: Identification of control criteria and options

At this stage, 8 general criteria were extracted based on a review of conducted studies conducted on the selection of control strategies (Ishaqi, Golmohammadi, & Khorram, 2012; Molaei Far et al., 2018; Sekhavati, Zadeh, Fam, & Zarandi, 2014). Subsequently, studies conducted on strategies for prevention or reducing the occupational violence against HWs were reviewed (Berry, McNeely, Beauregard, & Lagoe, 2013; Gallant-Roman, 2008; D. Gates et al., 2011; D. M. Gates, Ross, & McQueen, 2006; Gillespie et al., 2010; Henry & Ginn, 2002; Morphet et al., 2018; OSHA, 2015 ; Carol W. Runyan, 2001; Vladutiu et al., 2016; Wassell, 2009), and the most commonly used control strategies were extracted. The output of this stage constituted the input of the second and third research stages respectively. The criteria and options identified in the first step were then reviewed and edited by a panel of experts.

Stage 2: Weighing the criteria by the Fuzzy Analytic Hierarchy Process

In the present study, the Fuzzy Analytic Hierarchy Process (FAHP) was utilized to weigh the criteria for choosing control strategies. The FAHP is the upgraded version of the Analytic Hierarchy Process (AHP) in which fuzzy numbers are used instead of absolute values (Rokhsari & Sadeghi-Niaraki, 2015). The use of a fuzzy system increases the power of the method and is widely used in uncertainty conditions (Mamdani, 1974). Fuzzy theory, which is interpreted as inaccurate logic, is a part of operations research science in which quantitative

and qualitative data are converted through new ways (Chang, 1996). In a fuzzy system, linguistic variables are used instead of numbers; then linguistic scales are converted to triangular or trapezoidal numbers.

Since the measurement of some criteria was qualitative and subjective and because of the lack of numerical value for qualitative data, the evaluation of criteria was done on a linguistic scale. The triangular fuzzy numbers were used to show linguistic variables (Table 1).

Table 1: Linguistic Scale and corresponding fuzzy numbers for FAHP (Rajabi et al., 2018)

Linguistic Scale	Triangular fuzzy number	Reciprocal value of triangular fuzzy number
Just equal	(1, 1, 1)	(1, 1, 1)
Equally important	(0.5, 1, 1.5)	(0.66667, 1, 2)
Weakly important	(1, 1.5, 2)	(0.5, 0.66667, 1)
Strong more important	(1.5, 2, 2.5)	(0.4, 0.5, .66667)
Very Strong more important	(2, 2.5, 3)	(0.3333, 0.4, 0.5)
Absolute more important	(2.5, 3, 3.5)	(0.285714, 0.333, 0.4)

The geometric mean formula was used to combine the results of the expert group. Chang's extent analysis method, as the most widely used and simplest method, was applied to solve the FAHP calculations (Aggarwal & Singh, 2013; Chang, 1996). Details of FAHP calculations with the extent analysis method are presented in Appendix A.

One of the important issues in the FAHP method is the calculation of the consistency of expert opinions. To measure this, the pairwise comparison matrixes were evaluated by the Consistency Ratio (CR). The CR was obtained from dividing the Consistency Index (CI) by the random Consistency Index (RI) that was obtained from guidelines (Mazurek, 2017). If the calculated consistency ratio is less than 0.1, then the paired comparison and judgments are acceptable. Otherwise, the study continues until the desired consensus is reached. In this study, the second phase was repeated three times. For the first time, minor changes were made based on the opinions of the expert group on the selected criteria to reach a consensus of expert opinions.

(C) Prioritizing control strategies for occupational violence using the Fuzzy-ARAS method

After determining the weights of criteria in the previous step, control options (alternatives) of violence against the HWs were prioritized using the Fuzzy-ARAS method. This method is a relatively new technique that was developed by Zavadsk and Tursk who argued that complicated phenomena in the world could be understood by simple comparisons. This technique is a powerful way to choose the degree of utility of different options and is relatively simple (Turskis & Zavadskas, 2010). Due to the capabilities of this method, it has been used in several different studies (Keršulienė & Turskis, 2011, 2014; Turskis, Lazauskas, & Zavadskas, 2012). This study appears to be the first that conducted research with the F-ARAS method in the field of healthcare.

At this stage, the panel of experts was asked to evaluate the control strategies based on weighted criteria from the previous stage. To this end, the linguistic scales were used and converted to corresponding absolute scales (Table 2).

Table 2: Linguistic scale and related fuzzy numbers for F-ARAS

Linguistic Scale	Fuzzy number
Very Good (VG)	(7, 9, 9)
Good (G)	(5, 7, 9)
Fair (F)	(3, 5, 7)
Poor (P)	(1, 3, 5)
Very Poor (VP)	(1, 1, 3)

The optimality function and degree of the alternative utility were used to prioritize control strategies by the F-ARAS method. Higher values of these two indices for each of the control strategies indicated that they were better than the other control strategies. Details of F-ARAS calculations are provided in appendix A.

3- Findings

In this study, 8 criteria and 17 options for controlling violence against healthcare workers were identified and prioritized using the combination of FAHP and F-ARAS methods. Figure 2 indicates the results of calculating the relative weight of criteria for the selection of control measures using the FAHP method. Among 8 identified criteria, "safety", "efficiency", and "possibility for implementation" had the highest weights.

Figure 2. Relative weight for selection criteria of violence control measures using the FAHP method [about here]

The final results of the study indicated that among 17 investigated control strategies, “increasing number of security personnel”, “training staff on how to manage occupational violence”, and “prevention or restriction of the employee from work alone in high-risk area” were the best strategies for reducing or preventing occupational violence. Table 3 and Figure 3 present the results of prioritizing control strategies for occupational violence using the F-ARAS method.

Table 3: Identified control strategies for violence against healthcare workers and their values of the optimality function (S_i) by the F-ARAS method

Code	Alternatives (control measures)	S_i
A1	Elimination or substitution therapeutic procedures where the risk of violence is high	0.2773
A2	Using physical barriers (such as guards or door locks to reduce employee exposure to the hazard)	0.2911
A3	Installation of metal detectors	0.2825
A4	Installation of panic buttons or pagers	0.2989
A5	Improving lighting in indoor and outdoor areas	0.3172
A6	Increasing the number of exit routes in each space	0.3000
A7	Monitoring workplace using closed circuit videos	0.3137
A8	Providing impact-resistant and bulletproof glass at nurses' stations	0.2812
A9	Increasing the depth of counters at nurses' stations	0.2870
A10	Providing comfortable areas to reduce stress of patients and their companion	0.3108
A11	Elimination of items that could be used as weapons	0.3209
A12	Informing visitors about consequences of violence against HWs	0.3127
A13	Control of visitor entry	0.3187
A14	Prevention or restriction of employee from working alone in high risk areas	0.3233
A15	Post-incident tracking	0.3201
A16	Increasing number of security personnel	0.3523
A17	Training staff on how to manage occupational violence	0.3388

Figure 3: Prioritization results of control strategies based on the degree of utility using F-ARAS method [about here]

4- Discussion

The present study aimed to identify the criteria and options of control measures to prevent violence against HWs and rank them using a combination of FAHP and F-ARAS methods. The results obtained from the consensus of experts by the FAHP method indicated that the "safety" and "efficiency" criteria had the highest weights and should be prioritized when selecting of control measures of violence against HWs.

Results of ranking control measures based on the F-ARAS method also indicated that "increasing the number of security personnel," "training staff on how to manage occupational violence," and "preventing or restricting of employees from working alone in high-risk areas" had the highest impact on reducing or preventing occupational violence. As mentioned earlier, there are few studies on methods for controlling occupational violence against HWs, and there are no studies that systematically prioritize methods of controlling violence. Nonetheless, several studies have examined violence control strategies and the results of most of them are consistent with the present study. Stirling et al. found that increasing the number of security staff and improving security systems were effective for violence control (Stirling, Higgins, & Cooke, 2001). Zamanzadeh et al. also mentioned the limiting clients' access to hospitals as the most common approach for controlling violence against nurses (Zamanzadeh & Abdollahzadeh, 2007).

According to the results of the present study, increasing the number of security personnel and training staff were the best ways to control violence in healthcare centers. Many studies emphasized the role of education and increased awareness of staff and patients in reducing the incidence of violence or reducing consequences of violence in healthcare centers (Deans, 2004; Fernandes et al., 2002; D. Gates et al., 2011; Ishak & Christensen, 2002; Carol W Runyan, Zakocs, & Zwerling, 2000), although there were many contradictions in the effectiveness of training in previous studies (Wassell, 2009). In general, education should be provided to enhance the staff's knowledge, attitude, and skill for preventing or controlling verbal and physical violence (Hills et al., 2015). Training programs should be implemented, such as increasing the staff's knowledge about legal responsibilities, organizational procedures and policies, and strategies for interacting with patients to prevent violence (CDC, 2006). OSHA also mentions education as a key component of programs to address violence and emphasizes the need for educational programs in two ways, including general and specialized education for employees, managers, supervisors and security personnel in

healthcare centers. OSHA also recommends applying policies and procedures for preventing violence in the workplace, identifying risk factors and violent behavior, strategies for interacting with patients and managing violence in the workplace, training programs designed to familiarize employees with safety equipment such as alarm systems, and methods of protecting themselves and colleagues during the violent episodes (OSHA, 2015). Contrary to the above studies, the study by Havaei et al showed that nursing education was not an effective violence prevention strategy for nurses' perceptions of safety (Havaei et al., 2019).

Based on the consensus of experts with regard to all criteria for choosing a control strategy, administrative measures are more appropriate towards engineering (environmental) control measures of violence against the HWs. However, it is always recommended that administrative measures should be taken after the implementation of engineering measures in order to reduce the residual risk, or if engineering measures cannot be implemented (OSHA, 2015). Most studies have found that implementing environmental control measures was the most effective way to address workplace violence. However, these solutions seem to be more effective based on the availability of social and economic resources; and management practices are more effective than environmental measures to control the risk of violence in healthcare centers (Wassell, 2009). In fact, various aspects affecting the selection of the appropriate strategy such as cost, capability for implementation, and the comprehensiveness of the model should be taken into account when implementing a control strategy. Accordingly, most studies conducted on the control of violence in medical centers have suggested that the implementation of organizational and managerial interventions (including educational programs) is most effective in improving the control of violence against staff in healthcare centers (Barish, 2001; Flannery Jr et al., 1998; Henry & Ginn, 2002; Morrison & Love, 2003). Some studies have shown that the focus of organizational measures on patients, like employees, can reduce the incidence of violence in medical centers. In one of these studies, Kraus found that the patient management strategy reduced type II violence (violence of patients or relatives against employees) in medical centers (Kraus, Blander, & McArthur, 1995).

Despite the greater impact of managerial and organizational strategies over environmental solutions (engineering), the results of the present study indicated that roles of environmental strategies such as the removal of hazardous equipment, appropriate environmental designs, providing adequate lighting, using closed circuit videos, and implementing physical protections and barriers should not be ignored in the control of violence. Therefore, it is

suggested that environmental strategies be used along with administrative practice in violence control programs at workplaces. This result was consistent with the findings of several studies; in one of these studies, Arnetz found that intervention strategies such as increasing the number of security personnel, training staff, designing panic buttons, increasing the number of closed circuit video cameras, and modifying shifts to prevent staff from working alone in a unit were useful in reducing the rate of violence in the workplace (Arnetz et al., 2017). The results of a study by Hunter and Love indicated that an intervention plan including several environmental and administrative measures was very effective in reducing violence in hospitals (Hunter & Love, 1996).

5- Research Limitations

The type of violence control strategies may vary depending on the technical, economic, managerial, and cultural characteristics of the study location. Therefore, violence control measures investigated in this study may not be comprehensive and fully generalizable to healthcare centers in other regions.

6- Conclusion and practical suggestions

According to the results of the present study, measures should be taken to increase the number of security personnel and to train staff and clients of healthcare centers in order to control the occupational violence against HWs. Moreover, work schedules should be modified to prevent employees working alone in a unit and there should be control over entry and exit to the unit. Staff training programs should include ways to quickly identify violent and stressful behavior or identify any symptom or situation that may lead to violence in the workplace, strategies for the management and control of people during incidences of violence at work, and appropriate methods for dealing with patients and their relatives to prevent the occurrence of violence against HWs. To create a more effective violence control program, it is necessary to consider environmental solutions such as installing CCTV cameras, redesigning hazardous areas, and removing dangerous equipment that may be used violently.

7- Implications for nursing managers

Due to the diversity of quantitative and qualitative criteria, decision making is one of the most difficult responsibilities of managers, including nursing managers. Therefore, in recent years mathematical and computer science and new techniques such as multi-criteria decision making techniques have been used as a decision support system (DSS) to help managers make optimal decisions. In this study, we introduced the FAHP and F-ARAS methods to assist nursing managers in deciding on the optimal control strategies for violence against healthcare workers.

References:

- Aggarwal, R., & Singh, S. (2013). *AHP and extent fuzzy AHP approach for prioritization of performance measurement attributes*. Paper presented at the Proceedings of World Academy of Science, Engineering and Technology.
- Arnetz, J. E., Hamblin, L., Russell, J., Upfal, M. J., Luborsky, M., Janisse, J., & Essenmacher, L. (2017). Preventing patient-to-worker violence in hospitals: outcome of a randomized controlled intervention. *Journal of occupational and environmental medicine*, 59(1), 18. doi:doi: 10.1097/JOM.0000000000000909.
- Barish, R. C. (2001). Legislation and regulations addressing workplace violence in the United States and British Columbia. *American Journal of Preventive Medicine*, 20(2), 149-154.
- Berry, C., McNeely, A., Beauregard, K., & Lagoe, W. (2013). Workplace Violence Prevention Guidelines and Program for Health Care, Long Term Care and Social Services Workers. *N.C. Department of Labor. Occupational Safety and Health Program*.
- BLS. (2017). National census of fatal occupational injuries in 2017. *Department of Labor, Bureau of Labor Statistics*(<https://www.bls.gov/news.release/pdf/cfoi.pdf>).
- Cancela, J., Fico, G., & Waldmeyer, M. T. A. (2015). Using the analytic hierarchy process (AHP) to understand the most important factors to design and evaluate a telehealth system for Parkinson's disease. *BMC medical informatics and decision making*, 15(3), S7.
- CDC. (2006). *Centers for Disease Control and Prevention. Workplace violence prevention strategies and research needs*. Paper presented at the Conference Partnering in Workplace Violence Prevention Translating Research to Practice.
- Cezar, E. S., & Marziale, M. H. P. (2006). Occupational violence problems in an emergency hospital in Londrina, Paraná, Brazil. *Cadernos de Saúde Pública*, 22(1), 217-221. doi:DOI: /S0102-311X2006000100024
- Chang, D.-Y. (1996). Applications of the extent analysis method on fuzzy AHP. *European journal of operational research*, 95(3), 649-655.

- Accepted Article
- Cheraghi, M. A., Javanmardi, S., Zolfaghari, M., Bahrani, N., Akbari, A., & Javanmardi, Z. (2014). Predisposing Factors of Violence against Nurse in Emergency Department from Nurses', Patients', and accompanies' perspective. *Medycyna pracy*.
- Deans, C. (2004). Nurses and occupational violence: The role of organisational support in moderating professional competence. *Australian Journal of Advanced Nursing, The*, 22(2), 14.
- Dehnadi-Moghaddam, A., Asgharzadeh, A., S, H., Kouchakinejad-Eramsadati, L., kazemnejad-Leili, E., & Baghernia Hemmati, N. (2013). Frequency and Characteristics of Violence against Nurses of Emergency. *health and care, 14*(in persian), 62-68.
- Fernandes, C. M., Raboud, J. M., Christenson, J. M., Bouthillette, F., Bullock, L., Ouellet, L., & Moore, C. C. F. (2002). The effect of an education program on violence in the emergency department. *Annals of emergency medicine, 39*(1), 47-55.
- Flannery Jr, R. B., Hanson, M. A., Penk, W. E., Goldfinger, S., Pastva, G. J., & Navon, M. A. (1998). Replicated declines in assault rates after implementation of the Assaulted Staff Action Program. *Psychiatric Services, 49*(2), 241-243.
- Gacki-Smith, J., Juarez, A. M., Boyett, L., Homeyer, C., Robinson, L., & MacLean, S. L. (2009). Violence against nurses working in US emergency departments. *Journal of Nursing Administration, 39*(7/8), 340-349. doi:doi: 10.1097/NNA.0b013e3181ae97db.
- Gallant-Roman, M. A. (2008). Strategies and tools to reduce workplace violence. *AAOHN journal, 56*(11), 449-454.
- Garfi, M., Tondelli, S., & Bonoli, A. (2009). Multi-criteria decision analysis for waste management in Saharawi refugee camps. *Waste management, 29*(10), 2729-2739.
- Gates, D., Gillespie, G., Smith, C., Rode, J., Kowalenko, T., & Smith, B. (2011). Using action research to plan a violence prevention program for emergency departments. *Journal of Emergency Nursing, 37*(1), 32-39.
- Gates, D. M., Ross, C. S., & McQueen, L. (2006). Violence against emergency department workers. *The Journal of emergency medicine, 31*(3), 331-337.

- Gillespie, G. L., Gates, D. M., Miller, M., & Howard, P. K. (2010). Workplace violence in healthcare settings: risk factors and protective strategies. *Rehabilitation Nursing*, 35(5), 177-184.
- Havaei, F., Macphee, M., & Lee, S. E. (2019). The effect of violence prevention strategies on perceptions of workplace safety: A study of medical- surgical and mental health nurses. *Journal of advanced nursing*.
- Henry, J., & Ginn, G. O. (2002). Violence prevention in healthcare organizations within a total quality management framework. *Journal of Nursing Administration*, 32(9), 479-486.
- Hills, D. J., Ross, H. M., Pich, J., Hill, A. T., Dalsbø, T. K., Riahi, S., . . . Martínez-Jarreta, B. (2015). Education and training for preventing and minimising workplace aggression directed toward healthcare workers.
- Hsu, J. C., Hsieh, C.-Y., Yang, Y.-H. K., & Lu, C. Y. (2015). Net clinical benefit of oral anticoagulants: a multiple criteria decision analysis. *PloS one*, 10(4), e0124806.
- Hunter, M. E., & Love, C. C. (1996). Total quality management and the reduction of inpatient violence and costs in a forensic psychiatric hospital. *Psychiatric services (Washington, DC)*, 47(7), 751-754.
- Ishak, M., & Christensen, M. (2002). Achieving a better management for patients' aggressive behaviour: evaluation of a training program. *Journal of Occupational Health and Safety Australia and New Zealand*, 18(3), 231-238.
- Ishaqi, M., Golmohammadi, R., & Khorram, M. R. (2012). Prioritization of noise control methods in Hamedan Shine Co using the Analytical Hierarchical Process (AHP). *Occupational Health and Safety*, 3.
- Keršulienė, V., & Turskis, Z. (2011). Integrated fuzzy multiple criteria decision making model for architect selection. *Technological and Economic Development of Economy*, 17(4), 645-666.
- Keršulienė, V., & Turskis, Z. (2014). An integrated multi-criteria group decision making process: selection of the chief accountant. *Procedia-Social and Behavioral Sciences*, 110, 897-904.
- Kraus, J. F., Blander, B., & McArthur, D. L. (1995). Incidence, risk factors and prevention strategies for work-related assault injuries: a review of what is known, what needs to

be known, and countermeasures for intervention. *Annual Review of public health*, 16(1), 355-379.

Lanctôt, N., & Guay, S. (2014). The aftermath of workplace violence among healthcare workers: A systematic literature review of the consequences. *Aggression and violent behavior*, 19(5), 492-501.

Lanza, M. L., Zeiss, R. A., & Rierdan, J. (2006). Non-physical violence: a risk factor for physical violence in health care settings. *AAOHN journal*, 54(9), 397-402. doi:DOI: 10.1177/216507990605400903

Lin, Y.-H., & Liu, H.-E. (2005). The impact of workplace violence on nurses in South Taiwan. *International journal of nursing studies*, 42(7), 773-778. doi:DOI: 10.1016/j.ijnurstu.2004.11.010

Mamdani, E. H. (1974). *Application of fuzzy algorithms for control of simple dynamic plant*. Paper presented at the Proceedings of the institution of electrical engineers.

Mazurek, J. (2017). On inconsistency indices and inconsistency axioms in pairwise comparisons. *arXiv preprint arXiv:1703.05204*.

Moeini, Z., Fallahi, K., Hossaini, M., & Dalvandi, A. (2017). Nurse's knowledge, attitude and practice toward workplace violence in hospitals of isfahan university of medical sciences. *Health Promotion Management*(in persian).

Molaei Far, H., Rajabi, F., Zamanian, Z., Honarbakhsh, M., Ebrahimi, A., & Farhadi, P. (2018). Identifying and ranking the strategies of control exposure to heat stress in the rubber industry using a Delphi method and combination of Fuzzy AHP and TOPSIS approach. *Iran Occupational Health Journal*, 15(4), 21-33.

Moraveji, M., Soleymannejad, N., & Bazargan, M. (2012). Workplace violence against nurses working in hospitals of Zanjan province. *Iranian Journal of Health & Care*(persian).

Morphet, J., Griffiths, D., Beattie, J., Reyes, D. V., & Innes, K. (2018). Prevention and management of occupational violence and aggression in healthcare: A scoping review. *Collegian*.

Morrison, E. F., & Love, C. C. (2003). An evaluation of four programs for the management of aggression in psychiatric settings. *Archives of Psychiatric Nursing*, 17(4), 146-155.

- Mozafari, M., & Tavan, H. (2013). Survey of violence against nursing in the Ilam Training Hospitals at 2012. *Scientific Journal of Ilam University of Medical Sciences*(persian).
- Mühlbacher, A., Juhnke, C., & Kaczynski, A. (2016). Patients' Priorities in the Treatment of Neuroendocrine Tumours: An Analytical Hierarchy Process. *Gesundheitswesen (Bundesverband der Ärzte des Öffentlichen Gesundheitsdienstes (Germany))*, 78(10), 651-657.
- Nikathil, S., Olaussen, A., Gocentas, R. A., Symons, E., & Mitra, B. (2017). Workplace violence in the emergency department: A systematic review and meta analysis. *Emergency Medicine Australasia*, 29(3), 265-275.
- O'Connell, B., Young, J., Brooks, J., Hutchings, J., & Lofthouse, J. (2000). Nurses' perceptions of the nature and frequency of aggression in general ward settings and high dependency areas. *Journal of clinical Nursing*, 9(4), 602-610.
- Olivieri, A., Marchetti, M., Lemoli, R., Tarella, C., Iacone, A., Lanza, F., . . . Bosi, A. (2012). Proposed definition of 'poor mobilizer' in lymphoma and multiple myeloma: an analytic hierarchy process by ad hoc working group Gruppo Italiano Trapianto di Midollo Osseo. *Bone marrow transplantation*, 47(3), 342.
- OSHA. (2015). Occupational Safety and Health administration. Guidelines for Preventing Workplace Violence for Health Care & Social (available at: www.osha.gov (accessed: 15 Sep 2017)).
- Rafati Rahimzadeh, M., Zabihi, A., & Hosseini, S. (2011). Verbal and physical violence on nurses in hospitals of Babol University of Medical Sciences. *Journal of hayat*, 17(2), 5-11. doi:in persian
- Rajabi, F., Jahangiri, M., Molaeifar, H., Honarbakhsh, M., & Farhadi, P. (2018). Occupational stress among nurses and pre-hospital emergency staff: application of fuzzy analytic hierarchy process (FAHP) method. *EXCLI journal*, 17, 808.
- Rokhsari, S., & Sadeghi-Niaraki, A. (2015). Urban network risk assessment using Fuzzy-AHP and TOPSIS in GIS environment. *Iranian Journal of Operations Research*, 6(2), 73-86.
- Runyan, C. W. (2001). Moving forward with research on the prevention of violence against workers. *American Journal of Preventive Medicine*, 20(2), 169-172. doi:[https://doi.org/10.1016/S0749-3797\(00\)00294-4](https://doi.org/10.1016/S0749-3797(00)00294-4)

- Runyan, C. W., Zakocs, R. C., & Zwerling, C. (2000). Administrative and behavioral interventions for workplace violence prevention. *American Journal of Preventive Medicine*, 18(4), 116-127.
- Sekhavati, E., Zadeh, M. M., Fam, E. M., & Zarandi, A. F. (2014). Prioritizing methods of control and reduce noise pollution in Larestan cement Factory using analytical hierarchy process (AHP). *Toloo-e-behdasht*, 13(2), 156-167.
- Soheili, A., Mohammadpour, Y., Jafarizadeh, H., Habibzadeh, H., & Mehryar, H. (2014). Violence against nurses in emergency departments of Urmia university hospitals in 2013. *Journal of Urmia Nursing and Midwifery Faculty*, 12(9), 874-882. doi:in persian
- Spelten, E., Thomas, B., O'Meara, P. F., Maguire, B. J., FitzGerald, D., & Begg, S. J. (2017). Organisational interventions for preventing and minimising aggression directed toward healthcare workers by patients and patient advocates. *Cochrane Database of Systematic Reviews*(5).
- Stirling, G., Higgins, J. E., & Cooke, M. W. (2001). Violence in A&E departments: a systematic review of the literature. *Accident and emergency nursing*, 9(2), 77-85.
- Talas, M. S., Kocaöz, S., & Akgüç, S. (2011). A survey of violence against staff working in the emergency department in Ankara, Turkey. *Asian Nursing Research*, 5(4), 197-203. doi:DOI: 10.1016/j.anr.2011.11.001
- Turskis, Z., Lazauskas, M., & Zavadskas, E. K. (2012). Fuzzy multiple criteria assessment of construction site alternatives for non-hazardous waste incineration plant in Vilnius city, applying ARAS-F and AHP methods. *Journal of Environmental Engineering and Landscape Management*, 20(2), 110-120.
- Turskis, Z., & Zavadskas, E. K. (2010). A new fuzzy additive ratio assessment method (ARAS- F). Case study: The analysis of fuzzy multiple criteria in order to select the logistic centers location. *Transport*, 25(4), 423-432.
- Vladutiu, C. J., Casteel, C., Nocera, M., Harrison, R., & Peek- Asa, C. (2016). Characteristics of workplace violence prevention training and violent events among home health and hospice care providers. *American journal of industrial medicine*, 59(1), 23-30.

Accepted Article

Wassell, J. T. (2009). Workplace violence intervention effectiveness: A systematic literature review. *Safety Science*, 47(8), 1049-1055.

Zamanzadeh, V., & Abdollahzadeh, F. (2007). Nature of violence toward nurses working in hospitals. *Medical journal of Tabriz University of Medical Sciences and Health Services*, 29(2), 61-66.

Appendix A: Detailed description of FAHP and FARAS

A.1. FAHP

Equation A.1 shows a general form of fuzzy triangular numbers:

$$\mu_{\tilde{m}}(x) = \begin{cases} (x-l)/(m-l) & l \leq x \leq m \\ (u-x)/(u-m) & m \leq x \leq u \\ 0 & \text{other wise} \end{cases} \quad \text{Eq.A.1}$$

where l is the lower limit, u is the upper limit, and m is the median value. Therefore, fuzzy triangular numbers are shown as (l, m, u) .

The steps of the calculation of FAHP based on the Extent Analysis (EA) method in this study were as follows (Chang, 1996):

Step 1: The value of fuzzy combination value of i is expressed according to the Equation A.2:

$$S_i = \sum_{j=1}^m M_{gi}^j \quad \left[\sum_{i=1}^n \sum_{j=1}^m M_{ij} \right]^{-1} \quad \text{Eq.A.2}$$

The values of $\sum_{j=1}^m M_{gi}^j$ is calculated using the equations A.3 and A.4:

$$\sum_{j=1}^m M_{gi}^j = \left(\sum_{j=1}^m l_i \sum_{j=1}^m m_i \sum_{j=1}^m u_i \right) \quad \text{Eq.A.3}$$

$$\sum_{i=1}^n \sum_{j=1}^m M_{gi}^j = \left(\sum_{j=1}^m l_i \sum_{j=1}^m m_i \sum_{j=1}^m u_i \right) \quad \text{Eq.A.4}$$

The inverse of equation A.5 was calculated as follows:

$$\left[\sum_{i=1}^n \sum_{j=1}^m M_{gi}^j \right]^{-1} = \left(\frac{1}{\sum_{j=1}^m l_i}, \frac{1}{\sum_{j=1}^m m_i}, \frac{1}{\sum_{j=1}^m u_i} \right) \quad \text{Eq. A.5}$$

Step 2: If $M_2 = (l_2, m_2, u_2)$ and $M_1 = (l_1, m_1, u_1)$ are two fuzzy numbers, the probability degree is calculated according to Equation A.6:

$$V(M_1 \geq M_2) = y \geq x [\min(\mu_{M_1}(x), \mu_{M_2}(y))] \quad \text{Eq.A.6}$$

This equation can be expressed as follows:

$$V(M_2 \geq M_1) = hgt(M_1 \cap M_2) = \mu_{M_2}(d) = \left\{ \begin{array}{ll} 1, & \text{if } m_2 \geq m_1 \\ 0, & \text{if } l_1 \geq u_2 \\ \frac{l_1 - u_2}{(m_2 - u_2) - (m_1 - l_1)}, & \text{otherwise} \end{array} \right\} \quad \text{Eq. A.7}$$

Where, d is the length of the highest intersection between μ_{M_1} and μ_{M_2} .

Figure 4: The intersection between M_1 and M_2 [about here]

Step 3: The probability degree of a fuzzy point like M_i larger than point K is calculated using the Equation A.8:

$$V(M \geq M_1, M_2, \dots, M_K) = V [(M \geq M_1) \text{ and } (M \geq M_2) \text{ and } \dots (M \geq M_K)] = \min V (M \geq M_i), i = 1, 2, 3, \dots, k \quad \text{Eq.A.8}$$

Assuming that the following equation is true:

$$d'(A_i) = \min V (S_i \geq S_k) \quad \text{Eq. A. 9}$$

The weight of each vector is calculated according to the Equation A.10:

$$W' = (d'(A_1), d'(A_2), \dots, d'(A_n))^T \quad \text{Eq. A. 10}$$

Step 4: Finally, the normalized weight of each vector is calculated according to Equation A.11:

$$W = (d(A_1), d(A_1), \dots, d(A_n))^T \quad \text{Eq. A. 11}$$

Where, W is a non-fuzzy number.

A.1. F-ARAS

The main steps of the F-ARAS method to prioritize control strategies of the present study were as follows (Turskis & Zavadskas, 2010):

Step 1: Forming of a Fuzzy Decision Matrix

The general form of ARAS decision-making matrix is as follows:

$$\tilde{X} = \begin{bmatrix} \tilde{x}_{01} & \dots & \tilde{x}_{0j} & \dots & \tilde{x}_{0n} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ \tilde{x}_{i1} & \dots & \tilde{x}_{ij} & \dots & \tilde{x}_{in} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ \tilde{x}_{m1} & \dots & \tilde{x}_{mj} & \dots & \tilde{x}_{mn} \end{bmatrix}, \quad i = \overline{0, m}; j = \overline{1, n} \quad \text{Eq.A.12}$$

Where, m is the number of alternatives; n is the number of criteria; \tilde{x}_{ij} is the fuzzy value representing the i alternative based on j alternative; and \tilde{x}_{0j} is the optimal value of j criterion:

$$\tilde{x}_{0j} = \max_{\tilde{x}_{ij}, i} \text{, if } \max_{\tilde{x}_{ij}, i} \text{ is preferable, and}$$

$$\tilde{x}_{0j} = \min_{\tilde{x}_{ij}, i} \text{, if } \min_{\tilde{x}_{ij}, i} \text{ is preferable} \quad \text{Eq.A.13}$$

Step 2: Normalization of initial values of the decision matrix (\tilde{x}_{ij})

The following equation was used to normalize positive criteria:

$$\tilde{x}_{ij} = \frac{\tilde{x}_{ij}}{\sum_{i=0}^m \tilde{x}_{ij}} \quad \text{Eq.A.14}$$

Negative criteria were also normalized using the following equation:

$$\tilde{x}_{ij} = \frac{\tilde{x}_{ij}}{\tilde{x}_{ij}^*} \quad \tilde{x}_{ij} = \frac{\tilde{x}_{ij}}{\sum_{i=0}^m \tilde{x}_{ij}} \quad \text{Eq.A.15}$$

Step 3: Calculation of the normalized-weighted matrix (\tilde{x}_{ij}):

The following equation was used to calculate normalized-weighted matrix:

$$\tilde{x}_{ij} = \tilde{x}_{ij} \tilde{w}_j \quad i = \overline{0, m} \quad \text{Eq.A.16}$$

Where, w_j is the weight of each criterion and was calculated in the previous step using the FAHP.

Step 4: Determination the value of the optimality function and degree of the alternative utility

The value of the optimality i function is obtained according to the following equation:


$$\tilde{s}_i = \sum_{j=1}^n \tilde{x}_{ij}; \quad i = \overline{0, m} \quad \text{Eq.A.17}$$

Alternative with larger \tilde{s}_i is preferable to other alternatives. The following equation is used for defuzzification:


$$\tilde{s}_i = \frac{1}{3} (s_{i\alpha} + s_{i\beta} + s_{i\gamma}) \quad \text{Eq.A.18}$$

The degree of the alternative utility is calculated as follows:


$$k_i = \frac{s_i}{s_0} \quad i = \overline{0, m} \quad \text{Eq.A.19}$$




jonm_12989_f1.jpg



jonm_12989_f2.jpg



jonm_12989_f3.jpg



jonm_12989_f4.jpg