

University of Northern Colorado

Scholarship & Creative Works @ Digital UNC

2020 Undergraduate Presentations

Research Day 2020

4-2020

Analysis of Canabinoids in Natural and Synthetic Samples

Juan M. Chavez

Destinee Spurlin

Follow this and additional works at: https://digscholarship.unco.edu/ug_pres_2020

Analysis of Cannabinoids in Natural and Synthetic Samples

Juan M. Chavez and Destinee Spurlin

Mentors: Dr. Corina E. Brown and Dr. Richard M. Hyslop

Abstract

Recently, there has been an increased focus on the analysis of natural and synthetic cannabinoids. Several classes of cannabinoids exist, each with different chemical and pharmacological effects. The amount of each cannabinoid present varies with each strain of *Cannabis*, thus making accurate qualitative and quantitative is very important. Several commercial samples were analyzed by HPLC and GC for qualitative purposes.

Introduction

- Industrial hemp is a relatively new and rapidly expanding field.
- Production and sales of isolate and distillate requires potency analysis and some required qualitative analysis.
 - Production of $\Delta 9$ -THC free distillates and isolates
- Synthetic cannabinoids involves series of reactions to convert starting material to a desired material
 - Ex: Conversion of CBDV to CBD

Purpose

- Analyze synthetic CBD sample from Mile High Labs
 - Possible trace amounts of other cannabinoids

Methods






Column:	Luna Omega 5 μ m Polar C18 100 LC 150x4.6 mm
Mobile Phase A:	20 mM Ammonium Formate pH=3.2
Mobile Phase B:	HPLC-grade acetonitrile
Method Type:	Gradient
Flow Rate:	1.2 mL/min

Time=	Mobile Phase A	Mobile Phase B
0 min	40%	60%
9 min	5%	95%
12 min	5%	95%
16 min	40%	60%
20 min	40%	60%

Conclusion

- The samples Canna Pure Synthetic CBD and 98.5% CBD Isolate were shown to be the most pure samples of CBD.
 - $\Delta 9$ -THC detection
- The unknown synthetic sample contained the largest amounts of other minor cannabinoids
 - CBDV, $\Delta 9$ -THC, and other unknown compound detection
- Some of the cannabinoid standard solutions also had low detection levels of various cannabinoids.
 - Unknown origin
- GC results confirmed HPLC analysis

Results



Discussion

- Even though there were minor contaminants of $\Delta 9$ -THC, the isolate and synthetics could still be considered THC free.
 - <0.3% THC
- Quantification is required.

References

Bonn-Miller MO, Loflin MJE, Thomas BF, Marcu JP, Hyke T, Vandrey R. Labeling Accuracy of Cannabidiol Extracts Sold Online. *JAMA*. 2017;318(17):1708–1709. doi:10.1001/jama.2017.11909

Citti, C., Ciccarella, G., Braghiroli, D., Parenti, C., Vandelli, M. A., & Cannazza, G. (2016). Medicinal cannabis: Principal cannabinoids concentration and their stability evaluated by a high performance liquid chromatography coupled to diode array and quadrupole time of flight mass spectrometry method. *Journal of Pharmaceutical and Biomedical Analysis*, 128, 201–209. <https://doi.org/https://doi.org/10.1016/j.jpba.2016.05.033>