University of Northern Colorado

Scholarship & Creative Works @ Digital UNC

2020 Undergraduate Presentations

Research Day 2020

4-2020


The Birds and the Trees

Branden Lawson

Karina Sanchez

Lauryn Benedict

Follow this and additional works at: https://digscholarship.unco.edu/ug_pres_2020



The Birds and the Trees

Branden Lawson, Karina Sanchez, Dr. Lauryn Benedict Benedict Behavioral Ecology Lab, University of Northern Colorado, Greeley, Colorado



Objective / Prediction One

- ✓ Determine survival rate for urban and non-urban American Robin nests.
- ♣ Predicted that the survival rate of nestlings will be higher in Non-Urban habitats than in Urban habitats.

Objective / Prediction Two


- ♣ Collect data on tree characteristics in a 0.2 Hectare plot nests: Diameter at breast height, Canopy Cover, Number, Height.
- ✓ Predicted that non-urban locations would have an increase in tree collected characteristics.

Objective / Prediction Three

- ✓ Determine if the differences in tree characteristics between Urban and Non-urban locations affect the survival rate of American Robins.
- ✓ Predicted that survival rate will be lower in Urban locations vs. Nonurban location due to the difference in tree characteristics.


Introduction

Bird populations in North America have declined by 2.9 billion individuals since 1970.¹ The transformation of the earth's land surface due to urbanization is estimated to be 10-15%. This loss of habitat is due to many human enterprises that result in land transformation, biotic additions and losses, loss of biological diversity, as well as climate change.² It has been shown that the urban bird community is strongly influenced by vegetation, with the volume of native vegetation being most closely correlated with native bird density and species richness.³ American Robins (*Turdus* migratorius) are common in urban and non-urban areas and can serve as useful indicators of human activities' effects on wildlife.⁴ American Robins tend to nest in habitats that contain mature vegetation, large trees, and plenty of cover. Therefore, understanding the variation in survival between urban and non-urban locations is important to maintain habitat.




Survival Rate in urban locations was found to be 45.67% while in non-urban locations it was found to be 66.67%. These data were found to not be statistically significant.(Wilcoxon Rank-Sum test: Z = -0.32, p = 0.75).

Results




Principal Component Analysis allows us to Regression Analysis of survival rate as see that urban and non-urban study sites do not group together as we expected. As the number of trees within a plot increases so does the average cover within the plot. Additionally, as the average height of trees increases so does increase there is a decrease in survival. the average diameter at breast height.



predicted by the plot characteristics provides us with a negatively correlated relationship ($R^2 = -$ 0.12). As the average cover, diameter at breast height, height of trees, and number of trees

Methods

- ♣ Data collection occurred during the 2019 breeding season
- ✓ Nest monitoring occurred from clutch initiation to fledging.
- A circular plot (0.2 ha), with the nest at the center, was surveyed for tree density, tree height, the diameter of trees, and cover provided by the trees.



Twelve nesting sites that were categorized urban or non-urban based on distance from the city center of Greeley, Colorado.

Red = Urban

Yellow = Non-Urban

Conclusion

- ♣ The prediction that survival rate of nestlings will be higher in Non-Urban habitats than in Urban habitats could not be supported.
- ♣ The prediction that non-urban locations would have an increase in collected tree characteristics could not be supported.
- ♣ The prediction that survival rate will be lower in urban locations vs nonurban locations could not be supported.
- ♣ These results suggest that the American Robin, regardless of urban or non-urban habitat characteristics, are nesting in locations similar in number, diameter, and height of trees as well as in areas with the same amount of cover provided by trees.

References

- 1. Rosenberg, K. V. et al. 2019. Decline of the North American Avifauna. Science 365(6461)
- 2. Beaver, Donald. 1980. "Recovery of an American Robin Population after Earlier DDT Use." Journal of Field Ornithology 51(3): 220–28.
- 3. MillS, G. et al. 1991. "The Relationship between Breeding Bird Density and Vegetation Volume." The Wilson bulletin (Wilson Ornithological Society) 103(3): 468–79.
- 4. Vitousek, P.M. et al.. 1997. "Human Domination of Earth's Ecosystems." Science 277(5325): 494–99.



Branden Lawson Ecology, Evolution, and Conservation blawso@brandensbioblog.com Twitter: @brandensbioblog Linkedln: brandensbioblog