



## University of Dundee

### The Addition of Vitamin D Supplementation to Interferon and Ribavirin in the Treatment of Hepatitis C Virus Infection

Coats, Josh T.; Hapca, Adrian; Witham, Miles; Donnan, Peter; Heydtmann, Mathis; Bathgate, Andrew

*Published in:*  
Archives of Gastroenterology and Hepatology

*Publication date:*  
2018

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

*Citation for published version (APA):*

Coats, J. T., Hapca, A., Witham, M., Donnan, P., Heydtmann, M., Bathgate, A., Hayes, P. C., Fraser, A., Evans, M., Leen, C., Bramley, P., Barclay, S. T., Fox, R., & Dillon, J. F. (2018). The Addition of Vitamin D Supplementation to Interferon and Ribavirin in the Treatment of Hepatitis C Virus Infection: A Randomised Double-Blind PlaceboControlled Pilot Study. *Archives of Gastroenterology and Hepatology*, 1(2), 39-48. <https://www.sryahwapublications.com/archives-of-gastroenterology-and-hepatology/pdf/v1-i2/10.pdf>

#### General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## The Addition of Vitamin D Supplementation to Interferon and Ribavirin in the Treatment of Hepatitis C Virus Infection: A Randomised Double-Blind Placebo-Controlled Pilot Study

Josh T Coats<sup>1\*</sup>, Adrian Hapca<sup>2,3</sup>, Miles D Witham<sup>1,2</sup>, Peter T. Donnan<sup>2,3</sup>, Mathis Heydtmann<sup>4</sup>, Andrew Bathgate<sup>5</sup>, Peter C Hayes<sup>5</sup>, Andrew Fraser<sup>6</sup>, Morgan Evans<sup>7</sup>, Clifford Leen<sup>7</sup>, Peter Bramley<sup>8</sup>, Stephen T. Barclay<sup>9</sup>, Raymond Fox<sup>10,11</sup>, John F Dillon<sup>1</sup>

<sup>1</sup>Division of Molecular and Clinical Medicine, School of Medicine, University of Dundee, Ninewells Hospital and Medical School, Dundee, DD1 9SY, UK, <sup>2</sup>Tayside Clinical Trials Unit, Ninewells Hospital, Dundee, DD1 9SY, UK.

<sup>3</sup>Dundee Epidemiology and Biostatistics Unit (DEBU), Population Health Sciences, University of Dundee, MacKenzie Building, Kirsty Semple Way, Dundee, DD2 4BF, UK, <sup>4</sup>Royal Alexandra Hospital, Corsebar Road, Paisley, PA2 9PN, UK, <sup>5</sup>Royal Infirmary of Edinburgh, 51 Little France Crescent, Edinburgh, EH16, UK, <sup>6</sup>Aberdeen Royal Infirmary, Foresterhill, Aberdeen, AB25 2ZN, UK, <sup>7</sup>Western General Hospital, Crewe Road South, Edinburgh, EH4 2XU, UK, <sup>8</sup>Stirling Community Hospital, Livilands Gate, Stirling, FK8 2AU, UK, <sup>9</sup>Glasgow Royal Infirmary, 84 Castle Street, Glasgow, G4 0SF, UK, <sup>10</sup>Gartnavel General Hospital, 1053 Great Western Road, Glasgow, G12 0YN, UK, <sup>11</sup>The Brownlee Centre, 1053 Great Western Road, Glasgow, G12 0YN, UK.

*j.t.coats@dundee.ac.uk*

**\*Corresponding Author:** Josh T Coats, Division of Molecular and Clinical Medicine, School of Medicine, University of Dundee, Ninewells Hospital and Medical School, Dundee, DD1 9SY, UK.

### Abstract

**Background:** Vitamin D is known to be involved in the immune response to hepatitis C virus infection and preliminary data suggests that supplementation may improve response to treatment with interferon and ribavirin.

**Methods:** The ViaDUCT study was a pilot, multi-centre, randomised, double-blind, placebo-controlled, parallel group clinical trial. Participants with hepatitis C who were planned for treatment were recruited and were randomised to 100,000 units oral vitamin D3 supplementation or placebo each month whilst undergoing treatment with interferon and ribavirin-based regimens. The primary outcome was sustained virologic response at 12 weeks. Secondary outcomes were to identify the proportion of patients who dropped out and to assess for any increase in treatment-related toxicity. Study analyses were performed by modified intention to treat of the primary outcome which was compared between the two arms using logistic regression.

**Results:** 72 participants were randomised; 35 to vitamin D and 37 to placebo. The mean age of participants was 42.5 and 41.7 years respectively. The majority of trial participants were male - 25 (71%) in the vitamin D arm and 24 (65%) in the placebo arm. Following treatment, 60 (83%) attended for assessment of sustained virologic response at 12 weeks. Sustained virologic response was achieved in 82.9% (95%CI 67.3-91.9%) in the vitamin D arm and 73.0% (95%CI 57.0-84.6%) in the placebo arm. The odds of sustained virologic response at 12 weeks in the intervention arm was not significantly greater than in the placebo arm (adjusted odds ratio 1.74; 95%CI 0.43-6.97,  $p=0.44$ ). Adverse events were in keeping with the expected side effect profile of interferon and ribavirin based regimens.

**Conclusion:** Vitamin D supplementation did not significantly improve sustained virologic response when added to interferon and ribavirin-based treatment. The apparent effect size observed would require a larger trial to establish any true effect of this adjunctive therapy.

**Trial Registration:** ClinicalTrials.gov Identifier: NCT02053519, URL: <https://clinicaltrials.gov/ct2/show/NCT02053519>, East of Scotland Research Ethics committee: 13/ES/0116

**Keywords:** Vitamin D, Interferon, Ribavirin, Hepatitis C virus

## **INTRODUCTION**

Hepatitis C virus (HCV) infection is a major global health issue. Recent models estimate a global prevalence of 1% indicating that over 70 million individuals are currently affected with millions of new infections occurring annually[1]. Infection often leads to chronic hepatitis and ultimately may proceed to cirrhosis, hepatocellular carcinoma and death[2].

The therapeutic landscape for HCV has evolved dramatically in recent years advancing from interferon and ribavirin-based regimens to multiple, novel, direct-acting antiviral agents. These novel regimens are expensive and access remains problematic for many individuals affected by HCV[3]. For example, a course of sofosbuvir costs \$84,000 in the United States[4]. Many low and middle-income countries are, therefore, likely to continue to rely upon interferon and ribavirin based treatment protocols until generic manufacturing is permitted or access deals are agreed between governments and manufacturers [3] – a process that may take many years. Ways to maximise the efficacy of interferon/ribavirin regimes are therefore likely to be needed, especially in the short to medium term, if lower and middle income countries are to mount an affordable response to the public health challenge posed by hepatitis C infection.

Vitamin D is a term used to describe several fat-soluble compounds which play an important role in calcium homeostasis and an increasingly recognised role in immune function. Most of the required vitamin D is produced by the action of ultraviolet B radiation on the skin, with only a small contribution from dietary sources. Levels of circulating vitamin D metabolites are therefore often low in populations living at high latitudes, or in people with little exposure to sunlight due to other reasons, e.g. illness or conservative dress codes[5]. Associations between low circulating 25-hydroxyvitamin D (25OHD) levels and a wide range of illnesses have been noted, including malignancy, respiratory infection, autoimmune and cardio-metabolic diseases and notably, chronic liver disease including chronic HCV infection[6, 7].

The interplay between vitamin D and chronic liver disease has been further investigated in recent years. It has been demonstrated that 25OHD levels are lower in those with more severe liver dysfunction[8]. Interestingly it has also been shown that vitamin

D levels rise following successful clearance of HCV infection [9]. The T-cell response against the virus may also rely upon vitamin D suggesting that deficiency may impair the immune response [10, 11]. Given the key role of the liver in the 25-hydroxylation of vitamin D however, observational studies cannot dissect out whether low 25OHD levels are a contributor to, or an effect of hepatitis C infection.

Preliminary clinical data does however suggest that vitamin D status is associated with response to treatment for HCV. In a small study in Italy *Bitetto et al* found that low 25OHD levels were associated with a poorer response to interferon and ribavirin and that supplementation increased the likelihood of response[12]. Both *Abu-Mouch et al* and *Nimer et al* performed randomised controlled trials in Israeli cohorts to establish if vitamin D supplementation affected SVR[13, 14].

The *Abu-Mouch et al* trial randomised 72 individuals with HCV genotype 1 to treatment with interferon/ribavirin with or without vitamin D supplementation. They found that vitamin D supplementation increased SVR in comparison to the control group, 86% vs 42% ( $p < 0.001$ )[14]. In the *Nimer et al* trial, of 50 individuals with genotypes 2 and 3 infections, vitamin D supplementation was also associated with a significant increase in SVR, 95% vs 77% ( $p < 0.001$ ) [13]. SVR was assessed at 24 weeks in these trials as was the then standard of care.

None of these previous studies were placebo-controlled and none were performed in northern European populations – who are those most likely to have low 25OHD levels[5]. We therefore performed a randomised, placebo-controlled pilot trial in Scotland, UK to more rigorously assess the hypothesis that vitamin D supplementation would improve SVR when added to an interferon and ribavirin based treatment regimen.

## **METHODS**

### **General Study Design and Plan**

ViaDUCT was a pilot, multicentre, double blind, randomised, placebo controlled, parallel group trial to examine the effect of year-long course of monthly oral vitamin D3 treatment or placebo, on SVR rates in patients with HCV infection treated with interferon and ribavirin. As this was a pilot trial an important

## The Addition of Vitamin D Supplementation to Interferon and Ribavirin in the Treatment of Hepatitis C Virus Infection: A Randomised Double-Blind Placebo-Controlled Pilot Study

measure was to determine the number of eligible individuals consenting to enter a trial of this design as well as the rate of recruitment. The trial was approved by East of Scotland Research Ethics committee (13/ES/0116), and Clinical Trials Authorisation was given by the UK Medicines and Healthcare Regulatory Authority (EudraCT number 2013-003573-10). The trial was registered at [www.clinicaltrials.gov](http://www.clinicaltrials.gov), registration number NCT02053519

### Population

Individuals were recruited from viral hepatitis clinics in 5 health boards across the National Health Service in Scotland: NHS Tayside, NHS Greater Glasgow and Clyde, NHS Grampian, NHS Forth Valley and NHS Lothian. Inclusion criteria were: patients with known HCV infection confirmed by positive PCR, hepatitis C genotype 1 or 3, and planned to commence on standard HCV therapy, which at the time of trial recruitment was interferon/ribavirin based therapy.

The exclusion criteria were based upon the usual standard-of-care contraindications for treatment with interferon and ribavirin. In addition those at risk of, or known to have, hypercalcaemia were excluded due to the endocrine effects of vitamin D supplementation. The exclusion criteria were: hepatitis C genotype other than 1 or 3, contraindications to interferon or ribavirin therapy, estimated GFR  $<30\text{ml/min/1.73m}^2$  (by MDRD4 method) [15] currently decompensated liver disease characterised by ascites, encephalopathy or variceal bleeding, history of renal calculi, serum calcium  $<2.15$  or  $>2.60$  mmol/L, history of sarcoidosis or metastatic malignancy, hepatocellular carcinoma, taking  $>400$  international units of vitamin D per day, HIV positive, pregnancy, breastfeeding, of childbearing age and not on reliable contraception, or unable to provide written informed consent.

### Participant Recruitment

At the screening visit all participants had their medical history taken and gave written informed consent. Confirmation of HCV diagnosis by viral load by polymerase chain reaction (PCR) and viral genotype was taken from the last available values in the medical notes.

### Standard Treatment

Standard HCV therapy with interferon and ribavirin was commenced 1-4 weeks after the first dose of

vitamin D or placebo. Standard treatment for HCV was 24 weeks. Genotype 1 patients who did not respond to therapy at week 4 or 12 had all anti-viral therapy stopped. Other genotype 1 patients who did respond but did not become virus negative at 12 weeks of therapy were continued on a further 24 week course of standard therapy, total duration 48 weeks. Those who did not respond and stopped antiviral therapy did not continue to receive vitamin D or placebo for the planned 12 month course in this trial. Those participants who continued on 48 weeks of standard therapy received the same 12 months of vitamin D or placebo but returned at 3 and 6 months post-end of treatment for assessment.

Standard therapy for HCV changed as the trial was performed and therefore some patients received additional antivirals concomitantly with interferon and ribavirin. Genotype 1 patients at the outset of the study received interferon/ribavirin but later patients could also receive telaprevir, boceprevir, simeprevir or sofosbuvir at their prescribing physician's discretion. Similarly, Genotype 3 patients at trial outset received interferon/ribavirin but later patients could also receive sofosbuvir.

### Trial Treatment, Randomisation and Blinding

After successful confirmation of eligibility including safety, participants were randomised by the delegated nurse to either vitamin D3 or placebo in a double-blind fashion via a centrally controlled web-based GCP compliant randomisation system, run by Tayside Clinical Trials Unit (TCTU) to preserve allocation concealment. Randomisation was minimised for age ( $\leq 40$ ,  $> 40$ ) and genotype (1 or 3).

Oral vitamin D3 or matching placebo was administered once a month. Each dose of vitamin D3 or placebo was ingested by the participant in the presence of the study nurse. Participants were allowed to continue all their usual medication throughout the trial. The dose of vitamin D3 used was 100,000 units once a month (given as 5mls of Vigantol oil per dose [Merck KgAA]). Placebo was 5mls of Mygliol oil per dose; also provided by Merck KgAA; Mygliol is a medium-chain triglyceride used as the base oil in the manufacture of Vigantol. Vitamin D and placebo were identical on visual inspection and taste, and were dispensed from study bottles with no external indication of whether the bottle contained vitamin D or placebo. Participants,

research nurses recording outcomes, and healthcare teams thus remained blinded to treatment allocation.

### **Trial Outcomes**

The primary outcome of the trial was the absence of detectable virus in blood, using PCR to detect HCV RNA, at 12 weeks, termed sustained virologic response (SVR12). This is a recognised surrogate marker for the cure of infection. In this trial any record of a negative HCV RNA 12 weeks or more after completion of treatment was taken as achievement of SVR12. Conversely a positive result was taken as failure of SVR12. In the absence of an HCV RNA result, for the modified intention to treat analysis, the participant was assumed to have failed to achieve an SVR.

Secondary outcomes were to identify the proportion of eligible patients undergoing therapy for HCV infection who drop out and to assess the effect of supplementation with Vitamin D on adherence to standard therapy. The number of individuals recruited by each site and the rate of recruitment over the 18 month study period was also recorded (February 2014 to August 2015). This data was to establish recruitment variability across sites and help to inform a calculation regarding the required recruitment period for a future large study. A further secondary outcome was to assess for any increase in the expected toxicity of therapy.

### **Sample Size Calculation**

We estimated that a definitive trial would require 313 participants in each arm (assuming  $\alpha=0.05$  and  $\beta=0.80$ ) to have 80% power to show an improvement in SVR from 70% in the placebo group to 80% in the vitamin D group, at an alpha level of 0.05. The purpose of this pilot trial was not to detect this difference, but to estimate the likely effect size and recruitment rates to ensure that a future definitive trial was adequately powered. The sample size was therefore set at a maximum of 100 patients (50 randomised to each arm); this sample size was also deemed sufficient to demonstrate the ability of each of the study centres to recruit and retain subjects in the study, another key point for the pilot to address. This would enable the feasibility of a future larger study to be fully evaluated.

### **Statistical Analysis**

Study analyses were performed by modified intention to treat (mITT) of the primary outcome SVR at 12 weeks. The evaluable dataset were those randomised and receiving at least one dose of vitamin D3 or placebo, with those with missing values for SVR assumed to have not achieved SVR. The primary outcome (SVR12) was compared between the two arms utilising logistic regression stratified by centre with a parameter indicating arm of trial. The primary analysis was an adjusted analysis, adjusting for baseline age, sex, genotype, 25OHD level and viral load. Further analyses were performed without adjustment for baseline variables, and using a per-protocol approach (excluding those who did not attend for SVR measurement). Subgroup analyses were not performed in view of the small numbers in the trial; such analyses would carry a high risk of spurious positive findings.

A 2-sided p value of  $<0.05$  was taken as significant for all analyses. All analyses were conducted using SAS v9.3 by trial statisticians blind to treatment allocation.

### **RESULTS**

72 participants underwent randomisation and received the first dose of study medication. 35 were randomised to the vitamin D arm and 37 to the placebo arm. Flow through the trial is shown in Figure 1. 14 of the 35 (40%) in the vitamin D arm had a genotype 1 infection with the remainder having genotype 3 infection. 14 of the 37 (38%) in the placebo arm had genotype 1 infection with the remaining 23 having genotype 3 infection.

All received interferon/ribavirin therapy, however, within the vitamin D arm four genotype 3 participants also received sofosbuvir. Of genotype 1 participants, one received telaprevir and another boceprevir, three also received sofosbuvir and a further three received simeprevir. Within the placebo arm eight genotype 3 participants received sofosbuvir and among the genotype 1 participants, one received boceprevir and six received simeprevir. Of the 72 randomised participants, 60 (83%) had a blood test performed for HCV RNA 12 weeks after end of trial medication to allow confirmation of achievement of SVR12. Baseline details are given in Table 1.

## The Addition of Vitamin D Supplementation to Interferon and Ribavirin in the Treatment of Hepatitis C Virus Infection: A Randomised Double-Blind Placebo-Controlled Pilot Study

**Table 1.** Baseline characteristics of study participants

	Vitamin D	Placebo
N	35	37
Mean age (years) (SD)	42.5 (11.6)	41.7 (8.7)
Male sex (%)	25 (71%)	24 (65%)
Genotype 1 (%)	14 (40%)	14 (38%)
Genotype 3 (%)	21 (60%)	23 (62%)
Mean 25OHD level (nmol/L) (SD)	59 (24)	51 (21)
Mean HCV RNA level ( $\times 10^6$ IU/ml) (SD)	2.5 (5.4)	1.2 (1.4)

25OHD: 25-hydroxyvitamin D. HCV: Hepatitis C virus

### Changes in 25 OHD Levels across ViaDuct Trial

In the Vitamin D intervention group the mean change (averaged across all months) in 25OHD levels was +16.7 nmol/L (SD 39.5) compared to the placebo group where the mean change (averaged across all months) was -7.9 nmol/L (SD 21.8). The between-group difference was 24.7 nmol/L (95% CI 12.2 to 37.2;  $p < 0.001$  by Student's t-test)

### Primary Outcome

The proportion of positive outcome (SVR at 12 weeks) was 82.9% (95% CI 67.3 – 91.9%) in the vitamin D group, and 73.0% (95% CI 57.0 – 84.6%) in the placebo

**Table 2.** Primary outcome analyses (SVR12)

Variable		Vitamin D N=35	Placebo N=37	
SVR12 (mITT)	Positive (%)	6 (17.1)	10 (27.0)	
	Negative (%)	29 (82.9)	27 (73.0)	
SVR12* Per protocol	Positive (%)	1 (3.3)	3 (10.0)	
	Negative (%)	29 (96.7)	27 (90.0)	
Gender	Female (%)	10 (28.6)	13 (35.1)	
	Male (%)	25 (71.4)	24 (64.9)	
Genotype	1	Total achieving SVR		
		Interferon and ribavirin	4	5
		Interferon, ribavirin, telaprevir	1	0
		Interferon, ribavirin, boceprevir	1	0
		Interferon, ribavirin, sofosbuvir	3	0
		Interferon, ribavirin, simeprevir	3	6
	Total not achieving SVR		1	1
		Interferon and ribavirin	1	0
		Interferon, ribavirin, boceprevir	0	1
	Total genotype 1 for which SVR status unknown		1	2
3	Total achieving SVR		17	16
	Interferon and ribavirin	13	8	
	Interferon, ribavirin, sofosbuvir	4	8	
	Total not achieving SVR (interferon, ribavirin)		0	2
Total genotype 3 for which SVR status unknown		4	5	

\*12 missing SVR12 excluded

## The Addition of Vitamin D Supplementation to Interferon and Ribavirin in the Treatment of Hepatitis C Virus Infection: A Randomised Double-Blind Placebo-Controlled Pilot Study

### Secondary Outcomes

#### The Proportion of Eligible Patients Undergoing Therapy for HCV Infection Who Drop Out

In those patients consented to participate in the trial as described in the CONSORT diagram above 100

consented and only 72 started therapy. The major reason for this was not fulfilling entry criteria, the second commonest reason was failure to attend further appointments. 12 /72 (17%) consented participants failed to complete the trial. A breakdown of participant recruitment and retention by site is provided in table 3.

**Table 3. Recruitment by site**

Health board	No. randomised	Recruitment per month	Recruitment per 100,000 population	No. with SVR data available (%)
NHS Tayside	15	0.8	3.8	12 (80)
NHS Lothian	7	0.4	0.9	5 (71)
NHS Forth Valley	7	0.4	2.3	7 (100)
NHS Greater Glasgow and Clyde	39	2.2	3.0	33 (85)
NHS Grampian	4	0.2	0.8	3 (75)

#### The Effect of Supplementation with Vitamin D3 on Adherence to Standard HCV Therapy

The effect of supplementation with Vitamin D3 on adherence to standard HCV therapy was assessed using survival analysis and plotted using Kaplan-Meier plots. The survival analysis of time to drop-off or study completion (adherence) indicated no significant difference between placebo and drug (hazard ratio = 1.3; 95%CI 0.8 to 2.1; p=0.33).

#### Adverse Events

Adverse events in the trial are listed, broken down by intervention, in Table 4 by standard listing of numbers of each condition. Most events were non-serious and were related to expected side effects from interferon/ribavirin therapy. There was an excess of respiratory infections in the vitamin D arm, this is probably spurious and a reflection of small numbers, as other vitamin D trials have suggested that vitamin D may be protective against respiratory infections[16].

**Table 4. Adverse events**

Category	Vitamin D	Placebo
Blood and lymphatic system disorders	0	4
Ear and labyrinth disorders	0	1
Gastrointestinal disorders	16	21
Immune system disorders	0	1
Worsening Liver function tests	1	2
Metabolic disorders	2	4
Musculoskeletal disorders	2	4
Nervous system disorders	3	7
Psychiatric disorders	13	18
Urinary tract infection	4	0
Other renal/urological disorders	1	0
Reproductive system/breast disorders	2	1
Respiratory tract infection	11	2
Other Respiratory disorders	9	10
Dermatological disorders	18	17
Vascular disorders	2	1
Other Infections (excl respiratory and urinary infections)	2	6
Other	9	8
TOTAL	95	107
Deaths	0	0
Hospitalisations	1	2

## **DISCUSSION**

### **Key Findings**

We performed the first randomised, placebo-controlled parallel group trial to investigate the effects of vitamin D supplementation on response to interferon-based treatment for HCV. We achieved the aim of performing a pilot trial to assess the feasibility of recruitment to this trial design and to estimate the effect size of vitamin D supplementation to allow planning for a larger trial.

We observed no significant difference in SVR between the vitamin D and placebo groups which was not unexpected given the pilot nature of the trial. A trend towards improved SVR in the vitamin D group was observed (83% vs 73%,  $p=0.44$ ) and the groups were otherwise matched suggesting that this effect was independent of baseline characteristics.

The effect size had not been rigorously investigated prior to this study however our effect size was less than that seen in previous Israeli studies, which both demonstrated a statistically significant increase in SVR with vitamin D supplementation [13, 14]. This may be related to the more rigorous double-blind trial design which we adopted.

### **Strengths**

This trial was conducted in a group of patients known to have difficulty accessing healthcare and with a known high rate of drop-out from care [17]. It was therefore predicted that involving this population in a research study would be challenging. Despite this our study delivered reasonable recruitment and retention in a purely National Health Service (NHS) environment. Ultimately the drop-out rate observed in the study (16.7% dropout post-randomisation) was comparable to standard HCV care drop-out rates and reflects the real world nature and applicability of the study [18, 19].

### **Limitations**

We aimed to recruit 100 participants to the trial and ultimately 72 were enrolled and underwent randomisation. A key limitation to recruitment was the rapidly evolving therapeutic landscape with multiple new interferon-free treatments being approved and individual preference for these less toxic regimens. Extending the study to enable further

recruitment was not appropriate given the increased efficacy and reduced toxicity of these novel antivirals. We initially intended to assess for differences between individuals who declined trial involvement and those who consented. However, this was confounded by the adoption of interferon-free regimens during the study period and therefore these data has not been presented as they are no longer meaningful.

The use of interferon add-on therapies increased as the trial progressed and therefore multiple participants in both arms of the study received either telaprevir, boceprevir, sofosbuvir or simeprevir. Use of these agents complicates interpretation of our results, however it would have been unethical to withhold these from study participants given the dramatically increased SVR associated with these new agents. Equally the trend towards vitamin D supplementation improving SVR is of greater interest given the higher use of novel more effective antivirals in the placebo arm of our study.

60 of the 72 (83%) participants ultimately attended for post-treatment viral PCR to assess for SVR. It is therefore unknown if the treatment was effective in the other 12 individuals. It is common in HCV therapy for 5-10% of patients to drop out of therapy for patient related reasons, for a further 5% to have therapy stopped for safety side effects and for a further 5% to have therapy stopped due to lack of efficacy. Such patients are very difficult to keep in follow up to collect SVR data when the patient believes the therapy has not worked. Our observed dropout rate (17%) is therefore in keeping with real world published data [18, 19].

The quality of our data was also affected by a population of participants with variable attendance rates and chaotic lifestyles. Study visits often did not take place when planned and not all study procedures could be conducted at times planned in the protocol. In addition, all study activities were performed by NHS staff and not trained research assistants; an approach that reflected the pragmatic nature of the trial design and the realities of delivering a clinical service to a complex group of patients.

### **Generalisability**

This pilot study was not designed to provide generalizable conclusions on the efficacy of the addition of vitamin D to interferon and ribavirin-



## The Addition of Vitamin D Supplementation to Interferon and Ribavirin in the Treatment of Hepatitis C Virus Infection: A Randomised Double-Blind Placebo-Controlled Pilot Study

based HCV treatment. Our result does not warrant a change in clinical practice but helps to inform a future larger trial.

### Interpretation

Today the Scottish Medicines Consortium (SMC) has approved all-oral pan-genotypic regimens for HCV with SVR of over 97% [20]. Given this it is not surprising that it became increasingly challenging to recruit patients to a study and treatment which was rapidly becoming obsolete. However, it should be noted that the novel direct-acting antivirals are extremely expensive and funding will be a challenge in even the most highly-funded healthcare systems. The largest burden of HCV continues to be felt in lower and middle income countries (LMICs) and in these nations interferon and ribavirin are likely to be of continued importance for some years. An economical means to augment response to this regimen may, therefore, still provide meaningful benefit to both patients and to healthcare systems in LMICs.

Given the widespread adoption of direct-acting antivirals it is unlikely that a future large trial of vitamin D vs. placebo as an adjunct to interferon/ribavirin therapy will be conducted in any high-income country. In addition, using vitamin D as an adjunct to these novel therapies is not required given the near 100% response rates. However, such a trial could be performed in LMICs, could still be informative and could lead to a change in practice.

Based upon our data such a trial would require 540 participants (270 per group) to detect a similar difference in effect with 80% power, assuming  $\alpha=0.05$ . A larger number of participants would initially be required given the anticipated drop-out rates. Our trial enrolled 72 participants for which 60 (83%) have available SVR data. This 17% drop-out rate is comparable to that seen in other HCV treatment studies. Given this, if SVR data were to be obtained for 540 participants an initial enrolment and randomisation of 650 individuals would be required.

It is difficult to draw valid conclusions regarding recruitment given the previously discussed difficulties which we encountered. In total across the 5 Scottish health boards we recruited 4 individuals per month during the 18 month trial period. A predicted 158 month (13 year) recruitment period to enrol 632 participants from a majority of the Scottish population

in a larger study is unachievable. However if this were performed in a lower/middle-income country recruitment would be less affected by the widespread availability of interferon-free regimens.

The exact treatment regimen to be utilised in any future interferon/ribavirin study involving vitamin D vs. placebo is unclear and would be dependent upon the availability and cost of add-on direct acting antivirals which would boost overall efficacy.

In addition, recent evidence suggests that the monthly vitamin D bolus utilised in our study might be less effective than daily or weekly dosing and therefore a future trial would likely opt for more frequent dosing [16].

### CONCLUSION

Our study adds further credence to the hypothesis that vitamin D plays a crucial role in modulating the immune response to infection and inflammation. Whilst novel therapeutics have now replaced interferon/ribavirin regimens in high-income nations, interferon/ribavirin may continue to be an important therapeutic option in LMICs. Current evidence is insufficient to recommend adding vitamin D to interferon/ribavirin regimens, but our results support the case for larger trials of vitamin D supplementation as adjunctive therapy in patients with HCV treated with interferon/ribavirin. Such trials should be conducted in populations likely to be treated in this way – i.e. patients in LMICs.

### LIST OF ABBREVIATIONS

**25OHD** – 25-hydroxyvitamin D

**HCV** – hepatitis C virus

**LMIC** – lower and middle income countries

**NHS** – National Health Service

**PCR** – polymerase chain reaction

**SMC** – Scottish Medicines Consortium

**SVR** – sustained virologic response

### DECLARATIONS

#### Ethics Approval

The trial was approved by East of Scotland Research Ethics committee (approval number 138656), and Clinical Trials Authorisation was given by MHRA (EudraCT number 2013-003573-10) and ISRCTN number NCT 02053519.

## The Addition of Vitamin D Supplementation to Interferon and Ribavirin in the Treatment of Hepatitis C Virus Infection: A Randomised Double-Blind Placebo-Controlled Pilot Study

Written consent was provided by all participants prior to enrolment in the trial.

### Availability of Data and Materials

Raw data are available for collaborative, non-profit use on request from the authors.

### Acknowledgements

Tayside Clinical Trials Unit, University of Dundee, Dundee; in particular, James Flood, Margaret Band and Sarah Inglis, Clinical trial managers

Sheila Dickson, Liver Clinical Nurse Specialist, Glasgow Royal Infirmary

Stuart Kirkwood, Blood Borne Virus Clinical Nurse Specialist, Gartnavel General Hospital, Glasgow

Brian Stephens, Farsana Ahmed, Diane Knight and Jan Tait, Specialist nurses, Ninewells Hospital, Dundee

Lesley Briody, Lead Nurse, Wellcome Trust Clinical Research Facility, Edinburgh

Audrey Anderson, Anne Cameron-Burns, Specialist Nurses, Royal Alexandria Hospital, Paisley

Karen Prentice, Lorraine Moffat and Karen Gibb, Hepatology Specialist Nurses, Forth Valley Hepatology Service, Stirling.

### REFERENCES

- [1] Polaris Observatory HCVC. Global prevalence and genotype distribution of hepatitis C virus infection in 2015: a modelling study. *The Lancet Gastroenterology & hepatology*. 2017; 2(3): 161-76.
- [2] Shepard CW, Finelli L, Alter MJ. Global epidemiology of hepatitis C virus infection. *Lancet Infect Dis*. 2005; 5(9): 558-67.
- [3] Edwards DJ, Coppens DG, Prasad TL, Rook LA, Iyer JK. Access to hepatitis C medicines. *Bulletin of the World Health Organization*. 2015; 93(11): 799-805.
- [4] Zhang S, Bastian ND, Griffin PM. Cost-effectiveness of sofosbuvir-based treatments for chronic hepatitis C in the US. *BMC gastroenterology*. 2015; 15:98.
- [5] Rhein HM. Vitamin D deficiency is widespread in Scotland. *BMJ*. 2008;336(7659):1451.

- [6] Holick MF. Sunlight and vitamin D for bone health and prevention of autoimmune diseases, cancers, and cardiovascular disease. *Am J Clin Nutr*. 2004;80(6 Suppl):1678S-88S.
- [7] Cholongitas E, Theocharidou E, Goulis J, Tsochatzis E, Akriviadis E, Burroughs K. Review article: the extra-skeletal effects of vitamin D in chronic hepatitis C infection. *Alimentary pharmacology & therapeutics*. 2012;35(6):634-46.
- [8] Malham M, Jorgensen SP, Ott P, Agnholt J, Vilstrup H, Borre M, et al. Vitamin D deficiency in cirrhosis relates to liver dysfunction rather than aetiology. *World J Gastroenterol*. 2011;17(7):922-5.
- [9] Lange CM, Bojunga J, Ramos-Lopez E, von Wagner M, Hassler A, Vermehren J, et al. Vitamin D deficiency and a CYP27B1-1260 promoter polymorphism are associated with chronic hepatitis C and poor response to interferon-alfa based therapy. *J Hepatol*. 2011;54(5):887-93.
- [10] von Essen MR, Kongsbak M, Schjerling P, Olgaard K, Odum N, Geisler C. Vitamin D controls T cell antigen receptor signaling and activation of human T cells. *Nature immunology*. 2010;11(4):344-
- [11] Shen L. Vitamin D controls T cell activation: implication for causal association between vitamin D deficiency and fibrosis in chronic hepatitis C. *Hepatology*. 2010;52(5):1864.
- [12] Bitetto D, Fabris C, Fornasiere E, Pipan C, Fumolo E, Cussigh A, et al. Vitamin D supplementation improves response to antiviral treatment for recurrent hepatitis C. *Transplant international : official journal of the European Society for Organ Transplantation*. 2011;24(1):43-50.
- [13] Nimer A, Mouch A. Vitamin D improves viral response in hepatitis C genotype 2-3 naive patients. *World J Gastroenterol*. 2012;18(8):800-5.
- [14] Abu-Mouch S, Fireman Z, Jarchovsky J, Zeina AR, Assy N. Vitamin D supplementation improves sustained virologic response in chronic hepatitis C (genotype 1)-naive patients. *World J Gastroenterol*. 2011; 17(47): 5184-90.
- [15] Levey AS, Bosch JP, Lewis JB, Greene T, Rogers N, Roth D. A more accurate method to estimate glomerular filtration rate from serum creatinine:

## The Addition of Vitamin D Supplementation to Interferon and Ribavirin in the Treatment of Hepatitis C Virus Infection: A Randomised Double-Blind Placebo-Controlled Pilot Study

- a new prediction equation. Modification of Diet in Renal Disease Study Group. *Ann Intern Med.* 1999;130(6):461-70.
- [16] Martineau AR, Jolliffe DA, Hooper RL, Greenberg L, Aloia JF, Bergman P, et al. Vitamin D supplementation to prevent acute respiratory tract infections: systematic review and meta-analysis of individual participant data. *BMJ.* 2017;356:i6583.
- [17] Morrill JA, Shrestha M, Grant RW. Barriers to the treatment of hepatitis C. Patient, provider, and system factors. *Journal of general internal medicine.* 2005;20(8):754-8.
- [18] Cournot M, Glibert A, Castel F, Druart F, Imani K, Lauwers-Cances V, et al. Management of hepatitis C in active drugs users: experience of an addiction care hepatology unit. *Gastroenterologie clinique et biologique.* 2004; 28(6-7 Pt 1): 533-9.
- [19] Lieveld FI, van Vlerken LG, Siersema PD, van Erpecum KJ. Patient adherence to antiviral treatment for chronic hepatitis B and C: a systematic review. *Ann Hepatol.* 2013;12(3):380-91.
- [20] Dillon JF, Hayes P, Barclay S, Fox R, Fraser A. National Clinical Guidelines for the treatment of HCV in adults Health Improvement Scotland: NHS National Services Scotland; 2017 [cited 2017 07/17/17]. Available from: <http://www.hps.scot.nhs.uk/resourcedocument.aspx?resourceid=1598>.

**Citation:** JoshT Coats, Adrian Hapca, Miles D Witham, et al. *The Addition of Vitamin D Supplementation to Interferon and Ribavirin in the Treatment of Hepatitis C Virus Infection: A Randomised Double-Blind Placebo-Controlled Pilot Study. Archives of Gastroenterology and Hepatology.* 2018; 1(2): 39-48.

**Copyright:** © 2018 JoshT Coats, Adrian Hapca, Miles D Witham, et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.