



**British  
Geological Survey**  
NATURAL ENVIRONMENT RESEARCH COUNCIL

# Calcareous microfossils from laminated clay in an M1 cutting near Enderby Grange, Leicestershire

Internal Report IR/03/164



BRITISH GEOLOGICAL SURVEY

INTERNAL REPORT IR/03/164

# Calcareous microfossils from laminated clay in an M1 cutting near Enderby Grange, Leicestershire

Ian P. Wilkinson

The National Grid and other  
Ordnance Survey data are used  
with the permission of the  
Controller of Her Majesty's  
Stationery Office.  
Ordnance Survey licence number  
GD 272191/1999

*Key words*

Pleistocene, reworked  
microfaunas.

*Bibliographical reference*

WILKINSON, I. P.. 2003.  
Calcareous microfossils from  
laminated clay in an M1 cutting  
near Enderby Grange  
Leicestershire. *British  
Geological Survey Internal  
Report*, IR/03/164. 7pp.

## BRITISH GEOLOGICAL SURVEY

The full range of Survey publications is available from the BGS Sales Desks at Nottingham and Edinburgh; see contact details below or shop online at [www.thebgs.co.uk](http://www.thebgs.co.uk)

The London Information Office maintains a reference collection of BGS publications including maps for consultation.

The Survey publishes an annual catalogue of its maps and other publications; this catalogue is available from any of the BGS Sales Desks.

*The British Geological Survey carries out the geological survey of Great Britain and Northern Ireland (the latter as an agency service for the government of Northern Ireland), and of the surrounding continental shelf, as well as its basic research projects. It also undertakes programmes of British technical aid in geology in developing countries as arranged by the Department for International Development and other agencies.*

*The British Geological Survey is a component body of the Natural Environment Research Council.*

### **Keyworth, Nottingham NG12 5GG**

☎ 0115-936 3241 Fax 0115-936 3488  
e-mail: [sales@bgs.ac.uk](mailto:sales@bgs.ac.uk)  
[www.bgs.ac.uk](http://www.bgs.ac.uk)  
Shop online at: [www.thebgs.co.uk](http://www.thebgs.co.uk)

### **Murchison House, West Mains Road, Edinburgh EH9 3LA**

☎ 0131-667 1000 Fax 0131-668 2683  
e-mail: [scotsales@bgs.ac.uk](mailto:scotsales@bgs.ac.uk)

### **London Information Office at the Natural History Museum (Earth Galleries), Exhibition Road, South Kensington, London SW7 2DE**

☎ 020-7589 4090 Fax 020-7584 8270  
☎ 020-7942 5344/45 email: [bgs london@bgs.ac.uk](mailto:bgs london@bgs.ac.uk)

### **Forde House, Park Five Business Centre, Harrier Way, Sowton, Exeter, Devon EX2 7HU**

☎ 01392-445271 Fax 01392-445371

### **Geological Survey of Northern Ireland, 20 College Gardens, Belfast BT9 6BS**

☎ 028-9066 6595 Fax 028-9066 2835

### **Maclea Building, Crowmarsh Gifford, Wallingford, Oxfordshire OX10 8BB**

☎ 01491-838800 Fax 01491-692345

### *Parent Body*

### **Natural Environment Research Council, Polaris House, North Star Avenue, Swindon, Wiltshire SN2 1EU**

☎ 01793-411500 Fax 01793-411501  
[www.nerc.ac.uk](http://www.nerc.ac.uk)

# Contents

<b>Contents.....</b>	<b>i</b>
<b>Summary .....</b>	<b>i</b>
<b>1 Introduction.....</b>	<b>1</b>
<b>2 Faunal list.....</b>	<b>1</b>
2.1 Late Triassic-Early Jurassic: .....	1
2.2 Late Cretaceous:.....	1
2.3 Quaternary: .....	1
<b>3 Conclusions .....</b>	<b>2</b>

## Summary

Laminated clay, which on field evidence was considered to be of Pleistocene age, contained marine and brackish marine microfossils reworked from the Triassic-lower Jurassic, Upper Cretaceous and Quaternary. Fresh water ostracods or diatoms were no observed.

# 1 Introduction.

Whilst co-ordinating the palaeontological data in order to produce a palaeontological data set for the Leicester sheet (1:50K sheet 156) (WILKINSON, 2003) a sample (EGP8955) described as a 'laminated clay from below Chalky Boulder Clay and overlying glacial gravel', came to light (Register 90, page 36). The sample had been collected from the M1 cutting c.137m West of Enderby Grange and 640m E of Cailton Hayes Hospital (National Grid Reference SP 5438 9852). The register notes the presence of ostracods in the sample and a Pleistocene age was given to it. The implication is that the laminated clay represents a lacustrine or fluvial deposit and that the microfauna was *in situ*. However, the fauna examined comprises entirely reworked marine and brackish marine species.

## 2 Faunal list

### 2.1 LATE TRIASSIC-EARLY JURASSIC:

Foraminifera- *Eoguttulina* sp  
*Lingulina tenera tenera*  
 Ostracoda- *Ogmoconcha* sp

### 2.2 LATE CRETACEOUS:

Foraminifera- *Bolivinooides culverensis*  
*Loxostomum eleyi*  
*Gyroidinooides nitidus*  
*Dicarinella canaliculata*  
*Osangularia cordieriana*  
*Stensioeina* sp  
*Reusella szajnochae praecursor*  
*Marssonella trochus*  
*Stensioeina exsculpta exsculpta*  
*Stensioeina granulata incondita*  
*Whiteinella archaeocretacea*

### 2.3 QUATERNARY:

Foraminifera- *Elphidium* sp  
*Buccella frigida*  
 ?*Haynesina orbiculare*

### 3 Conclusions

The microfauna shows evidence of three ages.

The Rhaetic to early Jurassic is indicated by the presence of *Ogmoconcha* sp and *Lingulina tenera tenera*. These are long ranging taxa which are unlikely to be younger than the early Toarcian.

The late Cretaceous assemblage dominates the fauna and includes *Bolivinoidea culverensis*, a species indicative of the upper part of the *angulata* to *quadrata* macrofaunal zones (Lower and Middle Campanian). In terms of the lithostratigraphy of southern England, this species appears in the lower (but not basal) part of the Newhaven Chalk Formation and extends into the Culver Chalk Formation. *Stensioeina exsculpta exsculpta* became extinct in the basal *pilula* zone and is characteristic of the foraminiferal assemblages in the Seaford Chalk Formation and the lower part of the Newhaven Chalk Formation. The only part of the succession where both species are found together is in foraminiferal zone BGS19, which equates with the basal *pilula* zone, i.e. lower but not basal Newhaven Chalk.

The Quaternary element of the fauna is very minor. The Foraminifera listed above indicate cold waters and are derived from brackish to brackish marine water conditions, and are presumably derived from the North Sea. Had the sample represented a Pleistocene lacustrine or fluvial environment, fresh water ostracods and diatoms would have been expected. None was observed. There is no evidence, therefore, that the laminated clay is a fresh water deposit. However, on the basis of the Quaternary foraminifera, and despite the fact that they are reworked, the Pleistocene age assigned to the deposit is not unreasonable.

### 4 References

WILKINSON, I P 2000. A preliminary foraminiferal biozonation of the Chalk Group (In preparation for the Holostrat Project: Upper Cretaceous). *Internal Report of the British Geological Survey*, IR/00/13, 21pp.

WILKINSON, I P 2003. A data-base of palaeontological results pertaining to the Leicester Sheet (1:50K sheet 156). *Internal Report of the British Geological Survey*, IR/03/165, 21pp.