



UNIVERSITÀ
DEGLI STUDI
DI PADOVA

Università degli Studi di Padova

Padua Research Archive - Institutional Repository

Parabolic orbits of 2-nilpotent elements for classical groups

Original Citation:

Availability:

This version is available at: 11577/3341285 since: 2020-05-25T13:11:59Z

Publisher:

HELDERMANN VERLAG, LANGER GRABEN 17, LEMGO, GERMANY, 32657

Published version:

DOI:

Terms of use:

Open Access

This article is made available under terms and conditions applicable to Open Access Guidelines, as described at <http://www.unipd.it/download/file/fid/55401> (Italian only)

(Article begins on next page)

Parabolic orbits of 2-nilpotent elements for classical groups

Magdalena Boos¹, Giovanni Cerulli Irelli², Francesco Esposito³

Abstract

We consider the conjugation-action of the Borel subgroup of the symplectic or the orthogonal group on the variety of nilpotent complex elements of nilpotency degree 2 in its Lie algebra. We translate the setup to a representation-theoretic context in the language of a symmetric quiver algebra. This makes it possible to provide a parametrization of the orbits via a combinatorial tool that we call symplectic/orthogonal oriented link patterns. We deduce information about numerology. We then generalize these classifications to standard parabolic subgroups for all classical groups. Finally, our results are restricted to the nilradical.

1 Introduction

Let G be a classical complex group of rank n . Then G is either the general linear group $GL_n(K)$ or the symplectic group $SP_{2l}(K)$ or the orthogonal group $O_n(K)$, where $K = \mathbf{C}$. Let \mathfrak{g} be the corresponding Lie algebra.

The study of the adjoint action of (subgroups of) G on \mathfrak{g} and numerous variants thereof is a well-established and much considered task in algebraic Lie theory. Employing methods of geometric invariant theory, a classical topic is the study of orbits and their closures, which is also known as the *vertical* problem [11].

One famous example of a classification problem alike is the study of GL_n -conjugation (or SL_n -conjugation, this doesn't make a difference) on the variety of complex matrices of square-size n . A complete system of representatives up to conjugation is given by the Jordan canonical form [10] which dates back to the 19th century. This system of representatives is given by continuous parameters, the eigenvalues of the matrix, and discrete parameters. In order to determine the latter, it suggests itself to restrict the action to the nilpotent cone, namely to GL_n -conjugation on the set of nilpotent matrices. The number of conjugacy classes of nilpotent matrices is finite and can be described combinatorially by partitions of n .

¹Ruhr University Bochum, Faculty of Mathematics, 44780 Bochum, Germany. magdalena.boos-math@rub.de

²Sapienza University of Rome, Department SBAI, 00161 Rome, Italy. giovanni.cerulliirelli@uniroma1.it

³University of Padova, Department of Mathematics, 35121 Padova, Italy. esposito@math.unipd.it

One generalization of this setup is obtained by restricting the acting group from G to parabolic subgroups $P \subseteq G$. In particular, the Borel subgroup B is considered, then, and the question about a variety admitting only finitely many orbits is closely related to the concept of so-called spherical varieties [5]. One example of a parabolic action can be found in [9], where Hille and Röhrle prove a finiteness criterion for the number of orbits of parabolic conjugation on the unipotent radical of \mathfrak{g} .

Another adaption of the above setup is given by restricting the nilpotent cone \mathcal{N} of nilpotent matrices to certain subvarieties. For example, Melnikov parametrizes the Borel-orbits in the variety of 2-nilpotent elements in the nilradical \mathfrak{n} of $\mathfrak{g} = \text{Lie}(\text{GL}_n(K))$ in [14] which is inspired by the study of orbital varieties. A parametrization in the symplectic setup is published by Barnea and Melnikov in [2]. In [8], Gandini, Maffei, Möseneder Frajria and Papi consider the more general approach of B -stable abelian subalgebras of the nilradical of \mathfrak{b} in which they parametrize the B -orbits and describe their closure relations.

In this article, we consider the algebraic subvariety $\mathcal{N}(2)$ of 2-nilpotent elements of the nilpotent cone of \mathfrak{g} , namely

$$\mathcal{N}(2) = \mathcal{N}(2, G) = \{x \in \mathfrak{g} \mid x^2 = 0\}.$$

Every parabolic subgroup P of G acts on $\mathcal{N}(2)$. It is known that the number of orbits is always finite, since Panyushev shows finiteness for the Borel-action in [17]. In case $G = \text{GL}_n(K)$, a parametrization of the P -orbits and a description of their degenerations is given in [4] and [3] for each parabolic subgroup $P \subset G$.

Our first goal in this article is to prove in a different manner that there are only finitely many B -orbits in $\mathcal{N}(2)$ for the remaining classical groups, that is, for types B , C and D . We approach the problem in a way closely related to [4] from a quiver-theoretic point of view - but instead of translating to the representation variety of a quiver with relations of a special dimension vector, we translate the orbits to certain (isomorphism classes of) representations of a *symmetric* quiver with relations of a fixed dimension vector. In this setup we show that there are only finitely many of the latter which are parametrized by combinatorial objects which we call symplectic/orthogonal oriented link patterns, see Definitions 6.2 and 6.8.

For example the Borel-orbits of 2-nilpotent matrices in \mathfrak{o}_4 are parametrized by these five patterns:



The following five matrices give a system of representatives of these orbits.

$$\left(\begin{array}{cccc} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \end{array} \right) \quad \left(\begin{array}{cccc} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \end{array} \right) \quad \left(\begin{array}{cccc} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{array} \right) \quad \left(\begin{array}{cccc} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right) \quad \left(\begin{array}{cccc} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

Our second goal is to parametrize all orbits explicitly. The approach via a symmetric quiver makes it possible to classify the orbits by representations, and thus, by combinatorial data.

We afterwards generalize these results to parabolic subgroups $P \subset G$. To do so, we consider the isotropic flag corresponding to P and realize it as a representation M_P . The parabolic P equals the symmetric stabilizer of M_P and we obtain a classification of the P -orbits in $\mathcal{N}(2)$ as in the Borel case.

In the last section, we restrict our results to the action of P on the nilradical $\mathfrak{n}(2) := \mathcal{N}(2) \cap \mathfrak{n}$ of 2-nilpotent upper-triangular matrices in \mathfrak{g} and obtain complete parametrizations, here. In the symplectic case, the parametrization coincides with the parametrization by so-called symplectic link patterns of [2], even though the methods used to prove it are different.

Acknowledgments: The authors would like to thank Giovanna Carnovale for her input concerning the results and methods of this work. Furthermore, the first author thanks Martin Bender for many discussions about Lie-theoretical background.

2 Classical groups and Lie algebras

Let K be the field of complex numbers $K := \mathbf{C}$ and let n be an integer. We consider the complex classical groups, that is, the general linear group $\mathrm{GL}_n := \mathrm{GL}_n(K)$, the symplectic group $\mathrm{SP}_n := \mathrm{SP}_n(K)$, whenever $n = 2l$ for some integer l , and the orthogonal group $\mathrm{O}_n := \mathrm{O}_n(K)$. The corresponding Lie algebras are denoted by $\mathfrak{gl}_n := \mathfrak{gl}_n(K)$, $\mathfrak{sp}_n := \mathfrak{sp}_n(K)$ and $\mathfrak{o}_n := \mathfrak{o}_n(K)$.

In general, given a vector space V endowed with a non-degenerate bilinear form $\langle -, - \rangle$, let us denote by $\mathrm{Sym}(V)$ the group of symmetries of the vector space V which preserve $\langle -, - \rangle|_{V \times V}$. Then $\mathrm{Sym}(V)$ equals either the symplectic group $\mathrm{SP}(V)$ or the orthogonal group $\mathrm{O}(V)$, depending on whether $(V, \langle -, - \rangle)$ is symplectic or orthogonal). We define $\mathfrak{sym}(V) := \mathrm{Lie}(\mathrm{Sym}(V))$.

Let l be an integer, then we denote by $J = J_l$ the $l \times l$ anti-diagonal matrix with every entry on the anti-diagonal being 1:

$$J_l = \begin{pmatrix} 0 & \cdots & 1 \\ \vdots & \ddots & \vdots \\ 1 & \cdots & 0 \end{pmatrix}$$

It is easy to see (and well-known) that $J^{-1} = J$ and that the conjugate $J^T A J$ by J of the *transpose* ${}^T A$ of a matrix $A \in K^{l \times l}$ is given by "the transpose of A with respect to the anti-diagonal". For example, for $l = 2$, given $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$:

$$J^T A J = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} a & c \\ b & d \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} d & b \\ c & a \end{bmatrix}.$$

We set

$$\tilde{\varkappa} A := J^T A J.$$

In this notation, it is easy to write down the elements of the symplectic and orthogonal Lie algebras.

2.1 Symplectic group

Let V be an $n = 2l$ -dimensional complex vector space. Let us fix a basis of V and a bilinear form $F = F_V : V \times V \rightarrow K$, $F(v, w) = \langle v, w \rangle$, associated with the matrix (still denoted by F)

$$F = \begin{bmatrix} 0 & J_l \\ -J_l & 0 \end{bmatrix}. \quad (2.1)$$

The symplectic group SP_n consists of those matrices $A \in GL_n$ which preserve this bilinear form (i.e. $\langle Av, Aw \rangle = \langle v, w \rangle$); in other words A satisfies the equation

$${}^T A F A = F.$$

The Lie algebra \mathfrak{sp}_n of SP_n consists of those matrices $a \in \mathfrak{gl}_n$ which fulfill

$${}^T a F + F a = 0. \quad (2.2)$$

We write $a = \begin{bmatrix} A & B \\ C & D \end{bmatrix}$, where A, B, C, D are $l \times l$ -blocks, so that condition (2.2) translates into the following equations:

$$a = \begin{bmatrix} A & B = \tilde{\imath} B \\ C = \tilde{\imath} C & D = -\tilde{\imath} A \end{bmatrix}. \quad (2.3)$$

In particular, \mathfrak{sp}_n has dimension $l^2 + l(l+1) = l(n+1)$. The intersection of \mathfrak{sp}_n with the Borel subalgebra $\mathfrak{b}_n := \mathfrak{b}_n(K)$ of upper-triangular matrices is a solvable subalgebra of \mathfrak{sp}_n of dimension $l(l+1) = l^2 + l$. Since \mathfrak{sp}_n is a Lie algebra of type C_l , the number of positive roots is l^2 and the number of simple roots is l ; we hence see that $\mathfrak{b}(\mathfrak{sp}_n) := \mathfrak{sp}_n \cap \mathfrak{b}_n$ is a solvable subalgebra of maximal dimension and hence a Borel subalgebra. This is one of the advantages of working with the form F given by (2.1).

2.2 Orthogonal group

Let V be an n -dimensional complex vector space (where n can be even or odd). Let us fix a basis of V and let us choose the non-degenerate bilinear form on V associated with the matrix $F = J_n$. The orthogonal group O_n consists of those matrices $A \in GL_n$ for which ${}^T A F A = F$ holds true. The Lie algebra \mathfrak{o}_n consists of those matrices $a \in \mathfrak{gl}_n$ satisfying (2.2) which translates into the relation

$$a = -\tilde{\imath} a. \quad (2.4)$$

In particular, \mathfrak{o}_n has dimension $\frac{n(n-1)}{2}$. The intersection of \mathfrak{o}_n with the Borel subalgebra \mathfrak{b}_n of upper-triangular matrices, is a solvable subalgebra of \mathfrak{o}_n .

- If $n = 2l$, the dimension of such a solvable subalgebra is easily seen to be

$$\frac{n(n-1)}{2} - l(l-1) = l(2l-1) - l(l-1) = l^2.$$

Since \mathfrak{o}_n is a Lie algebra of type D_l , the number of positive roots is $l(l-1)$ and the number of simple roots is l ; we hence see that $\mathfrak{b}(\mathfrak{o}_n) := \mathfrak{o}_n \cap \mathfrak{b}_n$ is a solvable subalgebra of maximal dimension and hence a Borel subalgebra.

- Similarly, if $n = 2l + 1$, the dimension of $\mathfrak{b}(\mathfrak{o}_n) := \mathfrak{o}_n \cap \mathfrak{b}_n$ is easily seen to be

$$\frac{n(n-1)}{2} - (l(l-1) + l) = (2l+1)l - l^2 = l^2 + l.$$

Since \mathfrak{o}_n is a Lie algebra of type B_l , the number of positive roots is l^2 and the number of simple roots is l ; we hence see that $\mathfrak{b}(\mathfrak{o}_n)$ is a solvable subalgebra of maximal dimension and hence a Borel subalgebra.

As before, this is one of the benefits of working with the form F given by J_n .

3 Background on (symmetric) quiver representations

We include basic knowledge about the representation theory of finite-dimensional algebras via finite quivers [1] before introducing the notion of a symmetric quiver and discussing its representations. This theoretical background will be necessary later on to prove our main results.

A *finite quiver* \mathcal{Q} is a directed graph $\mathcal{Q} = (\mathcal{Q}_0, \mathcal{Q}_1, s, t)$, such that \mathcal{Q}_0 is a finite set of *vertices* and \mathcal{Q}_1 is a finite set of *arrows*, whose elements are written as $\alpha: s(\alpha) \rightarrow t(\alpha)$. The *path algebra* $K\mathcal{Q}$ is defined as the K -vector space with a basis consisting of all paths in \mathcal{Q} , that is, sequences of arrows $\omega = \alpha_s \dots \alpha_1$ with $t(\alpha_k) = s(\alpha_{k+1})$ for all $k \in \{1, \dots, s-1\}$; formally included is a path ϵ_i of length zero for each $i \in \mathcal{Q}_0$ starting and ending in i . The multiplication is defined as the concatenation of paths $\omega = \alpha_s \dots \alpha_1$ and $\omega' = \beta_t \dots \beta_1$, that is,

$$\omega \cdot \omega' = \begin{cases} \alpha_s \dots \alpha_1 \beta_t \dots \beta_1, & \text{if } t(\beta_t) = s(\alpha_1); \\ 0, & \text{otherwise.} \end{cases}$$

Let $\text{rad}(K\mathcal{Q})$ be the *path ideal* of $K\mathcal{Q}$, which is the (two-sided) ideal generated by all paths of positive lengths. An ideal $I \subseteq K\mathcal{Q}$ is called *admissible* if there exists an integer s with $\text{rad}(K\mathcal{Q})^s \subset I \subset \text{rad}(K\mathcal{Q})^2$. If this is the case for an ideal I , then the algebra $\mathcal{A} := K\mathcal{Q}/I$ is finite-dimensional.

We denote by $\text{rep}(K\mathcal{Q})$ the abelian K -linear category of all representations of \mathcal{Q} (which is equivalent to the category of $K\mathcal{Q}$ -modules). In more detail, the objects are given as finite-dimensional (K -)representations of \mathcal{Q} which, in more detail, are tuples

$$((M_i)_{i \in \mathcal{Q}_0}, (M_\alpha: M_i \rightarrow M_j)_{(\alpha: i \rightarrow j) \in \mathcal{Q}_1}),$$

where the M_i are K -vector spaces, and the M_α are K -linear maps. A *morphism of representations* $M = ((M_i)_{i \in \mathcal{Q}_0}, (M_\alpha)_{\alpha \in \mathcal{Q}_1})$ and $M' = ((M'_i)_{i \in \mathcal{Q}_0}, (M'_\alpha)_{\alpha \in \mathcal{Q}_1})$ consists of a tuple of K -linear maps $(f_i: M_i \rightarrow M'_i)_{i \in \mathcal{Q}_0}$, such that $f_j M_\alpha = M'_\alpha f_i$ for every arrow $\alpha: i \rightarrow j$ in \mathcal{Q}_1 .

Let us denote by $\text{rep}(\mathcal{A})$ the category of representations of \mathcal{Q} bound by I : For a representation M and a path ω in \mathcal{Q} as above, we denote $M_\omega = M_{\alpha_s} \cdot \dots \cdot M_{\alpha_1}$. A representation M is called *bound by I* , if $\sum_\omega \lambda_\omega M_\omega = 0$ whenever $\sum_\omega \lambda_\omega \omega \in I$. The category $\text{rep}(K\mathcal{Q}/I)$ is equivalent to the category of finite-dimensional \mathcal{A} -representations.

Let M be an \mathcal{A} -representation, let $B_i \subseteq \epsilon_i M$ be a K -basis of $\epsilon_i M$ for every $i \in \mathcal{Q}_0$ and let B be the disjoint union of these sets B_i . We define the *coefficient quiver* $\Gamma(M) := \Gamma(M, B)$ of M with respect to the basis B to be the quiver with exactly one vertex for each element of B , such that for each arrow $\alpha \in \mathcal{Q}_1$ and every element $b \in B_{s(\alpha)}$ we have

$$M_\alpha(b) = \sum_{c \in B_{t(\alpha)}} \lambda_{b,c}^\alpha c$$

with $\lambda_{b,c}^\alpha \in K$. For each $\lambda_{b,c}^\alpha \neq 0$ we draw an arrow $b \rightarrow c$ with label α . Thus, the quiver reflects the coefficients corresponding to the representation M with respect to the chosen basis B .

Given a representation $M \in \text{rep}(\mathcal{A})$, its *dimension vector* $\underline{\dim} M \in \mathbf{N}\mathcal{Q}_0$ is defined by $(\underline{\dim} M)_i = \dim_K M_i$ for $i \in \mathcal{Q}_0$. For a fixed dimension vector $\underline{d} \in \mathbf{N}\mathcal{Q}_0$, we denote by $\text{rep}(\mathcal{A}, \underline{d})$ the full subcategory of $\text{rep}(\mathcal{A})$ which consists of representations of dimension vector \underline{d} .

Let M and M' be two representations of \mathcal{A} . We denote by $\text{Hom}_{\mathcal{A}}(M, M')$ the space of homomorphisms from M to M' , by $\text{End}_{\mathcal{A}}(M)$ the set of endomorphisms and by $\text{Aut}_{\mathcal{A}}(M)$ the group of automorphisms of M in $\text{rep}(\mathcal{A})$.

For certain finite-dimensional algebras a convenient tool for the classification of the indecomposable representations (up to isomorphism) and of their homomorphisms is the *Auslander–Reiten quiver* $\Gamma(\mathcal{A})$ of $\text{rep}(\mathcal{A})$. Its vertices $[M]$ are given by the isomorphism classes of indecomposable representations of $\text{rep}(\mathcal{A})$; the arrows between two such vertices $[M]$ and $[M']$ are parametrized by a basis of the space of so-called irreducible maps $f: M \rightarrow M'$.

By defining the affine space $\mathbf{R}_{\underline{d}}(K\mathcal{Q}) := \bigoplus_{\alpha: i \rightarrow j} \text{Hom}_K(K^{d_i}, K^{d_j})$, one realizes that its points m naturally correspond to representations $M \in \text{rep}(K\mathcal{Q}, \underline{d})$ with $M_i = K^{d_i}$ for $i \in \mathcal{Q}_0$. Via this correspondence, the set of such representations bound by I corresponds to a closed subvariety $\mathbf{R}_{\underline{d}}(\mathcal{A}) \subset \mathbf{R}_{\underline{d}}(K\mathcal{Q})$.

The algebraic group $\text{GL}_{\underline{d}} = \prod_{i \in \mathcal{Q}_0} \text{GL}_{d_i}$ acts on $\mathbf{R}_{\underline{d}}(K\mathcal{Q})$ and on $\mathbf{R}_{\underline{d}}(\mathcal{A})$ via base change, furthermore the $\text{GL}_{\underline{d}}$ -orbits \mathcal{O}_M of this action are in bijection to the isomorphism classes of representations M in $\text{rep}(\mathcal{A}, \underline{d})$.

The notion of symmetry for a finite quiver comes into the picture as follows: A *symmetric quiver* is a pair (\mathcal{Q}, σ) where \mathcal{Q} is a finite quiver and $\sigma: \mathcal{Q}_0 \cup \mathcal{Q}_1 \rightarrow \mathcal{Q}_0 \cup \mathcal{Q}_1$ is an involution, such that $\sigma(\mathcal{Q}_0) = \mathcal{Q}_0$, $\sigma(\mathcal{Q}_1) = \mathcal{Q}_1$ and every arrow $i \xrightarrow{\alpha} j$ is sent to

the arrow $\sigma(j) \xrightarrow{\sigma(\alpha)} \sigma(i)$.

In this article, we represent the action of σ by adding the symbol $*$. For example,

$$1 \xrightarrow{a} 2 \xrightarrow{b} 3 \xrightarrow{b^*} 2^* \xrightarrow{a^*} 1^*$$

is the symmetric quiver (\mathcal{Q}, σ) with underlying quiver \mathcal{Q} being equioriented of type A_5 , such that σ acts on \mathcal{Q} by sending an element $x \in \mathcal{Q}_0 \cup \mathcal{Q}_1$ to x^* ; the vertex 3 is fixed by σ .

A *symmetric (K-)representation* of a symmetric quiver (\mathcal{Q}, σ) is a representation $M = (\{M_p\}_{p \in \mathcal{Q}_0}, \{M_\alpha\}_{\alpha \in \mathcal{Q}_1})$ in $\text{rep}(K\mathcal{Q})$ endowed with a non-degenerate bilinear form

$$\langle -, - \rangle : \bigoplus_{p \in \mathcal{Q}_0} M_p \times \bigoplus_{q \in \mathcal{Q}_0} M_q \rightarrow K,$$

such that:

(i) The equation

$$\langle -, - \rangle|_{M_p \times M_q} = 0 \quad (3.1)$$

holds true, unless $q = \sigma(p)$;

(ii) The equation

$$\langle M_\alpha(v), w \rangle + \langle v, M_{\sigma(\alpha)}(w) \rangle = 0 \quad (3.2)$$

holds true for every $v \in M_p$, $w \in M_{\sigma(q)}$ and for every arrow $p \xrightarrow{\alpha} q \in \mathcal{Q}_1$.

A representation $(M, \langle -, - \rangle)$ of a symmetric quiver (\mathcal{Q}, σ) is called *symplectic*, if the bilinear form is skew-symmetric and it is called *orthogonal*, if the bilinear form is symmetric.

Let (\mathcal{Q}, σ) be a symmetric quiver and let I be an ideal of $K\mathcal{Q}$, such that $\sigma \cdot I \subset I$. The involution σ induces an involution on the algebra $\mathcal{A} := K\mathcal{Q}/I$ and we can consider symplectic and orthogonal representations of the algebra \mathcal{A} : these are symplectic or orthogonal representations of \mathcal{A} which are annihilated by the ideal I .

We denote the categories of symmetric, symplectic and orthogonal representations by $\text{srep}(\mathcal{A})$ and make sure that it will always be clear from the context which one is meant. The restriction to the full subcategory of representations of a fixed dimension vector \underline{d} is denoted by $\text{srep}(\mathcal{A}, \underline{d})$. Analogously to the non-symmetric case, we associate a variety $\text{SR}_{\underline{d}}(\mathcal{A})$ to this category; and denote $M_x \in \text{srep}(\mathcal{A}, \underline{d})$ for $x \in \text{SR}_{\underline{d}}(\mathcal{A})$.

Let $(M, \langle -, - \rangle)$ and $(M', \langle -, - \rangle')$ be two representations in $\text{srep}(\mathcal{A})$. Let us denote by $\text{Hom}_{\text{srep}(\mathcal{A})}(M, M')$ the space of homomorphisms from M to M' and by $\text{Aut}_{\text{srep}(\mathcal{A})}(M)$ the group of automorphisms of M in $\text{srep}(\mathcal{A})$.

We have

$$\text{Aut}_{\text{srep}(K\mathcal{Q})}(M) = \text{Aut}_{K\mathcal{Q}}(M) \bigcap \prod_{p \neq \sigma(p)} \text{Sym}(M_p \oplus M_{\sigma(p)}) \times \prod_{p = \sigma(p)} \text{Sym}(M_p).$$

Let $\text{End}_{\text{rep}(\mathcal{A})}(M)$ be the Lie algebra of $\text{Aut}_{\text{rep}(\mathcal{A})}(M)$. An element $A = \{A_p\}_{p \in Q_0}$ of $\text{End}_{\text{rep}(\mathcal{A})}(M)$ is called a *symmetric endomorphism* of $(M, \langle -, - \rangle)$; it is an element of $\text{End}_{\mathcal{A}}(M)$ with the following extra conditions:

$$\begin{aligned} \text{If } M \text{ is symplectic: } & p \neq \sigma(p) \Rightarrow A_{\sigma(p)} = -\mathfrak{T}A_p \\ \text{If } M \text{ is orthogonal: } & p \neq \sigma(p) \Rightarrow A_{\sigma(p)} = \mathfrak{T}A_p \\ \text{Furthermore, } & p = \sigma(p) \Rightarrow {}^T A_p F_p + F_p A_p = 0. \end{aligned} \quad (3.3)$$

$$(3.4)$$

Remark 3.1. Conditions (3.3) follow by imposing the relation

$$\begin{bmatrix} {}^T A_p & 0 \\ 0 & {}^T A_{\sigma(p)} \end{bmatrix} \begin{bmatrix} 0 & J \\ \pm J & 0 \end{bmatrix} + \begin{bmatrix} 0 & J \\ \pm J & 0 \end{bmatrix} \begin{bmatrix} A_p & 0 \\ 0 & A_{\sigma(p)} \end{bmatrix} = 0.$$

Condition (3.4) means that A_p belongs to the Lie algebra of $\text{Sym}(V_p)$.

We hence have

$$\text{End}_{\text{rep}(KQ)}(M) = \text{End}_{KQ}(M) \bigcap \prod_{p \neq \sigma(p)} \text{sym}(V_p \oplus V_{\sigma(p)}) \times \prod_{p = \sigma(p)} \text{sym}(V_p).$$

4 B-orbits vs. isoclasses of symmetric representations

Let $G \in \{\text{SP}_n, \text{O}_n\}$ where $n = 2l$ in the symplectic case and $n \in \{2l, 2l + 1\}$ in the orthogonal case for some integer $l \in \mathbf{N}$ and let \mathfrak{g} be the corresponding symplectic or orthogonal Lie algebra. Let B be the standard Borel subgroup of G , that is, the subgroup of G of upper-triangular matrices which is obtained by intersecting the Borel subgroup of GL_n with G .

We consider the algebraic variety $\mathcal{N}(2)$ of 2-nilpotent elements of \mathfrak{g}

$$\mathcal{N}(2) = \mathcal{N}(2, G) = \{x \in \mathfrak{g} \mid x^2 = 0\}.$$

Then B acts on $\mathcal{N}(2)$ via conjugation and our first aim in this article is to prove by means of symmetric quiver representations that the action admits only a finite number of orbits. We thereby specify an explicit parametrization of the orbits.

4.1 Symmetric quiver setup

We define $\mathcal{A}(l)$ to be the algebra given by the quiver

$$Q_l : 1 \xrightarrow{a_1} 2 \xrightarrow{a_2} \dots \xrightarrow{a_{k-1}} k \xrightarrow{a_k} \omega \xrightarrow{a_k^*} k^* \xrightarrow{a_{k-1}^*} \dots \xrightarrow{a_2^*} 2^* \xrightarrow{a_1^*} 1^*$$

with relations $\alpha^2 = a_l^* a_l = 0$. Notice that the $2l$ vertices of Q_l are colored; the choice of the color will be clear in a few lines.

We consider the dimension vector

$$\mathbf{d}_\bullet = (d_1, \dots, d_l, d_\omega, d_{k^*}, \dots, d_{l^*}) = (1, 2, \dots, l-1, l, n, l, l-1, \dots, 2, 1)$$

and the variety $\text{SR}_{\mathbf{d}_\bullet}(\mathcal{A}(l))$. This variety is acted upon by the group

$$\text{GL}_{\text{sym}} := \text{GL}(d_1) \times \text{GL}(d_2) \times \dots \times \text{GL}(d_l) \times \text{Sym}(n)$$

where $\text{Sym}(n)$ denotes either the symplectic or the orthogonal group on a vector space of dimension n . Inside the variety $\text{SR}_{\mathbf{d}_\bullet}(\mathcal{A}(l))$ we consider the open subset $\text{SR}_{\mathbf{d}_\bullet}(\mathcal{A}(l))^0$ corresponding to the full subcategory $\text{srep}(\mathcal{A}(l), \mathbf{d}_\bullet)^0$ of $\text{srep}(\mathcal{A}(l), \mathbf{d}_\bullet)$ of those representations whose linear maps associated with the arrows a_i and a_i^* have maximal rank. For an element $x \in \text{SR}_{\mathbf{d}_\bullet}(\mathcal{A}(l))^0$, we have the natural notion of its GL_{sym} -stabilizer. We denote the corresponding stabilizer of the representation M_x in the category $\text{srep}(\mathcal{A}(l), \mathbf{d}_\bullet)$ by $\text{stab}_{\text{GL}_{\text{sym}}}(M_x)$.

Example 4.1. Let us consider the quiver \mathcal{Q}_2

$$1 \xrightarrow{a} 2 \xrightarrow{b} 3 \xrightarrow{b^*} 2^* \xrightarrow{a^*} 1^*$$

and the algebra $\mathcal{A}(2) = K\mathcal{Q}/(\alpha^2, b^*b)$. Let us consider the $\mathcal{A}(2)$ -representation M_0 given by

$$\begin{array}{ccccccc} 1 & \xrightarrow{a} & 2 & \xrightarrow{b} & 3 & & \\ & & & & & & \\ & & 2 & \xrightarrow{b} & 3 & & \\ & & & & 3 & \xrightarrow{b^*} & 2^* \\ & & & & 3 & \xrightarrow{b^*} & 2^* \xrightarrow{a^*} & 1^* \end{array} \quad (4.1)$$

and the $\mathcal{A}(2)$ -representation M'_0 given by

$$\begin{array}{ccccccc} 1 & \xrightarrow{a} & 2 & \xrightarrow{b} & 3 & & \\ & & & & & & \\ & & 2 & \xrightarrow{b} & 3 & & \\ & & & & 3 & & \\ & & & & 3 & \xrightarrow{b^*} & 2^* \\ & & & & 3 & \xrightarrow{b^*} & 2^* \xrightarrow{a^*} & 1^* \end{array} \quad (4.2)$$

In view of (3.2), in order for M to be symmetric, the arrows a, b of \mathcal{Q}_2 must act by 1 and the arrows a^*, b^* must act as -1 .

The symmetric structure of M_0 (that is, the choice of a non-degenerate bilinear form) is induced by the symmetric structure on the vector space at vertex 3. In the symplectic case, this bilinear form is given by the matrix

$$\begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{bmatrix};$$

and for the orthogonal case, it is defined by the anti-diagonal matrix with every entry on the anti-diagonal being 1 (see Section 2).

The symplectic space $\text{End}_{\text{rep}(\mathcal{A}(2))}(M_0)$ has dimension 6 and can be represented by the matrix

$$\left[\begin{array}{cc|cc} a & c & f & g \\ 0 & b & e & f \\ \hline 0 & 0 & -b & -c \\ 0 & 0 & 0 & -a \end{array} \right].$$

In orthogonal type, it is 4-dimensional and represented by

$$\left[\begin{array}{cc|cc} a & c & f & 0 \\ 0 & b & 0 & -f \\ \hline 0 & 0 & -b & -c \\ 0 & 0 & 0 & -a \end{array} \right].$$

In a similar way, we proceed for the orthogonal group \mathcal{O}_5 and look at the representation M'_0 . Then, as above, the stabilizer is given by

$$\left[\begin{array}{ccc|cc} a & c & e & f & 0 \\ 0 & b & d & 0 & -f \\ \hline 0 & 0 & 0 & -d & -e \\ 0 & 0 & 0 & -b & -c \\ 0 & 0 & 0 & 0 & -a \end{array} \right]$$

We have hence found the well known fact that the stabilizer of the complete standard flag is the Borel subgroup in both the symplectic and orthogonal setup. The Borel subgroups therefore equal the intersection of G with the standard Borel of upper-triangular matrices in GL_n .

Clearly, this example generalizes to larger n in a straight forward manner.

4.2 Translation

The translation from B -orbits in $\mathcal{N}(2)$ to the representation theory of a symmetric quiver is based on a theorem on associated fibre bundles which we recall for the convenience of the reader. Its origin can be found in [18].

Theorem 4.2. *Let G be an algebraic group, let X and Y be G -varieties, and let $\pi : X \rightarrow Y$ be a G -equivariant morphism. Assume that Y is a single G -orbit, $Y = Gy_0$. Define $H := \text{Stab}_G(y_0) = \{g \in G \mid g \cdot y_0 = y_0\}$ and $F := \pi^{-1}(y_0)$. Then X is isomorphic to the associated fibre bundle $G \times^H F$, and the embedding $\iota : F \rightarrow X$ induces a bijection between H -orbits in F and G -orbits in X preserving orbit closures.*

Corollary 4.3. *With the notation of Theorem 4.2, given a point $p \in F$, we have $\text{stab}_H(p) = \text{stab}_G(p)$*

Proof. Since H is a subgroup of G , $\text{Stab}_H(p) \subseteq \text{Stab}_G(p)$; viceversa, since $H \cdot p = G \cdot p \cap F$, the reversed inclusion also holds. \square

In view of Theorem 4.2, we can now prove the following key lemma, analogous to [3, Lemma 3.1].

Lemma 4.4. *There is a bijection between isoclasses of symplectic/orthogonal $\mathcal{A}(l)$ -representations in $\text{srep}(\mathcal{A}(l), \mathbf{d}_\bullet)^0$ and symplectic/orthogonal B -orbits in $\mathcal{N}(2)$. This bijection respects orbit closure relations and dimensions of stabilizers.*

Proof. Let \widetilde{Q}_l be the quiver obtained from Q_l by removing the loop α and let $\widetilde{\mathcal{A}(l)}$ be the corresponding symmetric algebra (also remove the relation α^2). By defining $\text{SR}_{\mathbf{d}_\bullet}(\widetilde{\mathcal{A}(l)})^0$ analogously to $\text{SR}_{\mathbf{d}_\bullet}(\mathcal{A}(l))^0$, we see that this variety is acted upon transitively by GL_{sym} and we denote the representation which is given by the complete standard flag by M_0 ; this is a generating point. The embedding $\widetilde{\mathcal{A}(l)} \subset \mathcal{A}(l)$ induces a GL_{sym} -equivariant projection

$$\pi : \text{SR}_{\mathbf{d}_\bullet}(\mathcal{A}(l))^0 \longrightarrow \text{SR}_{\mathbf{d}_\bullet}(\widetilde{\mathcal{A}(l)})^0$$

which is given by forgetting the linear map associated with the loop α . The fiber of π equals the variety $\mathcal{N}(2)$.

As we have seen before, the stabilizer of the symplectic/orthogonal representation M_0 is isomorphic to the Borel subgroup B of the symplectic/orthogonal group. Thus, Theorem 4.2 proves the claim. \square

We are hence left to classify the isomorphism classes of symplectic/orthogonal representations of $\mathcal{A}(l)$ of dimension vector \mathbf{d}_\bullet with maximal rank maps, which in view of Krull–Remak–Schmidt’s theorem is analogous to classifying the unique decompositions of elements of $\text{srep}(\mathcal{A}(l), \mathbf{d}_\bullet)^0$ into indecomposable symplectic/orthogonal representations (up to symmetric isomorphism). Let M and M' be two points of $\text{srep}(\mathcal{A}(l), \mathbf{d}_\bullet)^0$ which are contained in different orbits. Since $\pi(M) = \pi(M')$ under the morphism of the proof of Lemma 4.4, the only difference between them is given by the action of the loop α . This means that the only part of the coefficient quivers of M and M' which differs is the subquiver which represents the loop α .

5 Representation theory of $\mathcal{A}(l)$

In this section, we look at the (symmetric) representation theory of the algebra $\mathcal{A}(l)$ corresponding to the symmetric quiver Q_l . With these considerations, we are able to prove explicit parametrizations of the Borel-orbits in $\mathcal{N}(2)$ in Section 6.

5.1 Indecomposable symmetric $\mathcal{A}(l)$ -modules

The following proposition follows from [6, Section 3] by noticing that there are no band modules.

Proposition 5.1. *The algebra $\mathcal{A}(l)$ is a string algebra of finite representation type. In particular, the indecomposable $\mathcal{A}(l)$ -modules are string modules and their isoclasses are parametrized by words with letters in the arrows of Q_l and their inverses, avoiding relations.*

Let us give names to the indecomposable $\mathcal{A}(l)$ -modules (where $l + 1 := \omega$).

M_{ij} : For $1 \leq i \leq j \leq l + 1$, we denote by M_{ij} the string module associated with the word $a_i \cdots a_{j-1}$, i.e. it is the indecomposable module supported on vertices $i, i + 1, \dots, j$; its coefficient quiver is given by

$$i \longrightarrow i + 1 \longrightarrow \cdots \longrightarrow j - 1 \longrightarrow j$$

M_{ij}^* : For $1 \leq i \leq j \leq l + 1$, we denote by M_{ij}^* the string module associated with the word $a_{j-1}^* \cdots a_i^*$, i.e. it is the indecomposable module supported on vertices $j^*, (j - 1)^*, \dots, i^*$; its coefficient quiver is given by

$$j^* \longrightarrow (j - 1)^* \longrightarrow \cdots \longrightarrow (i + 1)^* \longrightarrow i^*$$

D_{ij}^+ : For $1 \leq i \leq j \leq l + 1$, we denote by D_{ij}^+ the indecomposable associated with the word $a_i a_{i+1} \cdots a_l \alpha^- a_l^- a_{l-1}^- \cdots a_j^-$; its coefficient quiver has the following form

$$\begin{array}{ccccccccccc} i & \longrightarrow & i + 1 & \longrightarrow & \cdots & \longrightarrow & j & \longrightarrow & j + 1 & \longrightarrow & \cdots & \longrightarrow & l & \longrightarrow & \omega \\ & & & & & & & & & & & & & & \uparrow \\ & & & & & & & & & & & & & & \omega \\ & & & & & & & & & & & & & & \downarrow \\ & & & & & & & & & & & & & & \omega \\ & & & & & & & & & & & & & & \downarrow \\ & & & & & & & & & & & & & & \omega \end{array}$$

D_{ij}^- : For $1 \leq i < j \leq l + 1$ we denote by D_{ij}^- the indecomposable associated with the word $a_i a_{i+1} \cdots a_l \alpha a_l^+ a_{l-1}^+ \cdots a_j^+$; its coefficient quiver is given by

$$\begin{array}{ccccccccccc} i & \longrightarrow & i + 1 & \longrightarrow & \cdots & \longrightarrow & j & \longrightarrow & j + 1 & \longrightarrow & \cdots & \longrightarrow & l & \longrightarrow & \omega \\ & & & & & & & & & & & & & & \downarrow \\ & & & & & & & & & & & & & & \omega \\ & & & & & & & & & & & & & & \downarrow \\ & & & & & & & & & & & & & & \omega \end{array}$$

C_{ij}^+ : For $1 \leq i \leq j \leq l + 1$ we denote by C_{ij}^+ the indecomposable associated with the word $(a_i^+)^- (a_{j-1}^+)^- \cdots (a_l^+)^- \alpha^- a_l^+ a_{l-1}^+ \cdots a_i^+$; its coefficient quiver is given by

$$\begin{array}{ccccccccccc} \omega & \longrightarrow & l^* & \longrightarrow & \cdots & \longrightarrow & j^* & \longrightarrow & (j - 1)^* & \longrightarrow & \cdots & \longrightarrow & (i + 1)^* & \longrightarrow & i^* \\ \downarrow & & & & & & & & & & & & & & \\ \omega & \longrightarrow & l^* & \longrightarrow & \cdots & \longrightarrow & j^* & & & & & & & & \end{array}$$

C_{ij}^- : For $1 \leq i < j \leq l+1$ we denote by C_{ij}^- the indecomposable module associated with the word $(a_j^*)^-(a_{j-1}^*)^- \cdots (a_i^*)^- \alpha a_i^* a_{i-1}^* \cdots a_1^*$; its coefficient quiver is given by

$$\begin{array}{ccccccccccc} \omega & \longrightarrow & l^* & \longrightarrow & \cdots & \longrightarrow & j^* & \longrightarrow & (j-1)^* & \longrightarrow & \cdots & \longrightarrow & (i+1)^* & \longrightarrow & i^* \\ \uparrow & & & & & & & & & & & & & & & \\ \omega & \longrightarrow & l^* & \longrightarrow & \cdots & \longrightarrow & j^* & & & & & & & & & \end{array}$$

Z_{ij}^+ : For $1 \leq i, j \leq l$ we denote by Z_{ij}^+ the indecomposable associated with the word $a_i a_{i+1} \cdots a_l \alpha^- a_l^* \cdots a_j^*$; its coefficient quiver is given by

$$\begin{array}{ccccccc} i & \longrightarrow & i+1 & \longrightarrow & \cdots & \longrightarrow & l & \longrightarrow & \omega \\ & & & & & & & & \uparrow \\ & & & & & & & & \omega & \longrightarrow & l^* & \longrightarrow & \cdots & \longrightarrow & j^* \end{array}$$

Z_{ij}^- : For $1 \leq i, j \leq l$ we denote by Z_{ij}^- the indecomposable associated with the word $a_i a_{i+1} \cdots a_l \alpha a_l^* \cdots a_j^*$; its coefficient quiver is given by

$$\begin{array}{ccccccc} i & \longrightarrow & i+1 & \longrightarrow & \cdots & \longrightarrow & l & \longrightarrow & \omega \\ & & & & & & & & \downarrow \\ & & & & & & & & \omega & \longrightarrow & l^* & \longrightarrow & \cdots & \longrightarrow & j^* \end{array}$$

Remark 5.2. All the modules above are non-isomorphic to each other, apart from $D_{l+1,l+1}^+ \simeq C_{l+1,l+1}^+$ and $M_{l+1,l+1} \simeq M_{l+1,l+1}^*$.

We consider the involution σ of \mathcal{Q}_l , which sends every vertex i to i^* , every arrow a to a^* and which fixes ω and α (here we use the convention that $(-)^{**} = (-)$). Then (\mathcal{Q}_l, σ) is a symmetric quiver and we can consider symmetric representations of $\mathcal{A}(l)$. The involution σ induces a duality on the category of representations of $\mathcal{A}(l)$ that we denote by ∇ (as in [7]).

Convention 5.3. Given an indecomposable $\mathcal{A}(l)$ -module M , we need to choose carefully the linear maps. Since we often work with its coefficient quiver, we fix one and for all a convention about these:

The arrows of the coefficient quiver of M colored with a_1, \dots, a_l act as 1, while the arrows colored with a_l^*, \dots, a_1^* act as -1 .

Every pair of two arrows $\omega_i \xrightarrow{\alpha_1} \omega_j$ and $\omega_{j^*} \xrightarrow{\alpha_2} \omega_{i^*}$ colored with ω (if they exist) has to satisfy the following conditions:

- For V to be orthogonal, α_1 acts as 1 and α_2 as -1 .
- For V to be symplectic, if $1 \leq i, j \leq l$ or $1 \leq i^*, j^* \leq l$, then α_1 acts as 1 and α_2 as -1 , otherwise α_1 and α_2 both act as 1.

5.2 Auslander–Reiten quiver of $\mathcal{A}(l)$

The algebra $\mathcal{A}(l)$ is a string algebra of finite representation–type, that is, it does only admit a finite number of isomorphism classes of indecomposable representations. Its Auslander–Reiten quiver can be obtained in several ways. We prefer to follow the treatment of Butler–Ringel [4] and get the following result.

Proposition 5.7. *The following are the Auslander–Reiten sequences of $\mathcal{A}(l)$:*

(i) *Auslander–Reiten sequences starting with M_{ij} :*

$$\begin{aligned} 0 &\longrightarrow M_{1,\omega} \longrightarrow Z_{1,1}^+ \longrightarrow M_{1,\omega}^* \longrightarrow 0, \\ 0 &\longrightarrow M_{i,\omega} \longrightarrow M_{i-1,\omega} \oplus Z_{i,1}^+ \longrightarrow Z_{i-1,1}^+ \longrightarrow 0, \text{ if } i > 1, \\ 0 &\longrightarrow M_{i,j} \longrightarrow M_{i,j-1} \oplus M_{i-1,j} \longrightarrow M_{i-1,j-1} \longrightarrow 0, \text{ if } i > 1 \text{ and } j \leq l, \\ 0 &\longrightarrow M_{i,i} = S_i \longrightarrow M_{i-1,i} \longrightarrow M_{i-1,i-1} \longrightarrow 0, \text{ if } i = j > 1. \end{aligned}$$

(ii) *Auslander–Reiten sequences starting with M_{ij}^* :*

$$\begin{aligned} 0 &\longrightarrow M_{1,j}^* \longrightarrow M_{2,j}^* \oplus M_{1,j+1}^* \longrightarrow M_{2,j+1}^* \longrightarrow 0, \text{ if } j \leq l-1, \\ 0 &\longrightarrow M_{1,l}^* \longrightarrow M_{2,l}^* \oplus P_\omega \longrightarrow C_{1,l}^- \longrightarrow 0, \\ 0 &\longrightarrow M_{1,\omega}^* \longrightarrow M_{2,\omega}^* \oplus C_{1,1}^+ \longrightarrow C_{1,2}^- \longrightarrow 0, \\ 0 &\longrightarrow M_{i,i}^* = S_{i^*} \longrightarrow M_{i+1,i}^* \longrightarrow M_{i+1,i+1}^* \longrightarrow 0, \text{ if } i < l, \\ 0 &\longrightarrow M_{l,l}^* = S_{l^*} \longrightarrow C_{1,l}^- \longrightarrow C_{1,\omega}^- \longrightarrow 0, \\ 0 &\longrightarrow S_\omega \longrightarrow M_{l,\omega} \oplus C_{1,\omega}^+ \longrightarrow Z_{l,1}^+ \longrightarrow 0, \\ 0 &\longrightarrow M_{i,\omega}^* \longrightarrow M_{i+1,\omega}^* \oplus C_{1,i}^+ \longrightarrow C_{1,i+1}^+ \longrightarrow 0, \text{ if } 1 < i < l, \\ 0 &\longrightarrow M_{l,\omega}^* \longrightarrow S_\omega \oplus C_{1,l}^+ \longrightarrow C_{1,\omega}^+ \longrightarrow 0, \\ 0 &\longrightarrow M_{i,j}^* \longrightarrow M_{i+1,j}^* \oplus M_{i,j+1}^* \longrightarrow M_{i+1,j+1}^* \longrightarrow 0, \text{ if } i > 1 \text{ and } j < l. \end{aligned}$$

(iii) *Auslander–Reiten sequences starting with D_{ij}^+ :*

$$\begin{aligned} 0 &\longrightarrow D_{1,\omega}^+ \longrightarrow D_{1,l}^+ \oplus S_\omega \longrightarrow M_{l,\omega} \longrightarrow 0, \\ 0 &\longrightarrow D_{i,\omega}^+ \longrightarrow D_{i-1,\omega}^+ \oplus D_{i,l}^+ \longrightarrow D_{i-1,l}^+ \longrightarrow 0, \text{ if } 1 < i \leq l, \\ 0 &\longrightarrow D_{1,j}^+ \longrightarrow D_{1,j-1}^+ \oplus M_{j,\omega} \longrightarrow M_{j-1,\omega} \longrightarrow 0, \text{ if } 1 < j \leq l, \\ 0 &\longrightarrow D_{i,j}^+ \longrightarrow D_{i-1,j}^+ \oplus D_{i,j-1}^+ \longrightarrow D_{i-1,j-1}^+ \longrightarrow 0, \text{ if } 1 < i \leq j \leq l. \end{aligned}$$

(iv) Auslander–Reiten sequences starting with D_{ij}^- :

$$\begin{aligned}
0 &\longrightarrow D_{1,\omega}^- \longrightarrow D_{1,l}^- \longrightarrow M_{l,l} = S_l \longrightarrow 0, \\
0 &\longrightarrow D_{i,\omega}^- \longrightarrow D_{i-1,\omega}^- \oplus D_{i,l}^+ \longrightarrow D_{i-1,l}^- \longrightarrow 0, \text{ if } 1 < i \leq l, \\
0 &\longrightarrow D_{i,i}^+ \longrightarrow D_{i-1,i}^- \oplus D_{i-1,i}^+ \longrightarrow D_{i-1,i-1}^+ \longrightarrow 0, \text{ if } 1 < i \leq \omega \\
0 &\longrightarrow D_{1,j}^- \longrightarrow D_{1,j-1}^- \oplus M_{j,l} \longrightarrow M_{j-1,l} \longrightarrow 0, \text{ if } 1 < j \leq l, \\
0 &\longrightarrow D_{i,j}^- \longrightarrow D_{i-1,j}^- \oplus D_{i,j-1}^- \longrightarrow D_{i-1,j-1}^- \longrightarrow 0, \text{ if } 1 < i < j \leq l.
\end{aligned}$$

(v) Auslander–Reiten sequences starting with C_{ij}^+ :

$$\begin{aligned}
0 &\longrightarrow C_{i,\omega}^+ \longrightarrow C_{i+1,\omega}^+ \oplus Z_{l,i}^+ \longrightarrow Z_{l,i+1}^+ \longrightarrow 0, \text{ if } 1 \leq i < l, \\
0 &\longrightarrow C_{l,\omega}^+ \longrightarrow C_{\omega,\omega}^+ \oplus Z_{l,l}^+ \longrightarrow D_{l,\omega}^+ \longrightarrow 0, \\
0 &\longrightarrow C_{i,j}^+ \longrightarrow C_{i+1,j}^+ \oplus C_{i,j+1}^+ \longrightarrow C_{i+1,j+1}^+ \longrightarrow 0, \text{ if } 1 < i \leq j \leq l.
\end{aligned}$$

(vi) Auslander–Reiten sequences starting with C_{ij}^- :

$$\begin{aligned}
0 &\longrightarrow C_{1,\omega}^- \longrightarrow Z_{l,1}^- = P_l \oplus C_{2,\omega}^- \longrightarrow Z_{l,2}^- \longrightarrow 0, \\
0 &\longrightarrow C_{i,\omega}^- \longrightarrow Z_{l,i}^- \oplus C_{i+1,\omega}^- \longrightarrow Z_{l,i+1}^- \longrightarrow 0, \text{ if } 1 \leq i \leq l, \\
0 &\longrightarrow C_{1,1}^- = P_\omega \longrightarrow C_{1,2}^- \oplus C_{1,2}^+ \longrightarrow C_{2,2}^+ \longrightarrow 0, \\
0 &\longrightarrow C_{i,j}^- \longrightarrow C_{i+1,j}^- \oplus C_{i,j+1}^- \longrightarrow C_{i+1,j+1}^- \longrightarrow 0, \text{ if } 1 < i \leq j \leq n.
\end{aligned}$$

(vii) Auslander–Reiten sequences starting with Z_{ij}^+ (note that $Z_{i,\omega}^+ = D_{i,\omega}^+$):

$$\begin{aligned}
0 &\longrightarrow Z_{1,j}^+ \longrightarrow Z_{1,j+1}^+ \oplus M_{j,\omega}^* \longrightarrow M_{j+1,\omega}^* \longrightarrow 0, \\
0 &\longrightarrow Z_{i,j}^+ \longrightarrow Z_{i,j+1}^+ \oplus Z_{i-1,j}^+ \longrightarrow Z_{i-1,j+1}^+ \longrightarrow 0, \text{ if } i > 1.
\end{aligned}$$

(viii) Auslander–Reiten sequences starting with Z_{ij}^- (note that $Z_{1j}^- = I_j^*$):

$$\begin{aligned}
0 &\longrightarrow Z_{i,1}^- = P_i \longrightarrow Z_{i-1,1}^- \oplus Z_{i,2}^- \longrightarrow Z_{i-1,2}^- \longrightarrow 0, \text{ if } i > 1, \\
0 &\longrightarrow Z_{i,j}^- \longrightarrow Z_{i-1,j}^- \oplus Z_{i,j+1}^- \longrightarrow Z_{i-1,j+1}^- \longrightarrow 0, \text{ if } 1 < i, j \leq l,
\end{aligned}$$

The resulting Auslander–Reiten quiver of $\mathcal{A}(l)$ has the shape of a "christmas tree"; its bottom part consists of pre–projective modules and its top consists of $l + 1$ periodic τ –orbits. The duality ∇ acts as a reflection through the vertical line formed by the self–dual $\mathcal{A}(l)$ –modules Z_{ii}^\pm and $D_{l+1,l+1}^+$. Figure 1 shows the Auslander–Reiten quiver of $\mathcal{A}(3)$.




Figure 1: Auslander–Reiten quiver of $\mathcal{A}(3)$

6 Parametrization of orbits

It is known by Panyushev [17] that B acts finitely on the variety $\mathcal{N}(2)$. We aim to prove explicit parametrizations of the orbits by means of symmetric representations and, thus, in a very combinatorial way. This way, we hope to be able to calculate e.g. degenerations in a follow-up article by means of the used representation-theoretic methods. We begin by discussing symplectic orbits in Subsection 6.1 and deduce orthogonal orbits in Subsection 6.2. In each type, we generalize the results to parabolic orbits in Section 7.

6.1 Orbits in type C

Let $G = \mathrm{SP}_n$, where $n = 2l$ for some integer l . We denote by B the standard Borel subgroup of G and consider the algebra $\mathcal{A}(l)$ and its symmetric representations as discussed in 5. Due to Lemma 4.4, we are interested in symplectic representations of dimension vector $\mathbf{d}_\bullet = (1, 2, \dots, l, 2l, l, \dots, 2, 1)$.

Let us begin with an example.

Example 6.1. *Figure 2 shows the complete list of isomorphism classes of symplectic*

representations in $\text{srep}(\mathcal{A}(2), \mathbf{d}_\bullet)^0$, where $n = 4 = 2l$ and B is the Borel subgroup. In more detail, the indecomposables are displayed by their coefficient quiver and then interpreted combinatorially by graphs on two vertices.

$M_0 = (M_{13} \oplus M_{13}^*) \oplus (M_{23} \oplus M_{23}^*)$				
$D_{12}^+ \oplus \nabla D_{12}^+$			$D_{12}^- \oplus \nabla D_{12}^-$	
$Z_{21}^+ \oplus \nabla Z_{21}^+$			$Z_{12}^- \oplus \nabla Z_{12}^-$	
$Z_{11}^+ \oplus (M_{23} \oplus M_{23}^*)$			$Z_{11}^- \oplus (M_{23} \oplus M_{23}^*)$	
$Z_{22}^+ \oplus (M_{13} \oplus M_{13}^*)$			$Z_{22}^- \oplus (M_{13} \oplus M_{13}^*)$	
$Z_{11}^+ \oplus Z_{22}^+$			$Z_{11}^- \oplus Z_{22}^-$	
$Z_{11}^+ \oplus Z_{22}^-$			$Z_{11}^- \oplus Z_{22}^+$	

Figure 2: Isomorphism classes of symplectic representations in $\text{srep}(\mathcal{A}(2), \mathbf{d}_\bullet)^0$ and their combinatorial interpretation

This observation leads us to the following definition.

Definition 6.2. A symplectic oriented link pattern (*solp* for short) of size l consists of a set of l vertices $1, 2, \dots, l$ together with a collection of oriented arrows between these vertices and a collection of oriented arrows with dots between these vertices, such that

(B) every vertex is touched by at most one arrow;

(SpOr) there are no loops of arrows without dots (that is, no arrows without dots from a vertex to itself).







We denote by Solp_l the set of solps of size l .

The definition of symplectic link patterns leads to a combinatorial parametrization of symplectic Borel-orbits in $\mathcal{N}(2)$.

Theorem 6.3. The B -orbits in the variety $\mathcal{N}(2) \subseteq \mathfrak{sp}_n$ are in bijection with the set Solp_l of solps of size l .

Proof. By Krull–Remak–Schmidt, there is a quite obvious bijection between the set of solps of size l and the set of symplectic representations in $\text{srep}(\mathcal{A}(l), \mathbf{d}_\bullet)^0$ up to isomorphism which maps an isomorphism class $[M]$ of a symplectic representation M to the subquiver of the coefficient quiver of M induced by M_α . Since this subquiver is completely determined by the vertices $1, \dots, l$ and the arrows By Lemma 4.4, the claim follows.

Let us state in detail, how the bijection works and how we can read of the representative 2-nilpotent matrix of a particular solp. For this, the following table gives a recipe; let $i < j$ and translate i^* and j^* via $(k^* = n - k + 1)$:

(Part of) solp	Indecomposable	(Part of) Matrix
i	$M_{i,l+1} \oplus M_{i,l+1}^*$	0
	$D_{i,j}^+ \oplus C_{i,j}^+$	$E_{i,j} - E_{j^*,i^*}$
	$D_{i,j}^- \oplus C_{i,j}^-$	$E_{j,i} - E_{i^*,j^*}$
	$Z_{i,j}^+ \oplus Z_{j,i}^+$	$E_{i,j^*} + E_{j,i^*}$
	$Z_{i,j}^- \oplus Z_{j,i}^-$	$E_{j^*,i} + E_{i^*,j}$
	$Z_{i,i}^+$	E_{i,i^*}
	$Z_{i,i}^-$	$E_{i^*,i}$

□

Since we have a combinatorial description, we can count the number of orbits.

Proposition 6.4. *Let s_l be the cardinality of Solp_l . Then the sequence $\{s_l\}$ is determined by*

- $s_0 = 1,$
- $s_1 = 3,$
- $s_l = 3s_{l-1} + 4(l-1)s_{l-2}.$

Proof. We divide the set Solp_l into the subset of symmetric link patterns where vertex 1 is not touched by any arrow and its complement. □

The sequence 1, 3, 13, 63, 345, 2043, ... of numbers of slps is classified in OEIS as A202837 [16].

Remark 6.5. For GL_n , the oriented link patterns considered in [4] only have to satisfy condition (B) that is, the 2-nilpotency conditions. We hence see that solps are special oriented link patterns as defined in [4]. This is not surprising; indeed the following fact is known by [12]: if two symplectic elements are conjugate under the Borel of GL_n , then they are conjugate under the Borel of SP_n , as well.

We finish the symplectic classification by describing all B -orbits for $SP(4)$ in detail.

Example 6.6. *The B -orbits in $\mathcal{N}(2) \subseteq \mathfrak{sp}_n$ are classified by the collection of solps of size 2. These are explicitly listed in the following table and the corresponding representative matrices in $\mathcal{N}(2)$ are displayed. This completes the example which we already considered in Figure 2.*

$\begin{pmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$
$\begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$	
$\begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$	

6.2 Orbits in types B and D

Let $G = O_n$, where $n \in \{2l, 2l + 1\}$. We denote by B the standard Borel subgroup of G and consider the algebra $\mathcal{A}(l)$ as discussed in Section 5. We are interested in orthogonal representations of dimension vector $\mathbf{d}_\bullet = (1, 2, \dots, l, n, l, \dots, 2, 1)$ by Lemma 4.4.

As before, we begin with an example.

Example 6.7. *Figure 3 shows the complete list of isomorphism classes of orthogonal representations in $\text{srep}(\mathcal{A}(2), \mathbf{d}_\bullet)^0$.*

The first table shows the isomorphism classes of orthogonal representations where $\mathbf{d}_\bullet = (1, 2, 4, 2, 1)$, i.e. it corresponds to O_4 .

The second table shows the isomorphism classes of orthogonal representations where $\mathbf{d}_\bullet = (1, 2, 5, 2, 1)$, i.e. it corresponds to O_5 .

This observation leads us to the following definition.




Figure 3: Isomorphism classes of orthogonal representations in $\text{srep}(\mathcal{A}(2), \mathbf{d}_\bullet)^0$ and their combinatorial interpretation

Definition 6.8. An orthogonal oriented link pattern (oolp for short) of size l consists of a set of l vertices $1, 2, \dots, l$ together with a collection of oriented arrows between these vertices and a collection of oriented arrows with dots with dots between these vertices, such that

(B) every vertex is touched by at most one arrow;

(SpOr) there are no loops of arrows without dots (that is, no arrows without dots from a vertex to itself).

(Or) there are no loops of arrows with dots (that is, no arrows with dots from a vertex to itself).

We denote by Oolp_l the set of oolps of size l .

Thus, an oolp is a solp without loops.

As in the symplectic case, the parametrization of the Borel-orbits in $\mathcal{N}(2)$ follows straight away.





Theorem 6.9. The B -orbits in the variety $\mathcal{N}(2) \subseteq \mathfrak{o}_n$, where $n \in \{2l, 2l + 1\}$, are in bijection with the set Oolp_l of oolps of size l .

Proof. In a similar manner to Theorem 6.3, there is a quite obvious bijection between the set Oolp_l of oolps of size l and the set of isoclasses of orthogonal representations

of $\mathcal{A}(l)$ in $\text{srep}(\mathcal{A}(l), \mathbf{d}_\bullet)^0$ which maps an isomorphism class $[M]$ of an orthogonal representation to a particular subquiver of the coefficient quiver of M induced by M_α .

In case n is even, this is done analogously to the symplectic case.

In case n is odd, the middle vertex of this particular subquiver is always determined as a fixed point as visualized in Figure 3. This is due to the fact that it corresponds to the direct summand $M_{\omega, \omega}$. The diagram representing M_α can thus be restricted to $2l$ vertices and we obtain the sought subquiver. This can be translated to an oolp as in the symplectic case again. In more detail, the translation of indecomposables to vertices / arrows in the oolp can be read off the following table - where also the translation to matrices in $\mathcal{N}(2)$ can be found.

(Part of) solp	Indecomposable	(Part of) Matrix
i	$M_{i,l+1} \oplus M_{i,l+1}^*$	0
	$D_{i,j}^+ \oplus C_{i,j}^+$	$E_{i,j} - E_{j^*,i^*}$
	$D_{i,j}^- \oplus C_{i,j}^-$	$E_{j,i} - E_{i^*,j^*}$
	$Z_{i,j}^+ \oplus Z_{j,i}^+$	$E_{i,j^*} - E_{j,i^*}$
	$Z_{i,j}^- \oplus Z_{j,i}^-$	$E_{j^*,i} - E_{i^*,j}$

As before, the claim follows from Lemma 4.4. □

Proposition 6.10. *Let o_l be the cardinality of Oolp_l . Then the sequence $\{o_l\}$ is determined by*

- $o_0 = 1$,
- $o_1 = 1$,
- $o_l = o_{l-1} + 4(l-1)o_{l-2}$.

Proof. We divide the set Oolp_l into the subset of oolps where vertex 1 is not touched by any arrow and its complement. □

The sequence 1, 1, 5, 13, 73, 281, 1741, ... which gives $\{o_l\}$ is classified in OEIS as A115329 [15].

Remark 6.11. As before, we see that oolps are special oriented link patterns. As in the symplectic case, this fact also follows from [12]: if two orthogonal elements are conjugate under the Borel of GL_n , then they are conjugate under the Borel of O_n , as well.

Again, we end the subsection by writing down the explicit classification of B -orbits in $\mathcal{N}(2)$ for the orthogonal group $\text{O}(4)$.

Example 6.12. Let $B \subset O(4)$, then the collection of oolps of size 2 and the corresponding matrices in $\mathcal{N}(2) \subset \mathfrak{o}_4$ are given in the following table:

$\overset{1}{\curvearrowright} \overset{2}{\curvearrowright}$	$\overset{1}{\curvearrowleft} \overset{2}{\curvearrowleft}$	$\overset{1}{\curvearrowright} \overset{2}{\curvearrowleft}$	$\overset{1}{\curvearrowleft} \overset{2}{\curvearrowright}$	$\overset{1}{\curvearrowright} \overset{2}{\curvearrowright}$
$\begin{pmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$

7 Generalization to parabolic actions

Since there are only finitely many Borel orbits in $\mathcal{N}(2)$, we know that every parabolic P acts finitely on the variety $\mathcal{N}(2)$, too. We aim to find explicit parametrizations for these parabolic orbits in case P is a standard parabolic subgroup of G .

For the rest of the article, we consider G to be either the symplectic group SP_{2l} in type C or the *special* orthogonal group SO_{2l} (type D) or SO_{2l+1} (type B).

7.1 Parabolic subgroups

We have seen before that the standard Borel subgroups of G equal the intersection of G with the standard Borel subgroup of GL_n . For standard parabolic subgroups, this is only true for types B and C; in type D there are standard parabolic subgroups $P \subset SO_n$ which are not given as the intersection of a standard parabolic subgroup of GL_n with SO_n .

Following Malle-Testermann in [13, Chapter 12], the parabolic subgroups of a classical group G are in bijection to the so-called totally isotropic flags. These are flags

$$V_1 \subset \dots \subset V_k,$$

such that every two elements $v, w \in V_i$ vanish according to the bilinear form of G , namely $(v, w) = 0$. Given such isotropic flag F , the parabolic P equals the stabilizer of F in G which we denote by $\text{stab}_G(F)$.

Let $F = V_1 \subset \dots \subset V_k$ be a totally isotropic flag, such that $P = \text{stab}_G(F)$ includes the standard Borel B . Then we define $d_i := \dim_K V_i$, the dimension vector $d_F = d_P = (d_1, \dots, d_k, n, d_k, \dots, d_1)$ and define the corresponding $\mathcal{A}(k)$ -representation $M_F = M_P$ of dimension vector d_F which represents the flag in a symmetric manner as usually.

Example 7.1. Let $n = 6$ and look at the totally isotropic flag

$$F_1 := (V_1 = \langle e_1 \rangle \subset V_2 = \langle e_1, e_2, e_3 \rangle)$$

Then F_1 corresponds to the $\mathcal{A}(2)$ -representation M_{F_1} of dimension vector $(1, 3, 6, 3, 1)$

with coefficient quiver

$$\begin{array}{ccccccc}
 \bullet & \xrightarrow{1} & \bullet & \xrightarrow{1} & \bullet & & \\
 & & \bullet & \xrightarrow{1} & \bullet & & \\
 & & \bullet & \xrightarrow{1} & \bullet & & \\
 & & & & \bullet & \xrightarrow{-1} & \bullet \\
 & & & & \bullet & \xrightarrow{-1} & \bullet \\
 & & & & \bullet & \xrightarrow{-1} & \bullet & \xrightarrow{-1} & \bullet
 \end{array} \tag{7.1}$$

Its stabilizer in G is given as the upper-block standard parabolic subgroup of GL_6 of block sizes $(1, 2, 2, 1)$, intersected with G .

Let now $n = 4$ and set

$$F_2 := (V_1 = \langle e_1, e_3 \rangle)$$

Then F_2 corresponds to the $\mathcal{A}(1)$ -representation M_{F_2} of dimension vector $(2, 4, 2)$ with coefficient quiver

$$\begin{array}{cccc}
 \bullet & \xrightarrow{1} & \bullet & \\
 & & \bullet & \xrightarrow{-1} & \bullet \\
 \bullet & \xrightarrow{1} & \bullet & \\
 & & \bullet & \xrightarrow{-1} & \bullet
 \end{array} \tag{7.2}$$

Its stabilizer in $\mathrm{SO}(4)$ is not given as the intersection of $\mathrm{SO}(4)$ with an upper-block standard parabolic subgroup of GL_4 . In fact,

$$\mathrm{Lie} \, \mathrm{stab}_{\mathrm{SO}(4)}(F_2) \cong \left\{ \left(\begin{array}{cccc} a & c & d & 0 \\ 0 & b & 0 & -d \\ e & 0 & -b & -c \\ 0 & -e & 0 & -a \end{array} \right) \mid a, b, c, d, e \right\} \supset \mathfrak{b}.$$

Let us fix a standard parabolic P now, together with the corresponding totally isotropic flag F_P and the representation $M_P := M_{F_P}$ of dimension vector \mathbf{d}_P . Our notions from Section 4 generalize to this setup by looking at the variety $\mathrm{SR}_{\mathbf{d}_P}(\mathcal{A}(k))$ which is acted upon by the group

$$\mathrm{GL}_{\mathrm{sym}}(P) := \mathrm{GL}(d_1) \times \mathrm{GL}(d_2) \times \cdots \times \mathrm{GL}(d_k) \times \mathrm{Sym}(n)$$

where $\mathrm{Sym}(n)$ denotes either the symplectic or the orthogonal group on a vector space of dimension n , as before. Again, the open subset $\mathrm{SR}_{\mathbf{d}_P}(\mathcal{A}(k))^0$ corresponds to the full subcategory $\mathrm{srep}(\mathcal{A}(k), \mathbf{d}_P)^0$ of $\mathrm{srep}(\mathcal{A}(k), \mathbf{d}_P)$ of maximal rank representations.

We define the block vector $\mathbf{b}_P = (d_1, d_2 - d_1, \dots, d_k - d_{k-1})$. In case of a standard parabolic which is induced by GL_n , this vector gives the explicit structure of the parabolic as can be seen in Example 7.1.

We obtain a translation as in Lemma 4.4 which can be proven in the same way.

Lemma 7.2. *There is a bijection between the isomorphism classes of symplectic/orthogonal $\mathcal{A}(k)$ -representations in $\mathrm{srep}(\mathcal{A}(k), \mathbf{d}_P)^0$ and symplectic/orthogonal P -orbits in $\mathcal{N}(2)$.*

7.2 Orbits in type C

We aim to classify the P -orbits in $\mathcal{N}(2) \subset \mathfrak{sp}_n$ by combinatorial objects and generalize the results of Subsection 6.1 in a straight forward manner.

Definition 7.3. An enhanced symplectic oriented link pattern (*esolp. for short*) of size k of type (b_1, \dots, b_k) consists of a set of k vertices $1, 2, \dots, k$ together with a collection of oriented arrows between different vertices, unoriented loops and oriented arrows with dots between arbitrary vertices.

Denote by x_i the number of sources of arrows, by y_i the number of targets of arrows, by z_i the number of unoriented loops and by l_i the number of loops with dots at vertex i . Then the following condition defines an esolp of such kind:

(Sp) $x_i + y_i + z_i - l_i \leq b_i$ holds for each vertex i .


Theorem 7.4. There is a natural bijection between the set of P -orbits in $\mathcal{N}(2) \subseteq \mathfrak{sp}_n$ and the set of esolps of size k and of type (b_1, \dots, b_k) .

Proof. In an analogous manner to Theorem 6.3, by Krull–Remak–Schmidt, there is a bijection between the set of esolps of size k of type (b_1, \dots, b_k) and the set of isomorphism classes of symplectic representations of $\mathcal{A}(k)$ in $\text{srep}(\mathcal{A}(k), \mathbf{d}_P)^0$ which maps an isomorphism class $[M]$ of a symplectic representation M to the subquiver of the coefficient quiver of M induced by M_α and then restricts the latter as follows: all vertices $i \in \{1, 2, \dots, l\}$ which correspond to the same step in the flag F , say to step $V_s \setminus V_{s-1}$ are glued together to vertex s of the esolp. This way, we obtain $2k$ vertices. We are again able to shrink them to k vertices as the table in Figure 7.2 shows - via this correspondence, we obtain a bijection between symplectic direct sum decompositions and esolps.

By Lemma 4.4, the claim follows. \square

Clearly, solps are special esolps: they are of size l and of type $(1, \dots, 1)$, such that we obtain the classification of Borel-orbits.

Example 7.5. Let P be the symplectic parabolic subgroup of block sizes $(4, 2)$, thus, $b_1 = 4$ and $b_2 = 2$. Then a symplectic representation of dimension vector $(4, 6, 12, 6, 4)$ is represented by a pattern of 12 coloured vertices which represents the map M_α of the representation, for example by



This pattern corresponds to the indecomposable direct sum decomposition

$$(D_{1,1}^+ \oplus C_{1,1}^+) \oplus Z_{1,1}^+ \oplus (Z_{1,2}^- \oplus Z_{1,2}^+) \oplus (M_{2,3} \oplus M_{2,3}^*)$$









Indecomposable	Multiplicity
$M_{s,k+1} \oplus M_{s,k+1}^*$	$b_s - x_s - y_s + l_s$ (symplectic) $b_s - x_s - y_s$ (orthogonal)
$D_{i,j}^- \oplus C_{i,j}^-$	number of arrows 
$D_{s,s}^+ \oplus C_{s,s}^+ \cong D_{s,s}^- \oplus C_{s,s}^-$	number of loops 
$D_{i,j}^+ \oplus C_{i,j}^+$	number of arrows 
$Z_{i,j}^- \oplus Z_{j,i}^-$	number of arrows 
$Z_{ij}^+ \oplus Z_{ji}^+$	number of arrows 
$Z_{s,s}^-$ (symplectic) $Z_{s,s}^- \oplus Z_{s,s}^-$ (orthogonal)	number of loops 
$Z_{s,s}^+$ (symplectic) $Z_{s,s}^+ \oplus Z_{s,s}^+$ (orthogonal)	number of loops 

Figure 4: Correspondence of indecomposables and esolps for $i < j < k+1$ and $s < k+1$

of a representation of the quiver Q_2 . The corresponding esolp is given by



We see that $b_1 - x_1 - y_1 + l_1 = 4 - 4 - 1 + 1 = 0$ and $b_2 - x_2 - y_2 + l_2 = 2 - 0 - 1 + 0 = 1$; these equations show that the multiplicity of the indecomposable $M_{1,3} \oplus M_{1,3}^*$ is zero and the multiplicity of the indecomposable $M_{2,3} \oplus M_{2,3}^*$ is one.

7.3 Orbits in type B and D

Let us consider a standard parabolic subgroup of SO_{2l+1} or SO_{2l} together with its totally isotropic flag F now. Then the classification of orbits is done analogously to the symplectic case, but in type D we have to be careful, since not every parabolic is obtained from a standard flag.

Definition 7.6. An enhanced orthogonal oriented link pattern (esolp. for short) of size k of type (b_1, \dots, b_k) consists of a set of k vertices $1, 2, \dots, k$ together with a collection of oriented arrows and oriented arrows with dots between these vertices.

Denote by x_i the number of sources of arrows, by y_i the number of targets of arrows and by z_i the number of unoriented loops at vertex i . Then the following condition defines an eoolp of such kind:

(SO) $x_i + y_i + 2z_i \leq b_i$ holds for each vertex i .

An esolp is an eoolp, if $b_i = 1$ implies that there are no loops at vertex i .

The classification of parabolic orbits follows similarly to the considerations in 7.4.

Theorem 7.7. There is a natural bijection between the set of P -orbits in $\mathcal{N}(2) \subseteq \mathfrak{g}$ and the set of eoolps of size k and of type (b_1, \dots, b_k) .

Proof. In type B, every standard parabolic subgroup is given as the stabilizer of a totally isotropic **standard** flag. Given a P -orbit $P.x$ in $\mathcal{N}(2)$ we look at the representation M_x via Lemma 4.4. Consider the coefficient quiver of M_x and restrict it to those n vertices which represent the loop M_α . Then the eoolp is obtained by identifying vertices according to the flag and then restricting to the vertices $1, \dots, k$ via the correspondence depicted in Figure 7.2 - similarly as in the symplectic case. Note that the vertex $k + 1$ is redundant, since it always represents the indecomposable $M_{\omega, \omega}$ (as in the proof of Theorem 6.9).

In type D, there are two cases to consider.

- (i) The flag F is a standard flag, such that P is the intersection of a standard parabolic of GL_n with SO_n .

In this case, the classification of the orbits is done similarly as in Theorem 7.4 with the correspondence table given in Figure 7.2.

- (ii) For every missing standard parabolic, the flag F is **not** a standard flag and P does not equal the intersection of a standard parabolic of GL_n with SO_n .

In this case, the classification is still done in the same mannner, since we deal with the Krull-Remak-Schmidt decomposition into unique indecomposables. Thus, Figure 7.2 is still a valid correspondence for the bijection between orbits and eolps and the combinatorial description via eoolps is just only dependent on the block vector of the flag (b_1, \dots, b_k) . \square

Clearly, oolps are special eoolps, they are of size l and of type $(1, \dots, 1)$, such that we obtain the classification of Borel-orbits.

Even though the classification of orbits for parabolics of non-standard totally isotropic flags in type D is done in a similar manner as in the other cases, these parabolics are more difficult to handle - especially if one is interested in partiular representative matrices of the P -orbits in $\mathcal{N}(2)$ - and we give them some more attention by calculating an example.

Example 7.8. *Let us consider $n = 6$ and the totally isotropic flag*

$$F = (V_1 = \langle e_1 \rangle \subset V_2 = \langle e_1, e_2, e_4 \rangle).$$

Then the coefficient quiver of M_F is

$$M_F = \begin{array}{ccccccc} \bullet & \xrightarrow{1} & \bullet & \xrightarrow{1} & \bullet & & \\ & & \bullet & \xrightarrow{1} & \bullet & & \\ & & & & \bullet & \xrightarrow{-1} & \bullet \\ & & \bullet & \xrightarrow{1} & \bullet & & \\ & & & & \bullet & \xrightarrow{-1} & \bullet \\ & & & & \bullet & \xrightarrow{-1} & \bullet \xrightarrow{-1} & \bullet \end{array} \quad (7.3)$$

and the isomorphism classes of indecomposables are listed in the table in Figure 5.

8 Restriction to the nilradical

If we restrict a parabolic action on $\mathcal{N}(2)$ to the nilradical $\mathfrak{n}(2)$ of 2-nilpotent upper-triangular matrices in the given Lie algebra, then we still have a parabolic action. The parametrization of the orbits can be obtained from our parametrizations of Sections 6 and 6 straight away. The action of the Borel subgroup in the symplectic case is parametrized in detail in [2], where Barnea and Melnikov also derive a description of the orbit closures and look at applications to orbital varieties in detail. In [8], Gandini, Maffei, Möseneder Frajria and Papi consider the more general approach of B -stable

Representation/Matrix	Representation/Matrix	Eoolp	Representation/Matrix	Esolp
 $E_{2,1} - E_{6,5}$	 $E_{4,1} - E_{6,3}$	 1 2	 0	1 2
 $E_{1,2} - E_{5,6}$	 $E_{1,4} - E_{3,6}$	 1 2	 $E_{3,2} - E_{5,4}$	1 2
 $E_{3,1} - E_{6,4}$	 $E_{5,1} - E_{6,2}$	 1 2	 $E_{2,3} - E_{4,5}$	1 2
 $E_{1,3} - E_{4,6}$	 $E_{1,5} - E_{2,6}$	 1 2		
 $E_{1,3} - E_{4,6}$	 $E_{1,5} - E_{2,6}$	 1 2		

Figure 5: Isomorphism classes of orthogonal representations in $\text{rep}(\mathcal{A}(2), \mathbf{d}_p)^0$ and their combinatorial interpretation

abelian subalgebras of the nilradical of \mathfrak{b} in which they parametrize the B -orbits and describe their closure relations.

Definition 8.1. A symplectic link pattern (*slp for short*) of size k is a symplectic oriented link pattern, such that every arrow is oriented from right to left and every loop is oriented counterclockwise. In the same way, the natural notion of orthogonal link pattern (*orlp for short*) enhanced symplectic and enhanced orthogonal link pattern is obtained.


The sets of (enhanced) symplectic and (enhanced) orthogonal link patterns are obtained by taking all such oriented patterns and deleting the orientation.

Corollary 8.2. There is a bijection between the parabolic orbits in the symplectic nilradical $\mathfrak{n}(2)$ (or orthogonal $\mathfrak{n}(2)$, resp.) and the set of enhanced symplectic (or orthogonal, resp.) link patterns.

Proof. The fact that all arrows are oriented from right to left corresponds equivalently to the nilpotent map at the loop being upper-triangular. Thus, these are directly the nilpotent elements contained in the nilradical of the particular Lie algebra. \square

We end the section with giving an example.

Example 8.3. For $l = 2$, we consider the Borel-action. Figures 2 and 3 show the possible patterns. Taking away the orientation is equivalent to only considering upper-triangular matrices. Thus, the following table gives a complete list of representatives of orbits for type C. The patterns which are marked green give a complete list of orthogonal B-orbits.



References

- [1] Ibrahim Assem, Daniel Simson, and Andrzej Skowroński. *Elements of the representation theory of associative algebras. Vol. 1*, volume 65 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 2006. Techniques of representation theory.
- [2] Nurit Barnea and Anna Melnikov. B-orbits of square zero in nilradical of the symplectic algebra. *Transform. Groups*, 22(4):885–910, 2017.
- [3] Magdalena Boos. Finite parabolic conjugation on varieties of nilpotent matrices. *Algebr. Represent. Theory*, 17(6):1657–1682, 2014.
- [4] Magdalena Boos and Markus Reineke. B-orbits of 2-nilpotent matrices and generalizations. In *Highlights in Lie Algebraic Methods (Progress in Mathematics)*, pages 147–166. Birkhäuser, Boston, 2011.
- [5] Michel Brion. Spherical varieties: an introduction. In *Topological methods in algebraic transformation groups (New Brunswick, NJ, 1988)*, volume 80 of *Progr. Math.*, pages 11–26. Birkhäuser Boston, Boston, MA, 1989.
- [6] M. C. R. Butler and Claus Michael Ringel. Auslander-Reiten sequences with few middle terms and applications to string algebras. *Comm. Algebra*, 15(1-2):145–179, 1987.
- [7] Harm Derksen and Jerzy Weyman. Generalized quivers associated to reductive groups. *Colloq. Math.*, 94(2):151–173, 2002.
- [8] Jacopo Gandini, Andrea Maffei, Pierluigi Möseneder Frajria, and Paolo Papi. The Bruhat order on abelian ideals of Borel subalgebras. 2018.

- [9] Lutz Hille and Gerhard Röhrle. A classification of parabolic subgroups of classical groups with a finite number of orbits on the unipotent radical. *Transform. Groups*, 4(1):35–52, 1999.
- [10] M. E. Camille Jordan. Sur la résolution des équations différentielles linéaires. *Oeuvres*, 4:313–317, 1871.
- [11] Hanspeter Kraft. *Geometrische Methoden in der Invariantentheorie*. Aspects of Mathematics, D1. Friedr. Vieweg & Sohn, Braunschweig, 1984.
- [12] Peter Magyar, Jerzy Weyman, and Andrei Zelevinsky. Symplectic multiple flag varieties of finite type. *J. Algebra*, 230(1):245–265, 2000.
- [13] Gunter Malle and Donna Testerman. *Linear algebraic groups and finite groups of Lie type*, volume 133 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2011.
- [14] Anna Melnikov. B -orbits in solutions to the equation $X^2 = 0$ in triangular matrices. *J. Algebra*, 223(1):101–108, 2000.
- [15] The On-Line Encyclopedia of Integer Sequences. Sequence a115329. published electronically at <http://oeis.org>, 2010.
- [16] The On-Line Encyclopedia of Integer Sequences. Sequence a202837. published electronically at <http://oeis.org>, 2010.
- [17] Dmitri I. Panyushev. Some amazing properties of spherical nilpotent orbits. *Math. Z.*, 245(3):557–580, 2003.
- [18] J.-P. Serre. Espaces fibrés algébriques. *Séminaire Claude Chevalley*, 3:1–37, 1958.