



Article

Feline *Leishmania* spp. Infection in a Non-Endemic Area of Northern Italy

Eva Spada ^{1,*}, Roberta Perego ^{1,*}, Fabrizio Vitale ², Federica Bruno ², Germano Castelli ², Gaia Tarantola ¹, Luciana Baggiani ¹, Sonia Magistrelli ³ and Daniela Proverbio ¹

¹ Dipartimento di Medicina Veterinaria (DIMEVET), Università degli Studi di Milan, via dell'Università 6, 26900 Lodi, Italy; gaiatarantola89@gmail.com (G.T.); luciana.baggiani@unimi.it (L.B.); daniela.proverbio@unimi.it (D.P.)

² Centro di Referenza Nazionale per le Leishmaniosi (C.Re.Na.L), Istituto Zooprofilattico Sperimentale (IZS) della Sicilia A. Mirri, Via G. Marinuzzi 3, 90129 Palermo, Italy; fabrizio.vitale@izssicilia.it (F.V.); federicabruno4486@gmail.com (F.B.); germanocastelli@gmail.com (G.C.)

³ Agenzia di Tutela della Salute di Milano (Dipartimento Veterinario), Corso Italia 19, 20122 Milan, Italy; SMagistrelli@ats-milano.it

* Correspondence: eva.spada@unimi.it (E.S); roberta.perego@unimi.it (R.P)

Received: 31 March 2020; Accepted: 05 May 2020; Published: 8 May 2020

Simple Summary: Leishmaniosis caused by *Leishmania infantum* is a parasitic disease of people and animals transmitted by sand fly vectors. Although dogs in south and central Italy are most affected, in recent decades feline leishmaniosis (FeL) has become an emerging disease. This study aimed to update information on the presence of FeL in stray cats in Milan, in northern Italy; to compare these results with previous studies performed in the same area; and to report aspects of *Leishmania* infection in cats. A total of 117 stray cats were included and 10 (8.6%) had *L. infantum* infection. The parasite was identified in popliteal lymph nodes in five cats and five had antibodies against *L. infantum*. Most infected cats were from a specific area of Milan. Increased gammaglobulins were seen in infected cats, which also had antibodies against the pathogens *Anaplasma phagocytophilum*, *Chlamydomphila felis*, and *Toxoplasma gondii*. There was a high prevalence of FeL in the surveyed area of northern Italy. Further studies are needed to understand if these infected cats are being imported from other areas, or if there is a real *Leishmania* focus in Milan. Investigation of the presence of sand fly vectors in Milan would be useful.

Abstract: Feline leishmaniosis (FeL) is an emerging vector-borne feline disease, with increasing numbers of cases reported and studies performed internationally. This study aimed to update the epidemiological status for FeL in stray cats in Milan, northern Italy; compare these results with previous studies in Northern Italy; and report clinicopathologic findings and coinfections in cats infected with *Leishmania* spp. A total of 117 cats were tested for *L. infantum* and retrovirus infection, hematological, and biochemical parameters. Demographic and clinical data were collected and FeL affected cats screened for selected coinfections. Overall, 10/117 (8.6%) cats tested positive for *L. infantum*: in five cats *L. infantum* DNA was found in popliteal lymph nodes and five were IFAT seropositive at titers from 1:80 to 1:160. Infected cats were concentrated in a specific area of Milan ($p = 0.0154$). No specific clinicopathologic abnormalities or retroviral infections were significantly linked to the infection, other than hypergammaglobulinemia ($p = 0.0127$). Seroreactivity to *Anaplasma phagocytophilum*, *Chlamydomphila felis*, and *Toxoplasma gondii* was found in some infected cats. A high prevalence of FeL was found in a non-endemic area of northern Italy and future studies should continually monitor this data to understand whether these cases are imported or if *Leishmania* vectors are present in this area.

Keywords: epidemiology; feline; indirect immunofluorescence test; infection; leishmaniosis; Northern Italy; polymerase chain reaction; prevalence; stray cats

1. Introduction

Feline leishmaniosis (FeL) is an emerging feline disease, with an overall estimated prevalence of 10%. Cases are being more frequently reported and numerous international research studies have been published in recent years [1,2]. Cats are naturally infected by, and are susceptible to the same *Leishmania* species that affect dogs and people, but most cases involve *Leishmania infantum* as in the countries in the Mediterranean basin [3].

A high seroprevalence rate (12.2%) of FeL is found in Southern European countries including Italy, Spain, Portugal, and Greece. However in Italy, both the overall seroprevalence (24%) and polymerase chain reaction (PCR) prevalence (21%) were found to be higher than in other countries [2].

In endemic regions such as Mediterranean countries, the subclinical form of feline *L. infantum* is common, whereas clinical illness is relatively uncommon [1,2,4,5]. Sandflies are the natural vectors of *Leishmania* spp. and may also feed on the blood of cats. Infected cats may therefore be urban reservoirs of *Leishmania* spp. and transmit the protozoan to other sandflies [6]. In addition, cats can be considered sentinel reservoir hosts at least in endemic areas of zoonotic visceral leishmaniasis [2]. They can therefore play a role as additional reservoir hosts of *L. infantum* and, in a “One Health” perspective, preventative measures should be taken in this species based on epidemiological data.

Over recent decades, many studies have confirmed that feline *Leishmania* infection is relatively common in areas where canine leishmaniosis (CanL) is endemic. The Lombardy region (except for a limited focus) is not considered endemic for *L. infantum* infection at the time of writing. However, previous studies found a seroprevalence up to 12.2% in stray cats from Northern Italy [7,8].

The aims of this study were (I) to update the epidemiological status for *L. infantum* infection in stray cats from Milan, Northern Italy using indirect fluorescent antibody test (IFAT) on serum and real-time polymerase chain reaction (RT-PCR) assays applied to a variety of biological samples; (II) to compare results with recent studies performed in owned and stray cats from the same, and from different, areas of Northern Italy; and (III) to report clinicopathological findings and coinfections of cats infected with *L. infantum* and to compare demographic, physical, and clinicopathologic findings between infected and uninfected cats to find significant associations with infectious status.

2. Materials and Methods

2.1. Study Population and Sample Collection

Stray cats in Milan city (45°28′38″28 N, 09°10′53″40 E), the capital of the Lombardy region in Northern Italy, involved in a trap–neutered and release (TNR) program were prospectively randomly sampled with no limitation for age, gender, and clinical status between June 2016 and December 2018. The TNR program was part of a national program to control stray pet populations under Italian National Law (law no. 281/1991). The project underwent ethical review and was given approval number OPBA_108_2018 by the University of Milan Animal Welfare Bioethical Committee.

Data on signalment including: breed, gender, age (estimated based on dentition, animals < 6 months of age were considered juvenile, whereas all others were considered adult) and colony of origin based on the nine municipalities of Milan city (zone 1–9) were collected for each cat.

During general anesthesia for neutering surgery the following clinical data were collected: rectal temperature; body condition score (BCS) evaluated by palpation of bone prominences and visual assessment of the animal’s silhouette with a score of 5/9 indicating normal BCS, 4–1/9 underweight, and 6–9/9 overweight [9]; data on clinical status based on physical and dermatological examination; cats were classified as either apparently healthy or unhealthy, when one or more of the following clinical abnormalities were present: fever, pale, or icteric mucous membranes, lymphadenopathy, cachexia, dehydration, gingivitis and/or stomatitis, emaciation, hepatomegaly, splenomegaly, dermal lesions, ocular lesions, signs of respiratory infections, and/or any other clinical findings indicating general illness.

The follow biological samples were collected: a 2.5 mL total volume blood sample collected by jugular or cephalic venipuncture into tubes with ethylenediaminetetraacetic acid (EDTA) anticoagulant (1 mL) and in plain collection tubes (1.5 mL) to obtain serum after centrifugation; fine needle aspirate from one popliteal lymph node; conjunctival swab from both eyes using sterile cotton swabs manufactured for bacteriological isolation, rubbed against the surface of the lower eyelid; dermal swabs from dermal lesions; and ectoparasites (ticks or fleas) when present.

2.2. Laboratory Analyses

EDTA whole blood samples were examined within 12 hours of collection for complete blood count (CBC). The remaining EDTA whole-blood samples, and oculoconjunctival and dermal swabs, and lymph node aspirates and ectoparasites were stored frozen at $-20\text{ }^{\circ}\text{C}$ and sent to the Experimental Zooprophyllactic Institute (IZS) of Sicily, National Reference Centre for Leishmaniosis (C.Re.Na.L.), where RT-PCR analysis was performed to determine the presence of *Leishmania spp.* DNA.

After centrifugation of blood collected in plain tubes one aliquot of serum sample was used to perform serology for retroviral infections and for selected coinfections, for measurements of serum protein content and for protein electrophoresis. An aliquot of each serum sample was stored frozen at $-20\text{ }^{\circ}\text{C}$ until sent to the IZS of Sicily where anti-*L. infantum* antibody titer was determined by immunofluorescence antibody test (IFAT).

2.2.1. Hematological Parameters

The following hematological parameters were assessed using an automated multiparameter hematology analyzer with software for animal samples (Cell-Dyn 3500 analyzer, Abbott, Rome, Italy): red blood cell (RBC) count, hemoglobin (Hb), hematocrit (HCT), mean cell volume (MCV), mean cell Hb (MCH), mean cell Hb concentration (MCHC), RBC distribution width (RDW), platelet (PLT) count, and white blood cell (WBC) count. Air-dried Wright-Giemsa-stained blood films were prepared (May-Grünwald-Giemsa MGG Quick Stain, Bio-Optica Milano Spa, Milan, Italy) and evaluated microscopically at oil immersion $\times 100$ magnification (Motic® BA 300, Milan, Italy) for the leukocyte differentiation (neutrophils, lymphocytes, eosinophils, basophils, and monocytes count) and the platelet count evaluation.

2.2.2. Serum Protein and Protein Electrophoresis

Serum total protein concentration was measured by spectrophotometry using the colorimetric biuret method (Hagen Diagnostica S.R.L., Florence, Italy) on a Cobas Mira Classics Roche automated chemistry analyzer (Roche S.p.A., Mannheim, Germany). Albumin and protein fractions were analyzed with Hydragel Kit $\beta 1$ - $\beta 2$ using a semiautomatic agarose gel electrophoresis system (Hydrasys, Sebia PN 1210, Issy-les-Moulineaux, France). Using the computer software Phoresis (Sebia PN 1210, Issy-les-Moulineaux, France) for Windows 2000 or XP Pro the electrophoretic curve for each sample was displayed. Protein fractions were determined based on the percentage of optical absorbance and the absolute concentration/dL was automatically calculated from the total serum protein concentration. Albumin to globulin (A/G) ratios were also calculated.

2.2.3. Detection of *L. infantum* by IFAT

Specific antibodies to *L. infantum* (WHO strain: MHOM/IT/80/IPT1) were detected using the IFAT against in-house cultured promastigotes following Office International des Epizooties (OIE) Terrestrial Manual protocol [10]. *Leishmania* strain was used as an antigen fixed on multispot microscope slides (Bio-Merieux, Marcy L'Etoile, France) in an acetone bath. The feline sera were prepared by serial 2-fold dilutions (1:40 to 1:5120) in phosphate buffered saline (PBS), pH 7.2, and added to the antigen-coated wells. The slides were incubated for 30 min at $37\text{ }^{\circ}\text{C}$. Positive and negative controls were included in each series of analyzed samples. The positive control consisted of a known titer serum of a cat with a positive cultural isolation. The negative control consisted of

serum from a cat testing negative to the culture. Fluoresceinated anti-cat immunoglobulin G (IgG) antibody (working anti-feline Anti-Cat IgG (whole molecule)—FITC antibody produced in goat, Sigma Aldrich, Saint Louis, MO, USA) was used (dilution 1:200 in PBS). The cut-off value for positivity was set at 1:80 according to the OIE Terrestrial Manual [10]. The slides were examined using a Leica DM 4000B fluorescence microscope (Leica, Heerbrugg, Switzerland).

2.2.4. Leishmania DNA Extraction and PCR Assays

EDTA whole-blood samples, oculoconjunctival swabs from both eyes, dermal swabs, ectoparasites, and popliteal lymph node aspirates were extracted by “PureLink® Genomic DNA Mini Kit” (Invitrogen, Carlsbad, CA, USA) according to the manufacturer’s instructions. Lymph node aspirates, oculoconjunctival and dermal swabs, and ectoparasites were resuspended in phosphate buffer saline (PBS) solution (200 µL) and homogenized.

The RT-PCR was carried out in a CFX96 Real-time System (Bio-Rad Laboratories s.r.l., Hercules, CA, USA) using TaqMan Master Mix (Applied Biosystems by ThermoFisher, Waltham, MA, USA) and performed as previously described [11]. The target DNA for amplification is a 116-bp fragment in the constant region of the kDNA minicircle. This is one of the kDNA minicircle families currently used to identify the *Leishmania* genus. The primer sequences were: U 5'-GGCGTTCTGCGAAAACCG-3'; D5'-AAAATGGCATTTCGGGCC-3'; while the associated probe was: 5'-TGGGTGCAGAAATCCCGTTCA- 3' 5'FAM and 3' black hole quencher (BHQ) labeled. The thermal cycling conditions comprised of an initial incubation for 2' at 50 °C for uracil-N-glycosylase activity. This step was followed by a 0' denaturation at 95 °C and 45 cycles at 95 °C for 15" and 60 °C for 1' each. Samples were amplified in a single 96-well plate. On each plate, a negative control was included. Each standard, each sample, and the negative control were analyzed in triplicate for each run. Cycle threshold (Ct) value was calculated for each sample by determining the point of the fluorescence value exceeding the threshold limit. A positive control containing genomic *L. infantum* DNA and a negative control without DNA were included. The parasitic DNA load was determined in each examined sample by comparison of the data with a specific standard curve based on the number of *Leishmania* per milliliter of extracted volume. Standard curves were prepared for both the *Leishmania* gene target and the internal positive control (IPC Applied Biosystems, ThermoFisher, Waltham, MA, USA). A stock solution of *L. infantum* DNA was obtained by extraction from 10⁹ promastigotes/mL. Ten fold serial dilution of the DNA stock solution were performed to obtain the six points of the curve spanning from 10⁶ to 10¹ DNA equivalent cells. The standard curve, calculated by independent experiments was linear over at least 6 log ranges of DNA concentration points with an average correlation coefficient of 0.988. The difference for each point of the curve was one log factor [11].

2.2.5. Detection of Coinfections

Serum from cats with *L. infantum* infection was subsequently tested for the IgG antibodies against *Rickettsia conorii*, *Ehrlichia canis*, *Anaplasma phagocytophilum*, *Chlamydomphila felis*, IgG, and immunoglobulin M (IgM) against *Toxoplasma gondii* and for feline retrovirus status.

IgG against *Rickettsia conorii*, *Ehrlichia canis*, and *Anaplasma phagocytophilum* were tested using IFAT canine commercial kits (Fluo *Rickettsia conorii*, Fluo *Ehrlichia canis*, Fluo *Anaplasma ph Dog*, Biopronix Product Line, Agrolabo Spa, Scarmagno, Turin, Italy.) The manufacturer’s protocol was followed for all serological tests, using anti-feline IgG ready to use (antibody concentration 7.5 µg/mL) instead of the canine ones provided by the kit. The samples were classified as positive if fluorescence was observed at a serum dilution of 1:64 or higher [12]. IgG specific to *Chlamydomphila felis* and IgG and IgM to *Toxoplasma gondii* were detected using a commercially available IFAT kit (Fluo *Chlamydomphila felis*, Fluo *Toxoplasma gondii*, Biopronix Product Line, Agrolabo Spa, Scarmagno, Turin, Italy). Titers ≥ 1:32 [13] and ≥ 1:64 [14] were considered seroreactive and, therefore, indicative of *C. felis* and *T. gondii* exposure, respectively. Positive and negative controls were included for each IFAT assay.

In all feline populations the presence of antibodies against p24 and gp40 antigens of feline immunodeficiency virus (FIV) and the presence of p27 antigen of feline leukemia virus infection (FeLV) were checked on serum samples using a commercial rapid enzyme-linked immunosorbent assay (ELISA) kit (SNAP® Combo Plus FeLV Ag/FIV Ab, IDEXX Laboratories, Europe).

2.3. Statistical Analysis

The data was analyzed using standard descriptive statistics and reported as mean \pm standard deviation (SD) or median and range, based on their distribution. Univariate analysis of categorical data using the Fisher's exact test or χ^2 analysis was performed to determine possible associations between *L. infantum* infection and the different variables collected. Differences between clinicopathological parameters in infected and uninfected cats were tested with a Student's *t*-test or Mann–Whitney test according to whether or not the data were normally distributed. Associations were described using a probability (*p*) value < 0.05 as statistically significant. Statistical analysis was performed using a commercially available software program (MedCalc Statistical Software version 19.1.3, MedCalc Software bv, Ostend, Belgium; 2019).

3. Results and Discussion

A total of 117 cats were randomly sampled during the study period. Demographic, physical, clinical data, and retrovirus status of the whole feline population is reported in Table 1. The median age for all cats was 2 years (range 4 months–10 years, SD \pm 2.26 years), rectal temperature was measured in 86/117 (73.5%) cats with a mean of 37.4 °C (range 35.0–39.3, SD \pm 0.83 °C).

For various technical reasons (insufficient volume of samples or no sample collected), not all data were available for all 117 cats. All 117 cats were tested with RT-PCR on conjunctival swabs, 102/117 were tested by IFAT for *L. infantum*, 109/117 cats were RT-PCR tested on whole blood, 115/117 on lymph node aspirate, 13/117 on ectoparasites, and 2/117 on dermal swabs from dermal lesions. For the same reasons only 104/117 whole blood samples were examined for CBC and 72/117 serum samples for total protein and protein electrophoresis.

Table 1. Demographic, physical, clinical data, and retroviral status of whole stray cat population, with both *Leishmania infantum* uninfected and infected cats. Due to insufficient volume of samples, only 95 cat samples were tested for of feline immunodeficiency virus (FIV) and feline leukemia virus infection (FeLV) infections.

Variables	Modalities	Whole Population (<i>n</i> = 117)	<i>Leishmania infantum</i> Infection Status		<i>p</i> Value
			Uninfected (<i>n</i> = 107)	Infected (<i>n</i> = 10)	
Age	Young	30 (25.6%)	25 (23.4%)	5 (50.0%)	0.0667
	Adult	87 (74.4%)	82 (76.6%)	5 (50.0%)	
Gender	Male	47 (40.2%)	40 (37.4%)	7 (70.0%)	0.0869
	Female	70 (59.8%)	67 (62.6%)	3 (30.0%)	
Reproductive status	Neutered	12 (10.3%)	10 (9.3%)	2 (20.0%)	0.2722
	Not-neutered	105 (89.7%)	97 (90.7%)	8 (80.0%)	
Breed	DSH	116 (99.1%)	106 (99.1%)	10 (100.0%)	1.0000
	Mixed Chartreux	1 (0.9%)	1 (0.93%)	0 (0.0%)	
Origin	Zone 1	12 (10.3%)	10 (9.3%)	2 (20.0%)	0.2722
	Zone 2	1 (0.9%)	1 (0.93%)	0 (0.0%)	1.0000
	Zone 3	2 (1.7%)	1 (0.93%)	1 (10.0%)	0.1643
	Zone 4	18 (15.4%)	18 (16.8%)	0 (0.0%)	1.0000
	Zone 5	17 (14.5%)	16 (15.0%)	1 (10.0%)	1.0000
	Zone 6	13 (11.1%)	12 (11.2%)	1 (10.0%)	1.0000
	Zone 7	27 (23.1%)	26 (24.3%)	1 (10.0%)	0.4499
	Zone 8	14 (12.0%)	13 (12.1%)	1 (10.0%)	1.0000

	Zone 9	7 (6.0%)	4 (3.7%)	3 (30.0%)	0.0154
	nr	6 (5.1%)	6 (5.6%)	0 (0.0%)	-
Body condition score	Normal	62 (53.0%)	56 (52.3%)	6 (60.0%)	0.7476
	Underweight	52 (44.4%)	49 (45.8%)	3 (30.0%)	0.5086
	Overweight	3 (2.6%)	2 (1.9%)	1 (10.0%)	0.2369
Health status	Healthy	97 (82.0%)	91 (85.0%)	6 (60.0%)	0.0665
	Unhealthy	20 (17.1%)	16 (15.0%)	4 (40.0%)	
Ectoparasites	Fleas	30 (25.6%)	29 (27.1%)	1 (10.0%)	0.4490
	Ticks	3 (2.6%)	3 (2.8%)	0 (0.0%)	1.0000
	Ear mites	8 (6.8%)	8 (7.5%)	0 (0.0%)	1.0000
Retrovirus status (n = 95)	Seropositive	14 (14.7%)	12/86 (14.0%)	2/9 (22.2%)	0.6167
	Seronegative	81 (85.3%)	74/86 (86.0%)	7/9 (77.8%)	
Retrovirus infection (n = 95)	FIV	6 (6.3%)	6/86 (7.0%)	0/9 (0.0%)	1.0000
	FeLV	6 (6.3%)	5/86 (5.8%)	1/9 (11.1%)	0.4590
	FIV+FeLV	2 (2.1%)	1/86 (1.2%)	1/9 (11.1%)	0.1814

DSH: domestic shorthair cat; nr: not reported; FIV: feline immunodeficiency virus; FeLV: feline leukemia virus. Results in bold indicate statistical significant p value < 0.05 at Fisher's exact test or χ^2 analysis.

Overall, 10 out of 117 stray cats sampled (8.6%) tested positive for *L. infantum* infection. *Leishmania* DNA was found in the popliteal lymph node aspirates of five cats. An additional five cats were IFAT seropositive with a titer of 1:160 (two cats) and 1:80 (three cats; Table 2 and 3). A further eight (7.8%) cats were IFAT seroreactive at a titer of 1:40, which is below the cut off of 1:80 considered positive for *L. infantum* infection in cats [4,10,15].

Table 2. Signalment and results of real-time polymerase chain reaction (RT-PCR) and immunofluorescence antibody test (IFAT) for *L. infantum* in the 10 infected domestic shorthair cat (DSH) stray cats.

n.	Gender	Age (years)	RT-PCR			IFAT Cut Off \geq 1:80
			Whole Blood	Popliteal Lymph Node	Conjunctival Swab	
1	F	2	negative	5 <i>Leishmania</i>/mL	negative	1:40
2	NF	4	negative	20 <i>Leishmania</i>/mL	negative	$<1:40$
3	M	1	negative	13 <i>Leishmania</i>/mL	negative	$<1:40$
4	M	1	negative	5 <i>Leishmania</i>/mL	negative	$<1:40$
5	F	0.6	negative	125 <i>Leishmania</i>/LI	negative	not tested
6	M	8	negative	negative	negative	1:160
7	M	1	negative	negative	negative	1:160
8	M	2	negative	negative	negative	1:80
9	NM	3	negative	negative	negative	1:80
10	M	1.1	negative	negative	negative	1:80

F: female; NF: neutered female; M: male; NM: neutered male; Positive results of RT-PCR and IFAT for *L. infantum* in bold.

Table 3 shows the results of other recent epidemiological studies performed in cats in Northern Italy. The first two studies [8,16] relate to the same population of stray cats and were performed in previous years by the authors of this study. The last two studies [17,18] relate to owned cat populations from cats surveyed in cities located in Northern Italy.

Table 3. Comparison of results of epidemiological studies for *L. infantum* infection of the current study with other recent studies performed in cats in northern Italy.

Variable	Spada et al 2014 [7]	Spada et al 2016 [8]	Current Study	Iatta et al 2019 [17]	Urbani et al 2020 [18]
Population	Stray	Stray	Stray	Owned	Owned
Geographic area	Milan	Milan	Milan	Northern Italy	Bologna

Sampling years	2008-2010	2014	2016-2018	2017-2018	2017	
Samples number	233	90	117	1543	152	
FeL overall prevalence	9.0%	12.2%	8.6%	1.6%	12.5%	
IFAT overall seropositivity (at $\geq 1:80$ cut off)	21/233 (9.0%)	11/90 (12.2%)	5/102 (4.9%)	20/1543 (1.3%)	18/152 (11.8%)	
IFAT titer	1:40	38 (16.3%)	16 (17.8%)	8/102 (7.8%)	0 (0.0%)	27 (17.8%)
	1:80	15 (6.4%)	11 (12.2%)	3/102 (2.9%)	20 (1.3%)	11 (7.2%)
	1:160	6 (2.6%)	0 (0.0%)	2/102 (2.0%)	0 (0.0%)	5 (3.3%)
	1:320	0 (0.0%)	0 (0.0%)	0/102 (0.0%)	0 (0.0%)	2 (1.3%)
PCR overall positivity	0 (0.0%)	2 (2.2%)	5/115 (4.4%)	5/1543 (0.3%)	1/150 (0.7%)	
PCR on	Whole blood	0 (0.0%)	1 (1.1%)	0/109 (0.0%)	5 (0.3%)	0/146 (0.0%)
	Conjunctiva	np	0 (0.0%)	0/117 (0.0%)	np	0/150 (0.0%)
	Lymph node	np	1 (1.1%)	5/115 (4.4%)	np	np
	Ectoparasites	np	np	0/13 (0.0%)	np	np
	Dermal lesion	np	np	0/2 (0.0%)	np	np
	Hair	np	np	np	np	1/150 (0.7%)

FeL: Feline leishmaniosis; IFAT: immunofluorescence antibody test; PCR: polymerase chain reaction; np: not performed.

Canine leishmaniosis has spread northwards in Italy in the last decades [19,20], but data on the epidemiology of CanL in the Lombardy region are scant and out of date, therefore difficult to compare with results of the current study. Cases of CanL are commonly diagnosed in the area where we performed our study, but the history of the affected dogs always reveals that they have visited or lived in areas endemic for CanL [21,22]. A canine serological survey on 313 dogs in a public animal shelter performed more than 15 years ago (2002–2003) in the urban area of Milan found a CanL seroprevalence of 3.4% [23]. Although the history of dogs in animal shelters is often unknown, some of these dogs may have come from areas that are endemic for *L. infantum* infection. However, it is unlikely that all *Leishmania* seropositive cats found in our study population were infected and imported from endemic areas. Therefore the results of this study highlight a high prevalence of FeL for a non-endemic area for *Leishmania* infection when compared to CanL, a prevalence similar to that found in cats in areas of Southern Italy endemic for CanL and human leishmaniosis [17] and similar to colony and shelter cats in central Italy in which recently a seroprevalence of 3.5% (at cut off $\geq 1:80$) was found [24]. The result of the current study is not surprising in the light of previous studies performed by the same authors in colony stray cats from Milan city. Stray cats in Milan have been found to be IFAT seroreactive for *L. infantum* since sample collection starting in 2008 [7]. However the FeL seroprevalence in this feline population has changed substantially since the first two studies were conducted and almost halved with respect to one study [7] and reduced to a third compared to the last study (Table 3) [8].

In general, the most common diagnostic test used in epidemiological studies is IFAT, however identification of *Leishmania* amastigotes in aspirated samples of bone marrow, spleen, and lymph node is specific and considered the gold standard method for diagnosing FeL [2,25]. Therefore, PCR is recommended preferentially over other diagnostic tests, especially when blood samples and other clinical samples contain a low parasitic burden [26]. *Leishmania* spp. DNA was found for the first time in stray cats in Milan in 2014 [8]. The current study found twice the prevalence of cats harboring *Leishmania* DNA with respect to the previous study [8], i.e., five cats in which parasite DNA was identified in the reticuloendothelial system by RT-PCR, confirming that these cats were indeed infected, some with a high parasitic load such as cat n. 5, a young entire female that had 125 *Leishmania*/mL in the popliteal lymph node aspirate.

As in previous studies we analyzed only stray cats, which may act as amplifying hosts and sentinels for vector-borne infections, some of which represent important zoonosis [27]. Stray cats receive no prophylaxis against fleas, ticks, and sand flies, which are potential vector of infections such as rickettsiosis, anaplasmosis, ehrlichiosis, and hematic mycoplasmosis, which have been identified to have a high prevalence in previous studies performed in stray cat populations in Milan [16,28]. Since stray cats are exposed to all vector-borne infections present in this area they can give

information as to the rate of these infections in a specific area. However, one of the problems with working with stray cat populations is the lack of anamnestic data that meant we could not exclude the possibility that the infected cats had been imported from other areas of Italy (or from other countries in southern Europe) where FeL is endemic. As previously described sporadic or rare occurrences of FeL in non-endemic areas can be the consequence of rehoming or moving cats, such as cases seen in Switzerland in cats imported from Spain [29,30].

Three RT-PCR-positive cats in this study had an IFAT titer <1:40, and one a titer of 1:40. Therefore, as previously shown [18], it seems that there was no agreement between the results obtained by IFAT and PCR. This is a common finding in cats from *L. infantum* endemic areas, where discordant results between serological and molecular techniques have been reported previously [7,24,31–34]. As previously hypothesized, the negative results obtained in the IFAT could reflect an ineffective immune system response or could be explained by the absence of antibody production during an early stage of the infection. For this reason, both direct and indirect techniques should be performed in epidemiological studies.

In comparison with other recent studies [17,18] performed in cats living in cities in Northern Italy (shown in Table 3), the rates of infection in cats surveyed varied widely. This may be due to the methodology used (IFAT versus PCR, type of samples analyzed with PCR), the geographic area of Northern Italy studied and the population under study (cat's life style, owned versus stray cats). Both previous studies were performed on owned cats in Northern Italy and had a seroprevalence at IFAT with $\geq 1:80$ cut off titer rates from 1.3% to 11.8% [17,18]. The last study [18] performed by Urbani et al. showed a very high seroprevalence (11.8%), more than double that found in the current study (4.9%), and this could be due to the fact the feline population surveyed comes from a city closer to the *L. infantum* endemic foci of central Italy and that the Emilia-Romagna region has been considered to be an endemic area for CanL [20]. PCR detection in these two studies ranged from 0.3% to 0.7%, which is lower than the results of the current study (4.4%). These studies used different primers in their RT-PCR [35,36] and different sensitivity of primers used could explain these differences. In addition, in feline epidemiological investigations most studies were performed on EDTA-blood and some positive cats were possibly missed because blood is not the most sensitive tissue for the detection of *L. infantum* DNA in dogs or cats [4,37]. The presence of *Leishmania* DNA in lymph node aspirates was the most common site for PCR identification of infection in the current study and detected the highest prevalence of infection for molecular testing when compared to all studies performed in cats from Northern Italy.

For the first time in the current study ectoparasites collected from cats in Northern Italy were analyzed by RT-PCR for presence of *Leishmania* DNA. *L. infantum* DNA was found in ticks and fleas removed from cats living in areas where canine leishmaniosis is endemic [38–40]. This opens the debate about the epidemiological role of ticks in feline leishmaniosis. For this reason we tested the ectoparasites we found in our cat population, but none of the 13 samples (11 fleas and 2 ticks) tested positive for the presence of *Leishmania* DNA at RT-PCR. A limitation is that since no ectoparasite was found on the infected cats we lack information on potential sources of infection in our feline infected population. In addition, only a limited number of ectoparasites were collected and analyzed.

The only statistically significant demographic factor linked to *L. infantum* infected cats was the origin from a colony located in area 9 of Milan city ($p = 0.0154$; Table 1). The fact that significantly more infected cats come from a specific area of Milan could be helpful to better understand whether there is a real endemic *Leishmania* focus in the city. For example, it may be useful to perform an entomological survey in this area to identify whether the classical vector of *Leishmania* spp., i.e., phlebotomine sand flies of genus *Phlebotomus* are present here. Until July 2019 there was no data on the presence of phlebotomine sand flies in the Milan area [41]. Sandflies included *Phlebotomus perniciosus* have previously been collected in the Lombardy region [20,41–43]. However, to the author's knowledge, no sandflies have been identified in the area of Milan from which the cats in this study originated.

Clinical data, results of CBC and protein profile related to the 10 infected cats are reported in Table 4, Table 5, and Table 6, respectively.

Table 4. Clinical data from ten *Leishmania infantum* infected stray cats from Northern Italy.

<i>n.</i>	Rectal Temperature (°C)	Mucous Membrane	Lymphadenomegaly	Respiratory Signs	Gastrointestinal Signs	Dermal Signs	Ocular Signs
1	38.4	n	np	np	np	np	np
2	37.7	n	Solitary	np	np	Bilateral temporal skin wounds	np
3	36.0	n	Solitary	np	Gingivostomatitis	np	np
4	37.4	n	Solitary	np	np	np	np
5	37.3	Pallor	np	np	Gingivostomatitis	np	np
6	37.8	Pallor	Solitary	Respiratory pulmonary rumors	Gingivostomatitis	np	Corneal ulcer and third eyelid lesion
7	nr	n	np	np	np	np	np
8	38.6	Pallor, subicterus	Generalized	np	np	np	np
9	nr	n	Solitary	np	np	Generalised scaling	np
10	37.4	n	np	np	np	np	np

BCS: body condition score; n: normal; np: not present.

Most infected cats showed solitary or generalized lymphadenomegaly and gingivostomatitis was the second most common clinical sign in the infected cats. When clinical signs of FeL are present, they are non-specific and frequently similar to those in other feline diseases and subclinical *L. infantum* feline infection is common [1,2,4,5,25,26,44,45]. The remaining subjects, 3/10 infected cats showed no clinical abnormalities and seemed to be apparently healthy and this corroborates the reports that FeL infected cats can be asymptomatic and that subclinical feline infection *L. infantum* is common [1,2,4,5,25,26,44–47].

Table 5. Complete blood count values in *L. infantum* infected stray cats from Northern Italy.

<i>n.</i>	RBCs (6560–11200 ×10 ³ /μL)	Hb (10.6– 15.6 g/dL)	Hct (31.7– 48.0%)	MCV (36.7–53.7 fl)	MCH (12.3–17.3 pg)	MCHC (30.1–35.6 g/dL)	RDW (16.7– 22.9%)	PLTs (175.0– 500.0 ×10 ³ /μL)	WBCs (4040– 18700/ μL)	Neutrophils (2300– 14000/μL)	Lymphocytes (800–6100/μL)	Monocytes (0–700/μL)	Eosinophils (0–1500/μL)	Basophils (0–100/μL)
1	5990	10.0	28.5	47.5	16.6	35	15.2	470	13600	11696	1088	408	408	0
2	7080	9.3	29.8	42.0	13.1	31.2	17.8	397	19500	14430	2340	390	2340	0
3	8370	11.6	42.2	50.0	13.8	27.5	19.2	321	8600	6708	516	172	1204	0
4	8010	12.1	38.6	48.0	15.1	31.3	17.3	444	12900	9159	2580	129	1032	0
5	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd
6	5470	8.9	29.0	60.0	16.2	27.2	16.2	393	19100	12033	4775	1719	573	0
7	8230	11.4	38.6	47.0	13.9	29.6	17.4	197	25400	20066	2286	1016	1778	254
8	8360	10.2	29.4	35.2	12.3	34.8	18.4	193	8580	5663	2574	0	343	0
9	8290	10.6	32.9	39.7	12.8	32.3	29.7	537	8250	5693	1485	577	495	0
10	8660	15.0	44.0	51.0	17.4	34.3	18.2	249	13800	10350	1656	276	1518	0

RBCs = red blood cells; Hb = hemoglobin; Hct = hematocrit; MCV = mean cell volume; MCH = mean cell hemoglobin; MCHC = mean cell hemoglobin concentration; RDW = red blood cell distribution width; WBCs = white blood cells; PLTs = platelets. In bold value outside the reference ranges; nd: not determined.

Table 6. Serum total protein, albumin and globulin content, and albumin to globulin ratio in *L. infantum* infected stray cats of Northern Italy.

<i>n</i>	Total Protein (6–8 g/dL)	Albumin (3.0–4.6 g/dL)	Alpha 1-globulins (0.1–0.5 g/dL)	Alpha 2-globulins (0.3–1.2 g/dL)	Beta 1-globulins (0.1–0.7 g/dL)	Beta 2-globulins (0.2–0.8 g/dL)	Gamma-globulins (0.3–0.8 g/dL)	A/G Ratio (0.8–1.6)
1	6.97	2.91	0.24	1.14	0.35	0.28	2.06	0.72
2	6.63	2.96	0.16	1.01	0.23	0.4	1.87	0.81
3	5.64	3.19	0.12	1.03	0.28	0.16	0.86	1.3
4	5.59	3.05	0.10	0.34	0.41	0.27	1.41	1.2
5	nd	nd	nd	nd	nd	nd	nd	nd
6	6.16	1.94	0.09	0.75	0.30	0.28	2.8	0.46
7	6.69	3.02	0.19	1.4	0.37	0.22	1.49	0.82
8	11.93	4.40	0.31	0.98	0.73	0.75	4.76	0.58
9	6.27	2.95	0.19	1.19	0.26	0.22	1.47	0.89
10	nd	nd	nd	nd	nd	nd	nd	nd

nd: not determined; A/G: albumin to globulin ratio. In bold values outside the reference ranges.

Mild to severe CBC abnormalities such as normocytic normochromic non-regenerative anemia, moderate to severe pancytopenia or leucocytosis may be seen in *L. infantum* infected cats [1,4,5,30,48,49]. Most infected cats in the current study had CBC parameters outside the reference range for healthy cats, however no significant statistical difference was reported in CBC median value between infected and uninfected cats.

While hyperproteinemia results were not significantly different between infected and uninfected cats in our population, statistically significant differences were found in protein fractions at serum electrophoresis. A statistically significant lower and higher concentration was found in alpha 2-globulins and gamma-globulins in infected compared to uninfected cats, respectively, as shown in Table 7. While median alpha 2-globulins remain in reference range in FeL infected cats, median gamma-globulins increased more in infected than in uninfected cats. Gammopathy are found in most cases described in the literature [1,4,5,18,30,48–52] with some cats showing also a normal proteinemia as in the feline population analyzed in this study. It can be hypothesized that in cats, as in dogs, the humoral immune response could also be activated by the production of immunoglobulins after infection, resulting in high concentrations of gamma-globulins. However polyclonal gammopathy occurs in many feline infectious and inflammatory diseases and is not specific for FeL. In addition, we should consider that both clinical, CBC and serum protein electrophoresis results may have been influenced by concurrent disease as well as retroviral or other non-diagnosed infections, which are common in stray feline populations. This could be supported by the fact that uninfected cats in our population also showed a median gamma-globulins content above the reference range for healthy cats.

Table 7. Comparison of total protein, albumin, globulin content, and albumin to globulin ratio between *L. infantum* infected and uninfected cats.

Parameter	<i>Leishmania infantum</i> Infection Status						<i>p</i> Value ^a
	Infected Cats			Uninfected Cats			
	<i>n</i>	Median	Average Rank	<i>n</i>	Median	Average Rank	
Total protein (6–8 g/dL)	8	6.4	45.9	64	5.9	35.3	0.1761
Albumin (3.0–4.6 g/dL)	8	2.9	49.0	64	2.7	34.9	0.0731
Alpha 1-globulins (0.1–0.5 g/dL)	8	0.1	24.7	64	0.2	37.9	0.0914
Alpha 2-globulins (0.3–1.2 g/dL)	8	1.0	21.6	64	1.2	38.3	0.0337
Beta 1-globulins (0.1–0.7 g/dL)	8	0.3	29.0	64	0.3	37.4	0.2818
Beta 2-globulins (0.2–0.8 g/dL)	8	0.2	38.6	64	0.2	36.2	0.7535
Gamma-globulins (0.3–0.8 g/dL)	8	1.6	53.8	64	1.0	34.3	0.0127
A/G ratio (0.8–1.6)	8	0.8	34.5	64	0.8	36.7	0.7743

A/G: albumin to globulin ratio. In bold where statistical significant p value <0.05 at Mann–Whitney test.

Some *L. infantum* infected cats were also retrovirus coinfecting and seropositive for the presence of antibodies (IgG) to *Anaplasma phagocytophilum* and *Toxoplasma gondii*. All tested cats were seropositive for *Chlamydomphila felis* as shown in IFAT results reported in Table 8. The prevalence of feline retroviruses in stray cats from our study area was slightly higher than rates detected in a previous study, 14.7% of the current study versus 10.4% in [53], and this was due to an increase in FeLV infections. This result was not surprising, as stray cats in Northern Italy are not routinely vaccinated against this retrovirus. A significant association between FIV and *L. infantum* infections has been found in previous studies [17,34,51,54] and also in stray cat populations in Milan investigated in a previous study [7]. FIV was previously reported to be the most frequent concomitant coinfection in FeL infected cats [1,5]. However, this significant association was not seen in the current study. In fact, although associations have been noted, immunosuppression induced by agents such as FIV does not always increase the risk for FeL, as in many studies there was no link between FIV infection and FeL infection [25,29,30,44,47,49,50,52,55–57].

Table 8. Coinfections, retrovirus status and seroreactivity to immunofluorescence antibody test in *L. infantum* infected stray cats in Northern Italy. Due to insufficient sample volume the end point titer for *Chlamydomphila felis* was not determined.

n.	<i>Anaplasma phagocytophilum</i>	<i>Ehrlichia canis</i>	<i>Rickettsia conorii</i>	<i>Toxoplasma gondii</i>		<i>Chlamydomphila felis</i>	FIV	FeLV
	(cut off 1:64)	(cut off 1:64)	(cut off 1:64)	IgG (cut off 1:64)	IgM (cut off 1:64)	(cut off 1:32)	Ab	Ag
1	negative	negative	negative	negative	negative	1:40	negative	negative
2	negative	negative	negative	negative	negative	1:80	negative	negative
3	negative	negative	negative	negative	negative	1:40	negative	negative
4	negative	negative	negative	negative	negative	>1:320	negative	negative
6	negative	negative	negative	1:128	negative	1:40	positive	positive
7	negative	negative	negative	negative	negative	1:40	negative	positive
8	1:64	negative	negative	negative	negative	>1:320	negative	negative
9	negative	negative	negative	negative	negative	>1:320	negative	negative

nd: not determined; IgG: immunoglobulin G; IgM: immunoglobulin M; FIV Ab: feline immunodeficiency virus antibodies; FeLV Ag: feline leukemia virus antigen. In bold positive results.

FeL infection does not exclude the possibility of concurrent diseases or coinfections. In previous studies the association between coronavirus, *Toxoplasma* and some vector-borne coinfections in cats by antibody and/or PCR positive to *L. infantum* has been explored [8,31,34,56,57]. Coinfections or comorbidities are frequently detected in stray cats [8,31]. The presence of one or more concomitant infection may influence the clinical presentation and presence of laboratory abnormalities, can contribute to a misrepresentation of clinical FeL, influence parasite burden or alter the progression of FeL.

4. Conclusions

The results of the present study highlight a stable FeL situation among the stray cats of the Milan city where the cats are exposed to, or infected by *L. infantum*. As stray cats do not receive any prophylactic measure other than neutering surgery and many unhealthy stray cats are hospitalized, FeL should be considered in the differential diagnosis list for cats showing hypergammabulinemia of unknown origin. Future studies on FeL infection in the same area of Northern Italy are needed to clarify whether FeL cases are the result of rehoming or movement of cats from *L. infantum* endemic areas or whether there is a real FeL focus in this region. Continual monitoring and further studies to investigate whether *Leishmania* vectors are present in this area are required.

Author Contributions: Conceptualization, E.S., D.P., R.P.; methodology, L.B., F.V., F.B., G.C., and G.T.; formal analysis, E.S., L.B., F.V., F.B., G.C.; investigation, E.S., L.B., F.V., F.B., G.C.; resources and samples E.S., G.T., S.M.; data curation, E.S., D.P., and R.P.; writing—original draft preparation, E.S.; writing—review and editing, E.S., D.P., R.P., F.V., G.T., F.B., G.C.; project administration, E.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding

Acknowledgments: . The ELISA tests for FIV and FeLV infections were donated by IDEXX. The IFATs kit for *Rickettsia conorii*, *Ehrlichia canis*, and *Anaplasma phagocytophilum* were donated by Agrolabo spa, Scarmagno, Turin, Italy.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Pennisi, M.G.; Persichetti, M.F. Feline leishmaniosis: Is the cat a small dog? *Vet. Parasitol.* **2018**, *251*, 131–137.
- Asfaram, S.; Fakhar, M.; Teshnizi, S.H. Is the cat an important reservoir host for visceral leishmaniasis? A systematic review with meta-analysis. *J. Venom. Anim. Toxins Incl. Trop. Dis.* **2019**, *25*, 1–10.
- Pennisi, M.G.; Hartmann, K.; Lloret, A.; Addie, D.; Belák, S.; Boucraut-Baralon, C.; Egberink, H.; Frymus, T.; Gruffydd-Jones, T.; Hosie, M.J.; et al. Leishmaniosis in cats: ABCD guidelines on prevention and management. *J. Feline Med. Surg.* **2013**, *15*, 638–642.
- Pennisi, M.G.; Cardoso, L.; Baneth, G.; Bourdeau, P.; Koutinas, A.; Miró, G.; Oliva, G.; Solano-Gallego, L. LeishVet update and recommendations on feline leishmaniosis. *Parasites Vectors* **2015**, *8*, 302..
- Fernandez-Gallego, A.; Bernabe, L.F.; Dalmau, A.; Esteban-Saltiveri, D.; Font, A.; Leiva, M.; Ortuñez-Navarro, A.; Peña, M.T.; Tabar, M.D.; Real-Sampietro, L.; et al. Feline leishmaniosis: Diagnosis, treatment and outcome in 16 cats. *J. Feline Med. Surg.* **2020**, online first.
- Maroli, M.; Pennisi, M.; Di Muccio, T.; Houry, C.; Gradoni, L.; Gramiccia, M. Infection of sandflies by a cat naturally infected with *Leishmania infantum*. *Vet. Parasitol.* **2007**, *145*, 357–360.
- Spada, E.; Proverbio, D.; Migliazzo, A.; Della Pepa, A.; Perego, R.; Bagnagatti De Giorgi, G. Serological and Molecular Evaluation of *Leishmania infantum* Infection in Stray Cats in a Nonendemic Area in Northern Italy. *ISRN Parasitol.* **2013**, 916376
- Spada, E.; Canzi, I.; Baggiani, L.; Perego, R.; Vitale, F.; Migliazzo, A.; Proverbio, D. Prevalence of *Leishmania infantum* and co-infections in stray cats in northern Italy. *Comp. Immunol. Microbiol. Infect. Dis.* **2016**, *45*, 53–58.
- Laflamme, D.P. Development and validation of a body condition scoring system for cats: A clinical tool. *Feline Pract.* **1997**, *25*, 13–18.
- Leishmaniosis. Available online: https://www.oie.int/fileadmin/Home/eng/Health_standards/tahm/3.01.11_LEISHMANIOSIS.pdf (accessed on 26 March 2020).
- Vitale, F.; Reale, S.; Vitale, M.; Petrotta, E.; Torina, A.; Caracappa, S. TaqMan-based detection of *Leishmania infantum* DNA using canine samples. *Ann. N. Y. Acad. Sci.* **2004**, *1026*, 139–143.
- Solano-Gallego, L.; Hegarty, B.; Espada, Y.; Llull, J.; Breitschwerdt, E. Serological and molecular evidence of exposure to arthropod-borne organisms in cats from northeastern Spain. *Vet. Microbiol.* **2006**, *118*, 274–277.

13. Di Francesco, A.; Donati, M.; Battelli, G.; Cevenini, R.; Baldelli, R. Seroepidemiological survey for *Chlamydomydia felis* among household and feral cats in northern Italy. *Vet. Rec.* **2004**, *155*, 399–400.
14. Dubey, J.P.; Lappin, M.R. Toxoplasmosis and Neosporidiosis. In *Infectious Diseases of the Dog and Cat*; Greene, C.E., Ed.; Saunders Elsevier: St. Louis, MO, USA, 2006; pp. 755–799.
15. Persichetti, M.F.; Solano-Gallego, L.; Vullo, A.; Masucci, M.; Marty, P.; Delaunay, P.; Vitale, F.; Pennisi, M.G. Diagnostic performance of ELISA, IFAT and Western blot for the detection of anti-*Leishmania infantum* antibodies in cats using a Bayesian analysis without a gold standard. *Parasites Vectors* **2017**, *10*, 119.
16. Spada, E.; Proverbio, D.; Galluzzo, P.; Della Pepa, A.; Bagnagatti De Giorgi, G.; Perego, R.; Ferro, E. Prevalence of haemoplasma infections in stray cats in northern Italy. *ISRN Microbiol.* **2014**, 298352.
17. Iatta, R.; Furlanello, T.; Colella, V.; Tarallo, V.D.; Latrofa, M.S.; Brianti, E.; Trerotoli, P.; Decaro, N.; Lorusso, E.; Schunack, B.; et al. A nationwide survey of *Leishmania infantum* infection in cats and associated risk factors in Italy. *PLoS Negl. Trop. Dis.* **2019**, *13*, 1–11.
18. Urbani, L.; Tirolo, A.; Salvatore, D.; Tumbarello, M.; Segatore, S.; Battilani, M.; Balboni, A.; Dondi, F. Serological, molecular and clinicopathological findings associated with *Leishmania infantum* infection in cats in Northern Italy. *J. Feline Med. Surg.* **2020**, online first.
19. Gramiccia, M.; Scalone, A.; Di Muccio, T.; Orsini, S.; Fiorentino, E.; Gradoni, L. The burden of visceral leishmaniasis in Italy from 1982 to 2012: A retrospective analysis of the multi-annual epidemic that occurred from 1989 to 2009. *Euro Surveill* **2013**, *18*, 20535.
20. Maroli, M.; Rossi, L.; Baldelli, R.; Capelli, G.; Ferroglio, E.; Genchi, C.; Gramiccia, M.; Mortarino, M.; Pietrobelli, M.; Gradoni, L. The northward spread of leishmaniasis in Italy: Evidence from retrospective and ongoing studies on the canine reservoir and phlebotomine vectors. *Trop. Med. Int. Health* **2008**, *13*, 256–264.
21. Perego, R.; Proverbio, D.; Bagnagatti De Giorgi, G.; Spada, E. Prevalence of dermatological presentations of canine leishmaniasis in a nonendemic area: A retrospective study of 100 dogs. *Vet. Med. Int.* **2014**, 374613.
22. Spada, E.; Proverbio, D. Canine leishmaniasis in non-endemic area: Incidence of positivity to IFAT test in dogs moved to endemic area. In Proceedings of the LVI National Congress SISVET, Giardini Naxos (ME), Italy, 26–28 September 2002; pp. 289–290.
23. Perego, R.; Proverbio, D.; Spada, E.; Ferro, E. Canine leishmaniasis: A sero-epidemiological survey by indirect fluorescence antibody test (IFAT) in 313 dogs at sanitary public kennel of Milan. In Proceedings of the LIX Congress SISVET, Viareggio (LU), Italy, 21–24 September 2005; pp. 273–274.
24. Morganti, G.; Veronesi, F.; Stefanetti, V.; Di Muccio, T.; Fiorentino, E.; Diaferia, M.; Santoro, A.; Passamonti, F.; Gramiccia, M. Emerging feline vector-borne pathogens in Italy. *Parasites Vectors* **2019**, *12*, 193.
25. Martín-Sánchez, J.; Acedo, C.; Muñoz-Pérez, M.; Pesson, B.; Marchal, O.; Morillas-Márquez, F. Infection by *Leishmania infantum* in cats: Epidemiological study in Spain. *Vet. Parasitol.* **2007**, *145*, 267–273.
26. Chatzis, M.K.; Andreadou, M.; Leontides, L.; Kasabalis, D.; Mylonakis, M.; Koutinas, A.F.; Rallis, T.; Ikononopoulos, J.; Saridomichelakis, M.N. Cytological and molecular detection of *Leishmania infantum* in different tissues of clinically normal and sick cats. *Vet. Parasitol.* **2014**, *202*, 217–225.
27. Day, M.J. One health: The importance of companion animal vector-borne diseases. *Parasites Vectors* **2011**, *4*, 2–7.
28. Spada, E.; Proverbio, D.; Galluzzo, P.; Della Pepa, A.; Perego, R.; Bagnagatti De Giorgi, G.; Ferro, E. Molecular study on selected vector-borne infections in urban stray colony cats in northern Italy. *J. Feline Med. Surg.* **2014**, *16*, 684–688.
29. Rufenacht, S.; Sager, H.; Müller, N.; Schaerer, V.; Heier, A.; Welle, M.M.; Roosje, P.J. Two cases of feline leishmaniasis in Switzerland. *Vet. Rec.* **2005**, *156*, 542–545.
30. Richter, M.; Schaarschmidt-Kiener, D.; Krudewig, C. Ocular signs, diagnosis and long-term treatment with allopurinol in a cat with leishmaniasis. *Schweiz Arch. Tierheilkd* **2014**, *156*, 289–294.
31. Miró, G.; Rupérez, C.; Checa, R.; Gálvez, R.; Hernández, L.; García, M.; Canorea, I.; Marino, V.; Montoya, A. Current status of *L. infantum* infection in stray cats in the Madrid region (Spain): Implications for the recent outbreak of human leishmaniasis? *Parasites Vectors* **2014**, *7*, 1–7.

32. Silaghi, C.; Knaus, M.; Rapti, D.; Kusi, I.; Shukullari, E.; Hamel, D.; Pfister, K.; Rehbein, S. Survey of *Toxoplasma gondii* and *Neospora caninum*, haemotropic mycoplasmas and other arthropod-borne pathogens in cats from Albania. *Parasites Vectors* **2014**, *7*, 62.
33. Montoya, A.; García, M.; Gálvez, R.; Checa, R.; Marino, V.; Sarquis, J.; Barrera, J.P.; Rupérez, C.; Caballero, L.; Chicharro, C.; et al. Implications of zoonotic and vector-borne parasites to free-roaming cats in central Spain. *Vet. Parasitol.* **2018**, *251*, 125–130.
34. Silva Vicente Sobrinho, L.; Nazaretian Rossi, C.; Peloi Vides, J.; Tozzi Braga, E.; Amélia Domingues Gomes, A.; Marçal Félix de Lima, V.; Helena Venturoli Perri, S.; Generoso, D.; Langoni, H.; Leutenegger, C.; et al. Coinfection of *Leishmania chagasi* with *Toxoplasma gondii*, Feline Immunodeficiency Virus (FIV) and Feline Leukemia Virus (FeLV) in cats from an endemic area of zoonotic visceral leishmaniasis. *Vet. Parasitol.* **2012**, *187*, 302–306.
35. Francino, O.; Altet, L.; Sánchez-Robert, E.; Rodriguez, A.; Solano-Gallego, L.; Alberola, J.; Ferrer, L.; Sánchez, A.; Roura, X. Advantages of real-time PCR assay for diagnosis and monitoring of canine leishmaniasis. *Vet. Parasitol.* **2006**, *137*, 214–221.
36. Ceccarelli, M.; Galluzzi, L.; Migliazzo, A.; Magnani, M. Detection and characterization of *Leishmania* (*Leishmania*) and *Leishmania* (*Viannia*) by SYBR green-based real-time PCR and high resolution melt analysis targeting kinetoplast minicircle DNA. *PLoS ONE* **2014**, *9*, e88845.
37. Lombardo, G.; Pennisi, M.G.; Lupo, T.; Migliazzo, A.; Capri, A.; Solano-Gallego, L. Detection of *Leishmania infantum* DNA by real-time PCR in canine oral and conjunctival swabs and comparison with other diagnostic techniques. *Vet. Parasitol.* **2011**, *184*, 10–17.
38. Salvatore, D.; Aureli, S.; Baldelli, R.; Di Francesco, A.; Tampieri, M.P.; Galuppi, R. Molecular evidence of *Leishmania infantum* in *Ixodes ricinus* ticks from dogs and cats In Italy. *Vet. Ital.* **2014**, *50*, 307–312.
39. Pennisi, M.G.; Persichetti, M.F.; Serrano, L.; Altet, L.; Reale, S.; Gulotta, L.; Solano-Gallego, L. Ticks and associated pathogens collected from cats in Sicily and Calabria (Italy). *Parasites Vectors* **2015**, *8*, 512.
40. Persichetti, M.F.; Solano-Gallego, L.; Serrano, L.; Altet, L.; Reale, S.; Masucci, M.; Pennisi, M.G. Detection of vector-borne pathogens in cats and their ectoparasites in southern Italy. *Parasites Vectors* **2016**, *9*, 247.
41. Plebotomine Maps. Available online: http://ecdc.europa.eu/en/healthtopics/vectors/vector-maps/Pages/VBORNET_maps_sandflies.aspx (accessed on 21 March 2020).
42. Morosetti, G.; Bongiorno, G.; Beran, B.; Scalone, A.; Moser, J.; Gramiccia, M.; Gradoni, L.; Maroli, M. Risk assessment for canine leishmaniasis spreading in the north of Italy. *Geospat. Health* **2009**, *4*, 115–127.
43. Maroli, M.; Gradoni, L.; Houry, C.; Bianchi, R.; Nicoli Aldini, R.; Gramiccia, M. Entomological survey on phlebotomine sandflies (Diptera: Psychodidae) in three sub-Alpine valleys of Lombardy region, northern Italy. *Parassitol.* **2006**, *48*, 159.
44. Vita, S.; Santori, D.; Aguzzi, I.; Petrotta, E.; Luciani, A. Feline leishmaniasis and ehrlichiosis: Serological investigation in Abruzzo region. *Vet. Res. Commun.* **2005**, *29*, 319–321.
45. Paşa, S.; Tetik Vardarli, A.; Erol, N.; Karakuş, M.; Töz, S.; Atasoy, A.; Balcioğlu, İ.C.; Emek Tuna, G.; Ermiş, Ö.V.; Ertabaklar, H.; et al. Detection of *Leishmania major* and *Leishmania tropica* in domestic cats in the Ege Region of Turkey. *Vet. Parasitol.* **2015**, *212*, 389–392.
46. Chatzis, M.K.; Leontides, L.; Athanasiou, L.V.; Papadopoulos, E.; Kasabalis, D.; Mylonakis, M.; Rallis, T.; Koutinas, A.F.; Andreadou, M.; Ikononopoulos, J.; et al. Evaluation of indirect immunofluorescence antibody test and enzyme-linked immunosorbent assay for the diagnosis of infection by *Leishmania infantum* in clinically normal and sick cats. *Exp. Parasitol.* **2014**, *147*, 54–59.
47. Baneth, G.; Nachum-biala, Y.; Zuberi, A.; Zipori-barki, N.; Orshan, L.; Kleinerman, G.; Shmueli-goldin, A.; Bellaiche, M.; Leszkowicz-mazuz, M.; Salant, H.; et al. *Leishmania* infection in cats and dogs housed together in an animal shelter reveals a higher parasite load in infected dogs despite a greater seroprevalence among cats. *Parasites Vectors* **2020**, *13*, 115.
48. Pennisi, M.G.; Venza, M.; Reale, S.; Vitale, F.; Lo Giudice, S. Case report of leishmaniasis in four cats. *Vet. Res. Commun.* **2004**, *28*, 363–366.
49. Brianti, E.; Celi, N.; Napoli, E.; Abbate, J.M.; Arfuso, F.; Gaglio, G.; Iatta, R.; Giannetto, S.; Gramiccia, M.; Otranto, D. Treatment and long-term follow-up of a cat with leishmaniasis. *Parasites Vectors* **2019**, *12*, 1–7.
50. Ozon, C.; Marty, P.; Pratlong, F.; Breton, C.; Blein, M.; Lelièvre, A.; Haas, P. Disseminated feline leishmaniasis due to *Leishmania infantum* in Southern France. *Vet. Parasitol.* **1998**, *75*, 273–277.

51. Poli, A.; Abramo, F.; Barsotti, P.; Leva, S.; Gramiccia, M.; Ludovisi, A.; Mancianti, F. Feline leishmaniosis due to *Leishmania infantum* in Italy. *Vet. Parasitol.* **2002**, *106*, 181–191.
52. Pimenta, P.; Alves-Pimenta, S.; Barros, J.; Barbosa, P.; Rodrigues, A.; Pereira, M.J.; Maltez, L.; Gama, A.; Cristóvão, J.M.; Campino, L.; et al. Feline leishmaniosis in Portugal: 3 cases (year 2014). *Vet. Parasitol. Reg. Stud. Rep.* **2015**, *1–2*, 65–69.
53. Spada, E.; Proverbio, D.; Della Pepa, A.; Perego, R.; Baggiani, L.; De Giorgi, G.B.; Domenichini, G.; Ferro, E.; Cremonesi, F. Seroprevalence of feline immunodeficiency virus, feline leukaemia virus and *Toxoplasma gondii* in stray cat colonies in northern Italy and correlation with clinical and laboratory data. *J. Feline Med. Surg.* **2012**, *14*, 369–377.
54. Akhtardanesh, B.; Moeini, E.; Shari, I.; Saberi, M.; Sadeghi, B.; Ebrahimi, M.; Otranto, D. Leishmania infection in cats positive for immunodeficiency virus and feline leukemia virus in an endemic region of Iran. *Vet. Parasitol. Reg. Stud. Rep.* **2020**, *20*, 100387.
55. Migliazzo, A.; Vitale, F.; Calderone, S.; Puleio, R.; Binanti, D.; Abramo, F. Feline leishmaniosis: A case with a high parasitic burden. *Vet. Derm.* **2015**, *26*, 69–70.
56. Sherry, K.; Miró, G.; Trotta, M.; Miranda, C.; Montoya, A.; Espinosa, C.; Ribas, F.; Furlanello, T.; Solano-Gallego, L. A Serological and Molecular Study of *Leishmania infantum* Infection in Cats from the Island of Ibiza (Spain). *Vector-Borne Zoonotic Dis.* **2011**, *11*, 239–245.
57. Marinho Dourado Coelho, W.; Francisco Talamini do Amarante, A.; de Carvalho Apolinário, J.; Marinho Dourado Coelho, N.; Marçal Felix de Lima, V.; Helena Venturoli Perri, S.; Denise Saraiva Bresciani, K. Seroepidemiology of *Toxoplasma gondii*, *Neospora caninum*, and *Leishmania* spp. infections and risk factors for cats from Brazil. *Parasitol. Res.* **2011**, *109*, 1009–1013.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).