

# Missed Study Visits and Subsequent HIV Incidence Among Women in a Predominantly Sex Worker Cohort Attending a Dedicated Clinic Service in Kampala, Uganda

Ivan Kasamba, MSc,<sup>a,b</sup> Stephen Nash, MSc,<sup>b,c</sup> Maryam Shahmanesh, PhD,<sup>d</sup> Kathy Baisley, MSc,<sup>b</sup> Jim Todd, MSc,<sup>b</sup> Onesmus Kamacooko, MSc,<sup>a</sup> Yunia Mayanja, MSc,<sup>a</sup> Janet Seeley, PhD,<sup>a,b</sup> and Helen A. Weiss, PhD<sup>b,c</sup>

**Background:** There is limited evidence on the relationship between sustained exposure of female sex workers (FSWs) to targeted HIV programmes and HIV incidence. We investigate the relationship between the number of missed study visits (MSVs) within each episode of 2 consecutively attended visits (MSVs) and subsequent HIV risk in a predominantly FSW cohort.

**Methods:** Women at high risk of HIV are invited to attend an ongoing dedicated clinic offering a combination HIV prevention intervention in Kampala, Uganda. Study visits are scheduled once every 3 months. The analysis included HIV-seronegative women with  $\geq 1$  follow-up visit from enrollment (between April 2008 and May 2017) to August 2017. Cox regression models were fitted adjusted for characteristics on sociodemographic, reproductive, behavioral, and sexually transmitted infections (through clinical examination and serological testing for syphilis).

**Findings:** Among 2206 participants, HIV incidence was 3.1/100 (170/5540) person-years [95% confidence interval (CI): 2.6 to 3.5].

Incidence increased from 2.6/100 person-years (95% CI: 2.1 to 3.2) in episodes without a MSV to 3.0/100 (95% CI: 2.2 to 4.1) for 1–2 MSVs and 4.3/100 (95% CI: 3.3 to 5.6) for  $\geq 3$  MSVs. Relative to episodes without a MSV, the hazard ratios (adjusted for confounding variables) were 1.40 (95% CI: 0.93 to 2.12) for 1–2 MSVs and 2.00 (95% CI: 1.35 to 2.95) for  $\geq 3$  MSVs ( $P$ -trend = 0.001).

**Conclusion:** Missing study visits was associated with increased subsequent HIV risk. Although several factors may underlie this association, the finding suggests effectiveness of targeted combination HIV prevention. But exposure to targeted interventions needs to be monitored, facilitated, and sustained in FSWs.

**Key Words:** HIV, HIV incidence, missed study visits, female sex workers, sub-Saharan Africa

(*J Acquir Immune Defic Syndr* 2019;82:343–354)

## INTRODUCTION

Combination of HIV-prevention programmes targeting sex workers is cost-effective, especially in settings with high HIV incidence and prevalence among sex workers.<sup>1,2</sup> However, when HIV prevention in sex work is not sustained in these settings, there can be rapid HIV acquisition and onward HIV transmission beyond female sex workers (FSWs) and their clients to the wider population.<sup>3</sup> In Uganda, a systematic review of epidemiological data among FSWs showed an estimated HIV prevalence in the range of 32%–52%, which is 4–7 times that in adult women in the general population.<sup>4,5</sup> Although there is sufficient evidence that supports effectiveness of multicomponent-targeted HIV interventions for FSWs, sustaining exposure to these interventions remains a major challenge.<sup>6–9</sup>

The WHO recommends routinely offering evidence-based comprehensive HIV-prevention services to key populations such as FSWs, which includes HIV counseling and testing (HCT), and diagnosis and treatment of sexually transmitted infections (STIs).<sup>10</sup> This regular exposure to services is likely to reinforce behavioral messages, to facilitate community empowerment, and to detect and treat STIs early among FSWs.<sup>8</sup> Furthermore, regular exposure to HCT can facilitate earlier initiation of antiretroviral therapy (ART) during the acute and early infectious phases of HIV.

Received for publication March 11, 2019; accepted July 8, 2019.

From the <sup>a</sup>MRC/UVRI and LSHTM Uganda Research Unit, Entebbe, Uganda; <sup>b</sup>London School of Hygiene and Tropical Medicine (LSHTM), London, United Kingdom; <sup>c</sup>MRC Tropical Epidemiology Group, London School of Hygiene and Tropical Medicine (LSHTM), London, United Kingdom; and <sup>d</sup>Institute for Global Health, University College London, London, United Kingdom.

Supported by the UK Medical Research Council (MRC) and the UK Department for International Development (DFID) under the MRC/DFID Concordat agreement (K012126/1). The clinic receives support from PEPFAR through the Centres for Disease Control and Prevention for the provision of clinical services. I.K. is jointly funded for a PhD by MRC Tropical Epidemiology Group, London School of Hygiene and Tropical Medicine (LSHTM), UK and UK Medical Research Council (MRC).

The authors have no conflicts of interest to disclose.

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's Web site ([www.jaids.com](http://www.jaids.com)).

Correspondence to: Ivan Kasamba, MSc, MRC/UVRI and LSHTM Uganda Research Unit, P.O. Box 49, Plot 51-59 Nakiwogo Road, Entebbe, Uganda (e-mail: [Ivan.Kasamba@lshtm.ac.uk](mailto:Ivan.Kasamba@lshtm.ac.uk)).

Copyright © 2019 The Author(s). Published by Wolters Kluwer Health, Inc. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial License 4.0 (CCBY-NC), where it is permissible to download, share, remix, transform, and buildup the work, provided it is properly cited. The work cannot be used commercially without permission from the journal.

Notwithstanding the limited coverage for FSW-targeted services across Africa, several factors exist at multiple levels that facilitate or deter FSWs from using services.<sup>11–13</sup> The most commonly identified contextual factors are stigma, discrimination, and criminalization and their associated consequences.<sup>14–16</sup> However, other contextual factors such as the high mobility among sex workers have been identified to delay the utilization of services.<sup>16,17</sup> These contextual factors typically act through the individual factors such as perceptions of HIV risk and importance of services as well as behavioral (sexual and substance abuse), reproductive, and biological characteristics (Fig. 1).<sup>14,15</sup> At the same time, these key determinants of service utilization are also important risk factors for HIV infection among FSWs.<sup>18–20</sup>

Although there is growing interest in improving service utilization among FSWs, it remains unclear how recent levels of service attendance are associated with subsequent HIV risk.<sup>14,19,20</sup> Using the proportions of study visits completed, research has linked reduced incidence of HIV and other STIs among FSWs to amount of exposure to targeted HIV interventions such as peer-mediated HIV education, condom promotion, risk-reduction interventions, and presumptive treatment for STIs.<sup>21–23</sup> However, assessment of the HIV risk associated with previous recent levels of service attendance may offer better evidence for risk associated with intervention exposure because an individual's level of attendance can vary over time and for different reasons. If we understand this relationship better, we could then inform the design and scale-up of appropriate interventions in sex work.

In a Kenyan FSW cohort with monthly clinic visits followed for a median of 16.2 months, a time-varying preceding gap in clinic attendance of  $\geq 60$  days was associated


with reduced HIV risk.<sup>24</sup> But this variable of preceding gap in clinic attendance in this study was only considered as a confounding variable and, thus, warrants further investigation as a main exposure of interest. As a main exposure, a different approach that considers the conceptual interrelationships with potential confounders and their relationship with HIV risk should be applied. Furthermore, instead of assigning the date of seroconversion as midpoint between the last-negative and first-positive test dates, we assigned a random date in this interval, thereby reducing on the artifactual clustering of seroconversion times in the middle of observation periods with missed scheduled testing.<sup>25,26</sup>

Therefore, the aim of this study was to examine the association between the number of missed study visits (MSVs) within episodes of 2 consecutively attended visits and subsequent HIV risk in a predominantly FSW cohort attending a dedicated clinic in Kampala, Uganda.

## METHODS

### Study Population

Between March 2008 and May 2009 (cohort-1), the Good Health for Women Project (GHWP) enrolled women acknowledging being FSWs or working in entertainment facilities such as bars, night clubs, or lodges (women at high risk of HIV infection) in Kampala, Uganda.<sup>27</sup> Participants were scheduled to attend the study clinic once every 3 months, which is consistent with the Uganda Ministry of Health guidelines for HCT of key populations.<sup>28</sup> From January 2013 (cohort-2), more participants were continuously enrolled into GHWP alongside cohort-1 participants.



**FIGURE 1.** Conceptual framework linking factors leading to MSVs and subsequent HIV incidence used to fit the Cox regression model within the GHWP. IEC, information education communication; SRH, sexual and reproductive health.

**TABLE 1.** HIV-Prevention Intervention Available at the GHWP by the Type of Intervention

Type of Intervention	Components	Delivery
Behavioral	Condom promotion, demonstration, and free condom distribution	Each visit
	Sexual counseling	Each visit
	Excessive alcohol use counseling including the AUDIT* questionnaire as part of the motivational interviewing	Each visit
	Knowledge of HIV status through HIV testing	Each visit and at moonlight outreaches to sex work hotspots
	If diagnosed with HIV, offered positive prevention strategies, treatment adherence counseling, and HIV care	Each visit and at ART refill visits
	Information sharing and health education	Each visit and at the group meetings held fortnightly for participants with upcoming visits
	General counseling services	Each visit
Biomedical	Syndromic management of sexually transmitted infections	Each visit
	Referrals to other collaborating service providers for participants in need of postexposure prophylaxis	As needed
Structural	Antiretroviral therapy (ART)† if diagnosed with HIV	As needed
	Access to a free sex worker–friendly dedicated clinic	As needed
	Sexual and reproductive health services including antenatal care and contraceptive provision	Every visit and as needed
	Free general health care and their children aged <5 years old	As needed
	Support by peer support network	As needed
	Outreach HIV counseling and testing, and supply of condoms and health information at the sex work hotspots	Two nights a week, with a team that includes peer educators, venue managers, and study staff
	Community mobilization involving police, venue managers and owners, and local authorities	Quarterly meetings
Protection services and referrals to community-based organizations in case of gender-based violence	As needed	

\*Alcohol Use Disorders Identification Test (AUDIT) was initiated in April 2012, but there was ongoing risk-reduction counseling.

†ART became available at the clinic in January 2013 (eligibility: CD4 cell counts <350 cells/μl, and from August 2014, for all diagnosed with HIV regardless of CD4 cell count), but before that participants with CD4 cell count below 250 cells/μl were referred to HIV care providers.

The procedures for recruiting, screening, enrolling, and following up of cohort-1 participants have been described in detail elsewhere.<sup>27</sup> But, identical procedures were followed for cohort-2. Briefly, outreach workers visited and assessed study eligibility for women at high risk of HIV infection who had been recruited by peer educators or contacted through evening community-outreach HCT sessions. Women were eligible if they were involved in commercial sex (self-identified FSWs or received money, goods, or other favors in exchange for sex) or employed in entertainment facilities within mapped sex work hotspots (clusters of bars, night clubs, lodges, and guest houses providing rooms for sex work or street spots frequented by sex workers). Those who were eligible were invited to attend the study clinic for rescreening and study enrollment.

**Targeted Intervention**

The GHWP intervention follows the WHO standard framework for HIV prevention in FSWs; aiming at reducing condomless sex, decreasing HIV transmission efficiency,

and empowering FSWs.<sup>29</sup> This intervention integrates sex worker–friendly HIV services with other sexual and reproductive health services and offers free general health care for participants and their children aged 5 years younger. Table 1 shows the detailed components of the intervention broadly classified as structural, behavioral, and biomedical.

At each visit, HIV-preventive behaviors were promoted to each participant, but the content of the counseling sessions was depended on each participant’s behavioral responses, outcomes of clinical examinations, and serological test results. Low-risk participants were encouraged to maintain behavior, whereas those with risky behaviors were supported to acknowledge the behavior and its possible consequences. In addition, high-risk participants were encouraged to take immediate steps to reduce the risk associated with behaviors such as harmful alcohol use or inconsistent condom use. Depending on their identified needs, participants were also counseled on a wide range of other physical and psychosocial issues including their general health, family, spiritual needs, and employment.

To ensure high retention, group meetings were held every 14 days at the study clinic with women who had scheduled visits within the following month. Participants with 2 or more consecutive MSVs were contacted by phone or traced in the community (either at their workplace or home) through a peer leader or the local council member of the respective area and then invited to attend the clinic at an appointed time.

## Data

At the clinic, trained nurse-counselors used structured questionnaires to collect data on sociodemographics, sexual behavior, reproductive health, and substance abuse (alcohol use and illicit drug use). Data were collected only at enrollment for age, education level, source of income (sex work alone or sex work and other jobs), place of client recruitment (street, bar, club or restaurant, and several avenues), and number of children. However, data collected at enrollment and follow-up visits included marital status and behavioral variables such as alcohol use, illicit drug use, number of sexual partners, number of paying sexual partners, frequency of condom use with clients, report of any paid sex in the last month, and contraceptive use and pregnancy. Pregnancy was determined using human chorionic gonadotropin urine sample tests or visible pregnancy.

A participant had an STI if he/she had an active syphilis infection (rapid plasma reagin titers  $\geq 1:8$ ) and positive *Treponema pallidum* hemagglutination assay based on serology tests conducted at 6-month intervals or had any of the STI symptoms of abnormal vaginal discharge, genital ulcer disease syndrome, or pelvic inflammatory disease. Between 2008 and 2010, endocervical and vaginal swabs were also used to detect *Neisseria gonorrhoeae*, *Chlamydia trachomatis*, *Mycoplasma genitalium*, and bacterial vaginosis.

## Outcome Variable

The primary outcome was HIV seroconversion, which was defined as a negative test result at a given visit attended followed by a positive result at the subsequent visit. Testing for HIV followed the Uganda MoH algorithm for HIV testing. According to this algorithm, HIV screening was performed using Determine HIV-1/2 (Abbott Diagnostics, Maidenhead, United Kingdom). Before 2013, reactive results were confirmed using 2 enzyme immunoassay tests for HIV screening (Vironostika Uniform II plus O, Murex HIV 1.2.0) and then a Western Blot Test. But from 2013, confirmation was done using Stat-Pak dipstick HIV-1/2 (Chembio Diagnostics), and discordant samples were further tested with the Uni-Gold HIV-2 (Trinity Biotech, Ireland). Participants were included in this analysis if they tested HIV-negative at enrollment and had at least one follow-up visit by 29th August 2017 (administrative censoring date). The total person-years (pyr) for each participant were calculated from their enrollment date to the earliest of (1) estimated date of seroconversion for seroconverters or (2) last known HIV-negative test date for nonseroconverters. The approach for estimating the date of

seroconversion is described under the statistical analysis section.

## Exposure Variable

The number of MSVs within each participant's episodes of 2 consecutively attended visits was used to define the exposure of interest. This exposure variable is a proxy for the extent of exposure to and engagement with the combined components of the HIV intervention offered at the clinic. We classified a scheduled study visit as missed if the participant did not attend the study clinic within 14 days before and 76 days after the scheduled visit date. At each follow-up visit, the number of scheduled visits that the participant consecutively missed since the previous visit attended was calculated, resulting in a time-varying exposure variable. However, because the exposure variable is likely to have sparse data for higher numbers of MSVs, we categorized the counts of MSV into 3 levels: (1) no missed visit, (2) 1–2 MSV, and (3)  $\geq 3$  MSV.

## Statistical Analysis

The total person-years of observation for each participant were split according to their number of episodes of attendance represented by the time period between 2 consecutively attended visits. Each episode was then assigned an appropriate category for the 3-level MSV variable defined previously. There would be increased uncertainty for the date of seroconversion during episodes with MSVs while estimating the person-years for seroconverters. To address this potential concern, the date of seroconversion was assigned as a random date from a uniform distribution bounded by the last-negative and first-positive test dates. In addition, the variability due to assigning a random date was accounted for by multiple imputations (200 imputations). For each set of imputed seroconversion dates, the HIV incidence rates and crude and adjusted hazard ratios (aHR) were calculated and combined using Rubin's rules.<sup>30</sup>

To facilitate identifying potential confounders for the association between MSV and HIV incidence, we examined factors associated with MSV using repeated-measures ordered logistic regression. In this model, the three-level MSV variable was fitted as the outcome. Within-woman clustering due to repeated measures was accounted for using the random-effects model. We also examined factors associated with HIV infection.

All models included an indicator variable for cohort (1 or 2) and age group a priori. The incidence analysis was performed in 3 stages: first, the association between MSV and HIV seroconversion was examined using Cox regression models that included the cohort indicator and age. Second, sociodemographic, behavioral, reproductive health, and STIs characteristics were added individually to the base model to assess whether the crude association persisted. Third, the final model was adjusted for variables that were likely confounders of the MSV and HIV seroconversion relationship (ie, at least a 10% change in the log-transformed MSV effects). As a sensitivity analysis, results from the final model were

**TABLE 2.** Comparing the Enrollment Characteristics for HIV-Negative Women With and Without a Follow-up Visit

Characteristic	Follow-up Status, n (%)		P
	At Least One Follow-up	No Follow-up Visit	
Number enrolled	2206	878	
Sociodemographic			
Age in years			<0.001
<24	891 (41.0)	459 (54.1)	
25–34	985 (45.3)	321 (37.8)	
35+	299 (13.7)	69 (8.1)	
Missing	31	29	
Highest education level			0.94
Less than primary	863 (41.3)	328 (42.0)	
Completed primary to incomplete O level	889 (42.6)	330 (42.3)	
Completed secondary ordinary level above	336 (16.1)	123 (15.7)	
Missing	118	97	
Marital status			0.01
Widowed/divorced	1381 (63.2)	484 (57.1)	
Currently married	148 (6.8)	67 (7.9)	
Never married	654 (30.0)	296 (34.9)	
Missing	23	31	
No. of children			<0.001
None	297 (13.7)	187 (22.4)	
One	542 (25.1)	239 (28.7)	
At least 2	1322 (61.2)	407 (48.9)	
Missing	45	45	
Source of income			0.01
Sex work alone	1227 (56.5)	523 (62.1)	
Sex work and other job	844 (38.8)	276 (32.8)	
No sex work	102 (4.7)	43 (5.1)	
Missing	33	36	
Where paying clients are recruited by participants			<0.001
Bar, club, or restaurant	1040 (50.9)	356 (45.4)	
Street	472 (23.1)	291 (37.1)	
Several avenues	532 (26.0)	138 (17.6)	
Missing	162	93	
Reproductive health			
Pregnancy status			0.02
Not pregnant	2048 (94.8)	812 (96.9)	
Pregnant	112 (5.2)	26 (3.1)	
Missing	46	40	
Current contraceptive use			<0.001
None or other	1285 (61.7)	531 (68.9)	
Oral cc	170 (8.2)	34 (4.4)	
Inject	514 (24.7)	180 (23.3)	
Pregnant	112 (5.4)	26 (3.4)	
Missing	125	107	
Behavioral characteristics			
Frequency of paid sex in the past 12 months			<0.001
Less than once a week/none	275 (12.6)	125 (14.6)	

**TABLE 2. (Continued)** Comparing the Enrollment Characteristics for HIV-Negative Women With and Without a Follow-up Visit

Characteristic	Follow-up Status, n (%)		P
	At Least One Follow-up	No Follow-up Visit	
At least once a week	735 (33.5)	210 (24.5)	
Daily	1181 (53.9)	522 (60.9)	
Missing	15	21	
No. of partners in the last month			0.07
<5	563 (27.0)	189 (24.3)	
5–19	499 (23.9)	170 (21.8)	
At least 20 or cannot remember	1022 (49.0)	420 (53.9)	
Missing	122	99	
No. of paying partners in the last month			0.10
<5	587 (28.2)	204 (26.3)	
5–19	488 (23.4)	162 (20.8)	
At least 20	1009 (48.4)	411 (52.9)	
Missing	122	101	
Condom use frequency with paid sex in the last month			0.09
Inconsistent	883 (42.2)	297 (38.0)	
Consistent (always)	969 (46.3)	397 (50.8)	
No paid sex missing	241 (11.5)	88 (11.3)	
Alcohol consumption frequency			0.46
Nondrinker	488 (22.8)	203 (24.9)	
Non-daily drinker	891 (41.6)	327 (40.2)	
Daily drinker	762 (35.6)	284 (34.9)	
Missing	65	64	
Binge drinking within the past 3 months			0.01
Nondrinker	489 (22.7)	201 (24.2)	
No bingeing	557 (25.8)	164 (19.7)	
Binged	1112 (51.5)	466 (56.1)	
Missing	48	47	
Illicit drug use in the past 3 months			0.45
Not used drugs in the past 3 months	1513 (72.5)	562 (72.1)	
Non-daily drug user	129 (6.2)	58 (7.4)	
Daily drug user	444 (21.3)	159 (20.4)	
Missing	120	99	
HIV testing history			<0.001
>1 yr ago or never	788 (36.5)	202 (24.3)	
7–12 months ago	309 (14.3)	95 (11.4)	
<6 months ago	1064 (49.2)	535 (64.3)	
Missing	45	46	
Sexually transmitted infections (STIs)*			
No STI	1349 (63.2)	592 (72.8)	<0.001
With STI	787 (36.8)	221 (27.2)	

\*STI variable was a composite variable combining data on STI symptoms from clinical examination and test results for STI for syphilis.

compared with those where analyses were restricted to women with at least 6 months of observation time. This is because HIV incidence peaked within the first 6 months, yet participants were less likely to miss scheduled visits during this period than later periods.<sup>26</sup> After performing an interaction of MSV with the cohort indicator variable in the final model adjusted for other confounders, results were also compared for the 2 cohorts.

Figure 1 shows the relationship between MSV and subsequent HIV risk and the relationships across factors controlled for in the regression model. Although behavioral factors and detection of STIs can influence attendance at the study clinic, attendance can also influence reported behavioral and STI outcomes. As a result, the Cox regression model adjusted for time-dependent variables at the preceding visits. Ordered categorical variables in the model were assessed for extralinear variability using likelihood ratio tests by comparing models with variables as categorical versus a linear fit. Analyses were performed using STATA 14 (StataCorp, College Station, TX).

## RESULTS

### Cohort Characteristics

Of the 3084 HIV-negative women enrolled at the clinic, 2206 (71.5%) attended  $\geq 1$  follow-up visit after enrollment: 93.5% (605/647) in cohort-1 and 65.9% (1601/2437) in cohort-2. Between the first enrollment (April 2008) and end of follow-up (Aug 2017), a total of 24,476 visits (cohort-1: 12,786, cohort-2: 11,690) had been scheduled by the last visit of each participant with  $\geq 1$  follow-up visit. Of these, 17,486 (71.4%) scheduled visits (cohort-1: 9,462, cohort-2: 8024) were attended. The median number of visits attended per participant was 15 [interquartile range (IQR): 8–22] in cohort-1 and 4 (IQR: 2–6) in cohort-2.

Table 2 compares enrollment characteristics for HIV-negative women with and without a follow-up visit. In the multivariable analysis (see Table 1, Supplemental Digital Content, <http://links.lww.com/QAI/B362>), the factors associated with having no follow-up include younger age, fewer children, street-based recruitment of clients, not on oral contraceptives, less frequent paid sex, and consistently using condoms with clients in the last month.

### HIV Incidence

Among women with  $\geq 1$  follow-up visit, we observed 5540 person-years and 170 (7.7%) seroconversions, giving an incidence rate of 3.1/100 person-years [95% confidence interval (CI): 2.6 to 3.5]. Overall incidence was similar in the 2 cohorts (cohort-1: 93 seroconversions; incidence = 3.1/100 person-years; cohort-2: 77 seroconversions; incidence = 3.0/100 person-years). HIV incidence declined after enrollment, from 3.6/100 person-years within the first year, 2.8/100 person-years in the second year, and 2.5/100 person-years beyond 2 years of follow-up.

Figure 2 shows that episodes of 2 consecutively attended visits during which HIV was diagnosed are longer

(have their distribution more skewed to the right) than those with HIV-negative test results at both attended visits. After restricting to episodes with  $\geq 3$  MSV, the median number of MSV within each participant's period of 2 consecutively attended visits was 4 MSVs (IQR: 3–6), whereas one person had the longest episode of 30 missed visits (Fig. 2). In examining the potential confounders for the association of MSV and HIV seroconversion (see Table 2, Supplemental Digital Content, <http://links.lww.com/QAI/B362>), there was evidence that the odds of being in higher categories of MSV than in lower categories were independently associated with: enrollment in cohort-2, longer follow-up, incomplete ordinary secondary education level, street-based recruitment of clients, being pregnant, reporting no paid sex within the last month, alcohol consumption, illicit drug use in past 3 months, and having no STI at the previous visit.

The number of MSVs seems to be positively correlated with HIV incidence. HIV incidence was 2.6/100 person-years (95% CI: 2.1 to 3.2) during episodes without missed visits, 3.0/100 person-years (95% CI: 2.2 to 4.1) in episodes with 1–2 MSV, and 4.3/100 person-years (95% CI: 3.3 to 5.6) in episodes with  $\geq 3$  MSV (*P* value for trend = 0.003; Table 3).

The number of MSVs remained correlated with subsequent increase in HIV risk even after taking account of differences in cohort of enrollment, age of participant, education level, having a stable sexual partner, location for recruiting clients, current type of contraceptive, the number of men with paid sex in the last month, frequency of condom use during paid sex, frequency of alcohol consumption, and having a current STI (Table 3).


In this multivariable model, episodes with 1–2 MSV were associated with 40% higher risk of HIV infection (aHR = 1.40, 95% CI: 0.93 to 2.12) than episodes without a MSV, but the evidence for this association was weak. However, episodes with  $\geq 3$  MSV had twice the risk of HIV infection (aHR = 2.00, 95% CI: 1.35 to 2.95) compared with that in episodes without a MSV. Furthermore, there was strong evidence in favor of the risk of HIV infection increasing with the increasing number of MSV within episodes; *P* value for trend = 0.001 and *P*-value for extralinear variability = 0.91.

The evidence for the association between MSV and increased HIV incidence remained when analyses were restricted to follow-up of  $>6$  months [*P* value for trend  $<0.001$ ; aHR = 1.61 (95% CI: 1.00 to 2.58) for 1–2 MSV, and aHR = 2.05 (95% CI: 1.31 to 3.22) for  $\geq 3$  MSV]. But the estimated effect for the 1–2 MSV category was stronger in this restricted analysis. Although there was no evidence that the effect of MSV differed by cohort (*P* value for interaction = 0.24), the results are also shown by cohort in Table 3, Supplemental Digital Content, <http://links.lww.com/QAI/B362>.

Other factors independently associated with increased risk of HIV incidence included the following: not having a stable sexual partner, recruiting clients from the street, not being pregnant, and inconsistent condom use with clients.

## DISCUSSION

Our findings demonstrate the importance of ensuring sustained exposure to combined HIV-prevention



**FIGURE 2.** Distribution for the number of MSVs within each participant’s episodes of 2 consecutively attended visits by HIV seroconversion status in the episodes.

interventions targeting women at high-risk of HIV infection in the prevention of HIV. Sustained exposure is not only likely to reinforce prevention interventions but also facilitate earlier initiation of ART among those recently infected with HIV. Participants attending the clinic had access to a range of services including HIV and other sexual health and reproductive services. At every visit, participants attended counseling sessions where they were encouraged to modify or maintain behaviors depending on their interview responses, results from clinical examinations, and serological test. Participants diagnosed with STIs were additionally given treatment. Therefore, MSV within episodes of consecutive attendances was a proxy for recent degree of exposure to combined HIV-prevention interventions.

Compared with episodes without MSV, those with 1–2 MSV had 40% higher risk of HIV infection, whereas those with  $\geq 3$  MSV had twice the risk. This dose–response relationship further supports the effectiveness of combination HIV-prevention interventions among FSWs for improved HIV-related outcomes such as consistent condom use with clients, earlier treatment of STIs, and consequently, lower HIV infection.<sup>6,7,31,32</sup> However, this effectiveness is dependent on good program adherence.

In addition, our findings are consistent with studies that have suggested a link between the amount of exposure to HIV interventions and improved HIV-related outcomes.<sup>21–23</sup> Most notable is a study in Côte d’Ivoire that conducted health talks, provided condoms, treated STIs at monthly clinic visits, and serological tests every 6 months.<sup>23</sup> In this study, HIV incidence was 4.6/100 person-years for women who attended 4 of the 5 scheduled visits before their last assessment and 13.0/100 person-years among those who attended fewer visits.<sup>23</sup> But a time-varying preceding gap in clinic attendance of  $\geq 60$  days was associated with reduced HIV risk in a Kenyan FSW cohort with monthly clinic visits.<sup>24</sup> Although the authors did not offer a reason for this association, it is likely that participants who missed the monthly visits may have had low-risk perception and, therefore, saw no need to

regularly attend visits. In addition, the variable indicating a preceding gap was correctly analyzed as a confounder, but where it is the main exposure of interest, the time-varying variables in that analysis represent potential mediating factors. Thus, these mediating factors underestimate the overall effect of the preceding gap.

In our study, the mechanisms through which MSV within episodes of consecutive visit attendances influence HIV infection may be explained by interconnected factors operating at the structural, behavioral, and biomedical levels.

### Structural Factors

HIV determinants such as high mobility, economic difficulties, and risky work environments of FSWs may underlie the risk associated with MSV. Although the ability to take risks (being a more risky individual) could influence an individual to be mobile in search of work; mobility itself can affect the risk of HIV if there is risk of HIV associated with the new environment/place (ie, the new place has high HIV prevalence, low condom use, and low ART coverage or adherence). A previous qualitative study in this cohort reported that participants commonly moved to fishing communities for extended periods in search of clients.<sup>15</sup> These communities have been linked with high HIV prevalence and incidence, yet the fishermen usually preferred condomless sex, thus increasing HIV vulnerabilities for the mobile participants.<sup>33</sup>

Other findings in this cohort are a signal to the potential economic pressures facing participants to not only deter them from accessing the much-needed clinic services but also affecting their ability to refuse sex with a client paying more for noncondom use. For example, when transport refund for participants was stopped in January 2013, the rate of MSV among those attending each visit increased substantially from 10% to 30%.<sup>26</sup> Furthermore, some 5% of participants reported lack of transport as a cause for missing 2 or more

**TABLE 3.** Association of HIV Seroconversion With the Number of Missed Visits Between Consecutively Attended Visits Based on the Cox Regression Model

Characteristic	New HIV Cases/pyr (Rate/100 pyr)	Crude Hazard Ratio (95% CI)	P	Adjusted Hazard Ratio (95% CI)	P
Overall (both cohort)	170/5540 (3.1)				
No. of visits missed between attendances					
No missed visit	81/3106 (2.6)	1	0.003*	1	0.001*
One/2 missed visit(s)	38/1250 (3.0)	1.18 (0.80 to 1.74)		1.40 (0.93 to 2.12)	
At least 3 missed visits	51/1180 (4.3)	1.77 (1.23 to 2.54)		2.00 (1.35 to 2.95)	
Sociodemographic					
Cohort of enrollment					
Cohort-1	93/3003 (3.1)	1	0.45	1	0.43
Cohort-2	77/2537 (3.0)	0.87 (0.61 to 1.24)		0.85 (0.56 to 1.27)	
Calendar period†					
Jan 2013–Dec 2014	46/1394 (3.3)	1	0.16		
Jan 2015–Aug 2017	65/2138 (3.0)	0.99 (0.64 to 1.55)			
Apr 2008–Dec 2012	59/2007 (2.9)	1.09 (0.70 to 1.70)			
Age in years					
<25	53/1562 (3.4)	1	0.49	1	0.37
25–34	92/2955 (3.1)	0.96 (0.68 to 1.35)		0.86 (0.60 to 1.24)	
At least 35	25/994 (2.5)	0.76 (0.46 to 1.24)		0.80 (0.48 to 1.33)	
With a stable sexual partner‡					
Yes	113/3959 (2.9)	1	0.01	1	0.003
No	57/1211 (4.7)	1.55 (1.13 to 2.14)		1.75 (1.23 to 2.50)	
Highest education level at enrollment					
Incomplete primary level	74/2361 (3.1)	1	0.03	1	0.24
Completed primary to incomplete O level	70/2267 (3.1)	0.97 (0.70 to 1.35)		1.06 (0.76 to 1.49)	
Completed secondary O level above	25/792 (3.2)	1.00 (0.63 to 1.57)		1.20 (0.75 to 1.92)	
Marital status§					
Widowed/divorced	112/3325 (3.4)	1	0.33		
Currently married	28/1150 (2.4)	0.77 (0.50 to 1.17)			
Never married	30/1044 (2.9)	0.81 (0.54 to 1.22)			
No. of children§					
None	21/583 (3.6)	1	0.41		
One	42/1247 (3.4)	0.94 (0.56 to 1.59)			
At least 2	107/3665 (2.9)	0.81 (0.51 to 1.30)			
Source of income§					
Sex work alone	74/2241 (3.3)	1	0.13		
Sex work and other job	73/2416 (3.0)	0.94 (0.68 to 1.31)			
No sex work	23/851 (2.7)	0.89 (0.54 to 1.45)			
Recruitment of clients§					
Bar, club, or restaurant	76/2717 (2.8)	1	0.01	1	0.004
Street	46/888 (5.2)	1.78 (1.23 to 2.57)		1.89 (1.27 to 2.80)	
Several avenues	48/1634 (2.9)	1.07 (0.75 to 1.54)		0.98 (0.68 to 1.42)	
Reproductive health					
Current contraceptive use§					
None or other	100/3160 (3.2)	1	0.001	1	<0.001
Oral cc	12/526 (2.3)	0.73 (0.40 to 1.33)		0.70 (0.38 to 1.28)	
Inject	56/1378 (4.1)	1.29 (0.93 to 1.80)		1.35 (0.96 to 1.88)	
Pregnant	2/397 (0.5)	0.16 (0.04 to 0.66)		0.13 (0.03 to 0.55)	
Behavioral characteristics					
No. of sexual partners§					
<5	77/2765 (2.8)	1	0.30		
5–19	32/1119 (2.9)	0.97 (0.64 to 1.49)			
At least 20 or cannot remember	61/1543 (4.0)	1.29 (0.90 to 1.84)			

(continued on next page)



**TABLE 3.** (Continued) Association of HIV Seroconversion With the Number of Missed Visits Between Consecutively Attended Visits Based on the Cox Regression Model

Characteristic	New HIV Cases/pyr (Rate/100 pyr)	Crude Hazard Ratio (95% CI)	P	Adjusted Hazard Ratio (95% CI)	P
No. of men the participant had paid sex within the last month§					
<5	79/2816 (2.8)	1	0.14	1	0.29
5–19	27/1089 (2.5)	0.84 (0.54 to 1.32)		0.70 (0.43 to 1.14)	
At least 20 or cannot remember	60/1522 (3.9)	1.29 (0.90 to 1.85)		0.96 (0.62 to 1.47)	
Condom use frequency with paid sex in the last month§					
Inconsistent	72/1616 (4.5)	1	0.01	1	0.002
Consistent (always)	55/2223 (2.5)	0.56 (0.40 to 0.80)		0.51 (0.35 to 0.74)	
No paid sex	43/1601 (2.7)	0.64 (0.43 to 0.96)		0.76 (0.48 to 1.20)	
Alcohol consumption frequency§					
Nondrinker	49/1870 (2.6)	1	0.30	1	0.41
Non-daily drinker	68/2202 (3.1)	1.17 (0.81 to 1.70)		1.17 (0.79 to 1.73)	
Daily drinker	53/1418 (3.7)	1.37 (0.92 to 2.04)		1.34 (0.87 to 2.07)	
Binge drinking in past 3 months§					
Nondrinker	48/1847 (2.6)	1	0.21		
No bingeing	58/1617 (3.6)	1.41 (0.95 to 2.07)			
Binged	64/2039 (3.1)	1.14 (0.78 to 1.67)			
Illicit drug use in the past 3 months§					
No	122/3949 (3.1)	1	0.34		
Yes	48/1260 (3.8)	1.18 (0.84 to 1.66)			
Sexually transmitted infections (STIs)§‡					
No STI	109/4033 (2.7)	1	0.01	1	0.12
With STI	61/1471 (4.1)	1.49 (1.09 to 2.05)		1.31 (0.93 to 1.85)	

Adjusted for: cohort of enrollment, age at follow-up in years, having a stable sexual partner (both married and unmarried), highest education level at enrollment, where clients are recruited from, current contraceptive use, condom use frequency with paid sex in the last month at follow-up, alcohol consumption frequency at follow-up, and presence of a sexually transmitted infection at follow-up.

\*Test for linear trend. Pyr—person-years of observation.

†The calendar period of January 2013–December 2014 was common to both cohorts (cohort-1 and cohort-2) because cohort-2 was initiated in January 2013 and was, therefore, used as the reference category.

‡STI variable was a composite variable combining data on STI symptoms from clinical examination and test results for STI for syphilis.

§Data collected at enrollment and follow-up study visits.

consecutive visits based on records of 342 HIV-negative women contacted between October and December 2015.

The social components of targeted HIV interventions such as peer outreach are not only critical in facilitating clinic attendance but are also associated with increased consistent condom use among FSWs.<sup>18,20</sup> Some of the participants with MSV might also have had limited engagement with peer educators as well as with outreach services, particularly participants working in isolation such as street or phone-based FSWs. Yet, our findings show that street-based FSWs were more likely to miss study visits and to acquire HIV infection than those who recruit clients from facilities such as bars or clubs. Among other reasons, street-based FSWs in this cohort as in other studies report a particularly higher risk of sexual or physical violence and exploitation by clients and law enforcers than venue-based FSWs.<sup>16,18,34</sup> Consequently, they are likely to be at a greater risk of STIs and HIV than those based in facilities. Furthermore, participants starting out in sex work are usually young and without social support networks to help them manage daily risks, including getting support for attending clinic visits.<sup>16</sup> These young participants are likely to be more isolated, more stigmatized, and to be highly vulnerable.<sup>34–37</sup> Therefore, targeted

programmes should work with potentially isolated groups of women at high risk of HIV (such as the highly mobile or young women) to develop better delivery models for services, which models could include scaling-up and intensifying mobile night clinics and intensifying peer outreach.<sup>20,38,39</sup>

### Behavioral Factors

Regular attendance of a dedicated clinic creates opportunities for reinforcing individual behavioral change messages among women at high risk of HIV. HIV prevention behavior can be achieved and sustained when participants are continually provided with appropriate information, are motivated to act, and enabled to obtain behavioral skills. These are key elements of the information–motivational–behavioral skill model that has been proposed for ART adherence but can also be applied for behavioral change among women at high risk of HIV.<sup>40</sup> In our study, trained counselors provide information on HIV prevention to participants through interactive sessions. Similar interactive sessions have been credited by FSWs in Zimbabwe for improving awareness of self-care as well as prevention and treatment of HIV and STIs.<sup>19</sup> Also, during

ongoing risk-reduction counseling sessions in our study, participants are motivated to improve or maintain healthy behaviors. This motivation may not only facilitate the adoption of healthy behaviors but also influence participants to regularly attend the clinic. Attending follow-up visits at the clinic was associated with higher parity, being on oral contraceptives, and having had an STI at the previous visit, which factors also reflect some underlying motivations related with the services offered at the clinic. According to the information–motivational–behavioral model, ongoing information on HIV prevention and motivation to stay healthy would act through acquired behavioral skills such as correct use of condoms to influence HIV incidence.

Therefore, participants who miss study visits are likely to also miss opportunities for reinforcing HIV risk-reduction behaviors. In addition, the HIV risk associated with MSV may increase where the participant's decision to attend the clinic is influenced by new intimate relationships. When participants start new intimate relationships, they might be afraid to associate with the clinic and they could be stopped by spouses from attending the dedicated clinic. In participants who missed  $\geq 2$  consecutive visits, 10% reported the reason as being in a new stable relationship or being stopped by a spouse. In this cohort as elsewhere, it is not uncommon for some of these new relationships to involve paying clients who transitioned into regular partnerships.<sup>16,34</sup> These partnerships have not only been associated with limited condom use which is usually seen as a sign of mistrust within intimate relationships across Africa but also with high HIV burdens.<sup>16,41</sup>

## Biomedical Factors

Regular attendance of a dedicated clinic enables timely access to biomedical interventions for prevention, treatment, and care. For example, 18% of all syndrome-based STIs might not have been detected if FSWs had not attended the clinic in a large study in Southern India on the impact of peer outreach on clinic utilization reported.<sup>20</sup> Similarly, the clinical examination at our study clinic is likely to facilitate earlier diagnosis and treatment of STIs for those regularly attending the clinic. As a result, earlier treatment would reduce the efficiency of STIs for HIV transmission.

Despite the substantial decline in HIV incidence after enrollment that was recently observed in this cohort, the incidence remained high and persisted over time.<sup>26</sup> HIV incidence also remained high during episodes with no MSV, although it was substantially lower in these episodes compared with those with at least 3 MSV. This persistently high HIV incidence despite combined HIV-prevention efforts may suggest presence of potential barriers in HIV control among women at high risk of HIV in this cohort. Among other factors, HIV control may be hindered by common barriers reported across studies in Africa such as accepting offers of sex without condoms for more money, hazardous alcohol use, and sexual and physical violence among women at high risk of HIV.<sup>18,34,42–44</sup> To further interrupt the spread of HIV through sex work, there is a need to strengthen targeted HIV-prevention efforts by integrating promising direct biomedical interventions such as pre-exposure prophylaxis, strengthening

risk-reduction interventions for substance abuse, and perhaps implementing mental health interventions.<sup>45–48</sup> These individual-level interventions should be further supported by structural interventions such as community empowerment that ensure targeted services are supplied and delivered to all women at high risk of HIV. Community empowerment conceptualized as an active involvement of sex workers, and the community in targeted activities could also reduce on the interconnected HIV risk factors for social stigmas, discrimination, and violence.<sup>6,8,49</sup> More importantly, there is need to monitor and strengthen regular engagement between women at high risk of HIV and service providers including their attendance at clinic services in the design and implementation of HIV-prevention interventions. Intensified peer outreach is associated with improved clinic utilization among FSWs and may be particularly important among isolated groups such as street-based FSWs.<sup>19–21</sup>

We note some key limitations. The measurement of MSV and seroconversion within the same episode could lead to some misclassification for the timing of seroconversions in the episodes with MSV if seroconversion occurred earlier before the missed visits. However, we assigned the timing of seroconversion as a random date rather than the commonly used midpoint to reduce on the potential for overestimation of HIV risk during episodes with MSVs.<sup>25,26</sup> Also, the number of events per predictor variable (EPV) in the multivariable model was relatively low (8.5) compared with the recommended minimum EPV of 10.<sup>50</sup> Although this low EPV can affect model-based inferences, the strong evidence for the association between MSV and HIV incidence in both the simple and multivariable models suggests that our conclusions could be valid. Similarly, data on some important risk factors such as duration of sex work were not collected, meaning this association could be confounded. Furthermore, the study did not fully record reasons for MSV and on a longitudinal basis, which would have facilitated better understanding of what underlies the HIV risk associated with MSV.

In conclusion, the need for scaling-up targeted combination HIV-prevention and treatment interventions for FSWs has been emphasized, but this study stresses the importance of monitoring, facilitating, and sustaining exposure to these services once women at high risk of HIV have been linked to services. Sustained exposure reinforces HIV prevention and facilitates earlier initiation of ART among those recently infected with HIV. Therefore, HIV programmes should include moving a wider scope of services to HIV-vulnerable women who are unable or unwilling to attend clinic services, particularly those working outside of established venues such as street-based FSWs.

## REFERENCES

1. Rinaldi G, Kiadaliri AA, Haghparast-Bidgoli H. Cost effectiveness of HIV and sexual reproductive health interventions targeting sex workers: a systematic review. *Cost Eff Resour Alloc*. 2018;16:63.
2. Vassall A, Pickles M, Chandrashekar S, et al. Cost-effectiveness of HIV prevention for high-risk groups at scale: an economic evaluation of the Avahan programme in south India. *Lancet Glob Health*. 2014;2:e531–e540.

3. Steen R, Wheeler T, Gorgens M, et al. Feasible, efficient and necessary, without exception—working with sex workers interrupts HIV/STI transmission and brings treatment to many in need. *PLoS One*. 2015;10:e0121145.
4. Muldoon KA. A systematic review of the clinical and social epidemiological research among sex workers in Uganda. *BMC Public Health*. 2015;15:1.
5. *Uganda Population-Based HIV Impact Assessment: 2016–2017*. Columbia University; 2018. Available at: [https://phia.icap.columbia.edu/wp-content/uploads/2018/07/3430%E2%80%A2PHIA-Uganda-SS\\_NEW\\_v14.pdf](https://phia.icap.columbia.edu/wp-content/uploads/2018/07/3430%E2%80%A2PHIA-Uganda-SS_NEW_v14.pdf). Accessed February 12, 2018.
6. Bekker LG, Johnson L, Cowan F, et al. Combination HIV prevention for female sex workers: what is the evidence? *Lancet*. 2015;385:72–87.
7. Chersich MF, Luchters S, Ntaganira I, et al. Priority interventions to reduce HIV transmission in sex work settings in sub-Saharan Africa and delivery of these services. *J Int AIDS Soc*. 2013;16:17980.
8. Kerrigan D, Kennedy CE, Morgan-Thomas R, et al. A community empowerment approach to the HIV response among sex workers: effectiveness, challenges, and considerations for implementation and scale-up. *Lancet*. 2015;385:172–185.
9. Shahmanesh M, Patel V, Mabey D, et al. Effectiveness of interventions for the prevention of HIV and other sexually transmitted infections in female sex workers in resource poor setting: a systematic review. *Trop Med Int Health*. 2008;13:659–679.
10. World Health Organization. *Consolidated Guidelines on HIV Prevention, Diagnosis, Treatment and Care for Key Populations*. Geneva, Switzerland: World Health Organization; 2014. Available at: <https://www.who.int/hiv/pub/guidelines/keypopulations/en/>. Accessed May 24, 2017.
11. Lafort Y, Greener R, Roy A, et al. Where do female sex workers seek HIV and reproductive health care and what motivates these choices? A survey in 4 cities in India, Kenya, Mozambique and South Africa. *PLoS One*. 2016;11:e0160730.
12. Nnko S, Kuringe E, Nyato D, et al. Determinants of access to HIV testing and counselling services among female sex workers in sub-Saharan Africa: a systematic review. *BMC Public Health*. 2019;19:15.
13. Ma PH, Chan ZC, Loke AY. The socio-ecological model approach to understanding barriers and facilitators to the accessing of health services by sex workers: a systematic review. *AIDS Behav*. 2017;21:2412–2438.
14. Lasater ME, Grosso A, Ketende S, et al. Characterising the relationship between migration and stigma affecting healthcare engagement among female sex workers in Lomé, Togo. *Glob Public Health*. 2019;1–14.
15. Davey C, Cowan F, Hargreaves J. The effect of mobility on HIV-related healthcare access and use for female sex workers: a systematic review. *Social Sci Med*. 2018;211:261–273.
16. Scorgie F, Chersich MF, Ntaganira I, et al. Socio-demographic characteristics and behavioral risk factors of female sex workers in sub-Saharan Africa: a systematic review. *AIDS Behav*. 2011;16:920–933.
17. Shannon K, Goldenberg SM, Deering KN, et al. HIV infection among female sex workers in concentrated and high prevalence epidemics: why a structural determinants framework is needed. *Curr Opin HIV AIDS*. 2014;9:174.
18. Shannon K, Strathdee SA, Goldenberg SM, et al. Global epidemiology of HIV among female sex workers: influence of structural determinants. *Lancet*. 2014;6736.
19. Ndori-Mharadze T, Fearon E, Busza J, et al. Changes in engagement in HIV prevention and care services among female sex workers during intensified community mobilization in 3 sites in Zimbabwe, 2011 to 2015. *J Int AIDS Soc*. 2018;21:e25138.
20. Krishnamurthy P, Hui SK, Shivkumar N, et al. Assessing the impact of peer educator outreach on the likelihood and acceleration of clinic utilization among sex workers. *PLoS One*. 2016;11:e0159656.
21. Luchters S, Chersich MF, Rinyiru A, et al. Impact of five years of peer-mediated interventions on sexual behavior and sexually transmitted infections among female sex workers in Mombasa, Kenya. *BMC Public Health*. 2008;8:143.
22. Steen R, Vuylsteke B, Decoito T, et al. Evidence of declining STD prevalence in a South African mining community following a core-group intervention. *Sex Transm Dis*. 2000;27:1–8.
23. Ghys PD, Diallo MO, Ettiegn-Traore V, et al. Effect of interventions to control sexually transmitted disease on the incidence of HIV infection in female sex workers. *AIDS*. 2001;15:1421–1431.
24. Graham SM, Raboud J, McClelland RS, et al. Loss to follow-up as a competing risk in an observational study of HIV-1 incidence. *PLoS One*. 2013;8:e59480.
25. Vandormael A, Dobra A, Bärnighausen T, et al. Incidence rate estimation, periodic testing and the limitations of the mid-point imputation approach. *Int J Epidemiol*. 2018;47:236–245.
26. Kasamba I, Nash S, Seeley J, et al. Human immunodeficiency virus infection attending a dedicated clinic in Kampala, Uganda: 2008–2017. *Sex Transm Dis*. 2019;46:407–415.
27. Vandepitte J, Bukonya J, Weiss HA, et al. HIV and other sexually transmitted infections in a cohort of women involved in high-risk sexual behavior in Kampala, Uganda. *Sex Transm Dis*. 2011;38:316–323.
28. Uganda Ministry of Health. *National HIV Testing Services Policy and Implementation Guidelines Uganda*. 2016. Available at: <http://health.go.ug/content/national-hiv-testing-services-policy-and-implementation-guidelines>. Accessed April 4, 2017.
29. World Health Organization. *Toolkit for Targeted HIV/AIDS Prevention and Care in Sex Work Settings*. Geneva, Switzerland: WHO; 2005.
30. Marshall A, Altman DG, Holder RL, et al. Combining estimates of interest in prognostic modelling studies after multiple imputation: current practice and guidelines. *BMC Med Res Methodol*. 2009;9:57.
31. Awungafac G, Delvaux T, Vuylsteke B. Systematic review of sex work interventions in sub-Saharan Africa: examining combination prevention approaches. *Trop Med Int Health*. 2017;22:971–993.
32. Prakash R, Bhattacharjee P, Blanchard A, et al. Effects of exposure to an intensive HIV-prevention programme on behavioural changes among female sex workers in Nairobi, Kenya. *Afr J AIDS Res*. 2018;17:99–108.
33. Seeley J, Nakiyingi-Miiri J, Kamali A, et al. High HIV incidence and socio-behavioral risk patterns in fishing communities on the shores of Lake Victoria, Uganda. *Sex Transm Dis*. 2012;39:433–439.
34. Mbonye M, Nakamanya S, Nalukenge W, et al. “It is like a tomato stall where someone can pick what he likes”: structure and practices of female sex work in Kampala, Uganda. *BMC Public Health*. 2013;13:741.
35. Lancaster KE, Cernigliaro D, Zulliger R, et al. HIV care and treatment experiences among female sex workers living with HIV in sub-Saharan Africa: a systematic review. *Afr J AIDS Res*. 2016;15:377–386.
36. Wanyenze RK, Musinguzi G, Kiguli J, et al. “When they know that you are a sex worker, you will be the last person to be treated”: perceptions and experiences of female sex workers in accessing HIV services in Uganda. *BMC Int Health Hum Rights*. 2017;17:11.
37. Dhana A, Luchters S, Moore L, et al. Systematic review of facility-based sexual and reproductive health services for female sex workers in Africa. *Glob Health*. 2014;10:46.
38. Lafort Y, Geelhoed D, Cumba L, et al. Reproductive health services for populations at high risk of HIV: performance of a night clinic in Tete province, Mozambique. *BMC Health Serv Res*. 2010;10:144.
39. Stadler J, Delany S. The “healthy brothel”: the context of clinical services for sex workers in Hillbrow, South Africa. *Cult Health Sex*. 2006;8:451–463.
40. Fisher JD, Fisher WA, Amico KR, et al. An information-motivation-behavioral skills model of adherence to antiretroviral therapy. *Health Psychol*. 2006;25:462.
41. Ngugi EN, Roth E, Mastin T, et al. Female sex workers in Africa: epidemiology overview, data gaps, ways forward. *SAHARA J*. 2012;9:148–153.
42. Beattie TS, Mohan HL, Bhattacharjee P, et al. Community mobilization and empowerment of female sex workers in Karnataka State, South India: associations with HIV and sexually transmitted infection risk. *Am J Public Health*. 2014;104:1516–1525.
43. Voeten HA, Egesah OB, Varkevissier CM, et al. Female sex workers and unsafe sex in urban and rural Nyanza, Kenya: regular partners may contribute more to HIV transmission than clients. *Trop Med Int Health*. 2007;12:174–182.
44. Robinson J, Yeh E. Transactional sex as a response to risk in Western Kenya. *Am Econ J Appl Econ*. 2011;3:35–64.
45. Deering KN, Amin A, Shoveller J, et al. A systematic review of the correlates of violence against sex workers. *Am J Public Health*. 2014;104:e42–e54.
46. McKinnon LR, Izulla P, Nagelkerke N, et al. Risk factors for HIV acquisition in a prospective Nairobi-based female sex worker cohort. *AIDS Behav*. 2015;19:2204–2213.

47. Li Q, Li X, Stanton B. Alcohol use among female sex workers and male clients: an integrative review of global literature. *Alcohol Alcohol*. 2010; 45:188–199.
48. Shahmanesh M, Wayal S, Cowan F, et al. Suicidal behavior among female sex workers in Goa, India: the silent epidemic. *Am J Public Health*. 2009;99:1239–1246.
49. Moore L, Chersich MF, Steen R, et al. Community empowerment and involvement of female sex workers in targeted sexual and reproductive health interventions in Africa: a systematic review. *Global Health*. 2014;10:47.
50. Vittinghoff E, McCulloch CE. Relaxing the rule of ten events per variable in logistic and Cox regression. *Am J Epidemiol*. 2007;165: 710–718.