

Perinatal or neonatal mortality among women who intend at the onset of labour to give birth at home compared to women of low obstetrical risk who intend to give birth in hospital: a systematic review and meta-analyses

Eileen K Hutton, PhD,^{1,2} Angela Reitsma, RM, MSc,² Julia Simioni, MSc,² Ginny Brunton, PhD,³ and Karyn Kaufman, DrPH²

1. Department of Obstetrics and Gynecology, Faculty of Health Sciences, McMaster University, Hamilton, Ontario, Canada
2. Midwifery Education Program, Faculty of Health Sciences, McMaster University, Hamilton, Ontario, Canada
3. EPPI-Centre, Department of Social Science, UCL Institute of Education, University College London, United Kingdom

Corresponding author: Professor Emerita Eileen K Hutton, McMaster University, 1280 Main Street West, HSC 4H24, Hamilton, Ontario, Canada, L8S 4K1, huttone@mcmaster.ca

Keywords: home childbirth, infant mortality, systematic review

Word Count: 3358/3000

SUMMARY (247/250)

Background: More women are choosing to birth at home in well-resourced countries. Concerns continue to be voiced that this out-of-hospital birth option contributes to higher perinatal and neonatal mortality. This systematic review and meta-analyses determine if risk of fetal or neonatal loss differs among low-risk women who begin labour intending to give birth at home compared to low-risk women intending to give birth in hospital.

Methods: In April 2018 we searched five databases from 1990 onward and used R to obtain pooled estimates of effect. We stratified by study design, study settings and parity. The primary outcome is any perinatal or neonatal death after the onset of labour. The study protocol is peer-reviewed, published and registered (PROSPERO No.CRD42013004046).

Findings: We identified 14 studies eligible for meta-analysis that included ~500,000 intended home births. Among nulliparous women intending a home birth in settings where midwives attending home birth are well-integrated in health services, the odds ratio (OR) of perinatal or neonatal mortality compared to those intending hospital birth was 1.07 (95% Confidence Interval [CI], 0.70 to 1.65); and in less integrated settings 3.17 (95% CI, 0.73 to 13.76). Among multiparous women intending a home birth in well-integrated settings, the estimated OR compared to those intending a hospital birth was 1.08 (95% CI, 0.84 to 1.38); and in less integrated settings was 1.58 (95% CI, 0.50 to 5.03).

Interpretation: The risk of perinatal or neonatal mortality was not different when birth was intended at home or in hospital.

Funding: Partial funding was received through an Association of Ontario Midwives open peer reviewed grant.

PANEL: RESEARCH IN CONTEXT

Evidence before this study

Although there is increasing acceptance for intended home birth as a choice for birthing women, controversy about its safety persists. The varying responses of obstetrical societies to intended home birth provide evidence of contrasting views. A Cochrane review of randomised controlled trials addressing this topic included one small trial and noted that in the absence of adequately sized randomised controlled trials on the topic of intended home compared to intended hospital birth, a peer reviewed protocol be published to guide a systematic review and meta-analysis including observational studies. Reviews to date have been limited by design or methodological issues and none has used a protocol published a priori.

Added value of this study

Individual studies are underpowered to detect small but potentially important differences in rare outcomes. This study uses a published peer-reviewed protocol and is the largest and most comprehensive meta-analysis comparing outcomes of intended home and hospital birth. We take study design, parity and jurisdictional support for home birth into account. Our study provides much needed information to policy makers, care providers and women and families when planning for birth.

Implications of all the available evidence

Women who are low risk and who intend to give birth at home do not appear to have a different risk of fetal or neonatal loss compared to a population of similarly low risk women intending to give birth in hospital.

INTRODUCTION

Birth has become the most common reason for hospital admission in well-resourced countries impacting healthcare costs;¹ however, it is unclear if hospitalisation for birth alters neonatal outcomes for women at low obstetrical risk. A small but growing number of women begin labour with the intention of giving birth at home² and research among this self-selected group consistently reports reduced obstetric interventions^{3,4}. It is uncertain however, if this reduction comes at the expense of neonatal wellbeing. In 2015, for example, two high-profile studies of out-of-hospital (home) birth reported contradictory findings regarding perinatal mortality and morbidity.^{3,4}

Acceptance of home birth as a choice for women is increasing,⁵ but controversy about safety persists. Quality evidence regarding outcomes associated with place of birth for low-risk pregnancies is urgently needed to inform parents, maternity care providers and policy makers. Because individual studies are underpowered to detect small but potentially important differences in rare outcomes, and randomised controlled trials are not feasible and do not contribute to these findings, a Cochrane review on this topic urged a careful systematic review and meta-analysis of cohort studies be undertaken to evaluate outcomes of intended home birth.⁶ Using our peer-reviewed, published, registered protocol⁷ (PROSPERO, <http://www.crd.york.ac.uk>, No.CRD42013004046) we undertook this systematic review and meta-analyses to determine if low-risk women who intend at the onset of labour to give birth at home are more or less likely to experience a fetal or neonatal loss compared to a cohort of similarly low-risk women who intend at the onset of labour to give birth in hospital.

METHODS

Methods reported previously in our published protocol⁷ were followed and are described briefly here.

Search strategy and Study Selection

The search included studies from 1990 onward and was completed on April 11, 2018 using Embase, Medline, AMED, CINAHL, and the Cochrane Library. Terms either as keywords or subject headings included: home delivery, home birth, home childbirth, and homebirth. Reference lists from review articles and all included studies were crosschecked. Two reviewers independently selected studies for full review if they had comparison groups of women at similarly low-risk, as defined within the study, for birth complications intending either to give birth in hospital or home; cohorts were defined by the intended location of birth rather than the actual location of birth; intention for a home birth was determined or reconfirmed at the onset of labour; parity was accounted for; and the study accounted for missing cases.

Data collection

Two reviewers independently collected data from the included studies using a detailed data abstraction form, compared their findings and reached consensus. Missing information was requested from authors of included studies as necessary. Whenever possible, findings were reported by parity sub-groups.

Because they answer somewhat different questions, we categorised studies into one of two study designs to reflect the assembly of birth cohorts. In all cases the comparison group included women intending hospital birth and deemed to be at low obstetrical risk. Studies designed to determine the safety of home birth in actual practice, included all intended home births in a given time frame, regardless of whether they would be considered eligible for home birth according to local standards. These ‘pragmatic’ design studies answer the research question: “Do women who intend at the onset of labour to give birth at home experience a higher or lower incidence of fetal or neonatal loss compared to women at low obstetric risk who intend at the onset of labour to give birth in hospital?” Other studies focused on outcomes of place of birth among women who met local selection standards for home birth thus assuring that only those of low obstetrical risk were included and answer the question: “Do women who intend to give birth at home and who meet their local eligibility criteria for home birth at the onset of labour experience a higher or lower incidence of fetal or neonatal loss compared to women who would have been eligible for home birth but intend at the onset of labour to give birth in hospital?” The latter study design may have resulted in the exclusion of, for example, any twin births or breech births that may have been intended and occurred at home, but that were not supported by local standards. We termed studies of this design ‘within standards.’ We stratified all analyses by study design in order to address both research questions. In addition, in order not to compromise power to find small differences, we conducted sensitivity analyses for all outcomes without stratification as described in the Sensitivity Analyses section below.

We hypothesised a priori⁷ that the degree of support for home birth and home birth care providers within the health care system where the study was carried out would act as an effect modifier of the relationship between intended place of birth and birth outcomes.⁸ We termed this context for home birth, described in detail elsewhere, as a ‘well-integrated’ versus ‘less well-integrated’ home birth environment.⁹ A well-integrated setting was described as a place where home birth practitioners: are recognised by statute within their jurisdiction; have received formal training; can provide or arrange care in hospital; have access to a well-established emergency transport system; and carry emergency equipment and supplies. Less well-integrated settings were those where one or more of these criteria are absent. Studies were categorised by an independent team of researchers⁹ based on information found within the study, from the study’s author via a questionnaire about the context of care at the time their study was undertaken, and from secondary publications such as policies or statements regarding home birth in the country where the study took place (Table 1).

Outcome

Our primary outcome is any perinatal or neonatal death after the onset of labour. If a study reported these data both including and excluding malformed infants, to minimise categorization bias we used data that included malformations in the primary analyses. Secondary outcomes include perinatal mortality (defined as stillbirth after the onset of labour or death to 7 completed days) and neonatal mortality (defined as death between 0 and 28 days of a live born baby). Where possible, we report perinatal mortality and neonatal mortality separately; and mortality rate excluding malformed infants. Additional neonatal outcomes included neonatal resuscitation, Apgar scores of less than 7 at one minute and less than 7 at 5 minutes, and admission to a neonatal intensive care unit (NICU). Definitions used by the authors for neonatal resuscitation and NICU admissions were recorded.

Because free standing birth centres cannot be considered to be a home or hospital setting, data from these out of hospital birth centres were not included. For studies that had more than one hospital comparison group, outcomes for the hospital groups were combined, provided that women in the groups being combined met eligibility criteria. If data for some or all outcomes could not be combined, we chose the comparison group most likely to minimise confounders; where the women were most like women choosing home birth, and the care providers were most like those providing care at home.

Risk of bias

Our study eligibility criteria ensured that the observational studies included in the review had a control group, used an intention-to-treat approach (analysed by intended place of birth at the onset of labour), and controlled for parity. Study quality was assessed using The Newcastle Ottawa Quality Assessment Scale for Cohort Studies (NOS).¹⁰ Risk of publication bias across studies was assessed through inspection of inverted funnel plots for the primary outcome.¹¹

Synthesis of results

Meta-analyses were conducted using the ‘metafor’ package in R statistical software version 3.3.1. Log odds ratios (OR) and corresponding sampling variances for each study were calculated using count data or ORs and confidence intervals. For studies using the home birth group as the reference category, the OR was first inverted. For studies that reported risk ratios, if count data were provided and it was possible to calculate an OR, this was done. If the adjusted risk ratio was the only method by which parity was accounted for, we were unable to combine these data with ORs from other studies. In this case, risk ratios were described separately. Data were then pooled by fitting a random-effects model and forest plots were created. Pooled ORs, 95% confidence intervals and measures of consistency (I^2) were calculated for each outcome within strata (Table 1).

Sensitivity analyses

We conducted a sensitivity analysis of our primary outcome excluding large datasets in order to determine if findings remained robust without those studies. We conducted sensitivity analyses for all outcomes without stratification by study design (pragmatic or within standards) to ensure that the smaller sample sizes resulting from stratification did not limit power and bias our analyses toward finding no difference.

Role of the funding source

The funder of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report. The corresponding author had full access to all the data in the study and had final responsibility for the decision to submit for publication.

RESULTS

The search, completed on April 11, 2018, provided 139 full text articles for review (Figure 1) and resulted in 23 cohort studies that met our predefined inclusion criteria for systematic review of intended place of birth. Two of these studies^{12,13} were excluded because they reported on data duplicated in other included studies and four studies because they did not include perinatal or infant outcomes.^{14–17} Of 17 studies eligible for systematic review of perinatal or infant outcomes, three provided no data either published or from study authors^{18–20} that could be included in a meta-analysis. Thus, the meta-analyses included 14 original cohort studies published between 1996 and 2017 that reported perinatal or neonatal outcomes for ~500,000-intended home births (Table 2). The precise number of births varies by analysis depending on the inclusion of one or the other of the large Dutch papers where there is likely considerable overlap in data. The included studies scored from 4 to 8 on the NOS. No randomised trials were found that included the outcomes of interest. A table of primary research studies excluded from this review can be found in Appendix 1.

Five included studies had more than one comparator group.^{20–23, 34} For the study by the Birthplace in England Collaborative Group, because all the women included were low obstetrical risk, we combined outcomes of the midwifery alongside unit (an in-hospital birthing unit) and the obstetrical unit.²³ For a variety of reasons the multiple hospital comparison groups in Janssen's papers could not be combined^{21,22} therefore we used the physician-attended hospital comparison group. For Davis et al, we used the primary unit comparison group.²⁰ For the paper by Homer et al we included home and hospital groups and excluded the birth centre group because it may have included out of hospital birth centres.

Integration and study design

The 17 studies included in the systematic review (14 included in the meta-analyses) described here took place in ten settings, which are illustrated in Table 1. Thirteen studies took place in six settings where midwives attending home birth were considered to be well-integrated into the

healthcare system (The Netherlands, England, Iceland, Canada, USA, New Zealand).^{18–30} Four studies took place in four settings where midwives attending home birth were considered to be less well-integrated into the healthcare system (Norway, Sweden, Japan, Australia)^{31–34} as described elsewhere.⁹ A pragmatic study design was used by ten studies,^{19,21,22,24,27–32} whereas seven studies included only those women who met local standards for home birth in their intended home birth cohorts.^{18,20,23,25,26,33,34}

Synthesis of results

There was no difference in the primary outcome between those who intended home and those who intended hospital birth when data from 13 studies were pooled (Figure 2 and Table 3). Among ten studies where midwives and homebirth were deemed to be well-integrated into the healthcare system, six reported results for all women (after accounting for parity through statistical adjustment or matching), and all used a pragmatic study design. The pooled OR (OR) was 0.89 (95% CI, 0.78 to 1.02). A sensitivity analyses that removed a large Dutch study (weighted 94.7% for this outcome),³⁰ remained non-significant for the primary outcome by intended place of birth for all women (after accounting for parity) (OR 1.16 (95% CI, 0.65 to 2.05)). Two studies reported findings as risk ratios and thus could not be included in any meta-analyses. The first, by van der Kooy et al¹⁹ reported outcomes similar to our pooled OR with a risk ratio (RR) of 0.80 (95% CI, 0.71 to 0.91). The second by Pang et al used a within standards design and reported a RR of 1.99 (95% CI, 1.06 to 3.73), which favoured intended hospital birth.¹⁸

Seven studies reported results stratified by parity in settings where midwives attending home birth were considered well-integrated into the healthcare system (three used a within standards design and four a pragmatic design). Regardless of study design and parity, no difference was found for the primary outcome between women who intended a home birth and those who intended a hospital birth. The pooled OR for nulliparous women was 1.07 (95% CI, 0.70 to 1.65) and for multiparous women 1.08 (95% CI, 0.84 to 1.38). The study by Pang et al reported a risk ratio for nulliparous women which favoured hospital birth (RR 2.73 (95% CI, 1.06 to 7.06)) but failed to report outcomes for multiparous women.¹⁸ Sensitivity analyses did not find statistically significant differences between intended home and hospital birth among nulliparous or multiparous women after excluding a large Dutch study (nulliparas OR 3.17 (95% CI, 0.73 to 13.76); multiparas OR 1.58 (95% CI, 0.50 to 5.03)).²⁵

In settings where home birth was less well-integrated into the health care system, two studies used a pragmatic design and one a within standards design. All three studies reported results stratified by parity. The pooled ORs for all three studies did not show a statistically significant difference between intended home and intended hospital birth regardless of parity (nulliparas OR 1.25 (95% CI, 0.58 to 2.67); multiparas OR 1.62 (95% CI, 0.78 to 3.35)).

We did not find any significant differences by intended place of birth in mortality outcomes exclusive of morbidity that is for: perinatal or neonatal mortality excluding malformed infants; perinatal or neonatal mortality including malformed infants; perinatal mortality; neonatal mortality (Table 3). These findings held true regardless of parity, degree of integration of midwives providing home birth care or study design.

An Apgar score less than seven at five minutes occurred less frequently among intended home births compared to hospital births in multiparas in settings where midwives were well-integrated (OR 0.76 (95% CI, 0.60 to 0.96)). This difference was not found among nulliparous women or when results were not stratified by parity. In addition, the study by Davis et al, that could not be combined due to use of risk ratios, reported no significant difference (for all women).²⁰ Two studies from settings where midwives were less well-integrated into the healthcare system reported this outcome, but the findings could not be combined in meta-analyses. Neither found a significant difference (Table 4).

An Apgar score less than seven at one minute was reported by two studies.^{21,22} Both took place in settings where midwives attending home birth were well-integrated and used a pragmatic design and neither stratified by parity. A meta-analysis of these studies found a statistically significant difference in favour of intended home birth compared to intended hospital birth (OR 0.73 (95% CI 0.63 to 0.83)) (Table 4).

Need for neonatal resuscitation was reported in five studies,^{21,22,27-29} where midwives attending home birth were well-integrated and where a pragmatic study design was used. A meta-analysis found no significant difference by intended place of birth for all women. Only one study stratified by parity and reported no difference amongst parity groups on need for neonatal resuscitation (Table 4).

Of seven studies that reported on NICU admissions all were settings where midwives attending home birth were well-integrated, but one could not be combined in meta-analysis.²⁰ Among nulliparous women, no difference in NICU admissions was found by intended place of birth, regardless of study design. Studies using a within standards study design reported fewer infant admissions to NICU born to multiparous women who intended home birth (OR 0.73 (95% CI, 0.65 to 0.83)). This finding held true when these studies were meta-analysed with the pragmatic studies (OR 0.73 (95% CI, 0.65 to 0.82)). This difference was also seen when stratification for parity was removed and all women were included (OR 0.83 (95% CI, 0.69 to 0.99)) (Table 4). Davis et al. reported no difference for NICU admission among all women by intended place of birth (RR 1.00 (95% CI, 0.66 to 1.50)).²⁰

Risk of bias across studies

Inverted funnel plots were created to assess for reporting bias across studies for our primary outcome, one for each strata of analysis, resulting in five plots. However, plots with fewer than ten studies are difficult to interpret¹¹ and the largest of our plots included only seven studies (Appendix 2).

DISCUSSION

This is the first systematic review and meta-analysis that has used a peer-reviewed, pre-published, registered protocol. Our results show that low risk women who intend to give birth at home when labour starts do not experience any increase in perinatal and neonatal mortality or morbidity compared to similarly low risk women who intend to give birth in a hospital. In order to fully understand any effects of intended place of birth and ensure that we are not missing any important pregnancy outcome related to mortality, we considered perinatal and neonatal mortality together as our primary outcome. In addition, combining these outcomes increases the number of events thus increasing the power of our study to determine any clinically important differences. Although jurisdictional variation exists in terms of how perinatal mortality is defined, the differences arise primarily in how the lower limits of viability are determined. Because our study included only those with low-risk pregnancies at the onset of labour, those with pre-term and in particular extreme preterm births are eliminated from the population under study. Furthermore, even with differences in definition of perinatal mortality between studies, the definition used within each study will be applied similarly to all study participants in that study thereby eliminating the likelihood of bias between the comparison groups in our study.

Other reviews have limited the scope of their review to deal with design differences or do not account for parity.³⁵ In our study we take a more inclusive approach and have included all relevant studies, dealing with differences in design by stratifying according to the approach taken, which we termed either pragmatic or within standards. In order to provide the most comprehensive understanding of the impact of choice of birthplace on perinatal outcomes, we present data both stratified and combined wherever possible to indicate where differences exist. Research design (pragmatic and within standards) did not have a significant impact on the results as findings were similar for the sub-set of women who met local criteria for choosing home birth, and for studies that included all women who intended a home birth, and thus may have included some who were non-compliant with local selection criteria. Outcomes were similar among nulliparous and multiparous women, and in settings where midwives providing home birth care were well-integrated into healthcare systems and where they were less well-integrated, although the data from integrated settings is more robust. There were no differences between intended home and intended hospital groups in other neonatal outcomes including NICU admission, Apgar scores, and the need for resuscitation. Although criteria may have varied across studies to determine NICU admission there should be no within study variance, thus providing confidence in our finding that there was no increased need for higher level care with births planned at home.

Select sub-groups showed results favouring home birth. The one exception to the primary outcome findings lies with the sole American study included in the review.¹⁸ This study could not be included in the meta-analyses; however, the findings indicated a significant increase in perinatal and neonatal mortality associated with intended home birth.

We applied pre-specified criteria to eligible studies to determine how well midwives attending home births were integrated into the health care system in that jurisdiction and presented this information in a separate publication.⁹ This categorisation was carefully undertaken using data from the studies, from questionnaires completed by authors of the included studies regarding care at the time that their study was completed as well as looking at standards and other jurisdictional documents. The highest quality studies came from large registries that were used in places where midwives providing home birth care were well-integrated into the health care system. Many home birth studies were ineligible for this review due to issues in study methodology such as: not including a control group of low-risk hospital births from the same region and time frame as the home births; not controlling for parity using matching, adjustment, or stratification; and excluding intended home births that transferred to hospital in labour from the home birth cohort thus potentially underestimating adverse outcomes. A table describing excluded studies provides transparency in our exclusion criteria (Appendix 1). We found that home birth studies occurring in less well-integrated settings were more often excluded perhaps due to inferior data collection practices. This could result in an unavoidable bias towards excluding studies most likely to have untoward outcomes.

Having fewer quality studies from less well-integrated settings resulted in loss of power and findings that were less precise and may introduce an unavoidable bias. Although we found no difference in mortality outcomes for intended home versus hospital births in less well-integrated settings there was a trend towards favouring hospital birth that is, perhaps, of interest. Generalisability of our findings should, therefore, be undertaken cautiously. We challenge readers to interpret the safety of home birth within a greater societal context and consider the integration of home birth practices within health care systems.

CONTRIBUTORS

All authors formulated the research objectives and designed the study methodology. GB conducted the literature search. AR and JS screened citations for eligibility and extracted data from eligible studies. JS conducted the data analyses. EKH, AR and JS prepared the initial manuscript draft. All authors edited the manuscript and approved the final version.

DECLARATION OF INTERESTS

We declare no competing interests.

ACKNOWLEDGEMENTS

This study was funded in part by a Mentored Midwifery Research Grant to Ms. Angela Reitsma from the Association of Ontario Midwives.

TABLES AND FIGURES

Table 1. Studies eligible for systematic review of perinatal and neonatal outcomes, stratified by degree of integration of home birth within the health care system and by study design.

		TYPE OF INTEGRATION INTO HEALTH SYSTEM	
		Well-integrated	Less well-integrated
STUDY DESIGN	Pragmatic (all women who intend home birth)	Halfdandottir 2015 ²⁷ Hutton 2009 ²⁸ Hutton 2015 ²⁹ Janssen 2002 ²¹ Janssen 2009 ²² van der Kooy 2011 ³⁰ van der Kooy 2017 ¹⁹ Wiegers 1996 ²⁴	Blix 2012 ³¹ Lindgren 2008 ³²
	Within standards (only women who meet criteria for birth at home)	Brocklehurst 2011 ²³ Davis 2011 ²⁰ de Jonge 2014 ²⁵ Hermus 2017 ²⁶ Pang 2002 ¹⁸	Hiraizumi 2013 ³³ Homer 2014 ³⁴

Figure 1. Flow diagram of study selection




Table 2. Description of included studies

Study	Data source & Time period	Method of accounting for parity	NOS Quality Score	Methods	Sample size	Setting	Outcomes Reported	Author questionnaire completed
Blix E, et al. 2012 ³¹	Home: Midwife's register, telephone interview, and midwife's birth protocols Hospital: Medical birth registry of Norway (MBRN) 1990-2007	Stratified	6	Pragmatic Retrospective cohort study	1631 home 16310 hospital	Norway (Midwives less well-integrated)	<u>1, 3-5, 8,</u> 11, 12, 15-18	yes
Birthplace in England Collaborative Group, 2011 ²³	Home: All NHS Trusts that provide home birth services OU: Random sample of 36 obstetric units within the NHS	Stratified and adjusted	7	Within standards Prospective cohort study 4 groups: Obstetric Unit, Alongside Midwifery Unit, Free-standing	16840 home 16710 ALU 19706 OU 11282 FSU Combined ALU and OU for comparison group	England (Midwives well-integrated)	<u>1, 2, 4, 8, 9,</u> 11, 12, 14-18	yes

Study	Data source & Time period	Method of accounting for parity	NOS Quality Score	Methods	Sample size	Setting	Outcomes Reported	Author questionnaire completed
	ALU: All NHS hospitals that have an alongside unit Data collection forms designed for this study 2008-2010			birth centre, Home				
Bolten N, et al. 2016 ¹⁴	DELIVER Study, recruited from 20 midwifery practices 2009-2011	Stratified	6	Within standards Prospective cohort study	2050 home 1445 hospital	Netherlands (Midwives well-integrated)	11, 12, 14, 16-18 No infant outcomes	yes
Davis D, et al. 2011 ²⁰	Midwifery Maternity Provider Organization Database 2006-2007	Adjusted	8	Within standards Retrospective cohort study	1830 home Primary unit 2877 Secondary hospital 7380 Tertiary hospital 4123 Used primary unit	New Zealand (Midwives well-integrated)	<u>8, 9</u> , 11, 14, 16, 18	no

Study	Data source & Time period	Method of accounting for parity	NOS Quality Score	Methods	Sample size	Setting	Outcomes Reported	Author questionnaire completed
					comparison group			
de Jonge A, et al. 2013 ¹⁵	LEMMoN Study database, National Perinatal database I, National Perinatal database II, National Neonatal Register 2004-2006	Stratified	8	Within standards Prospective cohort study	92333 home 54419 hospital	Netherlands (Midwives well-integrated)	11 No infant outcomes	yes
de Jonge A, et al. 2014 ²⁵	National Perinatal database I, National Perinatal database II, National Neonatal Register	Stratified	7	Within standards Retrospective cohort study	466112 home 276958 hospital 2000-2009	Netherlands (Midwives well-integrated)	<u>1,2,4,5,8,9</u>	yes

Study	Data source & Time period	Method of accounting for parity	NOS Quality Score	Methods	Sample size	Setting	Outcomes Reported	Author questionnaire completed
	2000-2009							
Halfdansdottir B, et al. 2015 ²⁷	Icelandic electronic birth registry and original midwife and doctor records extracted by study author using a structured item list. 2005-2009	Matched and Stratified	7	Pragmatic + Within standards Retrospective cohort study	307 home 921 hospital	Iceland (Midwives well-integrated)	<u>1,3,4,6,8-</u> 11,14-18	yes
Hermus M, et al. 2017 ²⁶	Midwifery practices using case report form developed for the study and linked with the Netherlands Perinatal Registry (Perined)	Stratified	6	Within standards Prospective cohort study	1086 home 701 hospital	Netherlands (Midwives well-integrated)	<u>1, 3, 4, 9-</u> 12, 14-18	yes

Study	Data source & Time period	Method of accounting for parity	NOS Quality Score	Methods	Sample size	Setting	Outcomes Reported	Author questionnaire completed
	2013							
Hiraizumi Y, et al. 2013 ³³	Japanese Red Cross Katsushika Maternity Hospital database 2007-2011	Presumed matched, equal proportion in groups by parity	7	Within standards Retrospective cohort study	168 home 123 hospital	Japan (Midwives less well-integrated)	<u>8</u> ,11-14,17,18	no
Homer C, et al. 2014 ³⁴	5 datasets in New South Wales. NSW Perinatal data collection NSW admitted patient data collection NSW register of congenital conditions NSW registry of births, deaths, and marriages Australian Bureau of	Stratified	7	Within standards Retrospective cohort study	735 home 221284 hospital 2000-2008 (birth centre outcomes excluded)	Australia (Midwives less well-integrated)	<u>1,2,4</u>	yes

Study	Data source & Time period	Method of accounting for parity	NOS Quality Score	Methods	Sample size	Setting	Outcomes Reported	Author questionnaire completed
	Statistics 2000-2008							
Hutton EK, et al. 2009 ²⁸	Ontario Midwifery Program dataset 2003-2006	Matched, Stratified	8	Pragmatic Retrospective cohort study	6692 home 6692 hospital 2003-2006	Ontario, Canada (Midwives well-integrated)	<u>1-3,5,6,8</u> -12, 14-18	yes
Hutton EK, et al. 2015 ²⁹	Ontario Midwifery Program dataset 2006-2009	Matched, Stratified	8	Pragmatic Retrospective cohort study	11493 home 11493 hospital	Ontario, Canada (Midwives well-integrated)	<u>1-6,8</u> ,10-12, 14-18	yes
Janssen P, et al. 2002 ²¹	Home: Home Birth Demonstration Project Hosp: British Columbia Perinatal Database Registry	Matched, Adjusted	6	Pragmatic Prospective and Retrospective cohort study	862 home 571 MW comparison 743 MD comparison Used MD comparison group	British Columbia, Canada (Midwives well-integrated)	<u>1,2,6-8</u> ,11-18	yes

Study	Data source & Time period	Method of accounting for parity	NOS Quality Score	Methods	Sample size	Setting	Outcomes Reported	Author questionnaire completed
	1998-1999							
Janssen P, et al. 2009 ²²	Home: BC Perinatal Database Registry + Rosters submitted to the College of Midwives of BC Hosp: BC Perinatal Database Registry 2000-2004	Matched, Adjusted	6	Pragmatic Retrospective cohort study	2899 home 4752 MW comparison 5331 MD comparison Used MD Comparison group	British Columbia, Canada (Midwives well-integrated)	1,2,4,6-8, 10-18	yes
Lindgren H, et al. 2008 ³²	Home: Home birth midwives reports, linked to Swedish Medical Birth Register Hosp: Swedish Medical Birth Register	Adjusted	6	Pragmatic Retrospective cohort study	897 home 11341 hospital	Sweden (Midwives less well-integrated)	1,2,4, 10-12, 16, 18	yes

Study	Data source & Time period	Method of accounting for parity	NOS Quality Score	Methods	Sample size	Setting	Outcomes Reported	Author questionnaire completed
	1992-2004							
Miller S, et al. 2012 ¹⁶	Midwives who chose to participate and report on their most recent nulliparous births. Not reported	Restricted to nulliparous	4	Within standards Retrospective cohort study	109 home 116 hospital	New Zealand (Midwives well-integrated)	11,12,14-18 No infant outcomes	yes
Nove A, et al. 2012 ¹⁷	St. Mary's Maternity Information System 1988-2000	Adjusted	8	Within standards Retrospective cohort study	5998 home 267874 hospital	England (Midwives well-integrated)	11 No infant outcomes	yes
Pang J, et al. 2002 ¹⁸	Washington State birth certificate data 1989-1996	Adjusted, Stratified	4	Within standards Retrospective cohort study	6133 home 10593 hospital	Washington state, USA (Midwives well-integrated)	<u>1,3,11</u> Data not available for meta-analysis	no
van der Kooy J, et al. 2011 ³⁰	Netherlands Perinatal Registry 2000-2007	Adjusted	7	Pragmatic Retrospective cohort study	402912 home 219105 hospital	Netherlands (Midwives well-integrated)	<u>1,2,4</u>	yes

Study	Data source & Time period	Method of accounting for parity	NOS Quality Score	Methods	Sample size	Setting	Outcomes Reported	Author questionnaire completed
van der Kooy J, et al. 2017 ¹⁹	Netherlands Perinatal Registry 2000-2007	Adjusted		Pragmatic Retrospective cohort study	402912 home 219105 hospital	Netherlands (Midwives well-integrated)	1 , 18 Data not available for meta-analysis	yes
Wiegers TA, et al. 1996 ²⁴	Questionnaires and the Birth Notification System 1990-1993	Stratified	6	Pragmatic Prospective and Retrospective cohort study	1140 home 696 hospital	Netherlands (Midwives well-integrated)	1,3,4,9,11,16-18	yes


Outcomes reported by included studies are listed in the table as follows. Outcomes reported in this manuscript are **bolded and underlined** in the table.

1. Any perinatal or neonatal mortality
2. Perinatal or neonatal mortality excluding malformations
3. Perinatal or neonatal mortality including malformations
4. Any perinatal mortality
5. Any neonatal mortality
6. Neonatal Resuscitation
7. Apgar <7 at 1 minute
8. Apgar <7 at 5 minutes
9. Admission to NICU
10. Maternal mortality
11. Postpartum hemorrhage


12. 3rd or 4th degree tear
13. Maternal infection
14. Oxytocin augmentation
15. Epidural
16. Episiotomy
17. Assisted vaginal delivery
18. Caesarean section

Figure 2. Forest plots showing meta-analyses for the primary outcome of perinatal or neonatal mortality.

a)




b)




c)

Perinatal or Neonatal Mortality: Midwives Integrated Setting, Not Stratified by Parity




d)

Perinatal or Neonatal Mortality: Midwives Not Well Integrated Setting, Nulliparous Women



e)

Perinatal or Neonatal Mortality: Midwives Not Well Integrated Setting, Multiparous Women



Note that for the Birthplace in England Collaborative Group study (2011), the hospital comparison group included data from the obstetrical unit and alongside midwifery unit. For the studies by Janssen et al (2002 and 2009), the physician-attended hospital group was used as the comparison.

Table 3. Summary of perinatal and neonatal mortality meta-analyses findings derived from Figure 2.

Strata	Number of studies	OR	95% CI	I ²
Primary Outcome: Perinatal or Neonatal Mortality (any)*†				
Midwives Well-integrated Setting				
Nulliparas	7	1.07	0.70, 1.65	18.9%
Within standards	3 ^{23,25,26}	1.30	0.47, 3.55	62.6%
Pragmatic	4 ^{24,27-29}	0.93	0.43, 1.99	0%
Multiparas	7	1.08	0.84, 1.38	0%
Within standards	3 ^{23,25,26}	1.04	0.80, 1.35	0%
Pragmatic	4 ^{24,27-29}	1.64	0.67, 4.04	0%
Not stratified by parity	-	-	-	-
Pragmatic	6 ^{21,22,27-30}	0.89	0.78, 1.02	0%
Midwives Less well-integrated Setting				
Nulliparas	3	3.17	0.73, 13.76	0%
Within standards	1 ³⁴	1.86	0.12, 29.97	n/a
Pragmatic	2 ^{31,32}	3.90	0.69, 21.95	0%
Multiparas	3	1.58	0.50, 5.03	0%
Within standards	1 ³⁴	1.41	0.09, 22.83	0%
Pragmatic	2 ^{31,32}	1.62	0.45, 5.78	0%
Outcome: Perinatal or Neonatal Mortality (excluding malformed infants)				
Midwives Well-integrated Setting				
Nulliparas	4	1.17	0.70, 1.97	41.0%
Within standards	2 ^{23,25}	1.52	0.48, 4.85	79.1%
Pragmatic	2 ^{28,29}	1.00	0.45, 2.23	0%
Multiparas	4	1.08	0.83, 1.40	0.5%
Within standards	2 ^{23,25}	1.04	0.80, 1.35	0%
Pragmatic	2 ^{28,29}	1.80	0.60, 5.37	0%
Not stratified by parity	-	-	-	-
Pragmatic	4 ^{21,22,28,29}	1.20	0.66, 2.18	0%
Midwives Less well-integrated Setting				
Nulliparas	-	-	-	-
Within standards	1 ³⁴	1.86	0.12, 29.97	n/a
Multiparas	-	-	-	-
Within standards	1 ³⁴	1.41	0.09, 22.83	n/a
Outcome: Perinatal or Neonatal Mortality (including malformed infants)*†				
Midwives Well-integrated Setting				
Nulliparas	-	-	-	-
Pragmatic	3 ^{24,27,29}	0.80	0.31, 2.03	0%
Multiparas	-	-	-	-
Pragmatic	3 ^{24,27,29}	1.52	0.53, 4.39	0%

Not stratified by parity	-	-	-	-
Pragmatic	4 ²⁷⁻³⁰	0.89	0.78, 1.01	0%
Midwives Less well-integrated Setting				
Nulliparas	-	-	-	-
Pragmatic	2 ^{31,32}	3.90	0.69, 21.95	0%
Multiparas	-	-	-	-
Pragmatic	2 ^{31,32}	1.62	0.45, 5.78	0%
Outcome: Perinatal Mortality†				
Midwives Well-integrated Setting				
Nulliparas	5	1.22	0.65, 2.27	39.1%
Within standards	2 ^{23,25}	1.52	0.48, 4.83	79.0%
Pragmatic	3 ^{24,27,29}	1.00	0.37, 2.70	0%
Multiparas	5	1.07	0.82, 1.38	0%
Within standards	2 ^{23,25}	1.06	0.82, 1.38	0%
Pragmatic	3 ^{24,27,29}	1.27	0.31, 5.27	0%
Not stratified by parity	-	-	-	-
Pragmatic	4 ^{22,27,29,30}	0.88	0.77, 1.01	0%
Midwives Less well-integrated Setting				
Nulliparas	3	3.58	0.82, 15.64	0%
Within standards	1 ³⁴	1.86	0.12, 29.97	n/a
Pragmatic	2 ^{31,32}	4.62	0.81, 26.37	0%
Multiparas	3	1.34	0.30, 5.91	0%
Within standards	1 ³⁴	1.41	0.09, 22.83	n/a
Pragmatic	2 ^{31,32}	1.31	0.23, 7.58	0%
Outcome: Neonatal Mortality				
Midwives Well-integrated Setting				
Nulliparas	2	0.96	0.71, 1.29	0%
Within standards	1 ²⁵	0.99	0.73, 1.34	n/a
Pragmatic	1 ²⁹	0.57	0.17, 1.95	n/a
Multiparas	2	1.08	0.74, 1.58	0%
Within standards	1 ²⁵	1.04	0.71, 1.54	n/a
Pragmatic	1 ²⁹	2.00	0.37, 10.92	n/a
Not stratified by parity	-	-	-	-
Pragmatic	2 ^{28,29}	1.07	0.50, 2.30	0%
Midwives Less well-integrated Setting				
Nulliparas	-	-	-	-
Pragmatic	1 ³¹	2.08	0.11, 38.66	n/a
Multiparas	-	-	-	-
Pragmatic	1 ³¹	0.68	0.09, 5.25	n/a

*Note that Pang et al (midwives well-integrated setting, within standards study design) reported: nulliparas RR 2.73 (95% CI, 2.06 to 7.06) and not stratified by parity RR 1.99 (95% CI, 1.06 to 3.73).¹⁸

†Note that van der Kooy et al 2017 (midwives well-integrated setting, pragmatic study design) reported: RR 0.80 (0.71, 0.91) (not adjusted for parity).¹⁹ Had we included these data, the van der Kooy et al 2011 data would have been excluded (for duplication).³⁰

Table 4. Summary of infant morbidity meta-analyses findings

Strata	Number of studies	OR	95% CI	I ²
Outcome: Apgar Score <7 at 5 Minutes*				
Midwives Well-integrated Setting				
Nulliparas	3	1.19	0.76, 1.87	83.2%
Within standards	2 ^{23,25}	1.12	0.69, 1.84	91.0%
Pragmatic	1 ²⁷	2.07	0.56, 7.57	n/a
Multiparas	3	0.76	0.60, 0.96	44.8%
Within standards	2 ^{23,25}	0.77	0.60, 0.99	64.1%
Pragmatic	1 ²⁷	0.33	0.04, 2.62	n/a
Not stratified by parity	6	0.87	0.74, 1.03	0%
Within standards	1 ²³	0.94	0.71, 1.25	n/a
Pragmatic	5 ^{21,22,27-29}	0.84	0.69, 1.03	0%
Midwives Less well-integrated Setting				
Nulliparas	-	-	-	-
Pragmatic	1 ³¹	0.14	0.01, 2.22	n/a
Multiparas	-	-	-	-
Pragmatic	1 ³¹	0.60	0.18, 1.95	n/a
Not stratified by parity	-	-	-	-
Within standards	1 ³³	1.10	0.18, 6.68	n/a
Outcome: Apgar Score <7 at 1 Minute				
Midwives Well-integrated Setting				
Not stratified by parity	-	-	-	-
Pragmatic	2 ^{21,22}	0.73	0.63, 0.83	0%
Outcome: Neonatal Resuscitation				
Midwives Well-integrated Setting				
Nulliparas	-	-	-	-
Pragmatic	1 ²⁷	4.20	0.91, 19.30	n/a
Multiparas	-	-	-	-
Pragmatic	1 ²⁷	0.85	0.28, 2.62	n/a
Not stratified by parity	-	-	-	-
Pragmatic	5 ^{21,22,27-29}	1.02	0.72, 1.43	0%
Outcome: NICU Admission†				
Midwives Well-integrated Setting				
Nulliparas	5	0.93	0.85, 1.02	0%
Within standards	3 ^{23,25,26}	0.94	0.85, 1.04	0%
Pragmatic	2 ^{24,27}	1.00	0.40, 2.48	77.9%
Multiparas	5	0.73	0.65, 0.82	0%
Within standards	3 ^{23,25,26}	0.73	0.65, 0.83	2.4%
Pragmatic	2 ^{24,27}	0.66	0.43, 1.03	0%
Not stratified by parity	3	0.83	0.69, 0.99	5.3%

Within standards	1 ²³	0.73	0.57, 0.94	n/a
Pragmatic	2 ^{27,28}	0.92	0.73, 1.17	0%

*Note that Davis et al (midwives well-integrated setting, within standards study design) reported: not stratified by parity RR 0.81 (95% CI, 0.39 to 1.68) (adjusted for maternal age, parity, ethnicity and smoking).²⁰

†Note that Davis et al (midwives well-integrated setting, within standards study design) reported: RR 1.00 (0.66 to 1.50) (adjusted for maternal age, parity, ethnicity and smoking).²⁰

APPENDICES

Appendix 1: Table of Studies Excluded from Systematic Review and Meta-Analysis

Appendix 2: Funnel Plots for the Primary Outcome

REFERENCES

- 1 Canadian Institute for Health Information. Giving Birth in Canada. The Costs. 2006. Available from: <https://secure.cihi.ca/estore/productFamily.htm?locale=en&pf=PFC421> (Accessed Jan 29, 2019)
- 2 Canadian Institute for Health Information. Giving Birth in Canada: Providers of Maternity and Infant Care. 2004. Available from: <https://secure.cihi.ca/estore/productFamily.htm?locale=en&pf=PFC348> (Accessed Jan 29, 2019)
- 3 Hutton EK, Cappelletti A, Reitsma AH, *et al.* Outcomes associated with planned place of birth among women with low-risk pregnancies. *CMAJ* 2015; **188**(5): e80-e90.
- 4 Snowden JM, Tilden EL, Snyder JS, Brian Q, *et al.* Planned Out-of-Hospital Birth and Birth Outcomes. *N Engl J Med* 2015; **373**: 2642–53.
- 5 Royal College of Obstetricians and Gynaecologists. Intrapartum care: Care of Healthy Women and their Babies During Childbirth. 2014. Available from: <http://www.nice.org.uk/guidance/CG190>. (Accessed Jan 29, 2019)
- 6 Olsen O, Jewell MD. Home versus hospital birth. *Cochrane Database Syst Rev* 2012; **9**: 32.
- 7 Hutton EK, Reitsma AH, Thorpe J, *et al.* Protocol: systematic review and meta-analyses of birth outcomes for women who intend at the onset of labour to give birth at home compared to women of low obstetrical risk who intend to give birth in hospital. *Syst Rev* 2014; **3**: 55.
- 8 Hutton EK. The Safety of Home Birth. *J Obstet Gynaecol Canada* 2016; **38**: 331–6.
- 9 Comeau A, Hutton EK, Simioni J, *et al.* Home birth integration into the health care systems of eleven international jurisdictions. *Birth* 2018; **45**(3): 311-321.
- 10 Wells G, Shea B, O’Connell D, *et al.* The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. *Ottawa Hosp Res Inst* 2013. Available from: http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp (Accessed Jan 29, 2019).
- 11 Sterne JAC, Sutton AJ, Ioannidis JPA, *et al.* Recommendations for examining and interpreting funnel plot asymmetry in meta-analyses of randomised controlled trials. *BMJ* 2011; **343**: d4002.
- 12 de Jonge de A, van der Goes BY, Ravelli AC *et al.* Perinatal mortality and morbidity in a nationwide cohort of 529,688 low-risk planned home and hospital births. *BJOG* 2009; **116**: 1177–84.
- 13 Davis D, Baddock S, Pairman S, *et al.* Risk of severe postpartum hemorrhage in low-risk

- childbearing women in new zealand: exploring the effect of place of birth and comparing third stage management of labor. *Birth* 2012; **39**: 98–105.
- 14 Bolten N, de Jonge A, Zwagerman E, *et al.* Effect of planned place of birth on obstetric interventions and maternal outcomes among low-risk women: A cohort study in the Netherlands. *BMC Pregnancy Childbirth* 2016; **16**: 329.
 - 15 de Jonge A, Mesman JA, Mannien J, *et al.* Severe adverse maternal outcomes among low risk women with planned home versus hospital births in the Netherlands: nationwide cohort study. *BMJ* 2013; **346**: f3263.
 - 16 Miller S, Skinner J. Are first-time mothers who plan home birth more likely to receive evidence-based care? A comparative study of home and hospital care provided by the same midwives. *Birth* 2012; **39**: 135–44.
 - 17 Nove A, Berrington A, Matthews Z. Comparing the odds of postpartum haemorrhage in planned home birth against planned hospital birth: results of an observational study of over 500,000 maternities in the UK. *BMC Pregnancy Childbirth* 2012; **12**: 130.
 - 18 Pang JWY, Heffelfinger JD, Huang GJ, *et al.* Outcomes of planned home births in Washington State: 1989-1996. *Obstet Gynecol* 2002; **100**: 253–9.
 - 19 van der Kooy J, Birnie E, Denktas S, *et al.* Planned home compared with planned hospital births: mode of delivery and Perinatal mortality rates, an observational study. *BMC Pregnancy Childbirth* 2017; **17**: 177.
 - 20 Davis D, Baddock S, Pairman S, *et al.* Planned Place of Birth in New Zealand: Does it Affect Mode of Birth and Intervention Rates Among Low-Risk Women? *Birth* 2011; **38**: 111–9.
 - 21 Janssen PA, Lee SK, Ryan EM, *et al.* Outcomes of planned home births versus planned hospital births after regulation of midwifery in British Columbia. *CMAJ* 2002; **166**: 315–23.
 - 22 Janssen PA, Saxell L, Page LA, *et al.* Outcomes of planned home birth with registered midwife versus planned hospital birth with midwife or physician. *CMAJ* 2009; **181**: 377–83.
 - 23 Brocklehurst P, Hardy P, Hollowell J, *et al.* Perinatal and maternal outcomes by planned place of birth for healthy women with low risk pregnancies: the Birthplace in England national prospective cohort study. *BMJ* 2011; **343**: d7400.
 - 24 Wiegers TA, Keirse MJ, van der Zee J, *et al.* Outcome of planned home and planned hospital births in low risk pregnancies: prospective study in midwifery practices in The Netherlands. *BMJ* 1996; **313**: 1309–13.
 - 25 de Jonge A, Geerts CC, van der Goes BY, *et al.* Perinatal mortality and morbidity up to 28 days after birth among 743 070 low-risk planned home and hospital births: A cohort study based on three merged national perinatal databases. *BJOG* 2014; **122**: 720–8.

- 26 Hermus MAA, Hitzert M, Boesveld IC, *et al.* Differences in optimality index between planned place of birth in a birth centre and alternative planned places of birth, a nationwide prospective cohort study in The Netherlands: results of the Dutch Birth Centre Study. *BMJ Open* 2017; **7**: e016958.
- 27 Halfdansson B, Smarason AK, Olafsdottir OA, *et al.* Outcome of planned home and hospital births among low-risk women in Iceland in 2005-2009: a retrospective cohort study. *Birth* 2015; **42**: 16–26.
- 28 Hutton EK, Reitsma AH, Kaufman K. Outcomes associated with planned home and planned hospital births in low-risk women attended by midwives in Ontario, Canada, 2003-2006: a retrospective cohort study. *Birth* 2009; **36**: 180–9.
- 29 Hutton EK, Cappelletti A, Reitsma AH, *et al.* Outcomes associated with planned place of birth among women with low-risk pregnancies. *CMAJ* 2016; 188(5): e80-90.
- 30 van der Kooy J, Poeran J, de Graaf JP, *et al.* Planned home compared with planned hospital births in the Netherlands: intrapartum and early neonatal death in low-risk pregnancies. *Obstet Gynecol* 2011; **118**: 1037–46.
- 31 Blix E, Huitfeldt AS, Oian P, *et al.* Outcomes of planned home births and planned hospital births in low-risk women in Norway between 1990 and 2007: A retrospective cohort study. *Sex. Reprod. Healthc.* 2012; **3**: 147–53.
- 32 Lindgren HE, Radestad IJ, Christensson K, *et al.* Outcome of planned home births compared to hospital births in Sweden between 1992 and 2004. A population-based register study. *Acta Obstet Gynecol Scand* 2008; **87**: 751–9.
- 33 Hiraizumi Y, Suzuki S. Perinatal outcomes of low-risk planned home and hospital births under midwife-led care in Japan. *J Obstet Gynaecol Res* 2013; **39**: 1500–4.
- 34 Homer CS, Thornton C, Scarf VL *et al.* Birthplace in New South Wales, Australia: An analysis of perinatal outcomes using routinely collected data. *BMC Pregnancy Childbirth.* 2014; **14**: no pagination.
- 35 Scarf VL, Rossiter C, Vedam S, *et al.* Maternal and perinatal outcomes by planned place of birth among women with low-risk pregnancies in high-income countries: A systematic review and meta-analysis. *Midwifery* 2018; **62**: 240–55.