

**UCC Library and UCC researchers have made this item openly available.
Please [let us know](#) how this has helped you. Thanks!**

Title	A plasmid-encoded putative glycosyltransferase is involved in hop tolerance and beer spoilage in <i>Lactobacillus brevis</i>
Author(s)	Feyereisen, Marine; Mahony, Jennifer; O'Sullivan, Tadhg; Boer, Viktor; van Sinderen, Douwe
Publication date	2020-01-21
Original citation	Feyereisen, M., Mahony, J., O'Sullivan, T. and van Sinderen, D. (2020) 'A Plasmid-Encoded Putative Glycosyltransferase Is Involved in Hop Tolerance and Beer Spoilage in <i>Lactobacillus brevis</i> ', <i>Applied and Environmental Microbiology</i> , 86(3), e02268-19, (10 pp). doi: 10.1128/AEM.02268-19
Type of publication	Article (peer-reviewed)
Link to publisher's version	https://aem.asm.org/content/86/3/e02268-19 http://dx.doi.org/10.1128/AEM.02268-19 Access to the full text of the published version may require a subscription.
Rights	© 2020 American Society for Microbiology. All Rights Reserved.
Embargo information	Access to this article is restricted until 6 months after publication by request of the publisher.
Embargo lift date	2020-07-21
Item downloaded from	http://hdl.handle.net/10468/9955

Downloaded on 2021-11-27T12:25:22Z

1 **Title: A plasmid-encoded putative glycosyltransferase is involved in hop tolerance and**
2 **beer spoilage in *Lactobacillus brevis*.**

3 **Running title: A glycosyltransferase involved in beer spoilage in *Lb. brevis***

4 Marine Feyereisen¹, Jennifer Mahony^{1,2}, Tadhg O’Sullivan³, Viktor Boer³ and Douwe van
5 Sinderen^{*1,2}

6 ¹School of Microbiology, University College Cork, T12 YT20 Cork, Ireland;
7 116221209@umail.ucc.ie (M.F.); J.Mahony@ucc.ie (J.M.)

8 ²APC Microbiome Ireland, University College Cork, T12 YT20 Cork, Ireland

9 ³HEINEKEN Global Innovation and Research, Heineken Supply Chain B.V, 2382 PH,
10 Zoeterwoude, The Netherlands; Tadhg.OSullivan@heineken.com (T.O.S.), Viktor Boer
11 viktor.boer@heineken.com (V.B.)

12 *Correspondence: d.vansinderen@ucc.ie; Tel.: +353 21 4901365

13

14 **Keywords:** Lactic acid bacteria, plasmid, resistance, HorA, cell wall polysaccharide, phage.

15 **Abstract**

16 *Lactobacillus brevis* beer-spoiling strains harbor plasmids that contain genes such as *horA*,
17 *horC* and *hitA*, which are known to confer hop tolerance. The *Lb. brevis* beer-spoiling strain
18 UCCLBBS124, which possesses four plasmids, was treated with novobiocin resulting in the
19 isolation of UCCLBBS124 derivatives exhibiting hop-sensitivity and an inability to grow in
20 beer. One selected derivative was shown to have lost a single plasmid, designated here as
21 UCCLBBS124_D, which harbors the *UCCLBBS124_pD0015* gene, predicted to encode a
22 glycosyltransferase. Hop tolerance and growth in beer was restored when
23 *UCCLBBS124_pD0015* was introduced in one of these hop-sensitive derivatives on a
24 plasmid. We hypothesize that this gene modifies the surface composition of the
25 polysaccharide cell wall conferring protection against hop compounds. Furthermore,
26 introduction of this gene *in trans* in *Lb. brevis* UCCLB521, a strain that cannot grow in and
27 spoil beer, was shown to furnish the resulting strain with the ability to grow in beer while its
28 expression also conferred phage-resistance. This study underscores how the acquisition of
29 certain mobile genetic elements plays a role in hop tolerance and beer spoilage for strains of
30 this bacterial species.

31 **Importance**

32 *Lactobacillus brevis* is a member of the lactic acid bacteria and is often reported as the
33 causative agent of food or beverage spoilage, in particular that of beer. Bacterial spoilage of
34 beer may result in product withdrawal or recall with concomitant economic losses for the
35 brewing industry. A very limited number of genes involved in beer spoilage have been
36 identified and primarily include those involved in hop resistance such as *horA*, *hitA*, and
37 *horC*. However, since none of these genes are universal, it is clear that there are likely (many)
38 other molecular players involved in beer spoilage. Here, we report on the importance of a
39 plasmid-encoded glycosyltransferase associated with beer spoilage by *Lb. brevis* that is

40 involved in hop tolerance. The study highlights the complexity of the genetic requirements to
41 facilitate beer spoilage and the role of multiple key players in this process.

42 Introduction

43 *Lactobacillus brevis* is a major threat for commercial and amateur brewers as strains
44 of this species are the predominant bacterial contaminants associated with beer spoilage (1).
45 Such *Lb. brevis* strains can grow in beer despite the presence of ethanol, low pH and the
46 depletion of oxygen and nutrients (2). Moreover, hop compounds added to beer for bitter
47 flavor development during the fermentation process also exert antibacterial activity through
48 the presence of *iso*- α -acids (1, 2). *Lb. brevis* beer-spoiling (BS) strains appear to have
49 acquired chromosomally- or plasmid-derived genetic content to survive and grow in beer (2).
50 *Lb. brevis* resistance to ethanol (up to 10 %) and pH lower than the optimal growth
51 conditions (pH 4-6) seems to be associated with chromosomal genes, possibly due to the
52 general stressors they represent (3, 4). However, *Lb. brevis* BS strains are also known to
53 harbor plasmids that are associated with their beer-spoilage phenotype and more specifically
54 with hop tolerance (5-8). Plasmid-derived genes that underpin hop-resistance in *Lb. brevis*
55 include *horA*, *horC*, *hitA* and *orf5_{ABBC45}* (1, 2). The genes *horA* and *horC* encode multidrug
56 transporter proteins driven by ATP and proton motive force (PMF), respectively, and were
57 identified as being involved in *iso*- α -acid extrusion from the bacterial cell (5, 7). The gene
58 *hitA* encodes a transmembrane protein involved in the transport of divalent cations such as
59 Mn²⁺ in exchange of protons released from hop bitter acids (8). The *orf5_{ABBC45}* gene was
60 identified in *Lb. brevis* BS strain ABBC45 which was unable to grow in beer after it had lost
61 a plasmid carrying this gene. The *orf5_{ABBC45}* gene encodes a predicted transmembrane protein
62 resembling a PMF-dependent multidrug transporter, which is presumed to be responsible for
63 *iso*- α -acid export (9).

64 However, these genes are not always indicative of BS ability as the presence of such
65 genes can be found among *Lb. brevis* strains that are unable to grow and consequently spoil
66 beer (designated here as NBS strains) (10). Indeed, *horA* is present in the *Lb. brevis* NBS

67 strain UCCLB556 (10). Moreover, genes identified as conferring hop-resistance are not
68 always simultaneously present in BS strains, e.g. the BS strain UCCLBBS124 carries
69 plasmids harboring *horA* and *horC*, however it does not possess *hitA* (10). Analysis of BS
70 strain *Lb. brevis* BSO 464 has highlighted the importance of plasmids and genes on mobile
71 genetic elements for bacterial growth in beer and beer spoilage ability (6). Recently, a gene
72 predicted to encode a glycosyltransferase was identified among BS strains responsible for
73 excess β -glucan formation (11). This gene is also present on the genome of *Lb. brevis* BS
74 strain UCCLBBS124, while it is absent in that of BS strain UCCLBBS449 (10). This
75 indicates that beer spoilage is not uniquely governed by the presence of a few genes, but
76 rather a combination of genes acting in concert to confer beer resistance to the strain. It also
77 suggests that other plasmid-encoded genes involved in beer spoilage are yet to be discovered.

78 In the present study we generated plasmid-cured derivatives of *Lb. brevis* BS strain
79 UCCLBBS124 using novobiocin. This approach has been successfully employed previously
80 to cure plasmids from lactic acid bacteria (LAB) isolates (6, 12). Plasmid-cured derivatives
81 were assessed for their ability to grow in the presence of hop and in beer. A derivative that
82 showed inability to grow in beer was selected and analyzed to ascertain which plasmids were
83 responsible for this phenotype. Bioinformatic analysis of the genetic content of such plasmids
84 revealed candidate genes required for growth in beer. These genes were used in
85 transformation experiments to revert the NBS phenotype.

86

87

88 **Results and Discussion**

89 *Derivatives with impaired growth in beer reveal loss of plasmid UCCLBBS124_D*

90 The beer-spoiling *Lb. brevis* strain UCCLBBS124 (abbreviated here as UCC124) possesses
91 four plasmids carrying genes of interest for bacterial beer spoilage (Table 3). Following
92 exposure to novobiocin, surviving *Lb. brevis* UCC124 cells were plated and fifty isolated
93 colonies (10) were randomly selected for further analysis. Thirty four of these fifty colonies
94 displayed impaired growth in beer. PCR-based identification of the hop-resistance gene *horA*
95 revealed the loss of this gene, located on plasmid UCCLBBS124_D (abbreviated here as
96 UCC124_D) in 33 out of the 34 isolates. One derivative, designated here as MB569, was
97 selected for genome sequencing, after which its sequence was compared to that of the WT,
98 confirming that plasmid UCC124_D had been lost from strain MB569.

99 *Tolerance of MB569 to iso- α -acids, ethanol and low pH*

100 The inability of strain MB569 to grow in beer highlights the apparent importance of plasmid
101 UCC124_D in conferring a beer spoilage phenotype on strain UCC124 (Figure 1). Beer is a
102 harsh environment incorporating a number of stresses such as low pH, lack of nutrients, and
103 the presence of ethanol and hop compounds. In order to understand which of these stresses
104 imposed a negative impact on growth of MB569, the WT strain and MB569 were grown in
105 MRS broth and mimicking conditions encountered in beer, e.g. pH4, 5.4 % ethanol, and 30
106 ppm *iso*- α -acids. Strain MB569 was shown to be capable of growth in MRS broth at neutral
107 pH and at pH 4, while it can also grow in the presence of ethanol comparable to the WT
108 strain (Figure 1). However, MB569 is incapable of growth in the presence of *iso*- α -acids
109 unlike the WT strain UCC124 (Figure 1). This indicates that plasmid-cured derivative
110 MB569 has lost the ability to spoil beer due to its sensitivity to the antimicrobial compounds
111 present in hops. Therefore, based on this phenotype and the finding that MB569 lacks

112 plasmid UCC124_D (when compared to its parental strain), it indicates that this plasmid is
113 linked to hop tolerance and thus contributing to the ability of strain UCC124 to cause beer
114 spoilage.

115 *Identification and functional annotation of genes present on plasmid UCC124_D*

116 Plasmid UCC124_D is 21 kb in size and is predicted to encompass 16 genes. Interestingly, a
117 7 kb region of this plasmid, contains six genes that are uniquely present among the plasmids
118 of *Lb. brevis* BS strains (Table 4) (10). In order to assess the possible role of these genes in
119 beer spoilage, the BS plasmid-specific genes *UCCLBBS124_pD0014* (abbreviated here as
120 *UCC124_D14*), encoding a predicted cytosine deaminase, *UCCLBBS124_pD0015* (renamed
121 *gtf_{D15}*), encoding a predicted glycosyltransferase, *UCCLBBS124_pD0016* (designated here as
122 *UCC124_horA*), which encodes HorA (Table 4), were individually cloned into plasmid
123 pNZ44 prior their transformation into NZ9000. The resulting plasmids were then introduced
124 into strain MB569 to determine the ability of the obtained recombinant strains to grow in beer
125 (where MB569 itself is unable to do so). Genes with locus tags *UCCLBBS124_pD0017*
126 (abbreviated as UCC124_D17), *UCCLBBS124_pD0018* (abbreviated as UCC124_D18) and
127 *UCCLBBS124_pD0019* (abbreviated as UCC124_D19) and encoding acyl-sn-glycerol-3-
128 phosphate acyltransferases and a glycosyltransferase (Table 4) were cloned together as a
129 cluster (as present in plasmid UCC124_D) in pNZ44 prior their introduction into NZ9000
130 and, subsequently, MB569.

131 Introduction of the genes *UCC124_D14*, *UCC124_horA*, *UCC124_D17*, *UCC124_D18* and
132 *UCC124_D19* in MB569 did not enable any obvious improvement of growth in the presence
133 of *iso- α -acid* (30 ppm) or beer (when compared to strain MB569) (data not shown).

134 Interestingly, expression of *gtf_{D15}* in MB569 was shown to confer a positive effect on its
135 ability to grow in MRS broth containing 30 ppm *iso- α -acids*, with a significant (P value <

136 0.05) growth increase after 72 h compared to the non-complemented strain or MB569
137 carrying the control plasmid pNZ44 (Figure 2A). When *Lb. brevis* MB569 pNZ44:*gtf_{D15}* was
138 cultivated in beer, it also exhibited an ability to grow in beer that was significantly better than
139 that of MB569 itself (P value < 0.05) (Figure 2B). Provision of *gtf_{D15}* *in trans* in MB569 did
140 not restore its growth in beer to the same level as the WT strain (i.e. the strain from which
141 MB569 was derived), but nonetheless allowed survival and growth in beer for this
142 recombinant strain across 96 h. MB569 and MB569 pNZ44 are still able to survive in the
143 presence of *iso-α*-acids or beer after culture for 72 h (Figure 2A and 2B) which might be due
144 to the presence of plasmid UCCLBBS124_C carrying the gene *horC* (Table 3). The *gtf_{D15}*
145 gene is predicted to encode a glycosyltransferase based on BLAST analysis and a HHPred
146 analysis (13) predicted the protein to belong to the glycosyl transferase family 8 associated
147 with cell wall glycosylation (99.9 % probability and E-value < 10⁻²⁸). Further sequence
148 scrutiny suggests that the Gtf_{D15} protein is a membrane-associated protein (TMHMM Server
149 2.0 (14)) with a predicted signal peptide in its N-terminus that may act as a membrane anchor
150 for the protein (<http://phobius.sbc.su.se/> (15)). These predictions suggest that GtfD15 is a cell
151 envelope-associated protein that confers protection against certain environmental stressors
152 such as hop compounds.

153 *Introduction of gtf_{D15} in NBS Lb. brevis* strains allows growth in beer

154 The introduction of *gtf_{D15}* in MB569 was shown to significantly improve growth of the strain
155 in MRS broth containing hop compounds (30 ppm *iso-α*-acids) and in beer indicating the
156 importance of this gene for beer spoilage by *Lb. brevis* strain UCC124. In order to assess the
157 potential growth-promoting effect of this gene for an NBS strain when inoculated in beer,
158 *gtf_{D15}* when cloned into pNZ44 (pNZ44:*gtf_{D15}*) was introduced into the NBS *Lb. brevis* strain
159 UCCLB521 (renamed here as UCC521) (Table 1). Remarkably, the presence of pNZ44:*gtf_{D15}*
160 in the NBS strain *Lb. brevis* UCC521 permitted the strain to grow significantly better (P

161 value < 0.05) in MRS broth containing 30 ppm *iso*- α -acids and in beer compared to the strain
162 carrying an empty plasmid which is incapable of survival or growth in these environments
163 (Figure 2C and D). These observations reinforce our results above and highlight the
164 significance of the *gtf_{D15}* gene in hop tolerance and beer spoilage. An alternative, though in
165 our opinion less likely explanation is that strains for which we obtained no or reduced CFUs
166 had entered a viable, but non-culturable (so-called VBNC) state as has been previously
167 observed for beer-passaged *Lb. brevis* strains (16).

168 Introduction of pNZ44:*gtf_{D15}* into ATCC 367, another NBS strain, did not allow improved
169 survival or growth in the presence of hop compounds or in beer (data not shown). This
170 suggests that a strain-specific mechanism and involvement of other genes that are absent in
171 ATCC 367 are responsible for increased hop tolerance. Among beer spoilage-related genes,
172 UCC521 possesses the *orf5_{ABBC45}* gene previously identified as involved in hop tolerance (9),
173 unlike ATCC 367 which does not harbor known genes involved in beer spoilage. Moreover,
174 UCC521, although a non-beer spoiler strain, was isolated from the brewery environment,
175 unlike ATCC 367 which was isolated from silage (10). UCC521 may have acquired genes
176 (such as *orf5_{ABBC45}*) or plasmids (UCC521 harbors five plasmids) throughout its presence in
177 the brewery environment, which confer hop tolerance when combined with *gtf_{D15}*. This
178 scenario has previously been observed, indicating that *Lb. brevis* strains can only survive and
179 grow in beer when multiple beer spoilage-related genes are present in a particular
180 combination (17). Selection pressures of the beer environment determine the genetic content
181 of beer-spoiling strains. The identification of diagnostic marker genes (DMGs) are important
182 in distinguishing BS from NBS strains, as well as predicting the ability of a given strain to
183 grow in beer (17, 18). In the study from Bergsveinson and Ziola, proposed DMGs were not
184 related to hop tolerance and no genes encoding glycosyltransferases were identified as
185 DMGs. However, a glycosyltransferase-encoding gene located on the plasmid of the BS *Lb.*

186 *brevis* strain BSO 464 which showed more than 99 % nt similarity to *gtf_{D15}* was described as
187 unique when compared to a NBS *Lb. brevis* strain KB290 and a BS *Pediococcus damnosus*
188 strain Pc344^T (17). From these observations and knowing that *gtf_{D15}* is highly prevalent in BS
189 *Lb. brevis* strains (Table 4), we propose to include this gene as a DMG to assess the beer
190 spoiling potential of *Lb. brevis* strains.

191 *Effect on phage sensitivity*

192 As demonstrated above *gtf_{D15}* was observed to play a role in hop and beer tolerance and is
193 predicted to encode a glycosyltransferase. Since the protein is predicted to be involved in
194 biosynthesis or modification of a cell surface-associated saccharidic polymer, the possible
195 role of this protein in bacteriophage infection was investigated. *Lb. brevis* strain UCC521 is
196 sensitive to *Lb. brevis* phages 3-521 and 521B (19). Plaque assays employing these phages
197 and *Lb. brevis* UCC521 harboring the empty vector pNZ44, or strain UCC521 containing
198 pNZ44:*gtf_{D15}* displayed similar EOP (Efficiency Of Plaquing) values with no significant
199 difference to the WT (Table 5). However, notable differences in plaque morphology were
200 observed, where plaques were faint and hard to distinguish on the bacterial lawn of UCC521
201 pNZ44:*gtf_{D15}*. Moreover, overnight incubation of the different strains with the two phages led
202 to complete lysis-in-broth of UCC521 and UCC521 containing pNZ44 with an approximately
203 1000-fold increase of phage titre after overnight propagation (Table 5). In contrast, UCC521
204 pNZ44:*gtf_{D15}* did not show visible lysis and was able to grow after overnight incubation with
205 just a ten-fold increase in phage numbers after overnight propagation (Table 5). These results
206 reinforce the role of the protein Gtf_{D15} in bacterial protection against diverse environmental
207 hazards such as hop compounds or bacteriophages.

208

209

210 **Conclusions**

211 In this study, we identified a novel genetic component required for beer spoilage and more
212 specifically for hop tolerance. This gene is located on plasmid UCC124_D of *Lb. brevis* BS
213 strain UCC124, validating the importance of plasmids to confer a beer spoilage phenotype.
214 Moreover, this gene had been highlighted previously as common among BS strains (10).
215 Genes required for hop tolerance have all been identified on plasmids (5, 7, 8), reinforcing
216 the importance of such mobile genetic elements in adaptation to the specific hurdles imposed
217 by the beer environment. A derivative of UCC124, MB569 showed impaired growth in beer
218 after the loss of plasmid UCC124_D and despite the presence of plasmids UCC124_B and
219 UCC124_C which carry several genes of interest in beer spoilage. Introduction of *gtf_{D15}* in
220 strain MB569 restored the hop tolerance phenotype of the strain which ultimately allowed it
221 to grow in beer. Similar results were observed when the gene was introduced into a NBS
222 strain confirming the notion that *gtf_{D15}* is required for the development of hop tolerance and
223 beer spoilage. Furthermore, this gene impacts on phage sensitivity of its host. This gene
224 seems a unique trait shared among BS strains of *Lb. brevis* and we propose *gtf_{D15}* as a DMG
225 for the detection of potential bacterial contamination of beer. The gene is predicted to encode
226 a glycosyltransferase and analysis of its topology suggests that it is a membrane-anchored
227 protein involved in the biosynthesis or modification of a cell surface-associated saccharidic
228 polymer. BS strains of *Lb. brevis* have been shown to increase higher molecular weight
229 lipoteichoic acids (LTA) in their cell wall, in the presence of hop bitter acids, thus believed to
230 confer resistance to the bacteria by enhancing the barrier functions of the cell wall and
231 preventing intrusion of hop compounds (20, 21). Moreover, lipoteichoic acids have been
232 described as phage receptors among lactobacilli phages as seen for *Lactobacillus delbrueckii*
233 phages LL-H and JCL1032 (22) but also for *Lactobacillus plantarum* ATCC8014-B2 (23).
234 Therefore, we speculate that this glycosyltransferase is involved in replacing alanine residues

235 with sugar residues on teichoic acids thereby changing their charge and preventing *iso- α -*
236 acids to penetrate the membrane as well as affecting phage adsorption and/or DNA injection.
237 This predicted glycosyltransferase shows only limited similarity (36 % amino acid similarity
238 in 20 % query cover) with the glycosyltransferase identified in a previous study as
239 responsible for β -glucan formation (11), and is thus believed to play a different role in beer
240 spoilage. Future studies will focus on defining the mechanism that underpins hop tolerance
241 and on determining how the genes identified to date (5, 7, 8) are linked to each other.
242 Moreover, located on the same plasmid as *gtf_{D15}* are genes predicted to encode a
243 glycosyltransferase and acyltransferases (Table 2) suggesting a common action on teichoic
244 acids with the acyltransferases involved in the acylation of alanine residues or the lipid
245 moiety of the lipoteichoic acids (24). Follow-up work may therefore focus on determining the
246 precise function of the glycosyltransferase (and other associated genes) in the modification of
247 the cell wall and/or cell surface. Another question to be addressed is if and how hop tolerance
248 is enhanced when these genes are present in a certain combination, and how such tolerance is
249 influenced by their expression level.

250

251

252 **Materials and Methods**

253 *Bacterial strains and cultivation media*

254 Bacterial strains used in this study are listed in Table 1. *Lb. brevis* strains were grown in
255 MRS broth (Oxoid Ltd., UK) at 30 °C while *Lactococcus lactis* NZ9000 was grown in M17
256 broth (Oxoid Ltd., UK) supplemented with 0.5 % glucose. 5 µg/mL chloramphenicol (Cm5)
257 was added to the medium when indicated.

258 *Plasmid curing and plasmid content analysis*

259 The overall experimental approach is presented in Figure S1. Plasmid curing of the BS strain
260 UCC124 was achieved using novobiocin treatment (25). A 1 % inoculum of a WT strain
261 overnight culture was used to inoculate 10 mL MRS broth containing 0.25 µg/mL
262 novobiocin. Cultures were incubated at 26 °C for 72 h. After incubation, cells were diluted
263 and plated on MRS agar. After 3 days of incubation at 26 °C, isolated colonies were
264 randomly selected and derivatives with impaired growth in beer (no growth observed after 72
265 h) were checked for the presence or loss of hop-resistance genes *horA*, *horC* and *orf5_{ABBC45}*
266 (Table 2). A derivative showing loss of hop-resistance gene was selected and sequenced
267 using Illumina sequencing technology. Paired-end sequence reads were generated using an
268 Illumina HiSeq2500 system (read length 2 x 250 bp). FASTQ sequence files were generated
269 using the Illumina Casava pipeline version 1.8.3. After Illumina sequencing the obtained
270 sequences were mapped back against the WT reference sequence to detect mutations by
271 single nucleotide polymorphism (SNP) or plasmid content loss. SNP analysis was performed
272 by aligning Illumina raw reads against a reference sequence using Bowtie2 V 2.3.5 (26). The
273 reads were then sorted using Samtools (27) and VarScan v2.3.9 was applied for the detection
274 of variants (28). A minimum allelic variation frequency cut-off of 0.25 was applied.

275 *Construction of plasmid vectors*

276 Genes of interest were amplified by PCR (Table 2) and cloned into the expression vector
277 pNZ44 (29). PCR products and pNZ44 plasmid DNA were digested with the appropriate
278 enzymes (Roche, USA) at 37 °C for at least 4 h, following the manufacturer's instructions
279 (Table 2). A ratio of (3:1) was applied for the ligation of the PCR product with pNZ44 using
280 T4 DNA ligase (Promega, USA). The mixture was incubated at room temperature for at least
281 4 hours prior to electrotransformation into *L. lactis* NZ9000 competent cells.

282 *Preparation of competent cells and electrotransformation*

283 Competent cells of *L. lactis* NZ9000 were prepared as previously described (30). Competent
284 cells of *Lb. brevis* UCC124 were prepared using an adapted version of a previously described
285 protocol (31): An overnight culture was transferred (1 % inoculum) to 10 mL MRS broth
286 containing 1 % glycine and incubated overnight at 30 °C. 5 mL of the overnight culture was
287 transferred to fresh MRS broth containing 1 % glycine (50 mL final volume) and cells were
288 grown to an OD_{600nm} of 0.6. Cells were harvested by centrifugation at 4,000 × *g* for 15 min at
289 4 °C and washed in ice-cold wash buffer (0.5 M sucrose, 10 % glycerol). The wash step was
290 repeated twice and the cells were finally resuspended in 200 µL wash buffer prior to storage
291 at -80 °C and/or electroporation (see below). All constructs were generated using *L. lactis*
292 NZ9000 as the cloning host, verified by sequencing after PCR amplification using the
293 primers pnz44F and pnz44R (Table 2) prior to their transfer into *Lb. brevis* strains.
294 Electrotransformation was performed using freshly prepared competent cells as described
295 above, where 45 µL of cells and 5 µL of plasmid construct were mixed into a pre-chilled 2
296 mm electroporation cuvette (Cell Projects, Kent, England) and subjected to electroporation at
297 1.5 kV (*Lb. brevis*) or 2.0 kV (*L. lactis*), 200 Ω, 25 µF. Following electroporation, 950 µL
298 recovery broth was added (MRS broth supplemented with 0.5 M sucrose and 0.1 M MgCl₂
299 (*Lb. brevis*) or GM17 broth supplemented with 20 mM MgCl₂ and 2 mM CaCl₂ (*L. lactis*)).
300 Cells were recovered at 30 °C for 3 h (*Lb. brevis*) or 2 h (*L. lactis*) prior to spread plating on

301 MRS (*Lb. brevis*) or GM17 (*L. lactis*) agar supplemented with Cm5. Presumed transformants
302 were purified on MRS agar + Cm5 and colonies were checked by sequencing after PCR
303 amplification using the primers pnz44F and pnz44R (Table 2) and applied to growth assays
304 as described below.

305 *Growth assays*

306 Growth profiles of the wild-type strain and its derivative were obtained by transferring an
307 overnight culture (1 % inoculum) to MRS broth, MRS broth supplemented with 30 ppm *iso*-
308 α -acids or beer (fresh Heineken lager 5 % ethanol, pH 4, 23 ppm *iso- α -acids). Cultures were
309 incubated at 30 °C for 72 hours. One mL of culture was retrieved after 24, 48, 72 and 96
310 hours, diluted in Ringer's solution and plated on MRS agar plates. Plates were incubated at
311 30 °C anaerobically for 48 hours prior to colony counting. The number of viable bacteria of
312 each strain was assessed after CFU/mL calculation. Non-inoculated controls were used in all
313 the experiments as blank measurements. These measurements were then subtracted from each
314 experimental condition to produce the values represented on growth curves. Statistical
315 differences were calculated using unpaired *t* test method (32).*

316 *Phage activity against Lb. brevis strains and transformants*

317 To assess phage sensitivity of *Lb. brevis* strains, transformants carrying genes of interest were
318 compared to the wild-type (WT) strain using plaque assays, as previously described (33). A
319 10 μ L volume of the appropriate phage dilution and 200 μ L of *Lb. brevis* culture were added
320 to 4 mL of soft agar supplemented with 10 mM CaCl₂, mixed and poured onto an MRS agar
321 plate supplemented with 10 mM CaCl₂ and 0.5% glycine. Plates were incubated at 30 °C
322 overnight and the resulting plaques were enumerated. Phage titre was determined as plaque-
323 forming units per mL (PFU/mL). The ability of phages to propagate and multiply within the
324 host cell was also tested. *Lb. brevis* strains were grown to early exponential phase (OD_{600nm} ~

325 0.25), at which point phages were added to the culture (T0) at a MOI (multiplicity of
326 infection) of 1, along with 10 mM CaCl₂. The mix was further incubated at 30 °C overnight
327 (T1). The number of phages present in the medium (i.e. following removal of bacterial cells
328 by centrifugation) at T1 was then determined by plaque assay. Phage propagation efficiency
329 on a given host was then determined by dividing the phage titre (PFU/mL) at T1 by the phage
330 titre (PFU/mL) at T0.

331 *GenBank accession numbers*

332 *Lb. brevis* UCCLBBS124: CP031169, *Lb. brevis* UCCLBBS124_A: CP031170, *Lb. brevis*
333 UCCLBBS124_B: CP031171, *Lb. brevis* UCCLBBS124_C: CP031172 and *Lb. brevis*
334 UCCLBBS124_D: CP031173.

335 **Author Contributions:** MF performed experiments and genomic analysis. DS, JM, TS and
336 VB provided materials and strains. MF, JM, TS and DS were involved in project design and
337 wrote the manuscript. All authors read and approved the final manuscript.

338 **Conflicts of Interest:** The authors declare that VB and TS are employees of Heineken.

339 **Funding:** Marine Feyereisen is the recipient of an Irish Research Council Enterprise
340 Partnership Scheme postgraduate scholarship (Ref. No. EPSPG/2015/7). Douwe van
341 Sinderen is supported by a Principal Investigator award (Ref. No. 450 13/IA/1953) through
342 Science Foundation Ireland (SFI). Jennifer Mahony is in receipt of a Starting Investigator
343 Research Grant (SIRG) (Ref. No. 15/SIRG/3430) funded by Science Foundation Ireland
344 (SFI).

345

346

347 **References**

- 348 1. Sakamoto K, Konings WN. 2003. Beer spoilage bacteria and hop resistance.
349 International Journal of Food Microbiology 89:105-124.
- 350 2. Suzuki K, Iijima K, Sakamoto K, Sami M, Yamashita H. 2006. A review of hop
351 resistance in beer spoilage lactic acid bacteria. Journal of the Institute of Brewing
352 112:173-191.
- 353 3. Wang C, Cui Y, Qu X. 2018. Mechanisms and improvement of acid resistance in
354 lactic acid bacteria. Archives of microbiology 200:195-201.
- 355 4. Makarova K, Slesarev A, Wolf Y, Sorokin A, Mirkin B, Koonin E, Pavlov A, Pavlova
356 N, Karamychev V, Polouchine N, Shakhova V, Grigoriev I, Lou Y, Rohksar D, Lucas
357 S, Huang K, Goodstein DM, Hawkins T, Plengvidhya V, Welker D, Hughes J, Goh
358 Y, Benson A, Baldwin K, Lee J, Díaz-Muñiz I, Dosti B, Smeianov V, Wechter W,
359 Barabote R, Lorca G, Altermann E, Barrangou R, Ganesan B, Xie Y, Rawsthorne H,
360 Tamir D, Parker C, Breidt F, Broadbent J, Hutkins R, O'Sullivan D, Steele J, Unlu G,
361 Saier M, Klaenhammer T, Richardson P, Kozyavkin S, Weimer B, Mills D. 2006.
362 Comparative genomics of the lactic acid bacteria. Proceedings of the national
363 academy of sciences of the United States of America 103:15611-6.
- 364 5. Sakamoto K, Margolles A, van Veen HW, Konings WN. 2001. Hop resistance in the
365 beer spoilage bacterium *Lactobacillus brevis* is mediated by the ATP-Binding
366 Cassette multidrug transporter HorA. Journal of Bacteriology 183:5371-5375.
- 367 6. Bergsveinson J, Baecker N, Pittet V, Ziola B. 2015. Role of plasmids in *Lactobacillus*
368 *brevis* BSO 464 hop tolerance and beer spoilage. Applied and environmental
369 microbiology 81:1234-41.

- 370 7. Iijima K, Suzuki K, Ozaki K, Yamashita H. 2006. HorC confers beer-spoilage ability
371 on hop-sensitive *Lactobacillus brevis* ABBC45cc. *Journal of Applied Microbiology*
372 100:1282-1288.
- 373 8. Hayashi N, Ito M, Horiike S, Taguchi H. 2001. Molecular cloning of a putative
374 divalent-cation transporter gene as a new genetic marker for the identification of
375 *Lactobacillus brevis* strains capable of growing in beer. *Applied Microbiology and*
376 *Biotechnology* 55:596-603.
- 377 9. Suzuki K, Sami M, Kadokura H, Nakajima H, Kitamoto K. 2002. Biochemical
378 characterization of horA-independent hop resistance mechanism in *Lactobacillus*
379 *brevis*. *International journal of food microbiology* 76:223-30.
- 380 10. Feyereisen M, Mahony J, Kelleher P, Roberts RJ, O'Sullivan T, Geertman J-MA, van
381 Sinderen D. 2019. Comparative genome analysis of the *Lactobacillus brevis* species.
382 *BMC Genomics* 20:416.
- 383 11. Fraunhofer ME, Geissler AJ, Wefers D, Bunzel M, Jakob F, Vogel RF. 2018.
384 Characterization of beta-glucan formation by *Lactobacillus brevis* TMW 1.2112
385 isolated from slimy spoiled beer. *International Journal of Biological Macromolecules*
386 107:874-881.
- 387 12. Karthikeyan V, Santosh SW. 2010. Comparing the efficacy of plasmid curing agents
388 in *Lactobacillus acidophilus*. *Beneficial microbes* 1:155-8.
- 389 13. Söding J, Biegert A, Lupas AN. 2005. The HHpred interactive server for protein
390 homology detection and structure prediction. *Nucleic Acids Res* 33.
- 391 14. Krogh A, Larsson B, von Heijne G, Sonnhammer EL. 2001. Predicting
392 transmembrane protein topology with a hidden Markov model: application to
393 complete genomes. *Journal of molecular biology* 305:567-80.

- 394 15. Käll L, Krogh A, Sonnhammer ELL. 2007. Advantages of combined transmembrane
395 topology and signal peptide prediction--the Phobius web server. *Nucleic acids*
396 *research* 35:W429-W432.
- 397 16. Liu J, Deng Y, Soteyome T, Li Y, Su J, Li L, Li B, Shirtliff ME, Xu Z, Peters BM.
398 2018. Induction and recovery of the viable but nonculturable state of hop-resistance
399 *Lactobacillus brevis*. *Frontiers in Microbiology* 9:2076-2076.
- 400 17. Bergsveinson J, Ziola B. 2017. Comparative genomic and plasmid analysis of beer-
401 spoiling and non-beer-spoiling *Lactobacillus brevis* isolates. *Canadian Journal of*
402 *Microbiology* 63:970-983.
- 403 18. Behr J, Geissler AJ, Schmid J, Zehe A, Vogel RF. 2016. The identification of novel
404 diagnostic marker genes for the detection of beer spoiling *Pediococcus damnosus*
405 strains using the BIAst Diagnostic Gene findEr. *PLOS ONE* 11:e0152747.
- 406 19. Feyereisen M, Mahony J, Lugli GA, Ventura M, Neve H, Franz CMAP, Noben J-P,
407 O'Sullivan T, Sinderen Dv. 2019. Isolation and characterization of *Lactobacillus*
408 *brevis* phages. *Viruses* 11:393.
- 409 20. Behr J, Gänzle MG, Vogel RF. 2006. Characterization of a highly hop-resistant
410 *Lactobacillus brevis* strain lacking hop transport. *Applied and Environmental*
411 *Microbiology* 72:6483-6492.
- 412 21. Yasui T, Yoda K. 1997. Purification and partial characterization of an antigen specific
413 to *Lactobacillus brevis* strains with beer spoilage activity. *FEMS Microbiology*
414 *Letters* 151:169-176.
- 415 22. Räsänen L, Schubert K, Jaakonsaari T, Alatossava T. 2004. Characterization of
416 lipoteichoic acids as *Lactobacillus delbrueckii* phage receptor components. *Journal of*
417 *Bacteriology* 186:5529-5532.

- 418 23. Wolin M, Douglas LJ. 1971. Cell wall polymers and phage lysis of *Lactobacillus*
419 *plantarum*. *Biochemistry* 10:1551-1555.
- 420 24. Kiriukhin MY, Neuhaus FC. 2001. d-alanylation of lipoteichoic acid: role of the d-
421 alanyl carrier protein in acylation. *Journal of bacteriology* 183:2051-2058.
- 422 25. Ruiz-Barba JL, Piard JC, Jiménez-Díaz R. 1991. Plasmid profiles and curing of
423 plasmids in *Lactobacillus plantarum* strains isolated from green olive fermentations.
424 *Journal of applied bacteriology* 71:417-421.
- 425 26. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nature*
426 *Methods* 9:357-359.
- 427 27. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,
428 Durbin R. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics*
429 25:2078-2079.
- 430 28. Koboldt DC, Zhang Q, Larson DE, Shen D, McLellan MD, Lin L, Miller CA, Mardis
431 ER, Ding L, Wilson RK. 2012. VarScan 2: somatic mutation and copy number
432 alteration discovery in cancer by exome sequencing. *Genome Research* 22:568-76.
- 433 29. McGrath S, Fitzgerald GF, van Sinderen D. 2001. Improvement and optimization of
434 two engineered phage resistance mechanisms in *Lactococcus lactis*. *Applied*
435 *environmental microbiology* 67:608-616.
- 436 30. van Pijkeren J-P, Britton RA. 2012. High efficiency recombineering in lactic acid
437 bacteria. *Nucleic acids research* 40:e76-e76.
- 438 31. Ahrné S, Molin G, Axelsson L. 1992. Transformation of *Lactobacillus reuteri* with
439 electroporation: studies on the erythromycin resistance plasmid pLUL631. *Current*
440 *microbiology* 24:199-205.
- 441 32. Kim TK. 2015. T test as a parametric statistic. *Korean journal of anesthesiology*
442 68:540-546.

- 443 33. Svensson U, Christiansson A. 1991. Methods for phage monitoring. Bulletin of the
444 international dairy federation 263:29-39.
- 445 34. Kuipers OP, de Ruyter PGGA, Kleerebezem M, de Vos WM. 1998. Quorum sensing-
446 controlled gene expression in lactic acid bacteria. Journal of biotechnology 64:15-21.
- 447
- 448
- 449
- 450

451 **Table 1.** Bacterial strains and plasmids used in this study.

Strain / Plasmid	Description	References
<i>Lb. brevis</i> strains		
UCCLBBS124 (UCC124)	Beer-spoiling strain isolated from spoiled beer keg (Singapore)	(10)
UCCLBBS449	Beer-spoiling strain isolated from unpasteurized spoiled beer (The Netherlands)	(10)
MB569	Non-beer spoiling strain derivative of UCCLBBS124	This study
UCCLB521 (UCC521)	Non-beer spoiling strain isolated from brewery environment (The Netherlands)	(10)
MB569 pNZ44	MB569 carrying pNZ44	This study
MB569 pNZ44: <i>gtfD15</i>	MB569 carrying pNZ44 with <i>gtfD15</i>	This study
UCCLB521 pNZ44: <i>gtfD15</i>	UCCLB521 carrying pNZ44 with <i>gtfD15</i>	This study
<i>L. lactis</i> strains		
NZ9000	Transformation host	(34)
Plasmids		
pNZ44	Transformation vector, chloramphenicol resistance gene	
pNZ44: <i>gtfD15</i>	pNZ44 harboring <i>gtfD15</i>	This study

452

453 **Table 2.** PCR primers used in this study. Incorporated restriction sites are indicated in capital
454 letters.

Primer name	Sequence (5' - 3')	Target	GenBank accession no.
<i>horAF</i>	cgcaactgaggctaactct	<i>horA</i> gene in UCCLBBS124	CP031173
<i>horAR</i>	ggcttgctatgctaggata	<i>horA</i> gene in UCCLBBS124	CP031173
<i>horCF</i>	gtatgcctaagtgcagt	<i>horC</i> gene in UCCLBBS124	CP031172
<i>horCR</i>	cattctctgcctctatac	<i>horC</i> gene in UCCLBBS124	CP031172
<i>orf5F</i>	ctggattgaggtgaggg	<i>orf5</i> gene in UCCLBBS124	CP031172
<i>orf5R</i>	gctgtaagggtagtgtattg	<i>orf5</i> gene in UCCLBBS124	CP031172
pNZ44F	aacaattgaaccatac	pNZ44 promoter	
pNZ44R	gaacgtttcaagccttg	pNZ44 MCS	
pD14F	aaaaaaCTGCAGgtccgaacagcgttcggatt	Gene <i>UCCLBBS124_pD0014</i> in UCCLBBS124_D	CP031173
pD14R	aaaaaaTCTAGAttaactctcgaatagtt	Gene <i>UCCLBBS124_pD0014</i> in UCCLBBS124_D	CP031173
pD15F	aaaaaaCCATGGgcggttgatatttatact	Gene <i>UCCLBBS124_pD0015</i> in UCCLBBS124_D	CP031173
pD15R	aaaaaaTCTAGAtcactcagtttcaattccc	Gene <i>UCCLBBS124_pD0015</i> in UCCLBBS124_D	CP031173
pD16F	aaaaaaCTGCAGgagccttgctatgctagg	Gene <i>UCCLBBS124_pD0016</i> in UCCLBBS124_D	CP031173
pD16R	aaaaaaTCTAGAtcaccgttgctcgt	Gene <i>UCCLBBS124_pD0016</i> in UCCLBBS124_D	CP031173
pD17-19F	aaaaaaCCATGGgggtagaatggtctgtt	Gene <i>UCCLBBS124_pD0017-19</i> in UCCLBBS124_D	CP031173
pD17-19R	aaaaaaTCTAGAttattgataatgaccagcaa	Gene <i>UCCLBBS124_pD0017-19</i> in UCCLBBS124_D	CP031173

455

456

457

458

459

460

461

462 **Table 3.** *Lb. brevis* UCC124 plasmids and genes of interest for beer spoilage.

463

UCC124 plasmids	Size (bp)	ORFs no.	Accession no.	Gene(s) of interest	References
UCCLBBS124_A (UCC124_A)	49,560	42	CP031170		
UCCLBBS124_B (UCC124_B)	23,078	20	CP031171	<i>gtf family 2</i>	(11)
UCCLBBS124_C (UCC124_C)	22,370	27	CP031172	<i>horB, horC, orf5</i>	(7, 9)
UCCLBBS124_D (UCC124_D)	20,971	16	CP031173	<i>horA</i>	(5)

464

465

466

467

468 **Table 4.** Presence and absence of genes of UCCLBBS124_D among *Lb. brevis* BS strains.

Gene	Predicted function	<i>Lb. brevis</i> BS strains						
		UCCLBBS124	UCCLBBS449	UCCLB95	TMW1.2108	TMW1.2111	TMW1.2112	TMW1.2113
<i>UCCLBBS124_pD0014 = UCC124_D14</i>	Cytosine deaminase	+	+	-	+	+	-	+
<i>UCCLBBS124_pD0015 = gtf_{D15}</i>	Glycosyltransferase family 8	+	+	-	+	+	+	+
<i>UCCLBBS124_pD0016 = UCC124_horA</i>	HorA	+	+	-	+	+	-	+
<i>UCCLBBS124_pD0017 = UCC124_D17</i>	Acyl-sn-glycerol-3-phosphate acyltransferase	+	+	-	+	+	+	+
<i>UCCLBBS124_pD0018 = UCC124_D18</i>	Glycosyltransferase family 8	+	+	-	+	+	+	+
<i>UCCLBBS124_pD0019 = UCC124_D19</i>	Acyl-sn-glycerol-3-phosphate acyltransferase	+	+	-	+	+	+	+

469

470

471

472

473

474

475


476 **Table 5.** Effect of phages 3-521 and 521B on *Lb. brevis* strain UCC521 and derivatives.

		<i>Lb. brevis</i> strains		
		UCC521	UCC521 pNZ44	UCC521 pNZ44:gtf _{D15}
	EOP (Efficiency Of Plaquing)	1.00	0.58 ± 0.29	0.64 ± 0.21
Phage 521B	Plaque morphology	Small clear plaques	Small clear plaques	Faint plaques
	Phage titre after O/N propagation* (PFU/mL)	2.90E+09	2.30E+09	3.00E+07
	EOP	1.00	1.52 ± 0.20	1.19 ± 0.19
Phage 3-521	Plaque morphology	Small clear plaques	Small clear plaques	Faint plaques
	Phage titre after O/N propagation* (PFU/mL)	4.30E+09	1.80E+09	4.80E+07


477 *Overnight propagation (O/N) was realized with a starting phage titre of 10⁶ PFU/mL (results

478 are average of triplicate assays).

479



480 **Figure 1.** Growth of the WT beer-spoiling strain *Lb. brevis* UCC124 (A) and its plasmid-
 481 cured derivative MB569 (B) in beer, MRS broth, MRS broth at pH4 and MRS broth
 482 supplemented with 5.4 % ethanol or 30 ppm *iso-α*-acids.



483 **Figure 2.** Number of viable bacteria (CFU/mL) of the WT BS strain *Lb. brevis* UCC124, the
 484 derivative MB569 +/- the empty plasmid pNZ44 and MB569 carrying the gene *gtf_{D15}* after
 485 growth in (A) MRS broth containing 30 ppm *iso*- α -acids for 72 h and (B) beer for 96 h
 486 ($P < 0.05$). CFU/mL of the WT BS strain *Lb. brevis* UCC124, the NBS UCC521 +/- the empty
 487 plasmid pNZ44 and the NBS UCC521 carrying the gene *gtf_{D15}* after growth in (C) MRS
 488 broth containing 30 ppm *iso*- α -acids and (D) beer for 96 h ($P < 0.05$).