

CORRECTION

Correction: Comparison of “Live High-Train Low” in Normobaric versus Hypobaric Hypoxia

Jonas J. Saugy, Laurent Schmitt, Roberto Cejuela, Raphael Faiss, Anna Hauser, Jon P. Wehrlin, Benjamin Rudaz, Audric Delessert, Neil Robinson, Grégoire P. Millet

The inspired fraction of oxygen (F_{iO_2}) for the normobaric hypoxic condition was incorrectly reported to be $15.8 \pm 0.8\%$. The correct F_{iO_2} value for the NH condition is $17.9 \pm 0.2\%$ and the correct inspired pressure of oxygen (P_{iO_2}) values were 111.1 ± 1.1 vs 111.5 ± 1.0 mmHg in normobaric and hypobaric hypoxia, respectively. The authors wish to clarify and confirm that the simulated altitude was perfectly matched to the real altitude of 2250m. The error appears only in the paper. This correction has no impact on the living altitudes between the two hypoxic conditions, as P_{iO_2} was strictly matched between conditions.

As a result, there are errors in [Fig 4](#). Please see the updated [Fig 4](#) here.



 OPEN ACCESS

Citation: Saugy JJ, Schmitt L, Cejuela R, Faiss R, Hauser A, Wehrlin JP, et al. (2015) Correction: Comparison of “Live High-Train Low” in Normobaric versus Hypobaric Hypoxia. PLoS ONE 10(7): e0133091. doi:10.1371/journal.pone.0133091

Published: July 14, 2015

Copyright: © 2015 Saugy et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.





Fig 4. A. Daily values of inspired pressure of oxygen (P_iO_2 in mmHg) during the Live High-Train Low (LHTL) camps for the normobaric hypoxia (NH) and hypobaric hypoxia (HH) groups. **B.** Mean values of night oxygen pulse saturation (S_pO_2). Data are presented in mean \pm standard error. Pre1-Pre2: measurements before the camps (1150 m, Prémanon, France); D01–D18: measurement during the camps (NH: hypoxic room in Prémanon, France; HH: Fiescheralp, Switzerland). # $P<0.05$, ### $P<0.001$ for differences between groups.

doi:10.1371/journal.pone.0133091.g001

Reference

1. Saugy JJ, Schmitt L, Cejuela R, Faiss R, Hauser A, Wehrli JP, et al. (2014) Comparison of “Live High-Train Low” in Normobaric versus Hypobaric Hypoxia. PLoS ONE 9(12): e114418. doi:[10.1371/journal.pone.0114418](https://doi.org/10.1371/journal.pone.0114418) PMID: [25517507](https://pubmed.ncbi.nlm.nih.gov/25517507/)