



# Time matters: genetic composition and evaluation of effective population size in temperate coastal fish species

Sara M. Francisco and Joana I. Robalo

MARE—Marine and Environmental Sciences Centre, ISPA—Instituto Universitário, Lisbon, Portugal

## ABSTRACT

**Background.** Extensive knowledge on the genetic characterization of marine organisms has been assembled, mainly concerning the spatial distribution and structuring of populations. Temporal monitoring assesses not only the stability in genetic composition but also its trajectory over time, providing critical information for the accurate forecast of changes in genetic diversity of marine populations, particularly important for both fisheries and endangered species management. We assessed fluctuations in genetic composition among different sampling periods in the western Portuguese shore in three fish species.

**Methods.** White seabream *Diplodus sargus*, sand smelt *Atherina presbyter* and shanny *Lipophrys pholis* were chosen, because of their genetic patterns in distinct ecological environments, insight into historical and contemporary factors influencing population effective size ( $N_e$ ), and degree of commercial exploitation. Samples were obtained near Lisbon between 2003 and 2014 and screened for genetic variation with mitochondrial and nuclear markers. Analyses included genealogies, genetic diversities, temporal structures and contemporary  $N_e$ .

**Results.** For mtDNA no temporal structure was detected, while for nDNA significant differences were recorded between some sampling periods for the shanny and the sand smelt. Haplotype networks revealed deep genealogies, with various levels of diversification. The shanny revealed a smaller  $N_e$ /generation when compared to the other species, which, in turn, revealed no evidence of genetic drift for most study periods. These results highlight the fact that temporal variations in genetic pool composition should be considered when evaluating the population structure of fish species with long distance dispersal, which are more vulnerable to recruitment fluctuations.

**Subjects** Aquaculture, Fisheries and Fish Science, Evolutionary Studies, Marine Biology, Population Biology

**Keywords** Effective population size, mtDNA, nDNA, Temporal structure, Temporal stability, Temporal method

## INTRODUCTION

Over the last four decades, we have assembled extensive knowledge on the genetic characterization of marine organisms, mainly concerning the spatial distribution and structuring of populations (e.g., *Knutsen et al., 2007; Plank et al., 2010; Robalo et al., 2013; Riginos et al., 2019; Verry et al., 2020*). With this information, it is now possible and relevant

Submitted 9 December 2019

Accepted 9 April 2020

Published 1 May 2020

Corresponding author

Sara M. Francisco,  
sara\_francisco@ispa.pt

Academic editor

María Ángeles Esteban

Additional Information and  
Declarations can be found on  
page 16

DOI 10.7717/peerj.9098

 Copyright

2020 Francisco and Robalo

Distributed under  
Creative Commons CC-BY 4.0

**OPEN ACCESS**

to understand how these patterns behave through time. Temporal monitoring assesses the genetic composition's stability and trajectories over time, providing critical information for the accurate forecast of changes in the genetic diversity of marine populations. This is increasingly important for both endangered and commercially exploited species, particularly in a context of severe broad-based anthropogenic influences and climate change (*Schwartz, Luikart & Waples, 2007; Cuveliers et al., 2011; Goldstien et al., 2013*).

Temporally replicated sampling also allows the estimate of  $N_e$ —effective population size, one of the most important parameters in both conservation and evolutionary biology (*Waples & Do, 2010; Hare et al., 2011*). From a genetic perspective,  $N_e$  is defined as the size of an ideal population that has the same rate of change in allele frequencies and heterozygosity as the observed population (*Wright, 1931*). Estimates of  $N_e$  may be used to assess the loss of genetic variation, increases in inbreeding, accumulation of mutations, and effectiveness of selection (*Waples & Do, 2010; Hare et al., 2011*). In the short to medium term, potential reductions in  $N_e$  may lead to inbreeding depression and/or drastic loss of genetic diversity (*Frankham, Bradshaw & Brook, 2014*), particularly in commercially overexploited populations, populations inhabiting highly degraded ecosystems or located near the margins of the species distribution. Loss of genetic diversity may take a long time to recover through mutation, thus limiting or impeding the adaptive response to environmental changes (*Lynch & Lande, 1998*).

Several approaches have been developed in the past decades to overcome difficulties in directly measuring  $N_e$  in the marine realm (e.g., *Wang, 2005; Luikart et al., 2010; Hare et al., 2011; Wang, Santiago & Caballero, 2016* for a review). Traditional demographic estimators based on Capture-Mark-Recapture (e.g., Jolly-Seber Model (*Jolly, 1965; Seber, 1965*), and multinomial approach in open populations (*Crosbie & Manly, 1985*)) have been complemented with genetic methods: single-sample  $N_e$  estimators (e.g., linkage disequilibrium method (*Hill, 1981*), heterozygote-excess method (*Pudovkin, Zaykin & Hedgecock, 1996*), sibship frequency (*Wang, 2009*), molecular coancestry (*Nomura, 2008*)) and two-sample  $N_e$  estimators (temporal change in allele frequencies (*Waples, 1989; Wang, 2001*)).

A frequently used genetic approach to measure contemporary changes in  $N_e$  has been the Temporal Method (*Nei & Tajima, 1981*). This method depends on estimates of allelic frequencies in two or more points in time, assuming that the observed genetic drift between two sampling moments will be more pronounced in small populations. *Jorde & Ryman (1995)* adjusted the method to incorporate the effects of age structure, demonstrating that the magnitude of changes in allelic frequencies was also determined by age-specific birth and survival rates, in addition to the actual size and the sampling interval. Later, these authors presented a direct extension of the previous model, allowing its application to temporal samplings of single cohorts born within a given number of years (*Jorde & Ryman, 2007; Jorde, 2012*).

Temporal genetic changes in coastal species have been linked to the stochastic nature of oceanographic conditions, including upwelling systems, fronts, eddies and sharp discontinuities in physicochemical variables (*Selkoe et al., 2006; White et al., 2010*). Moreover, the stability of allele frequencies in natural populations can be deeply influenced

by extreme events (anthropogenic and/or natural disturbing factors) (Allison *et al.*, 2003). Additionally, intrinsic biological and ecological factors associated with the species life cycle, also shape the genetic pattern of marine organisms, namely inshore-offshore spawning, pelagic larval duration and other early life-history traits (Riginos & Victor, 2001; Galarza *et al.*, 2009).

Genetic monitoring through time, specifically, the temporal method, has been successfully employed to evaluate the temporal stability and estimate  $N_e$  in several species, including some stocks with high economic significance such as the European hake *Merluccius merluccius* (Pita *et al.*, 2017), the brown trout *Salmo trutta* (Serbezov *et al.*, 2012) and the Atlantic cod *Gadus morhua* (Therkildsen *et al.*, 2010). In the present study, three coastal fishes were chosen to cover a wide range of contrasting environmental and ecological traits: the white seabream *Diplodus sargus* L. 1758 (Pisces: Sparidae); the sand-smelt *Atherina presbyter* Cuvier 1829 (Pisces: Atherinidae); and the shanny *Lipophrys pholis* L. 1758 (Pisces: Blenniidae).

Although they are not threatened—the three species are listed as least concerned—their population trend is unknown, and they can be locally vulnerable due to various reasons. The white-seabream is a commercially important species throughout European shores, wide-spread and locally abundant (Pollard *et al.*, 2014) and, although no major threats have been identified, some local overfishing may occur and lead to reduction in population size. The sand-smelt is commercially exploited as life bait for tuna and local coastal development can be a threat (Gon, 2015). In the long-term, climate change could be a problem for this temperate species. Finally, no major threats are found across the shanny's distribution range (Williams & Craig, 2014). This species can be used as an indicator for pollution monitoring due to several of its characteristics, including restricted home range and high sensitiveness to organic contaminants (Lima *et al.*, 2008).

These target species provide crucial insight into historical and present factors influencing effective population sizes and genetic patterns in distinct ecological environments and degrees of commercial interest. In this study, we assessed the changes in genetic composition in different points in time in the western Portuguese shore, using a mitochondrial and a nuclear marker (chosen due to their extensive use in the past decade). The specific objectives were: (1) to compare the inter-annual variation in genetic composition and structure of coastal species with contrasting traits, and (2) to test the temporal model and its potential for  $N_e$  estimation in populations with overlapping generations.

## MATERIALS & METHODS

### Target species

The white seabream *Diplodus sargus* is a coastal species in the north-eastern Atlantic ranging from Senegal to the Bay of Biscay, including the archipelagos of Canaries, Madeira and Azores, Mediterranean and Black Sea (Bauchot & Hureau, 1986). Eggs are planktonic, hatching after 3 days at 15–17 °C (Morato *et al.*, 2003). Larvae are also planktonic, settling after a pelagic larval duration (PLD) of 14–19 days (Di Franco *et al.*, 2011). Adults have considerable swimming ability and tend to remain near the coast. This is a commercially

exploited fish, with increasing landings per year in the last 40 years (FAO, 2019). Some degree of overfishing can be empirically inferred, even if stocks show no evidence of decline, since larger *D. sargus* are mainly spotted within Marine Protected Areas. Previous studies on the genetic structure of the white seabream revealed isolation-by-distance (IBD), suggesting genetic isolation over large geographic distances (e.g., Domingues et al., 2007). In contrast, allozyme studies reported significant divergences in cohorts sampled over a 6-month period in a Mediterranean population (Lenfant & Planes, 2002; Planes & Lenfant, 2002).

The sand smelt *Atherina presbyter* is an inshore marine species, also present in estuaries and coastal lagoons. Its distribution area comprises the British Isles and southern North Sea to the Canary Islands and Mauritania (Quignard & Pras, 1986), and has also been reported in the Azores (Santos, Porteiro & Barreiros, 1997). This species spawns in shallow waters, the eggs are demersal and attached to vegetation, and larvae hatch after 15–16 days with 6.7–7.5 mm TL at a temperature of 15 °C (Bamber, Henderson & Turnpenny, 1985). The larval stage is very short (the hatching larvae are well developed and ready to start exogenous feeding), which likely restricts passive dispersal. Migratory movements of adult sand-smelts along exposed shores are probably difficult, although they are active swimmers in the water column. Our current understanding of gene flow in *A. presbyter* indicates a structured population across its range, also showing IBD (Francisco et al., 2009). A previous mtDNA study revealed temporal stability in the genetic composition of the sand-smelt from the Portuguese west coast (Francisco & Robalo, 2015).

The shanny *Lipophrys pholis* is a rocky intertidal resident fish, very common in western European shores, ranging from Norway to Mauritania and from the Azores and Madeira to the entrance of the Mediterranean (Zander, 1986). The eggs are demersal, guarded by the male, and hatch after 16 days with 5.0 mm total length at a temperature of 17 °C (Almada et al., 1992; Faria et al., 2002). The larvae hatch in a well-developed stage and settle at 13–14 mm TL after a PLD of ca. 29 days at a temperature of 15.5–17.5 °C (Faria et al., 2002). Juveniles and adults show weak swimming capabilities and, consequently, restricted movements within the same rocky stretch (Faria, Almada & Goncalves, 1996). Previous molecular work on the shanny, using both nuclear and mitochondrial markers, strongly suggests panmixia across its distribution (Francisco, Vieira & Almada, 2006; Stefanni et al., 2006; Francisco et al., 2011). Contrary to *A. presbyter*, the mtDNA study of *L. pholis* detected significant genetic differentiation between some sampling years (Francisco & Robalo, 2015).

### Sampling scheme

Individuals of *L. pholis*, *D. sargus* and *A. presbyter* were sampled near Lisbon, in S. Pedro do Estoril (38°42'N, 9°22'W) and Fonte-da-Telha (38°34'N, 9°11'W) (Table 1) between 2003 and 2014. Both sampling locations are in the vicinity of marine protected areas and present heterogeneous rocky habitats mixed with sandy patches (e.g., Henriques, Gonçalves & Almada, 1999; Faria & Almada, 2001). They are known to harbour a relevant number of post-larvae and juveniles for several taxa, being important settlement and recruitment areas (e.g., Borges et al., 2009; Vinagre et al., 2018). Juveniles of each year were collected in intertidal rocky pools (*L. pholis*) and beach channels (*D. sargus* and *A. presbyter*). A small

**Table 1** Diversity measures for sampling periods of *Diplodus sargus*, *Atherina presbyter* and *Lipophrys pholis* based on the control region of the mitochondria: number of sequences ( $N$ ), number of haplotypes ( $N_h$ ), percentage of private haplotypes (%Ph), haplotype richness ( $R$ ), private allelic richness ( $pR$ ), haplotype diversity ( $h$ ), nucleotide diversity ( $\pi$ ) and mean number of pairwise differences ( $k$ ).

Species	Sampling period	$N$	$N_h$	%Ph	$R$	$pR$	$h$	$\pi$	$k$
<i>Diplodus sargus</i>	2006	20	19	68.42	20.000	16.008	0.995	0.034	13.058
	2009	39	30	70.00	16.441	11.55	0.974	0.032	12.093
	2011	27	22	59.09	15.379	9.476	0.983	0.033	12.479
	2014	95	73	79.45	16.431	11.643	0.993	0.029	11.135
	All	181	120	–	–	–	0.990	0.031	11.690
<i>Atherina presbyter</i>	2005	34	29	68.96	20.235	12.423	0.986	0.022	8.012
	2012	91	63	69.84	17.640	9.352	0.984	0.021	7.780
	2013	61	48	68.08	17.973	9.396	0.992	0.021	7.866
	2014	95	56	62.50	16.541	7.922	0.975	0.022	8.054
	All	281	155	–	–	–	0.984	0.021	7.931
<i>Lipophrys pholis</i>	2003	30	26	69.23	21.634	13.211	0.991	0.031	11.614
	2013	97	73	87.61	23.708	15.291	0.988	0.028	10.844
	2014	99	88	93.18	26.240	18.034	0.997	0.031	11.977
	All	226	171	–	–	–	0.995	0.030	11.522

sample of fin clip was collected, preserved in 96° ethanol and deposited in ISPA-IU/MARE collections. All sampling and handling of fish were conducted according to established animal welfare guidelines (ORBEA-ISPA, Animal Welfare Body) and following the relevant legislation, as none of the sampled species are endangered or protected in Portugal.

### DNA extraction, amplification and sequencing

Total genomic DNA was extracted from about 20 mg of tissue with REDEExtract-N-mp kit (Sigma-Aldrich) following manufacturer's instructions. Polymerase Chain Reaction (PCR) amplification was performed for two fragments, the mitochondrial control region (CR) and an intron of the nuclear *S7* ribosomal protein gene (*S7*), with the primer pairs: LPro1 (5'-ACT CTC ACC CCT AGC TCC CAA AG-3') and HDL1 (5'-CCT GAA GTA GGA ACC AGA TGC CAG-3') (CR for the three species) (Ostellari et al., 1996), S7RPEX1F (5'-TGG CCT CTT CCT TGG CCG TC-3') and S7RPEX2R (5'-AAC TCG TCT GGC TTT TCG CC-3') (first intron of the *S7* for the shanny and the white seabream), S7RPEX2F (5'-AGC GCC AAA ATA GTG AAG CC-3') and S7RPEX3R (5'-GCC TTC AGG TCA GAG TTC AT-3') (second intron of the *S7* for the sand-smelt) (Chow & Hazama, 1998). The PCR protocol was performed in a 20  $\mu$ l total reaction volume with 10  $\mu$ l of REDEExtract-N-amp PCR mix (Sigma-Aldrich), 0.8  $\mu$ l of each primer (10  $\mu$ M), 4.4  $\mu$ l of Sigma water and 4  $\mu$ l of template DNA. An initial denaturation at 94 °C for 3 min was followed by 35 cycles (denaturation at 94 °C for 30/45 s, annealing at 55/58 °C for 30/45 s, and extension at 72 °C for 1 min; values CR/*S7*, respectively) and a final extension at 72 °C for 10 min on a Bio-Rad MyCycler thermal cycler. The same primers were used for the sequencing reaction and PCR products were purified and sequenced at STABVIDA (Portugal, <http://www.stabvida.net>) and GATC (Germany, <http://www.gatc-biotech.com>).

Sequences were edited with Codon Code Aligner (<http://www.codoncode.com/index.htm>) and aligned with Clustal X2 (Larkin *et al.*, 2007). For S7 both strands of the same specimen were recovered, whenever possible, following the approach of Sousa-Santos *et al.* (2005). This approach takes advantage of the presence of indels in each nuclear marker and uses them to accurately reconstruct the individual haplotypes without the need for probabilistic estimation. Sequences were deposited in GenBank (Accession numbers MG992598–MG992888, MH024090–MH024357, MH030878–MH031272). Additional sequences from previous work (Francisco *et al.*, 2006; Francisco, Vieira & Almada, 2006; Domingues *et al.*, 2007; Francisco *et al.*, 2008; Francisco & Robalo, 2015) were retrieved from GenBank (Tables S1–S3 in Data S1).

### DNA analyses

The appropriate model of sequence evolution for the CR and S7 of each species was determined using the software jModeltest v.2.1.10 (Guindon & Gascuel, 2003; Durraba *et al.*, 2012), under the Akaike Information Criterion (AIC) (Nei & Kumar, 2000). Parsimony networks estimated with TCS version 1.21 (Clement, Posada & Crandall, 2000) were used to analyse relationships among haplotypes and to compute outgroup weights, based on parsimony methods (Templeton, Crandall & Sing, 1992). Visualization and network layout were edited with tcsBU (Múrias dos Santos *et al.*, 2016).

ARLEQUIN software package v.3.5 (Excoffier & Lischer, 2010) was used to estimate the genetic diversity ( $k$ —mean number of pairwise differences (Tajima, 1983);  $\pi$ —nucleotide diversity; and  $h$ —haplotype diversity (Nei, 1987)) within each sampling period, to perform analyses of molecular variance (AMOVA) (Excoffier, Smouse & Quattro, 1992) and to compute pairwise  $F_{ST}$ s. The  $\chi^2$  test (Salicru *et al.*, 1993) was used to assess the significance of differences in haplotype diversity among temporal samples. The software HP-Rare (Kalinowski, 2005) was used to estimate allelic richness  $R$  and private allelic richness  $pR$ , using rarefaction to correct for sample-size bias. Principal Coordinate Analysis (PCoA) was performed with GenAlEx 6.5 (Peakall & Smouse, 2006; Peakall & Smouse, 2012) to visualize the patterns of temporal genetic structure in a bi-dimensional space. The analyses of the S7 intron were also run in ARLEQUIN, after allowing the program to reconstruct the haplotypes present, using the ELB algorithm (Excoffier, Laval & Balding, 2003). The same software was used to perform the exact probability tests for deviations from the Hardy–Weinberg equilibrium (HWE) (Guo & Thompson, 1992).

Contemporary effective population size ( $N_e$ ) and genetic drift ( $F_s$ ) were estimated using TempoFs (Jorde & Ryman, 2007), under the temporal method of allele frequency shifts. The program reports  $F_s'$  (genetic drift corrected for sampling plan) and  $N_e$  per generation. For this approach, we used sampling plan II (individuals sampled before reproduction and not returned to the population; Waples, 1989) and a generation time of 2 yr for *L. pholis* (Milton, 1983; Faria, Almada & Goncalves, 1996), 2 yr for *D. sargus* (Morato *et al.*, 2003) and 1 yr for *A. presbyter* (Turnpenny, Bamber & Henderson, 1981).

## RESULTS


### Mitochondrial data

For *D. sargus*, a total of 181 CR sequences were obtained, comprising 120 haplotypes and 119 polymorphic sites (117 transitions, 15 transversions and two indels) (Table S1). For the seabream CRs, 16.67% of haplotypes were shared between sampling periods. For *A. presbyter*, 281 CR sequences were obtained corresponding to 155 haplotypes (368bp long fragment). Differences among haplotypes corresponded to 70 polymorphic sites (63 transitions, 19 transversions and six indels) (Table S2). For the sand-smelt, 16.13% of haplotypes were shared between sampling periods. A total of 226 shannies were sequenced for the CR (380 bp), corresponding to 171 distinct haplotypes (Table S3). A total of 108 polymorphic sites were found, corresponding to 105 transitions, 26 transversions and eight indels. Only 7.02% of haplotypes were shared among sampling periods. The generalized time-reversible (GTR) + invariable sites (I) (Tavare, 1986) was estimated as the optimal molecular evolutionary model for the CR of the three species.

The statistical parsimony networks constructed with the CR datasets revealed multiple levels of diversification and deep genealogies, seemingly without temporal structure (Fig. 1). The inferred ancestral haplotype for the CR of *D. sargus* included specimens collected in 2009, 2011 and 2014 (outgroup weight: 0.071). In this network, most of the haplotypes were not arranged in a star-like pattern, and some of the branches reached a maximum of 46 mutational steps, and 18 steps from the ancestral haplotype. For *A. presbyter*, the network built with the CR dataset presented several star-like patterns around haplotypes, including the haplotype inferred as ancestral (outgroup weight: 0.051), which comprised individuals from every sampling period. The network was dominated by two haplotypes shared by most sampling periods. In this network, also deep but with less diversification, some of the haplotypes differed 16 mutational steps from ancestor. For *L. pholis*, the CR network displayed several haplotypes, including the estimated ancestor, in the centre of star-like patterns. This ancestral haplotype included specimens collected in the three sampling periods (outgroup weight: 0.053), and some haplotypes differed as many as 17 mutations (Fig. 1). Interestingly, for all three species, haplotypes that initially had been merely inferred were sampled in more recent periods.

For the CR, genetic diversity indices were generally high and similar across years (Table 1). Salicru tests revealed non-significantly different levels of haplotype diversity among sampling years:  $\chi^2 = 1.557$  ( $p = 0.816$ ),  $\chi^2 = 4.342$  ( $p = 0.362$ ) and  $\chi^2 = 3.022$  ( $p = 0.388$ ), for *D. sargus*, *A. presbyter* and *L. pholis* respectively. The CR allelic richness and private allelic richness yielded distinct results for the three species, with no pattern for the seabream, increasing over sampling time in the shanny, and decreasing in the sand-smelt (Table 1).

Global genetic differentiation among sampling years was not significant for CR in any of the analysed species—AMOVA results (white seabream:  $F_{ST} = 0.003$ ,  $p = 0.250$ ; sand-smelt:  $F_{ST} = 0.002$ ,  $p = 0.312$ ; shanny:  $F_{ST} = 0.009$ ,  $p = 0.074$ ). Comparisons between periods were not significant in *D. sargus* and *A. presbyter*. However, for *L. pholis*, significant differentiation was detected between 2013 and 2014 samples ( $F_{ST} = 0.013$ ,  $p$



**Figure 1** Haplotype network for the CR of (A) *Diplodus sargus*, (B) *Atherina presbyter* and (C) *Lipophrys pholis*. The haplotype with the highest out group probability is displayed as a square, other haplotypes as circles. The area of the circles is proportional to each haplotype frequency. Colours refer to the year of sampling. In the case where haplotypes are shared among sampling periods, shading is proportional to the frequency of the haplotype in each period.

Full-size  DOI: [10.7717/peerj.9098/fig-1](https://doi.org/10.7717/peerj.9098/fig-1)



**Table 2** Estimates of contemporary effective population size and genetic drift for *Diplodus sargus*, *Atherina presbyter* and *Lipophrys pholis*. Mitochondrial control region (CR), nuclear S7 ribosomal protein gene (S7), estimated drift ( $F_s'$ ), estimated effective population size per generation ( $N_e/gen$ ).

Species	Marker	Sampling interval	$F_s'$	$N_e/gen$
<i>Diplodus sargus</i>	CR	2006–2009	0.029	102
		2009–2011	−0.005	∞
		2011–2014	0.005	426
	S7	2006–2009	0.022	68
		2009–2011	−0.008	∞
		2011–2014	−0.005	∞
<i>Atherina presbyter</i>	CR	2005–2012	−0.001	∞
		2012–2013	−0.001	∞
		2013–2014	0.001	1,282
	S7	2005–2012	0.352	10
		2012–2013	0.015	34
		2013–2014	−0.006	∞
<i>Lipophrys pholis</i>	CR	2003–2013	0.008	612
		2013–2014	0.010	94
	S7	2003–2013	0.534	5
		2013–2014	0.010	53

= 0.028). No species showed marked patterns of temporal genetic structure in CR and, therefore, the PCoA was not performed.

Estimates of effective population size based on the CR of the three species are given on Table 2, after correction for haploid data. *L. pholis* revealed a smaller  $N_e/generation$  when compared to the other species and *A. presbyter* showed no evidence of genetic drift for some of the study periods (negative values of  $F_s'$ ), which resulted in much higher  $N_e/generation$  ( $\sim inf$ ).

### Nuclear data

For *D. sargus*, 302 sequences of S7 were obtained with 308bp (corresponding to 151 individuals), comprising 99 haplotypes and 50 polymorphic sites (29 transitions, 21 transversions and 2 indels) (Table S1). Only 12.12% of haplotypes of S7 were shared between sampling periods for the seabream. The amplification of S7 ( $N = 286,572$  sequences) in *A. presbyter* resulted in a 201bp fragment, corresponding to 35 haplotypes (Table S2). A total of 25 polymorphic sites (6 transitions, 12 transversions and 7 indels) were found, and 57.14% of the haplotypes were shared between sampling periods. For the S7 of the shanny, a total of 360 sequences (180 individuals) were obtained (576 bp), comprising 24 distinct haplotypes (26.09% shared between sampling periods) (Table S3). The fragment yielded 20 polymorphic sites, corresponding to 9 transitions, 8 transversions and 3 indels. The unequal transitional substitution (TIM3) + invariable sites (I) + rate variation among sites (G) (Posada, 2003) was estimated as the optimal molecular evolutionary model for the S7 of the seabream. For the S7 of the sand-smelt, the selected model was the equal transitional

**Table 3** Diversity measures for sampling periods of *Diplodus sargus*, *Atherina presbyter* and *Lipophrys pholis* based on the S7: number of sequences ( $N$ ), number of haplotypes ( $N_h$ ), percentage of private haplotypes (%Ph), haplotype richness ( $R$ ), private allelic richness ( $pR$ ), haplotype diversity ( $h$ ), nucleotide diversity ( $\pi$ ), mean number of pairwise differences ( $k$ ) and test of deviations from the Hardy–Weinberg equilibrium observed/expected heterozygosity ( $Ho/He$ ).

Species	Sampling period	$N$	$N_h$	%Ph	$R$	$pR$	$h$	$\pi$	$k$	$Ho/He$
<i>Diplodus sargus</i>	2006	20	20	90.00	24.961	15.461	0.963	0.012	3.709	1.000/0.963
	2009	35	33	69.70	20.583	8.081	0.942	0.013	3.968	0.943/0.942
	2011	19	19	78.95	19.000	8.052	0.953	0.012	3.863	0.895/0.953
	2014	77	67	86.57	24.368	11.301	0.961	0.013	4.006	0.961/0.962
	All	151	125	–	–	–	0.981	0.013	3.948	–
<i>Atherina presbyter</i>	2005	38	4	50.00	4.000	2.070	0.242	0.002	0.380	0.034/0.033*
	2012	90	17	17.65	14.334	2.082	0.816	0.001	1.990	0.129/0.142*
	2013	64	19	21.05	16.223	4.262	0.743	0.009	1.772	0.099/0.099
	2014	94	26	26.92	17.830	4.605	0.770	0.010	2.063	0.102/0.115
	All	286	42	–	–	–	0.734	0.009	1.779	–
<i>Lipophrys pholis</i>	2003	20	7	51.71	7.000	4.637	0.360	0.002	1.121	0.400/0.360
	2013	76	13	53.85	7.162	2.608	0.757	0.004	2.534	0.750/0.757*
	2014	84	12	50.00	6.225	1.489	0.721	0.004	2.327	0.655/0.721*
	All	180	23	–	–	–	0.734	0.004	2.380	–

**Notes.**


\*Significant values of probability  $p$ .

substitution model (TIM1ef) + invariable sites (I) (Posada, 2003), and for the shanny was the unequal transitional substitution model (TIM3) + invariable sites (I) (Posada, 2003).

The S7 haplotype networks of the three species revealed shallower genealogies comparing to the CR (Fig. 2). The networks showed star-like patterns dominated by 2–3 very frequent haplotypes and no temporal structure was evident. The inferred ancestral haplotypes were shared among sampling periods, yielding outgroup weights of 0.067, 0.117 and 0.190, for *D. sargus*, *A. presbyter* and *L. pholis*, respectively.

The S7 genetic diversity indices of the three species are summarized in Table 3. The pattern of haplotypic diversity stability across years registered for the CR was only recovered for the S7 of *D. sargus* ( $\chi^2 = 1.289$ ,  $p = 0.863$ ). Salicru tests yielded significant differences for the haplotype diversity between sampling year for both *A. presbyter* ( $\chi^2 = 73.080$ ,  $p < 0.001$ ) and *L. pholis* ( $\chi^2 = 17.864$ ,  $p < 0.001$ ). Similarly, the Markov chain tests yielded significant deviations from the HWE for some years of the sand-smelt and shanny datasets ( $p < 0.001$ ) (Table 3).

For the S7 of the white seabream, the AMOVA results revealed no significant temporal structure among sampling years ( $F_{ST} = 0.003$ ,  $p = 0.204$ ). Pairwise  $F_{ST}$  comparisons were also non-significant. A different pattern was recovered for the S7 of the sand-smelt and the shanny, as significant global genetic differentiation was yielded ( $F_{ST} = 0.020$   $p < 0.001$  and  $F_{ST} = 0.067$   $p < 0.001$ , respectively). For *A. presbyter* significant differences were found between 2005 and every other period ( $p < 0.001$ ). For *L. pholis* pairwise  $F_{ST}$  was significant between 2003 and the more recent sampling times ( $p < 0.001$ ). The PCoA corroborated these patterns of temporal genetic structure, with the first axis explaining 94% and 99% of




**Figure 2** Haplotype network for the S7 of (A) *Diplodus sargus*, (B) *Atherina presbyter* and (C) *Lipophrys pholis*. The haplotype with the highest outgroup probability is displayed as a square, other haplotypes as circles. The area of the circles is proportional to each haplotype frequency. Colours refer to the year of sampling. In the case where haplotypes are shared among sampling periods, shading is proportional to the frequency of the haplotype in each period.

Full-size DOI: 10.7717/peerj.9098/fig-2

the observed variance, for the S7 of *A. presbyter* and *L. pholis*, respectively, and suggesting two groups associated with older vs recent sampling periods (Fig. 3).

For the S7 fragment, TempoFs yielded lower  $N_e$ /generation estimates for the shanny (Table 2). Both the white seabream and the sand-smelt revealed negative values of  $F_s'$



**Figure 3** Principal Coordinate Analysis for the S7 of (A) *Atherina presbyter* and (B) *Lipophrys pholis*.  
Full-size [DOI: 10.7717/peerj.9098/fig-3](https://doi.org/10.7717/peerj.9098/fig-3)

for some of the more recent sampling intervals, i.e., no evidence of genetic drift and, conversely, very high  $N_e$ /generation values ( $\sim\infty$ ) (Table 2).

## DISCUSSION

### Temporal genetic stability

Traditionally population genetic and phylogeographic studies have disregarded the temporal dimension, pooling samples based on their geographical origin and ignoring their collecting period (Viñas, Alvarado-Bremer & Pla, 2004; Charrier et al., 2006; Luttikhuisen et al., 2008; Gonzalez-Wanguemert et al., 2011; Stefanni et al., 2015). One of our main result is the subsequent appearance of inferred haplotypes, clearly reinforcing the importance of temporal evaluations. Similar results were also found using mitochondrial markers in several other species (e.g., *Genypterus capensis* (Henriques et al., 2017), *Prionace glauca* (Veríssimo et al., 2017), *Seriola lalandi* in the Pacific Ocean (Sepúlveda & González, 2017)).

Another important finding was the stability across years in these three ecologically distinct species, reinforcing the preliminary work by Francisco & Robalo (2015). This high degree of similarity was found in (a) the high genetic diversity, with little variation between sampling periods, especially true for the mitochondrial data (Table 1); (b) the deep genealogies with several diversification levels (Fig. 2); and (c) the absence of temporal structure for the mitochondrial data. In contrast, we detected temporal structure and some

marked differences among sampling years in both the shanny and the sand-smelt for S7 (Fig. 3). In fact, this differentiation between sampling periods was also detected for the CR of *L. pholis*, notwithstanding the absence of temporal genetic structure.

Stochasticity in larval survival and transport, due to variation in oceanographic conditions, and genetic differentiation among temporal samples are broadly documented in the literature (e.g., [Selkoe et al., 2006](#); [Selkoe et al., 2010](#)). *Lipohrys pholis* lays fewer eggs and has a longer planktonic larval stage when compared to both *A. presbyter* and *D. sargus*. Therefore, while its male-guarding behaviour can promote larval retention near rocky coasts, this is easily disrupted by atypical or severe events, when larvae may be transported very large distances if storm conditions prevail. Thus, the shanny's higher differentiation (marked differences among sampling periods in both markers and temporal structure for S7) might be closely related with its greater vulnerability to recruitment fluctuations. Conversely, the white seabream revealed a greater degree of temporal stability for both molecular markers. In contrast, allozyme studies in *D. sargus* in the Mediterranean found rapid genetic change within a population, assumed to be driven by genetic drift ([Lenfant & Planes, 2002](#); [Planes & Lenfant, 2002](#)). We suggest that these apparently contradictory findings may be due to the type of molecular marker. [Planes & Lenfant \(2002\)](#) associated the large variation in the white seabream's reproductive success with linkage disequilibrium and genetic relatedness shown in their allozyme data. In our study (mtDNA and nDNA) the absence of temporal structure is probably associated with *D. sargus*' higher fecundity ([Goncalves & Erzini, 2000](#); [Mouine et al., 2007](#)), longer spawning season ([Dias et al., 2016](#)) and relatively short PLD ([Di Franco et al., 2011](#)). Temporal stability has been described in several other marine species, such as *Solea solea* ([Cuveliers et al., 2011](#)), *Pagrus auratus* ([Bernal-Ramírez et al., 2003](#)), *Meganyctiphanes norvegica* ([Papetti et al., 2005](#)) and *Ammodytes marinus* ([Jiménez-Mena et al., 2019](#)).

Also noteworthy is the significant deviation from HWE in the sand-smelt and the shanny, the two species with most contrasting early life-history traits. Although unexpected for a relatively long PLD species like *L. pholis*, this result was previously reported in other marine organisms (e.g., [Chapman, Ball & Mash, 2002](#); [Karlsson & Mork, 2005](#); [Pérez-Portela et al., 2019](#)). Departure from HWE has been ascribed to natural selection, migration, Wahlund effect, null alleles, inbreeding and/or phenotypic assortative mating (e.g., [Addison & Hart, 2005](#); [Karlsson & Mork, 2005](#); [Gonzalez, Beerli & Zardoya, 2008](#); [Garnier-Géré & Chikhi, 2013](#)). Although we were not able to explain these results, we are aware that HW disequilibrium violates the assumptions behind temporal  $N_e$  estimation (e.g., [Nei & Tajima, 1981](#); [Serbezov et al., 2012](#); [Wang, Santiago & Caballero, 2016](#)).

### Estimates of contemporary effective population size

In the present work, the estimated population effective size per generation was smaller for the shanny than the other two species, corroborating preliminary findings by [Francisco & Robalo \(2015\)](#). The negative  $F_s'$  found for some of the more recent sampling intervals, for both white seabream and sand-smelt, evidenced no genetic drift (CR and S7). Conversely, the  $N_e$ /generation values for these species were very high ( $\sim\text{inf}$ ) (Table 2). Genetic stochasticity is most likely weaker in marine organisms with such large  $N_e$  ([Palstra &](#)

Ruzzante, 2008), as found in fish species like *S. solea* (Cuveliers et al., 2011) and *G. morhua* (Therkildsen et al., 2010) (see Marandel et al. (2019) for a review). Lower  $N_e$ /generation was found in fish species such as the herring *Clupea harengus* (Larsson et al., 2010), the silver seabream *P. auratus* (Hauser et al., 2002) and the thornback ray *Raja clavata* (Chevolot et al., 2008). The extremely low  $N_e$ /generation found in the first sampling interval of *L. pholis* and *A. presbyter* (S7) (Table 2) might be an underestimation due to statistical artefacts, immigration from other populations, or other factors (e.g., Palstra & Ruzzante, 2008).

Considering the target species and sampling scheme, the adjusted temporal method was, in our opinion, the best choice for estimating contemporary  $N_e$ . Not only is this method considered the least biased for species with overlapping generations (e.g., Jorde & Ryman, 2007; Waples & Yokota, 2007), the analysis of consecutive cohorts is also the best way to reduce  $N_e$  estimation bias (e.g., Jorde & Ryman, 1995; Luikart et al., 2010). Nevertheless, several factors can confound  $N_e$  estimates, including the aforementioned departure from HWE. For instance, temporal estimates of  $N_e$  may be biased by potential migratory movements within metapopulations that can override the effects of genetic drift (e.g., Palstra & Ruzzante, 2008; Waples & Do, 2010; Ryman et al., 2014). In fact, the assumption of complete isolation of the study population is frequently violated, and the resulting bias is generally of unknown magnitude (e.g., Ryman et al., 2014). In this study, this might result in biased  $N_e$  estimates for the sand-smelt and white seabream, for which IBD patterns have been recorded (Domingues et al., 2007; Francisco et al., 2009). Sampling strategy is another potential source of bias (e.g., Jorde & Ryman, 2007; Waples & Yokota, 2007; Luikart et al., 2010), including low sample size and sex-biased or age-biased sampling. In the present work, uneven sample size between years (considerably lower  $N$  for the first sampling point) likely affected our  $N_e$  estimates. Species life-history and reproductive strategy can also hamper interpretation of  $N_e$  estimates (e.g., Waples & Yokota, 2007). Sequential hermaphrodites, such like the protandrous *D. sargus* (Bauchot & Hureau, 1986), present sex ratios skewed towards the initial sex (Charnov & Bull, 1989). According to standard fixed-sex theory, this results in reduced  $N_e$  (Wright, 1938) due to greater genetic drift (Charlesworth, 2009). The very high  $N_e$ /generation values obtained for the white seabream seemingly contradict this prediction (Table 2). This expected lower  $N_e$  was also challenged in recent studies with protogynous species over historical timescales (Coscia et al., 2016), and in analysis of an eco-evolutionary model with ten hypothetical species (Waples, Mariani & Benvenuto, 2018).

Recently, the reliability of contemporary estimates of  $N_e$  has been challenged and questioned (e.g., Wang, Santiago & Caballero, 2016; Marandel et al., 2019). However, these reviews focused on single-point estimation approaches (Linkage-Disequilibrium methods), assuming discrete generations and, implicitly, reduced population sizes. Marandel et al. (2019) found significantly biased estimates in simulations of large populations of fish and therefore a need for extra-large sampling sizes (impracticable in most sampling schemes). Unlike the criticized studies, our work used a temporal method comprising extensions to estimate  $N_e$  in age-structured species (Jorde, 2012).

While the present study used only one geographical location and two genetic markers, these apparent limitations allowed a rigorous comparison with data from several species

collected in the same area over the past fifteen years and using the same two markers (mitochondrial control region and nuclear S7) (e.g., [Domingues et al., 2006](#); [Almada et al., 2012](#); [Robalo et al., 2013](#); [Francisco et al., 2014](#); [Stefanni et al., 2015](#); [Almada et al., 2017](#); [Pappalardo et al., 2017](#)), which facilitated comparison and calibration of genetic diversity results. Thus, it was possible to observe the high genetic diversity pattern found in several coastal fish populations in western Portugal (e.g., *D. vulgaris* ([Stefanni et al., 2015](#)) and *Labrus bergylta* ([Almada et al., 2017](#))).

## CONCLUSIONS

Genetic monitoring through time seeks to disentangle life-trait patterns affecting marine organisms, while deepening our ability to forecast changes in both genetic composition and diversity. This study assessed fluctuations in genetic composition among different sampling periods in the western Portuguese shore using a comparative approach in three species with distinct ecological and life-history characteristics. No temporal structure was detected for the mitochondrial marker, while for nDNA significant differences were recorded among some sampling periods for the shanny and the sand-smelt. The haplotype networks revealed deep genealogies, and one of our major findings was the repeated appearance of previously inferred haplotypes. The shanny revealed a smaller  $N_e$ /generation when compared to the other species but revealed no evidence of genetic drift for most study periods. These results clearly underline that temporal variation in gene pool composition should be considered when evaluating population structure of long larval dispersion fish species, which are more vulnerable to recruitment fluctuations. Comparison between the commercially and non-commercially explored species yielded no conclusive results.

Previous authors suggested increasing the number of time-point estimates (e.g., from pre-harvest and postharvest times around an overexploitation event) to circumvent difficulties in short-term prediction of  $N_e$  ([Pita et al., 2017](#)). Future studies with additional time-points covering a longer time span, additional sampling locations covering the species geographical distribution and additional markers would greatly improve the reliability of the present paper's results. Available tools include a combination of methods of demographic inference and large genomic datasets generated with RAD-seq (restriction site-associated DNA sequencing) (e.g., [Barbato et al., 2015](#)), as recently reported in critically endangered species (e.g., *Carcharias taurus* in [Reid-Anderson, Bilgmann & Stow, 2019](#)) and commercially important species (e.g., *Oncorhynchus kisutch* in [Barria et al., 2019](#); *Solea solea* in [Le Moan et al., 2019](#)). These genomic approaches, with their high-throughput sequencing methods, will likely improve our understanding of recent population demography ([Waples, 2016](#)). Applying and extending this framework to species with distinct features (e.g., life-history traits), conservation status and commercial importance would be of paramount importance to detect global patterns and predict the ability of species to adapt to future changes.

## ACKNOWLEDGEMENTS

We thank Frederico Almada, Ana Faria, Ana Lopes and Pedro Duarte Coelho for their help with field work. We are grateful to Per Erik Jorde for helpful suggestions on the temporal data analysis. We also thank André Levy for his valuable language editing.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This study was supported by Fundação para a Ciência e Tecnologia (FCT) Portugal, through the strategic project UID/MAR/04292/2019 granted to MARE, by ICES Science Fund of the project POPSIZE, and by an FCT fellowship (SFRH/BPD/84923/2012) granted to Sara M. Francisco. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:

Fundação para a Ciência e Tecnologia (FCT) Portugal, through the strategic project: UID/MAR/04292/2019.

ICES Science Fund of the project POPSIZE.

FCT fellowship: SFRH/BPD/84923/2012.

### Competing Interests

The authors declare there are no competing interests.

### Author Contributions

- Sara M. Francisco conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Joana I. Robalo conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.

### Animal Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

All sampling (fin clip) and handling of fish were conducted according to established animal welfare guidelines (ORBEA-ISPA, Animal Welfare Body—declaration 01/2019) and in accordance with the relevant legislation, as none of the sampled species is endangered or protected in Portugal.

### Field Study Permissions

The following information was supplied relating to field study approvals (i.e., approving body and any reference numbers):

The sampling took place in accordance with the relevant legislation, as none of the sampled species is endangered or protected in Portugal.



## DNA Deposition

The following information was supplied regarding the deposition of DNA sequences:

Sequences are available in GenBank: [MG992598–MG992888](#), [MH024090–MH024357](#), [MH030878–MH031272](#).

## Data Availability

The following information was supplied regarding data availability:

The complete list of specimens, sampling year, haplotype number and GenBank accession number for the mitochondrial control region and nuclear S7 ribosomal protein gene (intron 1 and 2) of the three target species (the white seabream *Diplodus sargus*, the sand-smelt *Atherina presbyter* and the shanny *Lipophrys pholis*) are available in the [Supplemental File](#).

## Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.9098#supplemental-information>.

## REFERENCES

- Addison JA, Hart MW. 2005.** Spawning, copulation and inbreeding coefficients in marine invertebrates. *Biology Letters* 1:450–453 DOI [10.1098/rsbl.2005.0353](#).
- Allison GW, Gaines SD, Lubchenco J, Possingham HP. 2003.** Ensuring persistence of marine reserves: catastrophes require adopting an insurance factor. *Ecological Applications* 13:8–24 DOI [10.1890/1051-0761\(2003\)013\[0008:EPOMRC\]2.0.CO;2](#).
- Almada VC, Almada F, Francisco SM, Castilho R, Robalo JI. 2012.** Unexpected high genetic diversity at the extreme northern geographic limit of *Taurulus bubalis* (Euphrasen, 1786). *PLOS ONE* 7:e44404 DOI [10.1371/journal.pone.0044404](#).
- Almada F, Francisco SM, Lima CS, Fitzgerald R, Mirimin L, Villegas-Ríos D, Saborido-Rey F, Afonso P, Morato T, Bexiga S, Robalo JI. 2017.** Historical gene flow constraints in a northeastern Atlantic fish: phylogeography of the ballan wrasse *Labrus bergylta* across its distribution range. *Royal Society Open Science* 4:160773 DOI [10.1098/rsos.160773](#).
- Almada VC, Goncalves EJ, Oliveira RF, Barata EN. 1992.** Some features of the territories in the breeding males of the intertidal blenny *Lipophrys pholis* (Pisces: Blenniidae). *Journal of the Marine Biological Association of the United Kingdom* 72:187–197 DOI [10.1017/S0025315400048876](#).
- Bamber RN, Henderson PA, Turnpenny AWH. 1985.** The early life history of the sand smelt (*Atherina Presbyter*). *Journal of the Marine Biological Association of the United Kingdom* 65:697–706 DOI [10.1017/S002531540005253X](#).
- Barbato M, Orozco-terWengel P, Tapio M, Bruford MW. 2015.** SNeP: a tool to estimate trends in recent effective population size trajectories using genome-wide SNP data. *Frontiers in Genetics* 6:109 DOI [10.3389/fgene.2015.00109](#).

- Barría A, Christensen KA, Yoshida G, Jedlicki A, Leong JS, Rondeau EB, Lhorente JP, Koop BF, Davidson WS, Yáñez JM. 2019.** Whole genome linkage disequilibrium and effective population size in a coho salmon (*Oncorhynchus kisutch*) breeding population using a high-density SNP array. *Frontiers in Genetics* **10**:498 DOI [10.3389/fgene.2019.00498](https://doi.org/10.3389/fgene.2019.00498).
- Bauchot M, Hureau J. 1986.** Sparidae. In: Whitehead P, Bauchot M, Hureau J, Nielsen J, Tortonese E, eds. *Fishes of the North-Eastern Atlantic and Mediterranean*. Paris: UNESCO, 883–907.
- Bernal-Ramírez JH, Adcock GJ, Hauser L, Carvalho GR, Smith PJ. 2003.** Temporal stability of genetic population structure in the New Zealand snapper, *Pagrus auratus*, and relationship to coastal currents. *Marine Biology* **142**:567–574 DOI [10.1007/s00227-002-0972-9](https://doi.org/10.1007/s00227-002-0972-9).
- Borges R, Vaz J, Serrão EA, Gonçalves EJ. 2009.** Short-term temporal fluctuation of very-nearshore larval fish assemblages at the Arrábida Marine Park (Portugal). *Journal of Coastal Research* **56**:376–380.
- Chapman R, Ball A, Mash L. 2002.** Spatial homogeneity and temporal heterogeneity of red drum (*Sciaenops ocellatus*) microsatellites: effective population sizes and management implications. *Marine Biotechnology* **4**:589–603 DOI [10.1007/s10126-002-0038-5](https://doi.org/10.1007/s10126-002-0038-5).
- Charlesworth B. 2009.** Effective population size and patterns of molecular evolution and variation. *Nature Reviews Genetics* **10**:195–205 DOI [10.1038/nrg2526](https://doi.org/10.1038/nrg2526).
- Charnov EL, Bull JJ. 1989.** Non-fisherian sex ratios with sex change and environmental sex determination. *Nature* **338**:148–150 DOI [10.1038/338148a0](https://doi.org/10.1038/338148a0).
- Charrier G, Chenel T, Durand JD, Girard M, Quiniou L, Laroche J. 2006.** Discrepancies in phylogeographical patterns of two European anglerfishes (*Lophius budegassa* and *Lophius piscatorius*). *Molecular Phylogenetics and Evolution* **38**:742–754 DOI [10.1016/j.ympev.2005.08.002](https://doi.org/10.1016/j.ympev.2005.08.002).
- Chevolot M, Ellis JR, Rijnsdorp AD, Stam WT, Olsen JL. 2008.** Temporal changes in allele frequencies but stable genetic diversity over the past 40 years in the Irish Sea population of thornback ray, *Raja clavata*. *Heredity* **101**:120–126 DOI [10.1038/hdy.2008.36](https://doi.org/10.1038/hdy.2008.36).
- Chow S, Hazama K. 1998.** Universal PCR primers for S7 ribosomal protein gene introns in fish. *Molecular Ecology* **7**:1247–1263.
- Clement M, Posada D, Crandall KA. 2000.** TCS: a computer program to estimate gene genealogies. *Molecular Ecology* **9**:1657–1659.
- Coscia I, Chopelet J, Waples RS, Mann BQ, Mariani S. 2016.** Sex change and effective population size: implications for population genetic studies in marine fish. *Heredity* **117**:251–258 DOI [10.1038/hdy.2016.50](https://doi.org/10.1038/hdy.2016.50).
- Crosbie SF, Manly BFJ. 1985.** Parsimonious modelling of capture-mark-recapture studies. *Biometrics* **41**:385–398 DOI [10.2307/2530864](https://doi.org/10.2307/2530864).
- Cuveliers EL, Volckaert FAM, Rijnsdorp AD, Larmuseau MHD, Maes GE. 2011.** Temporal genetic stability and high effective population size despite fisheries-induced

- life-history trait evolution in the North Sea sole. *Molecular Ecology* **20**:3555–3568 DOI [10.1111/j.1365-294X.2011.05196.x](https://doi.org/10.1111/j.1365-294X.2011.05196.x).
- Darriba D, Taboada GL, Doallo R, Posada D. 2012.** jModelTest 2: more models, new heuristics and parallel computing. *Nature Methods* **9**:772–772 DOI [10.1038/nmeth.2109](https://doi.org/10.1038/nmeth.2109).
- Di Franco A, De Benedetto G, De Rinaldis G, Raventos N, Sahyoun R, Guidetti P. 2011.** Large scale-variability in otolith microstructure and microchemistry: the case study of *Diplodus sargus sargus* (Pisces: Sparidae) in the Mediterranean Sea. *Italian Journal of Zoology* **78**:182–192 DOI [10.1080/11250003.2011.566227](https://doi.org/10.1080/11250003.2011.566227).
- Dias M, Roma J, Fonseca C, Pinto M, Cabral HN, Silva A, Vinagre C. 2016.** Intertidal pools as alternative nursery habitats for coastal fishes. *Marine Biological Research* **12**:331–344 DOI [10.1080/17451000.2016.1143106](https://doi.org/10.1080/17451000.2016.1143106).
- Domingues VS, Almada VC, Santos RS, Brito A, Bernardi G. 2006.** Phylogeography and evolution of the triplefin *Tripterygion delaisi* (Pisces, Blennioidei). *Marine Biology* **150**:509–519 DOI [10.1007/s00227-006-0367-4](https://doi.org/10.1007/s00227-006-0367-4).
- Domingues V, Santos R, Brito A, Alexandrou M, Almada V. 2007.** Mitochondrial and nuclear markers reveal isolation by distance and effects of Pleistocene glaciations in the northeastern Atlantic and Mediterranean populations of the white seabream (*Diplodus sargus*, L.). *Journal of Experimental Marine Biology and Ecology* **346**:102–113 DOI [10.1016/j.jembe.2007.03.002](https://doi.org/10.1016/j.jembe.2007.03.002).
- Excoffier L, Laval G, Balding D. 2003.** Gametic phase estimation over large genomic regions using an adaptive window approach. *Human Genomics* **1**:7 DOI [10.1186/1479-7364-1-1-7](https://doi.org/10.1186/1479-7364-1-1-7).
- Excoffier L, Lischer HEL. 2010.** Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Molecular Ecology Resources* **10**:564–567 DOI [10.1111/j.1755-0998.2010.02847.x](https://doi.org/10.1111/j.1755-0998.2010.02847.x).
- Excoffier L, Smouse PE, Quattro JM. 1992.** Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data. *Genetics* **131**:479–491.
- FAO. 2019.** FishStats; fisheries and aquaculture department. Available at <http://www.fao.org/fishery/> (accessed on 15 January 2019).
- Faria C, Almada V. 2001.** Microhabitat segregation in three rocky intertidal fish species in Portugal: does it reflect interspecific competition? *Journal of Fish Biology* **58**:145–159 DOI [10.1111/j.1095-8649.2001.tb00504.x](https://doi.org/10.1111/j.1095-8649.2001.tb00504.x).
- Faria C, Almada VC, Goncalves EJ. 1996.** Juvenile recruitment, growth and maturation of *Lipophrys pholis* (Pisces: Blenniidae), from the west coast of Portugal. *Journal of Fish Biology* **49**:727–730 DOI [10.1111/j.1095-8649.1996.tb00068.x](https://doi.org/10.1111/j.1095-8649.1996.tb00068.x).
- Faria C, Borges R, Gil F, Almada VC, Goncalves EJ. 2002.** Embryonic and larval development of *Lipophrys pholis* (Pisces: Blenniidae). *Scientia Marina* **66**:21–26 DOI [10.3989/scimar.2002.66n121](https://doi.org/10.3989/scimar.2002.66n121).
- Francisco SM, Almada VC, Faria C, Velasco EM, Robalo JI. 2014.** Phylogeographic pattern and glacial refugia of a rocky shore species with limited dispersal capability:

- the case of Montagu's blenny (*Coryphoblennius galerita*, Blenniidae). *Marine Biology* **161**:2509–2520 DOI [10.1007/s00227-014-2523-6](https://doi.org/10.1007/s00227-014-2523-6).
- Francisco SM, Cabral H, Vieira MN, Almada VC. 2006.** Contrasts in genetic structure and historical demography of marine and riverine populations of *Atherina* at similar geographical scales. *Estuarine Coastal and Shelf Science* **69**:655–661 DOI [10.1016/j.ecss.2006.05.017](https://doi.org/10.1016/j.ecss.2006.05.017).
- Francisco SM, Castilho R, Soares M, Congiu L, Brito A, Vieira MN, Almada VC. 2009.** Phylogeography and demographic history of *Atherina presbyter* (Pisces: Atherinidae) in the North-eastern Atlantic based on mitochondrial DNA. *Marine Biology* **156**:1421–1432 DOI [10.1007/s00227-009-1182-5](https://doi.org/10.1007/s00227-009-1182-5).
- Francisco SM, Congiu L, Stefanni S, Castilho R, Brito A, Ivanova PP, Levy A, Cabral H, Kiliass G, Doadrio I, Almada VC. 2008.** Phylogenetic relationships of the North-eastern Atlantic and Mediterranean forms of *Atherina* (Pisces, Atherinidae). *Molecular Phylogenetics and Evolution* **48**:782–788 DOI [10.1016/j.ympev.2007.12.009](https://doi.org/10.1016/j.ympev.2007.12.009).
- Francisco SM, Faria C, Lengkeek W, Vieira MN, Velasco EM, Almada VC. 2011.** Phylogeography of the shanny *Lipophrys pholis* (Pisces: Blenniidae) in the NE Atlantic records signs of major expansion event older than the last glaciation. *Journal of Experimental Marine Biology and Ecology* **403**:14–20 DOI [10.1016/j.jembe.2011.03.020](https://doi.org/10.1016/j.jembe.2011.03.020).
- Francisco SM, Robalo J. 2015.** Genetic structure and effective population size through time: a tale on two coastal marine species with contrasting life-history patterns. *Journal of Phylogenetics and Evolutionary Biology* **3**:155 DOI [10.4172/2329-9002.1000155](https://doi.org/10.4172/2329-9002.1000155).
- Francisco SM, Vieira MN, Almada VC. 2006.** Genetic structure and historical demography of the shanny *Lipophrys pholis* in the Portuguese coast based on mitochondrial DNA analysis. *Molecular Phylogenetics and Evolution* **39**:288–292 DOI [10.1016/j.ympev.2005.12.009](https://doi.org/10.1016/j.ympev.2005.12.009).
- Frankham R, Bradshaw CJA, Brook BW. 2014.** Genetics in conservation management: revised recommendations for the 50/500 rules, Red List criteria and population viability analyses. *Biological Conservation* **170**:56–63 DOI [10.1016/j.biocon.2013.12.036](https://doi.org/10.1016/j.biocon.2013.12.036).
- Galarza JA, Carreras-Carbonell J, Macpherson E, Pascual M, Roques S, Turner GF, Rico C. 2009.** The influence of oceanographic fronts and early-life-history traits on connectivity among littoral fish species. *Proceedings of the National Academy of Sciences of the United States of America* **106**:1473–1478 DOI [10.1073/pnas.0806804106](https://doi.org/10.1073/pnas.0806804106).
- Garnier-Géré P, Chikhi L. 2013.** Population subdivision, Hardy–Weinberg equilibrium and the Wahlund effect. In: *Encyclopedia of Life Sciences*. Hoboken: John Wiley & Sons, Ltd (Ed.) DOI [10.1002/9780470015902.a0005446.pub3](https://doi.org/10.1002/9780470015902.a0005446.pub3).
- Goldstien SJ, Inglis GJ, Schiel DR, Gemmell NJ. 2013.** Using temporal sampling to improve attribution of source populations for invasive species. *PLOS ONE* **8**:e65656 DOI [10.1371/journal.pone.0065656](https://doi.org/10.1371/journal.pone.0065656).
- Gon O. 2015.** *Atherina presbyter*. The IUCN Red List of Threatened Species 2015. DOI [10.2305/IUCN.UK.2015-4.RLTS.T194991A13489435.en](https://doi.org/10.2305/IUCN.UK.2015-4.RLTS.T194991A13489435.en).

- Goncalves JMS, Erzini K. 2000.** The reproductive biology of the two-banded sea bream (*Diplodus vulgaris*) from the southwest coast of Portugal. *Journal of Applied Ichthyology* **16**:110–116 DOI [10.1046/j.1439-0426.2000.00232.x](https://doi.org/10.1046/j.1439-0426.2000.00232.x).
- Gonzalez EG, Beerli P, Zardoya R. 2008.** Genetic structuring and migration patterns of Atlantic bigeye tuna, *Thunnus obesus* (Lowe, 1839). *BMC Evolutionary Biology* **8**:252 DOI [10.1186/1471-2148-8-252](https://doi.org/10.1186/1471-2148-8-252).
- Gonzalez-Wanguemert M, Froufe E, Perez-Ruzafa A, Alexandrino P. 2011.** Phylogeographical history of the white seabream *Diplodus sargus* (Sparidae): implications for insularity. *Marine Biology Research* **7**:250–260 DOI [10.1080/17451000.2010.499438](https://doi.org/10.1080/17451000.2010.499438).
- Guindon S, Gascuel O. 2003.** A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Systematics Biology* **52**:696–704 DOI [10.1080/10635150390235520](https://doi.org/10.1080/10635150390235520).
- Guo SW, Thompson EA. 1992.** Performing the exact test of Hardy-Weinberg proportion for multiple alleles. *Biometrics* **48**:361–372 DOI [10.2307/2532296](https://doi.org/10.2307/2532296).
- Hare MP, Nunney L, Schwartz MK, Ruzzante DE, Burford M, Waples RS, Ruegg K, Palstra F. 2011.** Understanding and estimating effective population size for practical application in marine species management. *Conservation Biology* **25**:438–449 DOI [10.1111/j.1523-1739.2010.01637.x](https://doi.org/10.1111/j.1523-1739.2010.01637.x).
- Hauser L, Adcock GJ, Smith PJ, Bernal-Ramirez JH, Carvalho GR. 2002.** Loss of microsatellite diversity and low effective population size in an overexploited population of New Zealand snapper (*Pagrus auratus*). *Proceedings of the National Academy of Sciences of the United States of America* **99**:11742–11747 DOI [10.1073/pnas.172242899](https://doi.org/10.1073/pnas.172242899).
- Henriques M, Gonçalves EJ, Almada VC. 1999.** The conservation of littoral fish communities: a case study at Arrábida coast (Portugal). In: Almada VC, Oliveira RF, Gonçalves EJ, eds. *Behaviour and conservation of littoral fishes*. Lisbon: ISPA, 473–519.
- Henriques R, Nielsen ES, Durholtz D, Japp D, Von der Heyden S. 2017.** Genetic population sub-structuring of kingklip (*Genypterus capensis*—Ophidiidae), a commercially exploited demersal fish off South Africa. *Fisheries Research* **187**:86–95 DOI [10.1016/j.fishres.2016.11.007](https://doi.org/10.1016/j.fishres.2016.11.007).
- Hill WG. 1981.** Estimation of effective population size from data on linkage disequilibrium. *Genetic Research* **38**:209–216 DOI [10.1017/S0016672300020553](https://doi.org/10.1017/S0016672300020553).
- Jiménez-Mena B, Le Moan A, Christensen A, Van Deurs M, Mosegaard H, Hemmer-Hansen J, Bekkevold D. 2019.** Weak genetic structure despite strong genomic signal in lesser sandeel in the North Sea. *Evolutionary Applications* **13**:376–387 DOI [10.1111/eva.12875](https://doi.org/10.1111/eva.12875).
- Jolly GM. 1965.** Explicit estimates from capture-recapture data with both death and immigration-stochastic model. *Biometrika* **52**:225–248 DOI [10.2307/2333826](https://doi.org/10.2307/2333826).
- Jorde PE. 2012.** Allele frequency covariance among cohorts and its use in estimating effective size of age-structured populations. *Molecular Ecology Resources* **12**:476–480 DOI [10.1111/j.1755-0998.2011.03111.x](https://doi.org/10.1111/j.1755-0998.2011.03111.x).
- Jorde PE, Ryman N. 1995.** Temporal allele frequency change and estimation of effective size in populations with overlapping generations. *Genetics* **139**:1077–1090.

- Jorde PE, Ryman N. 2007.** Unbiased estimator for genetic drift and effective population size. *Genetics* **177**:927–935 DOI [10.1534/genetics.107.075481](https://doi.org/10.1534/genetics.107.075481).
- Kalinowski ST. 2005.** hp-rare 1.0: a computer program for performing rarefaction on measures of allelic richness. *Molecular Ecology Notes* **5**:187–189 DOI [10.1111/j.1471-8286.2004.00845.x](https://doi.org/10.1111/j.1471-8286.2004.00845.x).
- Karlsson S, Mork J. 2005.** Deviation from HardyWeinberg equilibrium, and temporal instability in allele frequencies at microsatellite loci in a local population of Atlantic cod. *ICES Journal of Marine Science* **62**:1588–1596 DOI [10.1016/j.icesjms.2005.05.009](https://doi.org/10.1016/j.icesjms.2005.05.009).
- Knutsen H, Jorde PE, Albert OT, Hoelzel AR, Stenseth NC. 2007.** Population genetic structure in the North Atlantic Greenland halibut (*Reinhardtius hippoglossoides*): influenced by oceanic current systems? *Canadian Journal of Fisheries and Aquatic Sciences* **64**:857–866 DOI [10.1139/F07-070](https://doi.org/10.1139/F07-070).
- Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, Valentin F, Wallace IM, Wilm A, Lopez R, Thompson JD, Gibson TJ, Higgins DG. 2007.** Clustal W and Clustal X version 2.0. *Bioinformatics* **23**:2947–2948 DOI [10.1093/bioinformatics/btm404](https://doi.org/10.1093/bioinformatics/btm404).
- Larsson LC, Laikre L, André C, Dahlgren TG, Ryman N. 2010.** Temporally stable genetic structure of heavily exploited Atlantic herring (*Clupea harengus*) in Swedish waters. *Heredity* **104**:40–51 DOI [10.1038/hdy.2009.98](https://doi.org/10.1038/hdy.2009.98).
- Le Moan A, Jimenez-Mena B, Bekkevold D, Hemmer-Hanse J. 2019.** Fine scale population structure linked to neutral divergence in the common sole (*Solea solea*), a marine fish with high dispersal capacity. *BioRxiv*. DOI [10.1101/662619](https://doi.org/10.1101/662619).
- Lenfant P, Planes S. 2002.** Temporal genetic changes between cohorts in a natural population of a marine fish, *Diplodus sargus*. *Biological Journal of the Linnean Society* **76**:9–20 DOI [10.1046/j.1095-8312.2002.00041.x](https://doi.org/10.1046/j.1095-8312.2002.00041.x).
- Lima D, Santos MM, Ferreira AM, Micaelo C, Reis-Henriques MA. 2008.** The use of the shanny *Lipophrys pholis* for pollution monitoring: a new sentinel species for the northwestern European marine ecosystems. *Environment International* **34**:94–101 DOI [10.1016/j.ENVINT.2007.07.007](https://doi.org/10.1016/j.ENVINT.2007.07.007).
- Luikart G, Ryman N, Tallmon DA, Schwartz MK, Allendorf FW. 2010.** Estimation of census and effective population sizes: the increasing usefulness of DNA-based approaches. *Conservation Genetics* **11**:355–373 DOI [10.1007/s10592-010-0050-7](https://doi.org/10.1007/s10592-010-0050-7).
- Luttikhuizen PC, Campos J, Van Bleijswijk J, Peijnenburg KTCA, Van der Veer HW. 2008.** Phylogeography of the common shrimp, Crangon crangon (L.) across its distribution range. *Molecular Phylogenetics and Evolution* **46**:1015–1030 DOI [10.1016/j.ympev.2007.11.011](https://doi.org/10.1016/j.ympev.2007.11.011).
- Lynch M, Lande R. 1998.** The critical effective size for a genetically secure population. *Animal Conservation* **1**:70–72 DOI [10.1111/j.1469-1795.1998.tb00229.x](https://doi.org/10.1111/j.1469-1795.1998.tb00229.x).
- Marandel F, Lorange P, Berthelé O, Trenkel VM, Waples RS, Lamy J-B. 2019.** Estimating effective population size of large marine populations, is it feasible? *Fish and Fisheries* **20**:189–198 DOI [10.1111/faf.12338](https://doi.org/10.1111/faf.12338).

- Milton P. 1983.** Biology of littoral blennioid fishes on the coast of South-West England. *Journal of the Marine Biological Association of the United Kingdom* **63**:223–237 DOI [10.1017/S0025315400049912](https://doi.org/10.1017/S0025315400049912).
- Morato T, Afonso P, Lourinho P, Nash RDM, Santos RS. 2003.** Reproductive biology and recruitment of the white sea bream in the Azores. *Journal of Fish Biology* **63**:59–72 DOI [10.1046/j.1095-8649.2003.00129.x](https://doi.org/10.1046/j.1095-8649.2003.00129.x).
- Mouine N, Francour P, Ktari M-H, Chakroun-Marzouk N. 2007.** The reproductive biology of *Diplodus sargus sargus*; in the Gulf of Tunis (central Mediterranean). *Scientia Marina* **71**:461–469 DOI [10.3989/scimar.2007.71n3461](https://doi.org/10.3989/scimar.2007.71n3461).
- Múrias dos Santos A, Cabezas MP, Tavares AI, Xavier R, Branco M. 2016.** tcsBU: a tool to extend TCS network layout and visualization. *Bioinformatics* **32**:627–628 DOI [10.1093/bioinformatics/btv636](https://doi.org/10.1093/bioinformatics/btv636).
- Nei M. 1987.** *Molecular evolutionary genetics*. New York: Columbia University Press.
- Nei M, Kumar S. 2000.** *Molecular evolution and phylogenetics*. Oxford: Oxford University Press.
- Nei M, Tajima F. 1981.** Genetic drift and estimation of effective population size. *Genetics* **98**:625–640.
- Nomura T. 2008.** Estimation of effective number of breeders from molecular coancestry of single cohort sample. *Evolutionary Applications* **1**:462–474 DOI [10.1111/j.1752-4571.2008.00015.x](https://doi.org/10.1111/j.1752-4571.2008.00015.x).
- Ostellari L, Bargelloni L, Penzo E, Patarnello P, Patarnello T. 1996.** Optimization of single-strand conformation polymorphism and sequence analysis of the mitochondrial control region in *Pagellus bogaraveo* (Sparidae, Teleostei): rationalized tools in fish population biology. *Animal Genetics* **27**:423–427 DOI [10.1111/j.1365-2052.1996.tb00510.x](https://doi.org/10.1111/j.1365-2052.1996.tb00510.x).
- Palstra FP, Ruzzante DE. 2008.** Genetic estimates of contemporary effective population size: What can they tell us about the importance of genetic stochasticity for wild population persistence? *Molecular Ecology* **17**:3428–3447 DOI [10.1111/j.1365-294X.2008.03842.x](https://doi.org/10.1111/j.1365-294X.2008.03842.x).
- Papetti C, Zane L, Bortolotto E, Bucklin A, Patarnello T. 2005.** Genetic differentiation and local temporal stability of population structure in the euphausiid *Meganyctiphanes norvegica*. *Marine Ecology Progress Series* **289**:225–235 DOI [10.3354/meps289225](https://doi.org/10.3354/meps289225).
- Pappalardo AM, Francisco SM, Fruciano C, Lima CS, Pulvirenti V, Tigano C, Robalo JI, Ferrito V. 2017.** Mitochondrial and nuclear intraspecific variation in the rusty blenny (*Parablennius sanguinolentus*, Blenniidae). *Hydrobiologia* **802**:141–154 DOI [10.1007/s10750-017-3248-6](https://doi.org/10.1007/s10750-017-3248-6).
- Peakall R, Smouse PE. 2006.** Genalex 6: genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes* **6**:288–295 DOI [10.1111/j.1471-8286.2005.01155.x](https://doi.org/10.1111/j.1471-8286.2005.01155.x).
- Peakall R, Smouse PE. 2012.** GenALEx 6.5: genetic analysis in Excel. Population genetic software for teaching and research—an update. *Bioinformatics* **28**:2537–2539 DOI [10.1093/bioinformatics/bts460](https://doi.org/10.1093/bioinformatics/bts460).

- Pérez-Portela R, Wangensteen OS, Garcia-Cisneros A, Valero-Jiménez C, Palacín C, Turon X. 2019.** Spatio-temporal patterns of genetic variation in *Arbacia lixula*, a thermophilous sea urchin in expansion in the Mediterranean. *Heredity* **122**:244–259 DOI [10.1038/s41437-018-0098-6](https://doi.org/10.1038/s41437-018-0098-6).
- Pita A, Pérez M, Velasco F, Presa P. 2017.** Trends of the genetic effective population size in the Southern stock of the European hake. *Fisheries Research* **191**:108–119 DOI [10.1016/j.fishres.2017.02.022](https://doi.org/10.1016/j.fishres.2017.02.022).
- Planes S, Lenfant P. 2002.** Temporal change in the genetic structure between and within cohorts of a marine fish, *Diplodus sargus*, induced by a large variance in individual reproductive success. *Molecular Ecology* **11**:1515–1524 DOI [10.1046/j.1365-294X.2002.01521.x](https://doi.org/10.1046/j.1365-294X.2002.01521.x).
- Plank SM, Lowe CG, Feldheim KA, Wilson RR, Brusslan JA. 2010.** Population genetic structure of the round stingray *Urobatis halleri* (Elasmobranchii: Rajiformes) in southern California and the Gulf of California. *Journal of Fish Biology* **77**:329–340 DOI [10.1111/j.1095-8649.2010.02677.x](https://doi.org/10.1111/j.1095-8649.2010.02677.x).
- Pollard D, Russel B, Carpenter KE, Iwatsuki Y, Vega-Cendejas M, Jassim-Kawari A, Hartmann S, Alnazry H, Abdulqader E, Alam S, Bishop J, Hassan-Al-Khalf K, Kaymaram F. 2014.** *Diplodus sargus*. The IUCN Red List of Threatened Species 2014. DOI [10.2305/IUCN.UK.2014-3.RLTS.T170155A42736975.en](https://doi.org/10.2305/IUCN.UK.2014-3.RLTS.T170155A42736975.en).
- Posada D. 2003.** Using MODELTEST and PAUP\* to select a model of nucleotide substitution. *Current Protocols in Bioinformatics* **00**:6.5.1–6.5.14 DOI [10.1002/0471250953.bi0605s00](https://doi.org/10.1002/0471250953.bi0605s00).
- Pudovkin AI, Zaykin DV, Hedgecock D. 1996.** On the potential for estimating the effective number of breeders from heterozygote-excess in progeny. *Genetics* **144**:383–387.
- Quignard J-P, Pras A. 1986.** Atherinidae. In: Whitehead P, Bauchot M, Hureau J, Nielsen J, Tortonese E, eds. *Fishes of the North-Eastern Atlantic and Mediterranean*. Paris: UNESCO, 1207–1210.
- Reid-Anderson S, Bilgmann K, Stow A. 2019.** Effective population size of the critically endangered east Australian grey nurse shark *Carcharias taurus*. *Marine Ecology Progress Series* **610**:137–148 DOI [10.3354/meps12850](https://doi.org/10.3354/meps12850).
- Riginos C, Hock K, Matias AM, Mumby PJ, Van Oppen MJH, Lukoschek V. 2019.** Asymmetric dispersal is a critical element of concordance between biophysical dispersal models and spatial genetic structure in Great Barrier Reef corals. *Diversity and Distributions* **25**:1684–1696 DOI [10.1111/ddi.12969](https://doi.org/10.1111/ddi.12969).
- Riginos C, Victor BC. 2001.** Larval spatial distributions and other early life-history characteristics predict genetic differentiation in eastern Pacific blennioid fishes. *Proceedings of the Royal Society B* **268**:1931–1936 DOI [10.1098/rspb.2001.1748](https://doi.org/10.1098/rspb.2001.1748).
- Robalo JI, Lima CS, Francisco SM, Almada F, Bañon R, Villegas-ríos D, Almada VC. 2013.** Monitoring climate change impact on the genetic population structure: the case of the fivebeard rockling (*Ciliata mustela*, Linnaeus, 1758) in its southern limit of distribution. *Journal of Phylogenetics and Evolutionary Biology* **1**:1–4 DOI [10.4172/2329-9002.1000123](https://doi.org/10.4172/2329-9002.1000123).



- Ryman N, Allendorf FW, Jorde PE, Laikre L, Hössjer O. 2014. Samples from subdivided populations yield biased estimates of effective size that overestimate the rate of loss of genetic variation. *Molecular Ecology Resources* 14:87–99 DOI 10.1111/1755-0998.12154.
- Salicru M, Menendez ML, Morales D, Pardo L. 1993. Asymptotic distribution of  $(h, \varphi)$ -entropies. *Communications in Statistics—Theory and Methods* 22:2015–2031 DOI 10.1080/03610929308831131.
- Santos R, Porteiro F, Barreiros J. 1997. Marine fishes of the azores: annotated checklist and bibliography. *Arquipelago—Life and Marine Sciences Suppl* 1:1–268.
- Schwartz MK, Luikart G, Waples RS. 2007. Genetic monitoring as a promising tool for conservation and management. *Trends in Ecology and Evolution* 22:25–33 DOI 10.1016/J.TREE.2006.08.009.
- Seber GAF. 1965. A note on the multiple-recapture census. *Biometrika* 52:249–259 DOI 10.2307/2333827.
- Selkoe KA, Gaines SD, Caselle JE, Warner RR. 2006. Current shifts and kin aggregation explain genetic patchiness in fish recruits. *Ecology* 87:3082–3094 DOI 10.1890/0012-9658(2006)87[3082:CSAKAE]2.0.CO;2.
- Selkoe KA, Watson JR, White C, Horin TB, Iacchei M, Mitttarai S, Siegel DA, Gaines SD, Toonen RJ. 2010. Taking the chaos out of genetic patchiness: seascape genetics reveals ecological and oceanographic drivers of genetic patterns in three temperate reef species. *Molecular Ecology* 19:3708–3726 DOI 10.1111/j.1365-294X.2010.04658.x.
- Sepúlveda FA, González MT. 2017. Spatio-temporal patterns of genetic variations in populations of yellowtail kingfish *Seriola lalandi* from the south-eastern Pacific Ocean and potential implications for its fishery management. *Journal of Fish Biology* 90:249–264 DOI 10.1111/jfb.13179.
- Serbezov D, Jorde PE, Bernatchez L, Olsen EM, Vøllestad LA. 2012. Short-term genetic changes: evaluating effective population size estimates in a comprehensively described brown trout (*Salmo trutta*) population. *Genetics* 191:579–592 DOI 10.1534/genetics.111.136580.
- Sousa-Santos C, Robalo JI, Collares-Pereira MJ, Almada VC. 2005. Heterozygous indels as useful tools in the reconstruction of DNA sequences and in the assessment of ploidy level and genomic constitution of hybrid organisms. *DNA Sequence* 16:462–467 DOI 10.1080/10425170500356065.
- Stefanni S, Castilho R, Sala-Bozano M, Robalo JI, Francisco SM, Santos RS, Marques N, Brito A, Almada VC, Mariani S. 2015. Establishment of a coastal fish in the Azores: recent colonisation or sudden expansion of an ancient relict population? *Heredity* 115:527–537 DOI 10.1038/hdy.2015.55.
- Stefanni S, Domingues V, Bouton N, Santos RS, Almada F, Almada V. 2006. Phylogeny of the shanny, *Lipophrys pholis*, from the NE Atlantic using mitochondrial DNA markers. *Molecular Phylogenetics and Evolution* 39:282–287 DOI 10.1016/j.ympev.2005.07.001.

- Tajima F. 1983.** Evolutionary relationship of DNA sequences in finite populations. *Genetics* **105**:437–460.
- Tavare S. 1986.** Some probabilistic and statistical problems in the analysis of DNA sequences. *Lectures on Mathematics in the Life Sciences* **17**:57–86.
- Templeton AR, Crandall KA, Sing F. 1992.** Cladistic analysis of phenotypic associations with haplotypes inferred from restriction endonuclease mapping and DNA sequence data. 111. Cladogram estimation. *Genetics* **132**:619–633.
- Therkildsen NO, Nielsen EE, Swain DP, Pedersen JS. 2010.** Large effective population size and temporal genetic stability in Atlantic cod (*Gadus morhua*) in the southern Gulf of St. Lawrence. *Canadian Journal of Fisheries and Aquatic Sciences* **67**:1585–1595 DOI [10.1139/F10-084](https://doi.org/10.1139/F10-084).
- Turnpenny AWH, Bamber RN, Henderson PA. 1981.** Biology of the sand-smelt (*Atherina presbyter* Valenciennes) around Fawley power station. *Journal of Fish Biology* **18**:417–427 DOI [10.1111/j.1095-8649.1981.tb03783.x](https://doi.org/10.1111/j.1095-8649.1981.tb03783.x).
- Veríssimo A, Sampaio Í, McDowell JR, Alexandrino P, Mucientes G, Queiroz N, Da Silva C, Jones CS, Noble LR. 2017.** World without borders—genetic population structure of a highly migratory marine predator, the blue shark (*Prionace glauca*). *Ecology and Evolution* **7**:4768–4781 DOI [10.1002/ece3.2987](https://doi.org/10.1002/ece3.2987).
- Verry AJF, Walton K, Tuck ID, Ritchie PA. 2020.** Genetic structure and recent population expansion in the commercially harvested deep-sea decapod, *Metanephrops challengeri* (Crustacea: Decapoda). *New Zealand Journal of Marine and Freshwater Research* **54**:251–270 DOI [10.1080/00288330.2019.1707696](https://doi.org/10.1080/00288330.2019.1707696).
- Vinagre C, Mendonça V, Cereja R, Abreu-Afonso F, Dias M, Mizrahi D, Flores AAV. 2018.** Ecological traps in shallow coastal waters—Potential effect of heat-waves in tropical and temperate organisms. *PLOS ONE* **13**:e0192700 DOI [10.1371/journal.pone.0192700](https://doi.org/10.1371/journal.pone.0192700).
- Viñas J, Alvarado-Bremer J, Pla C. 2004.** Phylogeography of the Atlantic bonito (*Sarda sarda*) in the northern Mediterranean: the combined effects of historical vicariance, population expansion, secondary invasion, and isolation by distance. *Molecular Phylogenetics and Evolution* **33**:32–42 DOI [10.1016/j.ympev.2004.04.009](https://doi.org/10.1016/j.ympev.2004.04.009).
- Wang J. 2001.** A pseudo-likelihood method for estimating effective population size from temporally spaced samples. *Genetics Research* **78**:243–257 DOI [10.1017/S0016672301005286](https://doi.org/10.1017/S0016672301005286).
- Wang J. 2005.** Estimation of effective population sizes from data on genetic markers. *Philosophical Transactions of the Royal Society of London B* **360**:1395–1409 DOI [10.1098/rstb.2005.1682](https://doi.org/10.1098/rstb.2005.1682).
- Wang J. 2009.** A new method for estimating effective population sizes from a single sample of multilocus genotypes. *Molecular Ecology* **18**:2148–2164 DOI [10.1111/j.1365-294X.2009.04175.x](https://doi.org/10.1111/j.1365-294X.2009.04175.x).
- Wang J, Santiago E, Caballero A. 2016.** Prediction and estimation of effective population size. *Heredity* **117**:193–206 DOI [10.1038/hdy.2016.43](https://doi.org/10.1038/hdy.2016.43).
- Waples RS. 1989.** A generalized approach for estimating effective population size from temporal changes in allele frequency. *Genetics* **121**:379–391.

- Waples RS. 2016.** Life-history traits and effective population size in species with overlapping generations revisited: the importance of adult mortality. *Heredity* **117**:241–250 DOI [10.1038/hdy.2016.29](https://doi.org/10.1038/hdy.2016.29).
- Waples RS, Do C. 2010.** Linkage disequilibrium estimates of contemporary  $N_e$  using highly variable genetic markers: a largely untapped resource for applied conservation and evolution. *Evolutionary Applications* **3**:244–262 DOI [10.1111/j.1752-4571.2009.00104.x](https://doi.org/10.1111/j.1752-4571.2009.00104.x).
- Waples RS, Mariani S, Benvenuto C. 2018.** Consequences of sex change for effective population size. *Proceedings of the Royal Society B* **285**:20181702 DOI [10.1098/rspb.2018.1702](https://doi.org/10.1098/rspb.2018.1702).
- Waples RS, Yokota M. 2007.** Temporal estimates of effective population size in species with overlapping generations. *Genetics* **175**:219–233 DOI [10.1534/genetics.106.065300](https://doi.org/10.1534/genetics.106.065300).
- White C, Selkoe KA, Watson J, Siegel DA, Zacherl DC, Toonen RJ. 2010.** Ocean currents help explain population genetic structure. *Proceedings of the Royal Society B* **277**:1685–1694 DOI [10.1098/rspb.2009.2214](https://doi.org/10.1098/rspb.2009.2214).
- Williams J, Craig M. 2014.** *Lipophrys pholis*. The IUCN Red List of Threatened Species 2014. DOI [10.2305/IUCN.UK.2014-3.RLTS.T185180A1777432.en](https://doi.org/10.2305/IUCN.UK.2014-3.RLTS.T185180A1777432.en).
- Wright S. 1931.** Evolution in mendelian populations. *Genetics* **16**:97–159.
- Wright S. 1938.** Size of population and breeding structure in relation to evolution. *Science* **87**:430–431.
- Zander CD. 1986.** Blenniidae. In: Whitehead P, Bauchot M, Hureau J, Nielsen J, Tortonese E, eds. *Fishes of the North-Eastern Atlantic and Mediterranean*. Paris: UNESCO, 1096–1112.