

Blood-clotting enzymology. The activation of factor X by factor IXa

Citation for published version (APA):

van Dieijen, G., Rosing, J., Tans, G., Janssen-Claessen, G., Franssen, J., & Hemker, H. C. (1980). Blood-clotting enzymology. The activation of factor X by factor IXa: The effect of phospholipid and of antihemophilia factor VIII (F VIII) on the activation reaction. *Archives internationales de physiologie et de biochimie*, 88(3), B153-B154. <https://doi.org/10.3109/13813458009104857>

Document status and date:

Published: 01/01/1980

DOI:

[10.3109/13813458009104857](https://doi.org/10.3109/13813458009104857)

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

SOCIÉTÉ BELGE DE BIOCHIMIE
BELGISCHE VERENIGING VOOR BIOCHEMIE112^e RÉUNION, GENT, 23 FÉVRIER 1980

G. VAN DIEIJEN, J. ROSING, G. TANS, G. JANSSEN-CLAESSEN, J. FRANSSSEN and H. C. HEMKER (*Department of Biochemistry, University of Limburg, Maastricht, The Netherlands*).

Blood-clotting enzymology. The activation of factor X by factor IX_a ; the effect of phospholipid and of antihæmophilia factor VIII (F VIII) on the activation reaction.

The activation of bovine clotting factor X is mediated by the enzyme factor IX_a. Activation of factor X, *i.e.* the formation of the serine-protease X_a, is the result of the specific cleavage of a peptide bond in factor X. The rate of activation is strongly enhanced by the cofactor VIII_a and by a phospholipid (PL) surface, onto which the protein adsorbs and where the reaction is thought to occur.

In order to get more insight into the mechanism of the activation reaction and the role of PL, Ca²⁺ and factor VIII, we have determined the kinetic constants of factor X activation and the effects of PL and factor VIII on these constants. Without PL and factor VIII_a, K_m for factor X is 272 μM and V is 6 $\text{mmol X}_a \text{mol}^{-1} \text{IX}_a \text{min}^{-1}$. With PL (10 μM) the K_m is lowered to 0.06 μM and V is unchanged. In the presence of PL and factor VIII_a, K_m is 0.02 μM and V is 111 $\text{mol X}_a \text{mol}^{-1} \text{IX}_a \text{min}^{-1}$. The factor X concentration in plasma is 0.16 μM , so we conclude that PL in plasma serves to lower the K to 0.06 μM , and that factor VIII stimulates factor X activation because it increases V of X_a formation at least 10 000 times.

We further investigated the effect of PL concentration on the kinetic parameters with factor VIII_a left out. V for factor X_a formation does not change very much, but K_m for factor X increases from 0.06 μM at 10 μM PL to 1.35 μM at 260 μM PL. Earlier ROSING *et al.* (1980), studying the prothrombinase complex, found a similar relationship between PL concentration and the K_m for prothrombin.

This led to the hypothesis that it is the local substrate concentration at the phospholipid surface that determine the kinetics of its activation. An increased local concentration explains the dramatic decrease of K_m in the presence of PL. Further addition of PL lowers the substrate density at the PL surface with as consequence an increase of K_m .

The validity of this hypothesis can be tested if the binding parameters of substrate to PL are known. Recently, we found a new method to determine parameters of factor X binding to PL. This method is based on the observation that RVV-X (*i.e.* the factor-X-activating enzyme from Russell's viper venom) activates free factor X, but not factor X bound to PL. We found that PL vesicles consisting of equal amounts of phosphatidylserine and phosphatidylcholine (also used in our kinetic experiments) bind factor X with a dissociation constant (K_d) of 3.7×10^{-8} M. The concentration of factor-X-binding sites on these vesicles is 0.7 μM sites per 100 μM PL.

These data were used to convert K_m values from the kinetic experiments to K_m values expressed as factor X bound per PL surface area. The difference in K_m disap-

peared and one K_m value was found of $0.010 \pm 0.003 \mu M$ factor X/ μM PL for all PL concentrations. These results further support our hypothesis that the kinetics for X activation are governed by the density of the reactants bound to the PL surface.

Reference

ROSIK, J., TANS, G., GOVERS-RIEMSLAG, J. W. P., ZWAAL, R. F. A. & HEMKER, H. C. (1980) *J. Biol. Chem.*, in press.