

# Adsorption kinetics of protein mixtures

## Citation for published version (APA):

Cuypers, P. A., Willems, G. A., Hemker, H. C., & Hermens, W. T. (1987). Adsorption kinetics of protein mixtures: A tentative explanation of the Vroman effect. In E. F. Leonard, V. T. Turitto, & L. Vroman (Eds.), *Blood in Contact with Natural and Artificial Surfaces* (1 ed., pp. 244-252). Annals of the New York Academy of Sciences, Vol.. 516

## Document status and date:

Published: 28/12/1987

## Document Version:

Other version

## Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

## General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.umlib.nl/taverne-license](http://www.umlib.nl/taverne-license)

## Take down policy

If you believe that this document breaches copyright please contact us at:

[repository@maastrichtuniversity.nl](mailto:repository@maastrichtuniversity.nl)

providing details and we will investigate your claim.

5799

Reprinted from *Blood in Contact with Natural and Artific*  
Volume 516 of the *Annals of the New York Academy of Sciences*  
December 28, 1987

## Adsorption Kinetics of Protein Mixtures

### A Tentative Explanation of the Vroman Effect

PETER A. CUYPERS,<sup>a,b</sup> GEORGE M. WILLEMS,<sup>a</sup>  
H. COENRAAD HEMKER,<sup>c</sup> AND WIM TH. HERMENS<sup>a</sup>

<sup>a</sup>Department of Biophysics

<sup>c</sup>Department of Biochemistry

Biomedical Centre

University of Limburg

6200 MD Maastricht, The Netherlands

#### INTRODUCTION

When a surface is exposed to plasma, different proteins from the plasma will adsorb on the surface within several minutes. This adsorption from plasma or serum or protein mixtures depends on the concentration of the various proteins, the nature of the protein and surface, and the adsorption time. Despite numerous studies, the exact composition of the adsorbed protein layer from a multicomponent mixture such as plasma is not yet known.<sup>1-3</sup> Exposure of different surfaces to undiluted plasma results in a relatively limited adsorption of albumin, immunoglobulin, and fibrinogen compared with the adsorption of the same proteins from less concentrated solutions.<sup>4</sup> From experiments of Vroman<sup>5</sup> in which he used normal plasma and plasma lacking high molecular weight kininogen (HMWK), it was concluded that fibrinogen, initially adsorbed on hydrophilic surfaces exposed to normal plasma, is displaced by HMWK. These observations were confirmed by Brash.<sup>3</sup> Observations on the adsorption of proteins from various dilutions of plasma showed that no fibrinogen is adsorbed from concentrated plasma, adsorption followed by desorption of fibrinogen occurs for moderately diluted plasma, and a stable adsorbed fibrinogen monolayer is formed for plasma dilutions of about 5% or less.<sup>6-10</sup> The observation that a protein, abundantly present in plasma, predominates initially in the adsorbed protein layer only to be replaced thereafter by a scarce component of the plasma proteins with a higher affinity for the adsorbing surface fits qualitatively into classical binding theory. The dilution effects, however, seem unexplainable in this framework.

In recent studies on the adsorption kinetics of three different proteins on a double layer of phospholipids it was found that the value of the intrinsic adsorption rate constant,  $k_{on}^{int}$ , is strongly dependent on the surface concentration of the adsorbed proteins. The experiments showed that  $k_{on}^{int}$  decays exponentially with increasing surface concentration  $\Gamma$ . The present study shows that this behavior results in an interaction parameter between the adsorbed proteins that may in principle cause such phenomena as the replacement of fibrinogen by HMWK, and, more importantly, also explains the different types of adsorption from different dilutions of plasma.

<sup>b</sup>Present address: DSM Research BV, P.O. Box 18, 6160 MD Geleen, The Netherlands.

MATERIALS AND METHODS

Materials

A double layer of 1,2-dioleoyl-*sn*-glycero-2-phosphoserine (DOPS) was stacked on a chromium slide as described in refs. 11 and 16. Bovine prothrombin and human fibrinogen were used. The proteins were either obtained commercially or prepared according to established procedures.<sup>16</sup>

Adsorption and desorption experiments were performed by ellipsometry as described in detail previously.<sup>12-14</sup> The surface concentration of proteins was calculated from the measured refractive index and thickness according to the modified Lorenz-Lorentz equation.<sup>15</sup> Sorption experiments were performed in Tris-HCl buffer, pH 7.5, containing 0.1 M NaCl and 1.5 mM CaCl<sub>2</sub>.

Analysis of Sorption Kinetics

Introduction of  $\Gamma$ -dependent intrinsic adsorption and desorption rate constants, including the deviations from ideal Langmuir adsorption behavior, results in the following set of equations<sup>16,17</sup>

$$\frac{d}{dt}\Gamma(t) = k(\Gamma)_{on}^{app}(\Gamma_{max} - \Gamma(t))c_b - k(\Gamma)_{off}^{app}\Gamma(t) \tag{1a}$$

with

$$k(\Gamma)_{on}^{app} = k(\Gamma)_{on}^{int}D/(D + \delta k(\Gamma)_{on}^{int}(\Gamma_{max} - \Gamma)) \tag{1b}$$

and

$$k(\Gamma)_{off}^{app} = k(\Gamma)_{off}^{int}D/(D + \delta k(\Gamma)_{off}^{int}(\Gamma_{max} - \Gamma)), \tag{1c}$$

where  $\Gamma(t)$  = surface concentration ( $\mu\text{g}/\text{cm}^2$ ) dependent on time  $t$

$k(\Gamma)_{on}$  = adsorption rate function ( $\text{cm}^3/\mu\text{g} \cdot \text{s}$ )

$k(\Gamma)_{off}$  = desorption rate function ( $\text{s}^{-1}$ )

$c_b$  = protein concentration in the bulk ( $\mu\text{g}/\text{cm}^3$ )

$D$  = diffusion constant of the protein ( $\text{cm}^2/\text{s}$ )

$\delta$  = thickness of the unstirred layer (cm)

The observed values of  $k(\Gamma)_{on}^{app}$  and  $k(\Gamma)_{off}^{app}$  can be approximated by exponential relations (see FIGURE 2):

$$k(\Gamma)_{on}^{int} = k_{+1}e^{-\alpha\Gamma}$$

and

$$k(\Gamma)_{off}^{int} = k_{-1}e^{+\beta\Gamma}, \tag{2}$$

where  $\alpha$  and  $\beta$  are respectively the interaction constants for the adsorption and desorption of this particular protein.<sup>17</sup>

From the preadsorption experiments (see RESULTS) it was concluded that the total amount of adsorbed protein has to be taken into account in these relations; for instance,



FIGURE 1. Adsorption and desorption of prothrombin on a DOPS double layer.

for a mixture of two proteins

$$\Gamma = \Gamma_1 + \Gamma_2.$$

The equilibrium association constant  $K_a$  is defined by  $K_a = k_{on}^{int}/k_{off}^{int}$ . Using equations 1b, 1c, and 2,  $K_a$  can be written as:

$$K_a = k_{on}^{int}/k_{off}^{int} = k_{on}^{app}/k_{off}^{app} = Ke^{-\gamma\Gamma},$$

where  $\gamma = \alpha + \beta$  and  $K = k_+/k_-$  is the association constant in the limit of low surface coverage.

In equilibrium one has  $d/dt \Gamma(t) = 0$  and it follows from equation 1a that

$$\Gamma_{eq} = (k_{on}^{app}/k_{off}^{app}) (\Gamma_{max} - \Gamma_{eq})c_b = Ke^{-\gamma\Gamma_{eq}}(\Gamma_{max} - \Gamma_{eq})c_b.$$

For two components one then obtains:

$$\Gamma_{eq} = \Gamma_{1eq} + \Gamma_{2eq} = (K_1 e^{-\gamma_1 \Gamma_{eq}} c_{1b} + K_2 e^{-\gamma_2 \Gamma_{eq}} c_{2b})(\Gamma_{max} - \Gamma_{eq}). \quad (3)$$

Knowing the parameters  $K_1$ ,  $K_2$ ,  $\gamma_1$ ,  $\gamma_2$ ,  $c_{1b}$ ,  $c_{2b}$ , and  $\Gamma_{max}$ , this relation permits calculation of  $\Gamma_{eq}$ . Knowing  $\Gamma_{eq}$  one may then calculate  $\Gamma_{1eq}$  and  $\Gamma_{2eq}$ .

Instead of equation 3 one may also write:

$$\Gamma_{1eq}/\Gamma_{2eq} = [(K_1 c_{1b})/(K_2 c_{2b})] e^{-(\gamma_1 - \gamma_2)\Gamma_{eq}}. \quad (4)$$

This equation shows that the ratio of the adsorbed proteins is dependent on  $\Gamma_{eq}$ . If  $\Gamma_{eq}$  is




FIGURE 1. Measurement of adsorption and desorption of prothrombin, added 1600 s after preadsorption of fibrinogen (see text). The initial value of  $\Gamma = 0.335 \mu\text{g}/\text{cm}^2$  (at  $t = 0$ ) was calculated for the DOPS double layer.




FIGURE 2. The effect of surface concentration  $\Gamma$  on the value of  $k_{on}^{int}$  (logarithmic plot); O = without preadsorption of fibrinogen; \* = with preadsorption of  $0.08 \mu\text{g}/\text{cm}^2$  fibrinogen.

changed, for instance by changing the values of  $c_{1b}$  and  $c_{2b}$  by the same dilution factor, the ratio of the absorbed quantities of protein will also change.

RESULTS AND DISCUSSION

The Effect of Preadsorption of Fibrinogen on the Adsorption of Prothrombin

To investigate the effect of the presence of another absorbed protein on the absorption of prothrombin, a double layer of DOPS was stacked on a reflecting chromium surface and after 50 seconds  $20 \mu\text{g}/\text{ml}$  fibrinogen was added to the cuvette. The adsorption reaches a plateau value of  $0.1 \mu\text{g}/\text{cm}^2$  after 1000 s (FIGURE 1). The content of the cuvette was rinsed with buffer and a small desorption was observed. After 1600 s prothrombin adsorption was started by adding the protein to the cuvette (final concentration,  $20 \mu\text{g}/\text{ml}$ ). Within 100 seconds adsorption was completed at an adsorbed mass of  $0.18 \mu\text{g}/\text{cm}^2$ . After 2000 seconds the desorption of prothrombin was measured by again rinsing the cuvette with buffer.

The results of 10 prothrombin adsorptions with and without preadsorption of fibrinogen are shown in FIGURE 2. The effect of a preadsorption of fibrinogen ( $0.08 \mu\text{g}/\text{cm}^2$ ) on the adsorption of prothrombin is to lower the value of  $k_{on}^{int}$  by about a factor of ten and to cause a parallel shift to the left of about  $0.06 \mu\text{g}/\text{cm}^2$ . Comparing these values, it is concluded that the adsorbing prothrombin molecules, in first approximation, do not discriminate between preadsorbed fibrinogen or prothrombin. Apparently

the adsorption of one protein can be strongly influenced by the presence of another protein and in the case of a mixture of proteins the total adsorbed amount of proteins has to be taken into account. FIGURE 2 also shows the exponential relation between  $k_{on}^{int}$  and the surface concentration  $\Gamma$ , as has been observed before.<sup>16,17</sup>

### Computer Simulations

In order to investigate the consequences of this model for the adsorption of a protein mixture, we performed some computer simulations. The choice of the parameter values of the model was guided by the following considerations.

From previous experiments with adsorptions of prothrombin and fibrinogen mixtures a value of  $\Gamma_{max} = 0.3 \mu\text{g}/\text{cm}^2$  was estimated and for  $\Gamma_{eq}$  a value of  $0.25 \mu\text{g}/\text{cm}^2$  was chosen arbitrarily. If the first protein must show preferential adsorption for a highly diluted mixture and be displaced by the second protein for an undiluted mixture, the following requirements must be satisfied (cf. equation 4):

$$K_1 c_{b1} \gg K_2 c_{b2} \approx 1 \quad \text{and} \quad \frac{K_1 c_{b1}}{K_2 c_{b2}} e^{-(\gamma_1 - \gamma_2)0.25} \ll 1. \quad (5)$$

Therefore, the following numerical values were chosen:

$$K_1 c_{b1} = 10^4; K_2 c_{b2} = 10; \quad \frac{K_1 c_{b1}}{K_2 c_{b2}} e^{-(\gamma_1 - \gamma_2)0.25} = 10^{-2},$$

resulting in  $\gamma_1 - \gamma_2 = 45 \text{ (cm}^2/\mu\text{g)}$ .

For the dynamic simulations, values of  $k_{+1} = k_{+2} = 1 \text{ cm}^3/\mu\text{g} \cdot \text{s}$  were chosen. For proteins with molecular weights of 100,000, this value corresponds to an association rate constant of approximately  $10^8 \text{ M}^{-1}/\text{s}$ . Thus a high, and identical, value of  $k_1$  was chosen for both proteins, about one order of magnitude lower than the upper limit as predicted by Smoluchowsky theory. For the bulk concentrations protein values of  $c_{b1} = 1000 \mu\text{g}/\text{cm}^3$  and  $c_{b2} = 10 \mu\text{g}/\text{cm}^2$  were chosen, that is of the order of the plasma concentrations of fibrinogen and prothrombin, and the corresponding values of  $k_-$  thus became:  $k_{-1} = 0.1 \text{ s}^{-1}$  and  $k_{-2} = 1 \text{ s}^{-1}$ . A value of  $D/\delta = 10^{-3} \text{ cm/s}$  was chosen in Equation 1, in agreement with ref. 16.

It should be stressed, however, that the simulation results shown in FIGURES 3-5 are determined qualitatively by the conditions in equation 5 and do not depend critically on the other assumed parameter values. Also, the picture remains qualitatively unaltered if it is assumed that the effects of the adsorbed quantities of both species are not strictly additive. This is suggested, for instance, by the finding that preadsorption of fibrinogen is slightly less effective in lowering  $k_{on}$  for prothrombin than preadsorption of prothrombin itself. Such effects could be expected for proteins having different values of  $\gamma$  and could be simply accounted for by assuming a relation of the type  $\Gamma_{eq} = \Gamma_{1eq} + f\Gamma_{2eq}$ , with  $f$  as a weight factor.

The above estimated value of  $\gamma$ , the interaction constant, of about  $45 \text{ cm}^2/\mu\text{g}$  is in good agreement with the experimental values found. For the system tested until now  $\gamma$  values between  $20 \text{ cm}^2/\mu\text{g}$  and  $80 \text{ cm}^2/\mu\text{g}$  were found.<sup>17</sup>

### The Ratio of the Adsorbed Proteins as a Function of Time and Dilution

FIGURE 3 shows the sequence and ratio of the adsorption of the two proteins. Initially the protein with the highest  $K c_b$  value is adsorbed. At a surface concentration

CUYPERS *et al.*: ADSORPTION KINETICS

of about 0.15  $\mu\text{g}/\text{cm}^2$  and the corresponding maximum is reached at about 5 sec. It is strongly suggested that the protein with the highest adsorption capacity can replace the first protein.

The concentrations of the adsorbed proteins are presented in Figure 3. It is shown that, although the surface density changes from time to time, the total amount adsorbed remains constant. The effect of different values of the interaction constant  $\gamma$  for a constant velocity,  $v = 12.55 \text{ cm}^2/\text{sec}$ , is shown in Figure 4 for a constant velocity,  $v = 12.55 \text{ cm}^2/\text{sec}$ .

The importance of an "interaction constant" between protein molecules needs some comment, as has been pointed out recently by a number of authors. It describes the interaction between some real chemical physical




FIGURE 3. Sequential adsorption of proteins from a binary mixture. Parameter values are given in the text.

FIGURE 4. The effect of the interaction constant of protein on the final adsorbed amount of the protein (when  $v = 12.55 \text{ cm}^2/\text{sec}$ ).

and between proteins that influence their distribution. The effect of an interaction constant on the interaction constant (e.g., for albumin at pH = 7 and  $\gamma = 23$  and for  $\beta$ -globulin at pH = 6.0) strongly suggests the existence of electrostatic interactions. However, the data are not sufficient to explain such effect. Their self-aggregation and differences in molecular dimensions and configurations could largely influence the surface distribution of different proteins. It will be difficult to distinguish between electrostatic and other effects, but their net effect can be estimated by the experimental constants  $\alpha$  and  $\beta$  in our model.

The degree of displacement is not observed in our model. It is suggested that some special property of the BSA may be responsible for this displacement constant  $\gamma$  at physiological pH. This




FIGURE 4. The effect of dilution on the final surface concentrations of the proteins from a binary mixture. Parameter values are as in Figure 3.

of about  $0.16 \mu\text{g}/\text{cm}^2$  the exponential interaction term already influences  $\Gamma_{1\text{eq}}/\Gamma_{2\text{eq}}$  so strongly that the protein with the lower  $K_c$  value and a lower interaction constant  $\gamma$  can replace the first protein.

The concentrations of the adsorbed proteins as a function of dilution are presented in FIGURE 4. It is shown that, although the ratio of protein concentrations in the solution does not change upon dilution, there is a drastic change in the ratio of adsorbed proteins. The effect of different values of the interaction constants is shown in FIGURE 5 for a range of values  $\gamma_1 = 35\text{--}55 \text{ cm}^2/\mu\text{g}$ .

The significance of an "interaction constant" between protein molecules needs some comment, as has been pointed out recently.<sup>18</sup> In a narrow sense it describes the situation where some real chemical or physical, either attractive or repulsive, forces




FIGURE 5. The effect of the interaction constant of protein 1 on the final adsorbed amount. Other parameter values are as in FIGURE 3.

exist between molecules that influence their distribution. The effect of pH and calcium ions on the interaction constant (e.g., for albumin,  $\gamma = 76$  at pH = 7.5 and  $\gamma = 23$  at pH = 6.0) strongly suggests the existence of such forces, particularly the presence of electrostatic interactions. However, the finite dimensions of the molecules could be sufficient to explain such effect. Their self-penetration is impossible and differences in molecular dimensions and orientations could largely influence the surface distribution of different proteins. It will be difficult to differentiate between various physicochemical forces and spacing effects, but their net effects seem to be reasonably well described by the exponential constants  $\alpha$  and  $\beta$  in our model.

Fibrinogen displacement is not observed in HMWK-deficient plasma,<sup>5,9,10</sup> and this suggests some special property of the HMWK molecule resulting in a low value of the interaction constant  $\gamma$  at physiological pH. Indeed it is known that the light chain of



HMWK is responsible for the binding to negatively charged surfaces,<sup>19</sup> and this light chain has a rather unique amino acid composition with an average isoelectric point at pH 7.4. This implies that a large part of the molecule is uncharged at physiological pH, which could favor a low value of  $\gamma$ . It is tempting to speculate about such relations between the values of  $\gamma$  for different proteins and their amino acid composition and function. The same is true for a possible regulatory role of changes in the value of  $\gamma$  as a result of local triggers such as changes in pH or calcium concentrations. Such changes could cause altered compositions of a mixture of adsorbed proteins.

Sequential adsorption of proteins can be explained in a classical binding model by simply assuming that an abundant protein is displaced by a scarce protein with higher binding affinity. In such a model, however, dilution would not cause a qualitatively different behavior but would only make the process slower. A qualitative effect of dilution, as observed for fibrinogen, could still be explained in the framework of classical binding theory by assuming that, due to adsorption, the highly diluted bulk phases becomes depleted of scarce proteins.<sup>10</sup> The model presented in the present study, however, may also explain such a qualitative effect of dilution in those cases<sup>8</sup> where depletion of protein from the bulk phase did not occur.

### SUMMARY

A model for protein adsorption kinetics has been presented. This model includes diffusion-limited adsorption, adsorption and desorption rate constants  $k_{on}$  and  $k_{off}$ , which are dependent on the surface concentration, and an interaction term for the mutual influence of the adsorbed protein molecules. It has been shown that, in first approximation, values of  $k_{on}$  and  $k_{off}$  are exponential functions of the surface concentration. Assuming an adequate interaction term, it is possible to show with this model for a mixture of proteins that the ratio of the adsorbed proteins is strongly dependent on the overall surface concentration even if the ratio of the bulk concentrations of these proteins is kept constant. Differences in interaction terms for the different proteins offer a possible explanation for the peculiar behavior of plasma protein adsorption on a surface at different dilutions of the plasma, the so-called Vroman effect.

### ACKNOWLEDGMENT

We thank Dr. Thomas Horbett from the University of Washington, Seattle, and Dr. John Brash from the McMaster University, Hamilton, Ontario, for our stimulating discussions.

### REFERENCES

1. VROMAN, L., A. ADAMS, M. KLINGS & G. FISCHER. 1975. Fibrinogen, globulins, albumin and plasma at interfaces. *Adv. Chem. Ser.* **145**: 255.
2. VROMAN, L. & A. ADAMS. 1969. Finding with the recording ellipsometer suggesting rapid exchange of specific plasma proteins at liquid/solid interfaces. *Surf. Sci.* **16**: 438-466.
3. BRASH, J. L., S. UNİYAL & B. M. C. CHAN. 1981. Interactions of proteins in plasma with various foreign surfaces. *Artif. Organs* **5**(Suppl.): 475-477.
4. VROMAN, L., A. ADAMS, M. KLINGS, G. FISHER, P. C. MUNOZ & R. P. SOLENSKY. 1977. Reactions of formed elements of blood with plasma proteins at interfaces. *Ann. N.Y. Acad. Sci.* **283**: 65-76.

5. VROMAN L., A. L. ADAMS, G. C. FISCHER & P. C. MUNOZ. 1980. Interaction of high molecular weight kininogen, factor XII and fibrinogen in plasma at interfaces. *Blood* **55**: 156-159.
6. BREEMHAAR, W., E. BRINKMAN, D. J. ELLENS, T. BEUGELING & A. BANTJES. 1984. Preferential adsorption of high density lipoprotein from blood plasma onto biomaterial surfaces. *Biomaterials* **5**: 269-274.
7. BRASH, J. L. & P. TEN HOVE. 1984. Effect of plasma dilution on adsorption of fibrinogen to solid surfaces. *Thromb. Haemostas.* **51**(3): 326-330.
8. HORBETT, T. A. 1984. Mass action effects on competitive adsorption of fibrinogen from hemoglobin solutions and from plasma. *Thromb. Haemostas.* **51**(2): 174-181.
9. WOJCIECHOWSKI, P., P. TEN HOVE & J. L. BRASH. 1986. Phenomenology and mechanism of the transient adsorption of fibrinogen from plasma (Vroman effect). *J. Colloid Interface Sci.* **111**: 455-465.
10. VROMAN, L. & A. L. ADAMS. 1986. Adsorption of proteins out of plasma and solutions in narrow spaces. *J. Colloid Interface Sci.* **111**: 391-402.
11. BLODGETT, C. 1935. Films built by depositing successive monolayers on a solid surface. *J. Am. Chem. Soc.* **57**: 1007-1022.
12. CUYPERS, P. A. 1976. Dynamic ellipsometry: Biochemical and biomedical applications. Ph.D. thesis. University of Limburg, Maastricht, the Netherlands.
13. CUYPERS, P. A., W. TH. HERMENS & H. C. HEMKER. 1977. Ellipsometric study of protein films on chromium. *Ann. N.Y. Acad. Sci.* **283**: 77-85.
14. KOP, M. J. M. M., P. A. CUYPERS, T. LINDHOUT, H. C. HEMKER & W. TH. HERMENS. 1984. The adsorption of prothrombin to phospholipid monolayers quantitated by ellipsometry. *J. Biol. Chem.* **259**: 13993-13998.
15. CUYPERS, P. A., J. W. CORSEL, M. P. JANSSEN, J. M. M. KOP, W. TH. HERMENS & H. C. HEMKER. 1983. The adsorption of prothrombin to phosphatidylserine multilayers quantitated by ellipsometry. *J. Biol. Chem.* **258**: 2426-2431.
16. CORSEL, J. W., G. M. WILLEMS, J. M. M. KOP, P. A. CUYPERS & W. TH. HERMENS. 1986. The role of intrinsic binding rate and transport rate in the adsorption of prothrombin, albumin and fibrinogen to phospholipid bilayers. *J. Colloid Interface Sci.* **111**: 544-553.
17. CUYPERS, P. A., G. M. WILLEMS, J. M. M. KOP, J. W. CORSEL, M. P. JANSSEN & W. TH. HERMENS. 1986. Kinetics of protein sorption on phospholipid membranes measured by ellipsometry. *In* Proteins at Interfaces. J. L. Brash & T. A. Horbett, Eds. American Chemical Society, Washington, D.C.
18. KAROLCZAK, M. 1984. The specification of the condition leading to interpretation of adsorption isotherms in the terms of excess electrochemical free energy of mixing and some implications of the interpretations. *J. Colloid Interface Sci.* **97**: 284-290.
19. KELLERMANN, J., F. LOTTSPEICH, A. HENSCHEN & W. MÜLLER-ESTERL. 1986. Completion of the primary structure of human high molecular mass kininogen. *Eur. J. Biochem.* **154**: 471-478.