

Risk factors, ageing and signal transduction

Citation for published version (APA):

ten Haaf, J. A., & Jolles, J. (1990). Risk factors, ageing and signal transduction: special reference to metals and inositol phospholipid metabolism. In *From Gene to Man* (pp. 182-185). Stichting Gerontologie en Geriatrie.

Document status and date:

Published: 01/01/1990

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

RISK FACTORS, AGING, AND SIGNAL TRANSDUCTION: SPECIAL REFERENCE TO METALS AND INOSITOL PHOSPHOLIPID METABOLISM

J.A. ten Haaf and J. Jolles

Department of Neuropsychology and Psychobiology, University of Limburg, P.O. Box 616, 6200 MD Maastricht

ABSTRACT

The effects of several metal ions on the metabolism of (poly-) phosphoinositides in rat brain membranes were studied. It appeared that divalent as well as trivalent ions exerted differential effects on the phosphorylation of the phosphoinositides, suggesting a role of this polyphosphoinositide (pPI) system in the healthy functioning of the brain, in neurological disorders related to age, and in brain dysfunctions due to exposure to neurotoxic metals.

INTRODUCTION

The pPI metabolism is an intermediate system involved in the process of trans membrane neuronal signal transduction. Therefore, the study of the pPI metabolism may lead to a better understanding of processes, such as ageing, that affect signal transduction. Receptor stimulation leads to phospholipase C (PLC) activation subsequently cleaving pPI into the well known second messengers diacylglycerol (DAG) and inositolphosphates. On the other hand, the pPIs are interconverted into each other by membrane-bound, specific kinases and phosphomonoesterases. DAG is rapidly phosphorylated into phosphatidic acid (PA) by cytosolic DAG-kinase, after which in the endoplasmic reticulum pPI is reconstituted from PA-CMP and inositol. In our laboratory the effects of risk factors - known to be likely involved in processes such as mental retardation and/or neuronal degradation- on this pPI metabolism are studied in relation to ageing. Several metal ions play an essential role as "micro-" and "macro-minerals" (present in the mammalian brain in μmolar and mmolar concentrations respectively) in the living organism. More specifically, Mg^{++} is essential for the kinases and monoesterases involved in the pPI metabolism, Ca^{++} for PLC activity, and Mn^{++} and Mg^{++} are essential for the transferases. Several metal ions not belonging to these essential elements are known for their (neuro-)toxic effects: These include Cd^{++} , Hg^{++} , Pb^{++} , and Al^{+++} . The present study reports on the differential effects of several metal ions on the PI metabolism. Especially those metals receive attention which are implicated in aging processes and neurodegenerative diseases, such as for instance Zn^{++} and Alu^{+++} .

EXPERIMENTAL PROCEDURE

Following the approach of Jolles et al. (1980), a lysed crude mitochondrial/synaptosomal fraction from rat brain homogenate was prepared and phosphorylation started by the addition of ATP, using radiolabeled ATP. After termination of the reaction and subsequent lipid extraction, thin layer chromatography revealed phosphate incorporation in PI-phosphate (PIP), PI-bis phosphate (PIP2) and PA.




Fig. 1 Effect of Ni^{++} on inositollipid phosphorylation. Abscissa: log concentration of cation during incubation, B: no cation present. Ordinate: phosphate incorporation as percentage of control level.

RESULTS

Incubation of the homogenate in the presence of monovalent metal ions (K^+ , Li^+ in a dose range of $1 \mu\text{M}$ to 1mM) prior to ATP addition had no effect (data not shown), whereas ($1 \mu\text{M}$ to 1mM) Mn^{++} , Co^{++} , Ni^{++} , Cu^{++} , Zn^{++} (in Fig. 1, the dose response curve of Ni^{++} is given), and Cd^{++} , Hg^{++} , Pb^{++} (Fig. 2, Pb^{++}) exerted differential effects on PI metabolism. Al^{+++} and Au^{+++} inhibited PIP, PIP2 and PA formation dose-dependently (Fig. 3, Al^{+++}).




Fig. 2 Effect of Pb^{++} on inositollipid phosphorylation. For legends, see Fig.1.




Fig. 3 Effect of Al^{+++} on inositollipid phosphorylation. For legends, see Fig.1.

DISCUSSION

At which level may these ions interfere with pPI metabolism? Cations may form complexes with ATP, selectively interact with monoesterases, PLC, or with one or more of the involved kinases, interfere with transferase activity. We have good indications that in our assay system no PLC activity is present under basal conditions. This is the more likely since no Ca^{++} is present during the incubation. Pb^{++} stimulates PA, and PIP and PIP2 are inhibited (Fig. 4), this could point to a stimulation of PLC activity. This is in agreement with findings reported in literature that Pb^{++} can "imitate" Ca^{++} at the biochemical level. The chemical parameters of a certain cation -such as ion radius, coordination number, or character of binding with chelators- can not simply explain the differential effects we found with most metal ions. As far as we tested, metal ions can only be grouped on charge with respect to the effects found: Monovalent ions are not effective, bivalent ions exert differential effects, and trivalent ions only inhibited phosphate incorporation in PIP, PIP2, an PA. Further research will be directed to uncover possible mechanisms. The possibility that metal ions are involved in neurodegenerative processes such as Alzheimer disease is recovering increasingly interest, whereas a causal relation to certain syndromes is amply documented (eg Aluminum encephalopathy). Based upon the results presented here, follow-up research is now being performed with the same ions in brains of old rats and of Alzheimer brain material.

REFERENCES

- Jolles J., H. Zwiers, C. van Dongen, P. Schotman, K.W.A. Wirtz and W.H. Gispen, Modulation of brain polyphosphoinositide metabolism by ACTH-sensitive protein phosphorylation, *Nature* 286 (1980) 623-625.
- Ten Haaf J.A., T. van Teunenbroek, M. Markerink and J. Jolles, Differential effects of metal ions on the metabolism of phosphatidylinositol in rat brain membrane preparation. In: Proceedings of the Giuseppe Porcellati Foundation International Meeting on Neurochemical Aspects of Phospholipid Metabolism, May 26-28 Perugia, Italy, 1988, 61.