

The Ramanujan Journal (2020) 51:329–332
<https://doi.org/10.1007/s11139-018-0091-y>



A note on the Moll–Arias de Reyna integral

M. L. Glasser^{1,2} 

Received: 1 May 2018 / Accepted: 12 September 2018 / Published online: 13 February 2019
 © Springer Science+Business Media, LLC, part of Springer Nature 2019

Abstract

The Moll–Arias de Reyna integral

$$\int_0^\infty \frac{dx}{(x^2 + 1)^{3/2}} \frac{1}{\sqrt{\varphi(x) + \sqrt{\varphi(x)}}} \quad \text{where } \varphi(x) = 1 + \frac{4}{3} \left(\frac{x}{x^2 + 1} \right)^2$$

is generalized and several values are given.

Keywords Definite integral · Elliptic integral · Elliptic modulus · Algebraic integrand

Mathematics Subject Classification Primary 33E05, 33E20

1 Introduction

We define

$$f(a, b) = \int_0^\infty \frac{dx}{(x^2 + 1)^a} \frac{1}{\sqrt{\varphi(x) + \sqrt{\varphi(x)}}} \quad (1.1)$$

where

$$\varphi(x) = 1 + 4b^{-2}u^2, \quad u = \frac{x}{x^2 + 1}. \quad (1.2)$$

Partial financial support is acknowledged to the Spanish Junta de Castilla y León (VA057U16) and MINECO (Project MTM2014-57129-C2-1-P).

✉ M. L. Glasser
 larryg@clarkson.edu

¹ Departamento de Física Teórica, Atómica y Óptica, Facultad de Ciencias, University of Valladolid, Valladolid 40007, Spain

² Department of Physics, Clarkson University, Potsdam, NY 13676, USA

The value $f(3/2, \sqrt{3}) = \frac{\pi}{2\sqrt{6}}$ appeared as entry 3.248.5 in [5] and was shown to be incorrect by Moll et al. [1]. Unable to find the correct evaluation, the editors decided to take this entry out of later editions of the table [6]. The exact value

$$f(3/2, \sqrt{3}) = \frac{\sqrt{3}-1}{2} \Pi\left(\frac{\pi}{2}, 3-\sqrt{3}, 3^{-1/2}\right) - 6^{-1/2} F\left(\sin^{-1} \sqrt{2-\sqrt{3}}, 3^{-1/2}\right) \tag{1.3}$$

was recently provided in a mathematical *tour de force* by Arias de Reyna [2]. The aim of the present note is to provide further values of (1.1) and to suggest that the incorrect value in [5] is not merely a misprint within the scope of the parametrization (1.2).

2 Calculation

Factor $\sqrt{\varphi(x)}$ from the denominator of the integrand of (1.1), multiply the numerator and denominator by $\sqrt{\sqrt{\varphi(x)}-1}$, change the integration variable to u (note that the range of x must be divided into $[0, 1] \cup [1, \infty)$) and set $s = 2u$ to obtain

$$f(a, b) = 2^{-a} b \int_0^1 \frac{ds}{s\sqrt{1-s^2}} \left\{ [1 + \sqrt{1-s^2}]^{a-1} + [1 - \sqrt{1-s^2}]^{a-1} \right\} \sqrt{1 - \frac{b}{\sqrt{b^2+s^2}}} \tag{2.1}$$

Since both quadratic surds can be rationalized by the elliptic substitution $s = \text{cn}(\kappa, x)$ for a suitable modulus, $f(a, b)$ should be expressible in terms of elliptic integrals for integer and half-integer values of a , even in the trigonometric case $\kappa = 0$. For example, it is clear that

$$f(2, b) = \frac{1}{2} f(1, b) \tag{2.2}$$

and with the substitution $t = b/\sqrt{b^2+s^2}$

$$f(1, b) = k \int_{\kappa}^1 \frac{dt}{(t+1)\sqrt{(1-t)(t^2-k^2)}}, \quad k = \frac{b}{\sqrt{b^2+1}}. \tag{2.3}$$

which is a complete elliptic integral of the third kind which can be manipulated into standard form [4]

$$f(1, b) = \frac{\kappa}{\sqrt{k+1}} \Pi\left(\frac{\pi}{2}, \alpha^2, \kappa\right)$$

with

$$\alpha^2 = \frac{\sqrt{b^2 + 1} + b}{2\sqrt{b^2 + 1}}, \quad \kappa = \sqrt{b^2 + 1} - b. \tag{2.4}$$

For $a = 3, 4, 5, \dots$, $f(a, b)$, with $x = s^2$, can be seen to be a multiple of $f(1, b)$ plus an integral of the form

$$\int_0^1 \frac{P(x)}{\sqrt{1-x}} \sqrt{1 - \frac{b}{\sqrt{b^2 + x}}} dx, \tag{2.5}$$

where P is a polynomial. Such an integral can be transformed into a sum of elliptic integrals by the substitutions $x \rightarrow 1 - x^2, x \rightarrow \sqrt{b^2 + 1} \sin t$. For example,

$$f(3, b) = \frac{1}{2}f(1, b) - \frac{b^2}{4k} \int_k^1 \sqrt{\frac{x(x-k)}{1-x^2}} dx. \tag{2.6}$$

For $a = 3/2$ (2.4) yields

$$f(3/2, b) = \frac{b}{4} \int_0^{\pi/2} [\csc(t/2) + \sec(t/2)] \sqrt{1 - \frac{b}{\sqrt{b^2 + \sin^2 t}}} dt. \tag{2.7}$$

This can be further simplified by the substitutions $\sin t = b \sin u, \sin u = x, \sqrt{1 + x^2} = 1/y$ to

$$f(3/2, \sqrt{3}) = \frac{3}{\sqrt{8}} \sum_{\pm} \int_{\sqrt{3}/2}^1 \frac{dy}{\sqrt{y(1+y)(4y^2-3)(y \pm \sqrt{4y^2-3})}} \tag{2.8}$$

which may be reduced further to

$$f(2/3, \sqrt{3}) = \frac{3^{1/4}}{2} \int_0^{1/\sqrt{3}} \frac{dx}{X^{3/2} \sqrt{4-3X^2}} \frac{\sqrt{X-2x} + \sqrt{X+2x}}{\sqrt{X+2/\sqrt{3}}} \tag{2.9}$$

with $X = \sqrt{x^2 + 1}$ and which offers an alternative approach to (1.3).

3 Discussion

Very recently a preprint by Blaschke [3] has appeared pointing out that if the nested square root in (1.1) is replaced by the three halves power the value $\pi/2\sqrt{6}$ is obtained. Thus the error in [5] is indeed merely a misprint. Nevertheless, we examine the possibility of reproducing this value by a specific choice of (a, b) in (1.1). To keep things

simple, take $b = i$, so that $b\sqrt{s^2 + b^2} = (1 - s^2)^{-1/2}$ thus eliminating one of the surds in (1.1). Then one has

$$\begin{aligned} f(a, i) &= 2^{-a} \int_0^1 \frac{ds}{s\sqrt{1-s^2}} \left\{ [1 + \sqrt{1-s^2}]^{a-1} + [1 - \sqrt{1-s^2}]^{a-1} \right\} \\ &\quad \times \sqrt{\frac{1}{\sqrt{1-s^2}} - 1} \\ &= 2^{1-a} \int_0^1 \left[\frac{(1+x^2)^{a-2}}{\sqrt{1-x^2}} + \frac{(1-x^2)^{a-1/2}}{\sqrt{1+x^2}} \right] dx \\ &= 2^{-a} \pi \left[{}_2F_1(1/2, 2-a; 1; -1) + \frac{\Gamma(1-a)}{\sqrt{\pi}\Gamma(a-1/2)} \right]. \end{aligned}$$

On solving $f(a, i) = \pi/2\sqrt{6}$ numerically one finds the two values

$$f(0.701935\dots, i) = f(11.8052\dots, i) = \frac{\pi}{2\sqrt{6}}.$$

Acknowledgements The author thanks Victor Moll for informing him of [2,3].

References

1. Amdeberhan, T., Moll, V.: The integrals in Gradshteyn and Ryzhik. Part 14: an elementary evaluation of entry 3.411.5. *Scientia* **19**, 97–103 (2010)
2. Arias-de Reyna, J.: True value of an integral in Gradshteyn and Ryzhik's table (2018). [arxiv:1801.0964v1](https://arxiv.org/abs/1801.0964v1) [math.CA]
3. Blaschke, P.: Hypergeometric form of fundamental theorem of Calculus (2018). [arxiv:1808.04837v1](https://arxiv.org/abs/1808.04837v1) [math.CA]
4. Byrd, P.F., Friedman, M.D.: *Handbook of Elliptic Integrals for Engineers and Scientists*, 2nd edn. Springer, Berlin (1971)
5. Gradshteyn, I.S., Ryzhik, I.M.: In: Jeffrey, A., Zwillinger, D. (eds.) *Table of Integrals, Series, and Products*, 6th edn. Academic Press, New York (2000)
6. Gradshteyn, I.S., Ryzhik, I.M.: In: Zwillinger, D., Moll, V. (eds.) *Table of Integrals, Series, and Products*, 8th edn. Academic Press, New York (2015)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.