

2020

Marked increase in community-associated methicillin-resistant *Staphylococcus aureus* infections, Western Australia, 2004–2018

Lauren E. Bloomfield

The University of Notre Dame Australia, lauren.bloomfield@nd.edu.au

G W. Coombs

S Tempone

P K. Armstrong

Follow this and additional works at: https://researchonline.nd.edu.au/med_article



Part of the [Medicine and Health Sciences Commons](#)

This article was originally published as:

Bloomfield, L. E., Coombs, G. W., Tempone, S., & Armstrong, P. K. (2020). Marked increase in community-associated methicillin-resistant *Staphylococcus aureus* infections, Western Australia, 2004–2018. *Epidemiology and Infection*, *Early View Online First*.

Original article available here:

[10.1017/S0950268820000849](https://doi.org/10.1017/S0950268820000849)

This article is posted on ResearchOnline@ND at https://researchonline.nd.edu.au/med_article/1152. For more information, please contact researchonline@nd.edu.au.



This is the author's version of the following article, as accepted for publication.

Bloomfield, L.E., Coombs, G.W., Tempone, S., and Armstrong, P.K. (2020) Marked increase in community-associated methicillin-resistant *Staphylococcus aureus* infections, Western Australia, 2004-2018. *Epidemiology and Infection, Online First*. doi: 10.1017/S0950268820000849

This article was published in *Epidemiology and Infection*, 23 April, 2020.

Published version available online at: -

<https://doi.org/10.1017/S0950268820000849>

Title: Marked increase in community-associated methicillin-resistant *Staphylococcus aureus* infections, Western Australia, 2004 – 2018

Authors: L.E. BLOOMFIELD^{1,2,3}, G.W. COOMBS^{4,5}, S. TEMPONE¹, P.K. ARMSTRONG¹

1. Communicable Disease Control Directorate, Perth, Western Australia

2. Edith Cowan University, Perth, Western Australia

3. The University of Notre Dame, Fremantle, Western Australia

4. PathWest Laboratory Medicine, Perth, Western Australia

5. Murdoch University, Perth, Western Australia

Disclaimers: Nil

Corresponding Author: Lauren Bloomfield, School of Medicine, The University of Notre Dame Australia, PO Box 1225 Fremantle WA 6959, AUSTRALIA; email: lauren.bloomfield@nd.edu.au

This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI 10.1017/S0950268820000849.

Epidemiology & Infection is published by Cambridge University Press.

Summary

This study presents enhanced surveillance data from 2004 – 2018 for all community-associated methicillin-resistant *Staphylococcus aureus* (CA-MRSA) specimens collected in Western Australia (WA), and describes the changing epidemiology over this period. A total of 57,557 cases were reviewed. Annual incidence rates increased from 86.2 cases per 100,000 population to 245.6 per 100,000 population (IRR = 2.9, CI₉₅ 2.7 – 3.0). The proportion of isolates carrying Panton-Valentine leukocidin (PVL)-associated genes increased from 3.4% to 59.8% (χ^2 test for trend 7021.9, $p < 0.001$). The emergence of PVL-positive, “Queensland CA-MRSA” (ST93-IV) and “WA 121” (ST5-IV) accounted for the majority of increases in CA-MRSA across the study period. It is unclear why some clones are more prolific in certain regions. In WA, CA-MRSA rates increase as indices of temperature and humidity increase after controlling for socioeconomic disadvantage. We suggest climatic conditions may contribute to transmission, along with other socio-behavioural factors. A better understanding of the ability for certain clones to form ecological niches and cause outbreaks is required.

Marked increase in community-associated methicillin-resistant *Staphylococcus aureus* infections, Western Australia, 2004 – 2018

BACKGROUND

Staphylococcus aureus is the most commonly isolated bacterial pathogen in humans and is an important cause of skin and soft-tissue infections (SSTIs), pneumonia, septic arthritis, endocarditis, osteomyelitis, foreign-body infections, and sepsis [1]. Methicillin-resistant *S. aureus* (MRSA) isolates are resistant to penicillins and other β -lactam antibiotics and in the past were confined mainly to health care environments [2]. However, over the last 30 years there has been an increase in the number of MRSA infections in persons without the usual healthcare-associated risk factors. An increased recognition of new MRSA clones, commonly referred to as community-associated MRSA (CA-MRSA) in order to distinguish them their healthcare-associated counterparts, are responsible for such infections [2].

In Australia, CA-MRSA infections first emerged from the northern Kimberley region in Western Australia (WA) in the early 1990s, mainly associated with Aboriginal people living in remote communities [3]. Since that time, the heavy burden of staphylococcal disease and an increasing prevalence of CA-MRSA in these populations across northern Australia has been noted [4].

Different CA-MRSA strains have been demonstrated to have found ecological niches, with different regions experiencing outbreaks largely comprising different dominant strains [1], such as USA300 in North America, ST80-IV 'European clone' in countries of the Gulf Cooperation Council [5] the ST30-IV 'South Western Pacific clone' in New Zealand [2]. While the global emergence and spread of CA-

MRSA has previously been well-documented [6], Australia has been no exception [7], with ST93-IV “Queensland CA-MRSA” noted as the predominant clone in the Australia for over a decade [8].

Western Australia has adopted an aggressive approach to MRSA management, with stringent policy requirements for the screening, reporting and management of cases introduced in the 1980s [9]. Colonisation or infection with MRSA has been a notifiable condition in WA since 1982, and since 1997 all MRSA isolates have been referred to the PathWest Gram-positive typing Laboratory (GPTL) where isolates are characterised as either healthcare-associated MRSA (HA-MRSA) or CA-MRSA based on molecular markers [10].

Since 2003, GPTL has employed multilocus sequence typing (MLST) and staphylococcal chromosomal cassette *mec* (SCC*mec*) typing which has enabled the identification of novel clones not previously reported. This study presents 15 years of enhanced surveillance data from specimens collected in WA, and describes the changing epidemiology and burden of disease over this period.

CA-MRSA and HA-MRSA

Phenotypically, CA-MRSA clones are generally resistant to fewer non- β -lactam antibiotics than HA-MRSA clones [11]. Clinically, CA-MRSA infections generally occur in younger, otherwise healthy patients and are associated predominantly with SSTIs, but may also cause severe clinical syndromes such as necrotizing pneumonia and sepsis [1, 12]

In WA, CA-MRSA clones are frequently responsible for infections detected in the hospital setting, including those classified as nosocomial infections [13]. Consequently, classification of CA-MRSA and HA-MRSA clones based solely on hospitalisation and risk factors for healthcare may lead to frequent differential misclassification [11] with molecular typing methods considered a more accurate means of identifying and classifying CA-MRSA clones.

Community-associated MRSA may be differentiated from HA-MRSA clones by genotypic features such as multilocus sequence typing (MLST), having a relatively smaller *SCCmec* elements (typically type IV and V), and by the frequent presence of the Panton-Valentine leucocidin (PVL) *lukF-PV* and *lukS-PV* associated genes [11]. Typing data from GPTL are used to direct patient management and appropriate infection control responses within WA healthcare facilities.

PVL status

The protein coded by the *lukF-PV* and *lukS-PV* PVL-associated genes mediates leukocyte and tissue destruction, and it has been suggested that its presence may cause more severe SSTIs than PVL-negative (PVL-) isolates [12, 14]. However, controversy remains regarding whether observed increases in disease severity are due to the presence of the PVL toxin, or if it is a marker for other virulence factors [1]. Conflicting evidence from animal and human studies has led to inconclusive findings of the exact role of PVL in disease severity; however, evidence suggests that the effects may be strain-specific [1].

Although PVL-positive (PVL+) isolates do not appear to be related to increases in bacteraemia or mortality [12], a demonstrated propensity to cause more invasive SSTIs would place a significant burden on healthcare services if the incidence of these strains increased, with more complex procedures required to treat and manage these infections.

METHODS

Genotyping

In WA the classification of CA-MRSA and HA-MRSA clones is determined by a combination of the molecular analysis of the seven housekeeping genes sequence types using MLST, and the *SCCmec*

type using multiplex PCR [15]. Prior to 2014, all CA-MRSA (regardless of PVL status) was individually typed. Increases in case numbers and subsequent workload, combined with the low applicability of the results to patient management, led to a decision to group all PVL- CA-MRSA under one typing result. This classification was applied retrospectively to all data, and therefore all PVL- CA-MRSA are grouped together. Typing data for PVL+ isolates is available for the full study period.

Epidemiology

Cases were extracted with the following inclusion criteria:

- Collection date between 1 January 2004 and 31 December 2018
- Persons with a WA address (classified by residential postcode) at the time a sample was collected
- Sample was positive for a CA-MRSA clone identified by GPTL
- Sample was identified as a clinical specimen (screening specimens excluded)
- Specimen was not a duplicate (specimen of the same strain collected from the same patient)

Estimated resident population (ERP) figures from 2004 – 2017 were obtained from the Australian Bureau of Statistics for each region. Regional ERP for 2018 was estimated based on linear population growth trends for the period 2015-2017.

Study population

WA comprises approximately 2.6 million km² and is commonly divided into eight major geographical regions. Each of these regions varies widely on a number of factors, including geography and climate, population density, proportion of Aboriginal residents, and indices of rurality and remoteness. Regional areas of the state, particularly northern and easterly regions, contain populations with some of the highest indices of socio-economic disadvantage in Australia [16]. The

population of WA is approximately 2.6 million people, with almost three-quarters of the population residing in metropolitan Perth [17].

Regions were further categorised based on their climatic properties, using Bureau of Meteorology data for climate zones based on temperature and humidity [18].

Regions were assigned to climate zones based on having a majority of regional landmass and/or population within a particular band. Zone 1 (hot, humid summer) contained the northernmost Kimberley region, Zone 3 (hot dry summer, mild winter) contained the northern-central Pilbara and Midwest regions, Zone 4 (hot dry summer, cold winter) contained the central Wheatbelt and Goldfields regions, and Zone 5 (warm summer, cold winter) contained the southerly Metropolitan, Southwest and Great Southern regions (Figure 1). Deciles taken from the ABS Indices of Relative Socio-Economic Advantage and Disadvantage (IRSAD) were applied based on patient postcode.

Statistical analysis

Data analysis was performed using R version 3.5.2 (R Foundation for Statistical Computing; Vienna, Austria). Annual incidence rates per 100,000 population were calculated. A Wilcoxon rank sum test was used to assess difference in continuous variables. Multivariable logistic regression was used to assess age, sex and climatic region of residence as independent risk factors for PVL+ CA-MRSA. Cases with missing data for sex and age group were deleted listwise. Negative binomial regression models were selected due to over dispersion of annual count data, and were used to estimate year-adjusted regional incidence rate ratios (IRRs) and associated 95% confidence intervals (CI₉₅).

RESULTS

A total of 57,557 cases of CA-MRSA reported to GPTL between 01 January 2004 and 31 December 2018 were extracted from the database in January 2019. While overall, the majority (57%) of cases reported over the 15-year period were PVL- CA-MRSA, assessment of annual trends demonstrates the proportion of PVL+ CA-MRSA increased from 3.4% to 59.8%, comprising the majority of isolates since 2015 (chi-square test for trend 7021.9, $p < 0.0001$) (Table 1).

Incidence rates by climatic region and PVL status

Between 2004 and 2018, annual incidence rates of CA-MRSA in WA increased almost three-fold, from 86.2 cases per 100,000 population to 245.6 per 100,000 population (IRR = 2.9, CI_{95} 2.7 – 3.0). At the commencement of the monitoring period, there were some differences noted in the incidence rates of CA-MRSA in the hotter climatic zones, with significantly higher total rates in Zone 1 (IRR = 3.2, CI_{95} 2.5 – 3.9) and Zone 3 (IRR = 1.7, CI_{95} 1.4 – 2.0) compared with the cooler Zone 5 (reference category) in the south of the state. The differences noted at this time are entirely attributable to PVL- strains, with very few cases of PVL+ CA-MRSA ($n = 58$) reported within the state in 2004 (Figure 2).

The IRR of all CA-MRSA increased significantly as indices of temperature and humidity increased across climatic zones (Table 2). After adjusting for year in the negative binomial model, a trend was identified for both PVL- and PVL+ CA-MRSA. Using Zone 5 (warm summer, cold winter) as a reference, the IRR for all CA-MRSA was over 9-fold higher in total in the northerly region of the state (Zone 1), compared the reference (Zone 5).

The median age for PVL+ cases was significantly lower at 26.7 years, compared with 54.5 years for PVL- cases ($p < 0.001$). Multivariable logistic regression adjusting for collection year and IRSAD

demonstrated that male sex, younger age, and residence in hotter, more humid regions were positively associated with infection with PVL+ CA-MRSA, compared to other strains (Table 3).

Molecular Biology

The emergence of two PVL+ clones, “Queensland CA-MRSA” (ST93-IV) and “WA121” (ST5-IV), account for the majority of increases in CA-MRSA across the study period. Proportionally, Queensland CA-MRSA accounted for 1.6% of all CA-MRSA strains in 2004; this increased to 37.9% of the total by 2018. Case numbers for this clone increased across the state, and is now the single most prolific strain in WA. As shown in Figure 3, the case numbers of this clone are disproportionately higher in Zone 1, which has comparable counts to Zone 5, despite Zone 5 having a population approximately 66 times the size.

The emergence of WA121 occurred in 2011 with 5 cases of this previously undetected clone occurring in this year. This number increased rapidly to 73 cases in 2012. Since this time, year-on-year increases in WA121 have been observed, in 2018 accounted for 740/6,287 (11.8%) of all reported CA-MRSA. The same regions in which Queensland CA-MRSA has proliferated again account the majority of increase in case numbers of WA121.

Greater diversity among PVL+ isolates is noted in the southerly, cooler, more densely populated regions, with South West Pacific (SWP) CA-MRSA (ST30-IV) and the PVL+ ST22-IV representing a sizeable minority within these zones. While PVL- CA-MRSA rates increased overall during this period, proportionally PVL- isolates declined from 97% of the total to 40%. This demonstrates that rather than replacement, PVL+ isolates have had an additive effect on the over rates of CA-MRSA in WA over last 15 years.

DISCUSSION

This study reports a sustained increase in clinical CA-MRSA rates in WA, with an apparent commencement of an outbreak of PVL+ MRSA in regional areas commencing 2009. Reports of increases in CA-MRSA are occurring globally, including Europe, the Asia-Pacific region, and North America [1, 19, 20]. The introduction and proliferation of Queensland CA-MRSA in the last 15 years has dominated the landscape of CA-MRSA in WA, with this clone alone accounting for almost 40% of all CA-MRSA cases in the most recent reporting period.

In addition, the recent emergence of WA121 appears to follow a similar pattern as the Queensland CA-MRSA, with a sudden emergence and rapid proliferation in remote communities. Given the demonstrated ability for a newly introduced clone to add to the burden of disease in the community, the emergence of a new PVL+ clone in these regions is of particular concern. This same clone rapidly emerged in New Zealand between 2006 – 2011 [21] to become to dominant strain in that country during this period.

The emergence and spread of CA-MRSA, particularly in the Kimberley region, has previously been documented [9, 22]. The precise reason for the emergence and spread of PVL+ strains during this period is not currently known; however, a disproportionate burden of PVL+ CA-MRSA among younger, Aboriginal Australians is well established [7, 23-25]. Reasons for the increased number of cases within these populations are complex, and centre around environmental and behavioural determinants, such as domestic overcrowding, comorbidities (especially the presence of other skin conditions), hygiene and antibiotic use [23, 24]. Although Aboriginal status was not available as part of the dataset, the increase in case numbers associated with regions and remote areas with a high proportion of Aboriginal residents [26] suggests that Aboriginal people are likely over-represented in these data.

Queensland CA-MRSA has previously been implicated as a major contributor to the burden of disease in Aboriginal populations, suggesting particular genetic factors unique to this clone may have facilitated its spread in this population across Australia [24, 27]. The epidemiology of PVL+ USA300 within North America closely mirrors the WA experience with Queensland CA-MRSA in many ways – a rapid emergence and subsequent dominance of a particular well-adapted clone [1], which is now responsible for an increasing number of healthcare-associated infections [28].

Despite numerous incursions of USA300 observed over the surveillance period, this strain has not proliferated in WA. Similarly, despite small-scale outbreaks of Queensland CA-MRSA in the United Kingdom [29] has not resulted in the same widespread dissemination in this country [1]. In fact, this clone has had apparently very limited spread, even within the Asia-Pacific region [19]. Reasons why some strains seem to dominate in certain regions and not others remain unclear, however this suggests that there are other factors, outside of a susceptible population and apparently fit strain, which allow establishment and transmission in a certain region.

The rate of PVL+ CA-MRSA in the highly populous Zone 5 has increased, but is still lower than that of PVL- CA-MRSA. Rates of CA-MRSA in other regions that also contain areas of high disadvantage have not increased as markedly as within the state's north. This suggests that a combination of socio-behavioural and environmental factors may be reducing the ability of PVL+ CA-MRSA to proliferate within more temperate geographic locations within the state.

Climatic influences on MRSA SSTIs have previously been explored [30], although a paucity of evidence exists. The data collected by Sahoo and colleagues suggest a combination of temperature and humidity; namely, average weekly maximum temperatures above 33°C and an average weekly relative humidity between 55% and 78% was associated with an increase in *S. aureus* associated SSTIs [30]. Average maximum temperature and humidity data across climate bands published by the

Bureau of Meteorology suggests that optimum conditions for colonisation and spread of MRSA are present in northerly climate zones of the state for a large proportion of the year.

Further work linking transmission to climatic factors is recommended to better describe this phenomenon. Factors outside of the presence of PVL-associated genes and the presence of high-risk populations may help to explain the geographic differences in prolific CA-MRSA clones, and could at least partially account for the apparent ecological niche that seemingly encourages transmission of some strains within a region, while limiting spread of others.

Limitations of this retrospective, population-based study include the inability to account for changes in the healthcare provider behaviour over the 15-year study period, in particular the propensity to send specimens for culture and typing. It is reasonable to assert that a rapid rise of specific PVL+ isolates within a region, causing SSTIs in increasing number and severity, raised the profile of CA-MRSA among local clinicians and drove up testing numbers to an extent.

The observed increases in PVL- CA-MRSA observed mainly in the Kimberley region may, indeed, be largely driven by an increased propensity to test for CA-MRSA. Increased testing, however, should not have any impact on the proportion of PVL+ isolates comprising the total number of specimens, suggesting that the increase in PVL+ cases in regional WA is not entirely an artefact of increased testing, but instead represents a true outbreak within the state.

Despite this limitation, the increased rates of PVL- disease, combined with an almost 50% reduction in these clones proportionally demonstrates that PVL+ clones have not replaced PVL-clones, but rather added to the burden of disease within the state. While information on individual PVL- clones is not available, the relatively minor comparative increase in rates of these clones overall suggests

that it is unlikely that one or more particular outbreak clones were responsible for driving this, as was observed with PVL+ clones.

An additional limitation was that Aboriginality was not available as an indicator for this dataset, prohibiting further investigation of this as a risk factor. In place of this variable, IRSAD was used a proxy measure, to account for the relative socio-economic disadvantage prevalent in remote communities. After controlling for IRSAD in the regression model, still there were still significantly increased proportions of CA-MRSA identified in the northern zones.

Public health management of this current outbreak remains a challenge. The suggested propensity for CA-MRSA, particularly PVL+ clones, to spread in hotter and more humid areas, particularly among vulnerable communities, suggests WA may expect similar increases in incidence rates across a number of regions in future. Countries with climatic conditions favourable for colonisation and spread of CA-MRSA should adopt proactive mitigation strategies, particularly in areas where vulnerable populations reside.

Further work to evaluate the impact of this outbreak on the healthcare system within affected regions could provide evidence of the increased burden on individual patients that results from severe SSTIs, and further quantify the economic impact to healthcare systems. Such evidence could be used as further justification for increased prevention efforts, both to halt the current outbreak and to prevent emergence of virulent strains in future.

Acknowledgements: Nil

Financial support: Nil

Conflicts of interest: None

REFERENCES

1. David MZ, Daum RS. Community-associated methicillin-resistant *Staphylococcus aureus*: epidemiology and clinical consequences of an emerging epidemic. *Clinical Microbiology Reviews*. 2010;23(3):616-87.
2. Lakhundi S, Zhang K. Methicillin-Resistant *Staphylococcus aureus*: Molecular Characterization, Evolution, and Epidemiology. *Clinical Microbiology Reviews*. 2018;31(4).
3. Udo EE, Pearman JW, Grubb WB. Genetic analysis of community isolates of methicillin-resistant *Staphylococcus aureus* in Western Australia. *The Journal of Hospital Infection*. 1993;25(2):97-108.
4. Tong SY, Bishop EJ, Lilliebridge RA, Cheng AC, Spasova-Penkova Z, Holt DC, et al. Community-associated strains of methicillin-resistant *Staphylococcus aureus* and methicillin-susceptible *S. aureus* in indigenous Northern Australia: epidemiology and outcomes. *The Journal of Infectious Diseases*. 2009;199(10):1461-70.
5. Udo EE. Community-acquired methicillin-resistant *Staphylococcus aureus*: the new face of an old foe? *Medical principles and practice : International Journal of the Kuwait University, Health Science Centre*. 2013;22 Suppl 1:20-9.
6. Wong JW, Ip M, Tang A, Wei VW, Wong SY, Riley S, et al. Prevalence and risk factors of community-associated methicillin-resistant *Staphylococcus aureus* carriage in Asia-Pacific region from 2000 to 2016: a systematic review and meta-analysis. *Clinical Epidemiology*. 2018;10:1489-501.
7. Agostino JW, Ferguson JK, Eastwood K, Kirk MD. The increasing importance of community-acquired methicillin-resistant *Staphylococcus aureus* infections. *The Medical Journal of Australia*. 2017;207(9):388-93.
8. Williamson DA, Coombs GW, Nimmo GR. *Staphylococcus aureus* 'Down Under': contemporary epidemiology of *S. aureus* in Australia, New Zealand, and the South West Pacific. *Clinical microbiology and infection : the official publication of the European Society of Clinical Microbiology and Infectious Diseases*. 2014;20(7):597-604.
9. Riley TV, Rouse IL. Methicillin-resistant *Staphylococcus aureus* in Western Australia, 1983-1992. *The Journal of Hospital Infection*. 1995;29(3):177-88.
10. Coombs GW, Monecke S, Pearson JC, Tan HL, Chew YK, Wilson L, et al. Evolution and diversity of community-associated methicillin-resistant *Staphylococcus aureus* in a geographical region. *BMC Microbiology*. 2011;11:215.
11. David MZ, Glikman D, Crawford SE, Peng J, King KJ, Hostetler MA, et al. What is community-associated methicillin-resistant *Staphylococcus aureus*? *J Infect Dis*. 2008;197(9):1235-43.
12. Jahamy H, Ganga R, Al Raiy B, Shemes S, Nagappan V, Sharma M, et al. *Staphylococcus aureus* skin/soft-tissue infections: the impact of SCCmec type and Pantone-Valentine leukocidin. *Scand The Journal of Infectious Diseases*. 2008;40(8):601-6.
13. WA Department of Health. Infection Prevention and Control of Methicillin-resistant *Staphylococcus aureus* (MRSA) in Western Australian Healthcare Facilities (Version 1). Perth: WA Department of Health; 2013.
14. Shallcross LJ, Fragaszy E, Johnson AM, Hayward AC. The role of the Pantone-Valentine leucocidin toxin in staphylococcal disease: a systematic review and meta-analysis. *The Lancet Infectious Diseases*. 2013;13(1):43-54.
15. Coombs GW, Pearman JW, Robinson O. Western Australian Methicillin-Resistant *Staphylococcus aureus* (MRSA) Epidemiology and Typing Report. July 1 2017 to June 30 2018 Perth: WA Department of Health; 2018.
16. Australian Bureau of Statistics. IRSAD Interactive Map 2018 [Available from: <https://www.abs.gov.au/ausstats/abs@.nsf/Lookup/by%20Subject/2033.0.55.001~2016~Main%20Features~IRSAD%20Interactive%20Map~16>].

17. Australian Bureau of Statistics. Western Australia: Regional Data Summary 2018 [Available from:
http://stat.abs.gov.au/itt/r.jsp?RegionSummary®ion=5&dataset=ABS_REGIONAL_ASGS&eoconcept=REGION&measure=MEASURE&datasetASGS=ABS_REGIONAL_ASGS&datasetLGA=ABS_NRP9_LGA®ionLGA=REGION®ionASGS=REGION.
18. Bureau of Meteorology. Climate classification maps 2016 [Available from:
http://www.bom.gov.au/jsp/ncc/climate_averages/climate-classifications/index.jsp.
19. Huh K, Chung DR. Changing epidemiology of community-associated methicillin-resistant *Staphylococcus aureus* in the Asia-Pacific region. *Expert Review of Anti-Infective Therapy*. 2016;14(11):1007-22.
20. Li M, Diep BA, Villaruz AE, Braughton KR, Jiang X, DeLeo FR, et al. Evolution of virulence in epidemic community-associated methicillin-resistant *Staphylococcus aureus*. *Proceedings of the National Academy of Sciences of the United States of America*. 2009;106(14):5883-8.
21. Williamson DA, Roberts SA, Ritchie SR, Coombs GW, Fraser JD, Heffernan H. Clinical and molecular epidemiology of methicillin-resistant *Staphylococcus aureus* in New Zealand: rapid emergence of sequence type 5 (ST5)-SCCmec-IV as the dominant community-associated MRSA clone. *PLoS One*. 2013;8(4):e62020.
22. Dailey L, Coombs GW, O'Brien FG, Pearman JW, Christiansen K, Grubb WB, et al. Methicillin-resistant *Staphylococcus aureus*, Western Australia. *Emerging Infectious Diseases*. 2005;11(10):1584-90.
23. Tong SY, McDonald MI, Holt DC, Currie BJ. Global implications of the emergence of community-associated methicillin-resistant *Staphylococcus aureus* in Indigenous populations. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2008;46(12):1871-8.
24. Harch SAJ, MacMorran E, Tong SYC, Holt DC, Wilson J, Athan E, et al. High burden of complicated skin and soft tissue infections in the Indigenous population of Central Australia due to dominant Panton Valentine leucocidin clones ST93-MRSA and CC121-MSSA. *BMC Infectious Diseases*. 2017;17(1).
25. Boan P, Tan HL, Pearson J, Coombs G, Heath CH, Robinson JO. Epidemiological, clinical, outcome and antibiotic susceptibility differences between PVL positive and PVL negative *Staphylococcus aureus* infections in Western Australia: a case control study. *BMC Infectious Diseases*. 2015;15:10.
26. Australian Bureau of Statistics. Aboriginal and Torres Strait Islander Population, 2016 2016 [Available from:
<https://www.abs.gov.au/ausstats/abs@.nsf/Lookup/by%20Subject/2071.0~2016~Main%20Features~Aboriginal%20and%20Torres%20Strait%20islander%20Population%20Article~12>.
27. van Hal SJ, Steinig EJ, Andersson P, Holden MTG, Harris SR, Nimmo GR, et al. Global Scale Dissemination of ST93: A Divergent *Staphylococcus aureus* Epidemic Lineage That Has Recently Emerged From Remote Northern Australia. *Frontiers in Microbiology*. 2018;9:1453.
28. Moore CL, Hingwe A, Donabedian SM, Perri MB, Davis SL, Haque NZ, et al. Comparative evaluation of epidemiology and outcomes of methicillin-resistant *Staphylococcus aureus* (MRSA) USA300 infections causing community- and healthcare-associated infections. *International Journal of Antimicrobial Agents*. 2009;34(2):148-55.
29. Ellington MJ, Ganner M, Warner M, Boakes E, Cookson BD, Hill RL, et al. First international spread and dissemination of the virulent Queensland community-associated methicillin-resistant *Staphylococcus aureus* strain. *Clinical microbiology and infection : the official publication of the European Society of Clinical Microbiology and Infectious Diseases*. 2010;16(7):1009-12.
30. Sahoo KC, Sahoo S, Marrone G, Pathak A, Lundborg CS, Tamhankar AJ. Climatic factors and community - associated methicillin-resistant *Staphylococcus aureus* skin and soft-tissue

TABLES

Table 1 CA-MRSA cases, WA, 2004-2018

	PVL Negative CA-MRSA	PVL Positive CA-MRSA	Total		
	n (%)	n (%)	N		
TOTAL	32868 (57.6%)	24689 (42.4%)	57,557		
Sex	FEMALE	17263 (60.4%)	28,727		
	MALE	15540 (54.7%)	28,745		
	UNKNOWN	65 (76.5%)	85		
	2004	1649 (96.7%)	58 (3.3%)	1,707	
Year	2005	1573 (94.1%)	101 (5.9%)	1,674	
	2006	1693 (92%)	153 (8%)	1,846	
	2007	1718 (83.8%)	356 (16.2%)	2,074	
	2008	1916 (78.9%)	540 (21.1%)	2,456	
	2009	2077 (75.8%)	712 (24.2%)	2,789	
	2010	1933 (71.7%)	811 (28.3%)	2,744	
	2011	2289 (67.6%)	1156 (32.4%)	3,445	
	2012	2239 (62.5%)	1450 (37.5%)	3,689	
	2013	2364 (56.1%)	2006 (43.9%)	4,370	
	2014	2675 (51.8%)	2511 (48.2%)	5,186	
	2015	2769 (46.2%)	3157 (53.8%)	5,926	
	2016	2644 (42%)	3698 (58%)	6,342	
	2017	2802 (40.3%)	4220 (59.7%)	7,022	
	2018	2527 (41.2%)	3760 (58.8%)	6,287	
	Age group	0 - 9 years	3891 (41.1%)	5380 (58.9%)	9,271
		10 -19 years	2306 (34.9%)	4211 (65.1%)	6,517
		20 - 29 years	3025 (40%)	4442 (60%)	7,467
30 - 39 years		3272 (45.4%)	3898 (54.6%)	7,170	
40 - 49 years		3145 (51.4%)	3013 (48.6%)	6,158	
50 - 59 years		3041 (61.6%)	1955 (38.4%)	4,996	
60 - 69 years		3225 (76.1%)	1024 (23.9%)	4,249	
70 - 79 years		3475 (89.4%)	431 (10.6%)	3,906	
80+ years		7475 (95.9%)	334 (4.1%)	8,970	
UNKNOWN		13 (93.3%)	1 (6.7%)	14	
Climate Zone	ZONE 1	2570 (28.4%)	6402 (71.6%)	8,972	
	ZONE 3	3579 (45.9%)	4264 (54.1%)	7,843	
	ZONE 4	2187 (55.8%)	1796 (44.2%)	3,983	

ZONE 5 24532 (67.4%) 12227 (32.6%) 36,759

Table 2 Year-adjusted incidence rate ratios, by climate zone

Climate Zone	PVL ⁻ CA-MRSA		PVL ⁺ CA-MRSA		All CA-MRSA	
	IRR*	CI ₉₅	IRR*	CI ₉₅	IRR*	CI ₉₅
ZONE 1	5.38	4.70 - 6.15	15.74	11.70 - 21.16	9.11	7.22 - 11.49
ZONE 3	2.37	2.08 - 2.70	3.49	2.60 - 4.68	2.90	2.31 - 3.63
ZONE 4	1.30	1.13 - 1.48	1.74	1.29 - 2.34	1.41	1.12 - 1.77
ZONE 5	1.00	ref	1.00	ref	1.00	ref

*adjusted for year of specimen collection

Table 3 Factors associated with a PVL+ CA-MRSA infection

		aOR*	CI ₉₅
Sex	FEMALE	1.00	ref
	MALE	1.27	1.22 - 1.32
Climate Zone	ZONE 1	2.24	2.07 - 2.41
	ZONE 3	1.33	1.25 - 1.42
	ZONE 4	1.10	1.02 - 1.20
	ZONE 5	1.00	ref
	0 - 9 years	20.20	17.92 - 22.84
Age group	10 -19 years	31.22	27.60 - 35.43
	20 - 29 years	25.05	22.19 - 28.36
	30 - 39 years	19.76	17.50 - 22.37
	40 - 49 years	15.13	13.38 - 17.16
	50 - 59 years	10.80	9.52 - 12.28
	60 - 69 years	5.52	4.84 - 6.32
	70 - 79 years	2.46	2.12 - 2.87
	80+ years	1.00	ref

*adjusted for age group, sex, climate zone, IRSAD and year

FIGURES





Figure 1 Regions and climatic zones, Western Australia, adapted from Bureau of Meteorology [18]



Key: Zone 1 – hot, humid summer, Zone 3 – hot dry summer, mild winter, Zone 4 – hot dry summer, cold winter, Zone 5 – warm summer, cold winter

Figure 2 Incidence rates, CA-MRSA, Western Australia, 2004 – 2018



Key: Bengal Bay CA-MRSA (ST772-V), Taiwan CA-MRSA (ST59-V), WA121 (ST5-V), Queensland CA-MRSA (ST93-IV), USA300 (ST8-IV), South West Pacific (SWP) (ST30-IV)

Figure 3 PVL-positive CA-MRSA clones, by climatic region, 2004 – 2018