

**CORRELATION OF MATERNAL SERUM  
FETUIN/ $\alpha_2$ -HS-GLYCOPROTEIN CONCENTRATION WITH  
MATERNAL INSULIN RESISTANCE AND ANTHROPOMETRIC  
PARAMETERS OF NEONATES IN NORMAL PREGNANCY AND  
GESTATIONAL DIABETES**

László Kalabay<sup>1</sup>, Károly Cseh<sup>2a</sup>, Attila Pajor<sup>3</sup>, Éva Baranyi<sup>4</sup>, György M.  
Csákány<sup>5</sup>, Zsolt Melczer<sup>3</sup>, Gábor Speer<sup>6</sup>, Margit Kovács<sup>1</sup>, György Siller<sup>2b</sup>,  
István Karádi<sup>1</sup>, and Gábor Winkler<sup>7</sup>

1. 3<sup>rd</sup> Department of Internal Medicine Semmelweis University, Faculty of  
Medicine

2. 1<sup>st</sup> Department of Internal Medicine<sup>a</sup> and Department of Urology<sup>b</sup> Károlyi  
Hospital

3. 2<sup>nd</sup> Department of Obstetrics and Gynecology, Semmelweis University,  
Faculty of Medicine

4. Department of Diabetology, Semmelweis University, Faculty of Health  
Sciences

5. 2<sup>nd</sup> Department of Obstetrics and Gynecology, Semmelweis University,  
Faculty of Health Sciences

6. 1<sup>st</sup> Department of Internal Medicine Semmelweis University

7. 2<sup>nd</sup> Department of Internal Medicine, Szent János Hospital, Budapest,  
Hungary

**Corresponding author:** László Kalabay MD, PhD, 3<sup>rd</sup> Department of Internal Medicine, Faculty of Medicine, Semmelweis University, Kútvölgyi út 4., Budapest, Hungary, H-1125 Tel.: (36)-1-355-1122, Fax: (36)-1-355-8251, e-mail: [kalasz@kut.sote.hu](mailto:kalasz@kut.sote.hu)

**Short running title:**  $\alpha_2$ -HS glycoprotein in gestational diabetes

**ABSTRACT**

**Objective:** Human fetuin/ $\alpha_2$ -HS-glycoprotein (AHSG) is a 49 kD serum and tissue protein, which is a natural inhibitor of insulin receptor signaling. We investigated serum AHSG levels during pregnancy and whether the protein is involved in insulin resistance observed in healthy pregnant women and patients with gestational diabetes.

**Design:** One hundred and four healthy pregnant women and 23 of their neonates, 30 patients with gestational diabetes and their neonates and 30 healthy age-matched non-pregnant females as a control group were investigated in a case-control cross-sectional study.

**Methods:** Serum AHSG was determined by radial immunodiffusion.

**Results:** We observed an increase of serum AHSG concentration in the second and third trimester. Gestational diabetes patients had significantly higher AHSG levels than healthy pregnant women and non-pregnant controls. There was a highly significant positive correlation between serum AHSG concentration and indirect parameters of insulin resistance, i.e. tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), leptin, C-peptide, and C-peptide/blood glucose ratio. There was also a negative correlation between maternal AHSG, TNF- $\alpha$ , leptin levels and head circumference, body length and body weight of newborns.

**Conclusion:** AHSG, TNF- $\alpha$  and leptin may contribute to insulin resistance during normal pregnancy and gestational diabetes. AHSG along with these cytokines may also negatively regulate the neonatal skeletal development.

## Introduction

Human fetuin/ $\alpha_2$ -HS-glycoprotein (AHSG) is a 49 kD serum and tissue protein, which plays a role in the host defense and bone metabolism [1, 2]. The protein is one of the major components of the non-collagenous bone matrix, especially in the fetal age. The main source of its serum isoform is the liver, where its phosphorylated variant decreases the signal transduction of the insulin receptor; hence, it may contribute to the cellular insulin resistance [3, 4, 5]. The tumor necrosis factor (TNF) system [6] and leptin [7] were also found to contribute to insulin resistance in obesity and Type 2 diabetes, and may influence the insulin secretory capacity of the  $\beta$ -cells, too. Both cytokines may also be involved in the intrauterine bone development [8, 9].

Progressive increase in insulin resistance has repeatedly been demonstrated during the course of the normal pregnancy [10]. Decreased insulin sensitivity is even more pronounced in patients with gestational diabetes [11]. The status of hepatic insulin sensitivity in pregnancy and gestational diabetes is less clear. Studies on pregnant rats described an increased hepatic insulin resistance [12]. Because of its established role in signal transduction of the insulin receptor we studied the potential contribution of AHSG in insulin resistance accompanying normal pregnancy and GDM.

It can be suspected that some of the neonatal complications of diabetic foetopathy (e.g. alterations in body composition and anthropometrical parameters) may be explained by the pathophysiological influence of the

TNF-system and leptin. Therefore, in a case-control cross-sectional study we measured maternal serum AHSG, TNF- $\alpha$ , soluble TNF-receptor-1 and -2 (sTNFR-1 and sTNFR-2) and leptin concentrations in healthy pregnant women being in different trimesters of the pregnancy and in patients with gestational diabetes mellitus (GDM). The relationship between indirect parameters of maternal insulin resistance (fasting C-peptide level, C-peptide/fasting blood glucose ratio) and the concentration of the above mentioned proteins were studied. Different anthropometric parameters of the neonates (body length, body weight and head circumference) were compared to the maternal cytokine and AHSG levels.

### **Subjects and Methods**

One hundred and four healthy pregnant women (35 in the first, 31 in the second, and 38 in the third trimester), 30 GDM patients with high fasting C peptide levels in the 20-40<sup>th</sup> gestational weeks and 30 healthy age-matched non-pregnant females as a control group were investigated (Table 1). All participants gave their written informed consent. The diagnostic criteria of GDM were stated by a 75 g oral glucose tolerance test (OGTT) according to the WHO classification [13] To maintain normoglycemia all of these patients required intensified insulin treatment (3 or more short acting shots, bedtime intermediar insulin).

Anthropometric parameters (body weight, length and head circumference) of 30 neonates (13 boys, 17 girls) of mothers with GDM, 23 newborns (11 boys, 12

girls) delivered by healthy pregnant women between the 38<sup>th</sup>-40<sup>th</sup> week in the third trimester have been analyzed in correlation with the maternal clinical and laboratory parameters. All deliveries of the mothers with GDM also succeeded between the 38<sup>th</sup>-40<sup>th</sup> gestational weeks. Table 2 shows the anthropometric parameters of neonates.

Serum AHSG concentration was determined by radial immunodiffusion using a monospecific goat anti-human AHSG antibody (IgG fraction, Incstar, Cat No. 81931, 13.7 mg/ml, in a final concentration of 84 µl/11.5 ml gel, mean coefficient of variance, IACV: 3.6%, mean interassay coefficient of variance, IECV: 6.2%). Tumor necrosis factor- $\alpha$  (Sigma, St.Louis, MI, USA, IACV: 4.8%, IECV: 6.7%), sTNFR-1 (Bender MedSystem, Vienna, Austria, IACV: 1.9%, IECV: 8.6%), sTNFR-2 (Bender MedSystem, Vienna, Austria, IACV: 1.4%, IECV: 2.0%), leptin (DRG International, USA, IACV: 4.6%, IECV: 6.6%) concentrations were determined by ELISA, serum fasting C-peptide concentration by RIA (Biodata, Rome, Italy, IACV: 5.6%, IECV: 7.3%, normal fasting range 0.66-2.50 ng/ml). HbA<sub>1c</sub> was measured by HPLC (BioRad, normal value at non-diabetics 4.3-5.8%), serum fructosamine by Boehringer (Mannheim, Germany) automatic analyzer kit (normal value at non-pregnant non-diabetics 185-280 µmol/l). The statistical analysis was performed by the Mann-Whitney linear correlation (Spearman), and multivariate analysis using the SPSS v10 statistical program.

## Results

### Serum AHSG concentrations in non-pregnant and pregnant women and in patients with gestational diabetes mellitus

Compared to non-pregnant women there was an increase of serum AHSG concentration during the first, second, and third trimesters of pregnancy (Figure 1). In patients with GDM significantly elevated AHSG levels were observed compared to non-pregnant controls and of healthy pregnant women at any trimester (Figure 1).

Significantly elevated ( $p < 0.01$ ) serum TNF- $\alpha$ , sTNFR-1, sTNFR-2, leptin and C-peptide levels were found in GDM patients as compared to healthy non-pregnant controls and healthy pregnant females at any trimester (Table 3). In healthy pregnant females these values were also significantly higher ( $p < 0.01$ ) in the third trimester compared to the first and second ones and to the non-pregnant controls (Table 3).

### Correlation between maternal serum AHSG concentration and the indirect parameters of insulin resistance

There was a significant positive correlation between serum AHSG concentration and the indirect parameters of maternal insulin resistance (fasting C-peptide concentration and C-peptide/ Blood glucose ratio) in GDM patients and in healthy pregnant women (Table 4). Maternal serum AHSG concentration was also found to be in a linear positive correlation with the maternal serum TNF- $\alpha$  and leptin values in the GDM group and in the healthy pregnant groups. These correlations were not observed in non-pregnant controls (Table 4).

Multivariate analysis showed that TNF- $\alpha$  had a significant effect on AHSG concentrations both in GDM ( $r = 0.781$ ,  $p < 0.0001$ ) and in normal pregnancy ( $r = 0.737$ ,  $p = 0.004$ ). In this analysis, AHSG alone did not have a significant effect on indirect parameters of insulin resistance (Cp and Cp/Blood sugar ratio), however, in combination with TNF- $\alpha$  and leptin it had a significant effect in GDM (adjusted  $r^2 = 0.575$ ,  $p = 0.001$ ) and in normal pregnancy (adjusted  $r^2 = 0.823$ ,  $p < 0.001$ ).

Correlation between maternal serum AHSG concentration and the anthropological parameters of newborns

The anthropological parameters of neonates (both sexes) delivered by the GDM mothers were slightly decreased as compared to those delivered by healthy pregnant women (Table 2).

Body length and head circumference of the newborns were found to be in a significant negative correlation with the maternal AHSG concentration in GDM (Table 5). In healthy pregnant women a significant negative linear correlation was calculated with all three anthropological parameters of the newborns (Table 5). Among body length (bl), head circumference (hc), body weight (bw) of the newborns and maternal serum TNF- $\alpha$  ( $r_{hc} = -0.4857$ , 95% C.I. =  $-0.7254 - -0.1411$ ) and leptin ( $r_{bl} = -0.4376$ , 95% C.I. =  $-0.6951 - -0.0807$ ,  $p = 0.0156$ ;  $r_{bw} = -0.3706$ , 95% C.I. =  $-0.6513 - -0.0007$ ,  $p = 0.0438$ ) levels significant negative correlations have been calculated in the GDM group. In healthy pregnant women a significant negative linear correlation was observed only between



maternal leptin concentration and the head circumference of the neonates ( $r = -0.6001$ , 95% C.I.:  $-0.8283 - -0.2010$ ,  $p = 0.0026$ ).

Multivariate analysis showed that maternal serum AHSG concentration alone did not have a significant influence neither on head circumference nor on body length of neonates in GDM and normal pregnancy. However, in combination with TNF- $\alpha$ , leptin and C-peptide AHSG had a significant effect on these parameters in GDM (adjusted  $r^2 = 0.563$ ,  $p = 0.002$ ), but not in normal pregnancy (adjusted  $r^2 = -0.069$ ,  $p = 0.583$ ).

## **Discussion**

The increase of maternal serum AHSG concentration during pregnancy has been observed but not analyzed and interpreted in details [14]. AHSG is considered as a negative acute phase reactant, the concentration of which decreases during infections, trauma and liver cirrhosis [15, 16, 17]. It is tempting to speculate the biological significance of this protein in clinical settings where its concentration increases, i.e. in pregnancy.

The aim of the present study was to investigate the role of AHSG in maternal insulin resistance in GDM and in normal pregnant women. The phosphorylated variant of AHSG was described among the first proteins, which interfere with the intracellular signaling of the insulin receptor [1, 2, 3, 4, 5]. No relationship between serum AHSG levels and any parameters of insulin resistance has been

described so far. Only the phosphorylated form of AHSG is able to inhibit insulin signaling *in vitro*, the dephosphorylated form is inactive [3, 5]. AHSG isolated from human serum still retains some inhibitory activity on insulin receptor signaling, yet this activity is much less than that of recombinant AHSG produced in the baculovirus expression vector system [3]. The presence of phosphorylated AHSG in human serum (approximately 20% of the total) has been recently been demonstrated in healthy persons [18].

During the course of normal pregnancy increasing insulin resistance occurs, which is even more pronounced in GDM. Previously we found that the progressive elevation of the fasting C-peptide and C-peptide/Blood glucose ratio are easily measurable parameters of progressive insulin resistance in normal pregnancy and in GDM (due to ethical reasons clamp or other *in vitro* models for exact estimation of insulin resistance were not performed in our subjects) [19]. Therefore we considered these states as a model to study the contribution of different parameters (e.g. TNF-system, leptin and also AHSG) to the transitory insulin resistance.

We found significant positive correlations among the TNF-system, leptin, AHSG, and the indirect parameters of insulin resistance in both healthy pregnant women and GDM patients. These data suggest the contribution of the elevated AHSG levels to insulin resistance. The liver is the main site of AHSG synthesis, however in pregnancy the foetoplacental unit can be an other source of the protein. This is supported by the elevation of the protein concentration during the course of pregnancy, especially in marked in the third trimester. In adults

AHSG can be expressed not only in the liver but in other organs, thus in the endometrium. In addition, autoantibodies against endometrial AHSG and transferrin have been demonstrated in patients with endometriosis [20]. An increased hepatic synthesis during pregnancy, however, cannot be ruled out. Our observations may suggest that this increased synthesis of AHSG in the liver contributes to the decreased hepatic insulin sensitivity during normal pregnancy and GDM. TNF- $\alpha$  may also have a regulatory role on the synthesis of the protein.

The negative correlation between maternal serum leptin, TNF- $\alpha$  and head circumference of the newborns may raise the possibility that these cytokines can serve as negative regulators of the neonatal bone development, as it was suggested earlier in animal models [8, 9]. The negative regulatory role of AHSG on bone development has also been postulated, e.g. AHSG prevented bone resorption and hydroxylapatite formation in vitro [21, 22]. The biological significance of AHSG in the regulation in human neonatal bone development has not been clarified, however, several data from animal studies including mice knocked out for AHSG gene raise the possibility that the protein may prevent unwanted calcification [22]. The negative correlation between the AHSG concentrations measured in maternal serum during normal pregnancy and GDM and the head circumference of the neonates further supports the hypothesis of the negative regulatory role of the protein in human neonatal bone development. Further studies are necessary to the exact role of AHSG in this process.

In conclusion, AHSG, TNF- $\alpha$  and leptin in combination may contribute to insulin resistance during normal pregnancy and GDM. These cytokines and AHSG may also negatively regulate the neonatal skeletal development.

### **Acknowledgment**

Authors thank Professor George Füst (3<sup>rd</sup> Department of Medicine, Faculty of Medicine, Semmelweis University) for his technical support and critical review of the manuscript. This work was supported by a grant of Ministry of Health Hungary, ETT 03/2000, OTKA T025449, and FKFP-0106/2000.

### **References**

1. Dickson IR, Poole AR, Veis A: Localization of plasma  $\alpha_2$ HS glycoprotein in mineralising human bone. *Nature* 1975 **256** 430-432.
2. Arnaud P, Miribel L, Emerson DL:  $\alpha_2$ -HS glycoprotein. *Methods in Enzymology* 1988 **163** 431-441.
3. Srinivas PR., Wagner AS, Reddy LV, Deutsch DD, Leon MA, Goustin AS et al. Serum  $\alpha_2$ -HS-glycoprotein is an inhibitor of the human insulin receptor at the tyrosine kinase level. *Molecular Endocrinology* 1993 **7** 1445-1455.

4. Kahn CR. Causes of insulin resistance. *Nature* 1995 **373** 384-385.
  
5. Kalabay L, Chavin K, Lebreton JP, Robinson KA, Buse MG, Arnaud P. Human recombinant  $\alpha$ 2-HS glycoprotein is produced in insect cells as a full length inhibitor of the insulin receptor tyrosine kinase. *Hormone and Metabolic Research* **30** 1-6.
  
6. Hotamisligil GS. Mechanisms of TNF- $\alpha$  induced insulin resistance. *Experimental and Clinical Endocrinology and Diabetes* 1999 **107** 119-125.
  
7. Frühbeck G, Salvador J. Relation between leptin and the regulation of glucose metabolism. *Diabetologia* 2000 **43** 3-12.
  
8. Suda T, Takahashi N, Udagawa N, Jimi E, Gillespie MT, Martin TJ. Modulation of osteoclast differentiation and function by the new members of the tumor necrosis factor receptor and ligand families. *Endocrine Reviews* 1999 **20** 345-357.
  
9. Ducy P, Amling M, Takeda S, Priemel M, Schillink F, Bell FT et al. Leptin inhibits bone formation through a hypothalamic relay: a central control of bone mass. *Cell* 2000 **100** 197-207.
  
10. Buchanan TA. Intermediary Metabolism During Pregnancy: Implications for Diabetes Mellitus. In *Diabetes Mellitus. Fundamental and Clinical Text*, pp

677-684. Eds D LeRoith, SI Taylor, JM Olefsky. Philadelphia:  
Lippincott-Raven, 1996

11. Ratner RE. Gestational Diabetes Mellitus. In *Diabetes Mellitus. Fundamental and Clinical Text*, pp 710-715. Eds D LeRoith, SI Taylor, JM Olefsky. Philadelphia: Lippincott-Raven, 1996

12. Rossi G, Sherwin RS, Penzias AS, Lapaczewski P, Jacob RJ, Shulman GI et al. Temporal changes in insulin resistance and secretion in 24-h-fasted conscious pregnant rats. *American Journal of Physiology* 1993 **265** E845-E851.

13. Alberti KG, Zimmet PZ. Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus provisional report of a WHO consultation. *Diabetic Medicine* 1998 **15** 539-553.

14. Cleve H, Dencker H. Quantitative variations of the group-specific component (Gc) and of barium- $\alpha_2$ -glycoprotein of human serum in health and disease. In *Proteins and Biological Fluids* Vol. 14, pp 273-276. Ed H Peeters Oxford: Pergamon Press, 1966.

15. Lebreton JP, Joisel F, Raoult JP, Lannuzel B, Rogez JP, Humbert G. Serum concentration of human  $\alpha_2$  HS glycoprotein during the inflammatory process. *Journal of Clinical Investigation* 1979 **64** 1118-1129.

16. Kalabay L, Cseh K, Benedek S, Fekete S, Masszi T, Herjeczki K et al. Serum  $\alpha_2$ -HS glycoprotein concentration in patients with hematological malignancies. *Annals of Hematology* 1991 **63** 264-269.
17. Kalabay L, Jakab L, Fekete B, Prohászka Z, Benkő Zs, Teledgy L et al. Human fetuin/ $\alpha_2$ HS-glycoprotein level as a novel indicator of liver cell function and short-term mortality in patients with liver cirrhosis and liver cancer. *European Journal of Gastroenterology and Hepatology* 2002 **14** 1-6.
18. Haglund AC, Ek B, Ek P. Phosphorylation of human plasma  $\alpha_2$  Heremans-Schmid glycoprotein (fetuin) in vivo. *Biochemical Journal* 2001 **357** 437-446.
19. Winkler G, Salamon F, Harnos G, Salamon D, Speer G, Szekeres O et al. Elevated serum tumor necrosis factor-alpha concentrations and -bioactivity in Type 2 diabetics and patients with android type obesity. *Diabetes Research and Clinical Practice* 1998 **42** 169-174.
20. Pillai S, Zhou GY, Arnaud P, Jiang HX, Butler WJ, Zhang HM. Antibodies to endometrial transferrin and alpha2-Heremans-Schmid (HS) glycoprotein in patients with endometriosis. *American Journal of Reproductive Immunology* 1996 **35** 483-494.

21. Colclasure GC, Lloyd WS, Lamkin M, Gonnermann W, Troxler RF, Offner GD et al. Human serum  $\alpha_2$ HS glycoprotein modulates in vitro bone resorption. *Journal of Clinical Endocrinology and Metabolism* 1988 **66** 187-192.

22. Jahnen-Dechent W, Schinke T, Trindl A, Müller-Esterl W, Sablitzky F, Kaiser S et al. Cloning and targeted deletion of the mouse fetuin gene. *Journal of Biological Chemistry* 1997 **272** 31496-31503.



Table 1. Clinical and laboratory parameters of healthy pregnant women, patients with gestational diabetes mellitus patients and non-pregnant controls (mean  $\pm$  SD)

	<b>GDM patients</b>	<b>Healthy pregnant women</b>	<b>Non-pregnant women</b>
n	30	104	30
Age (years)	28.0 $\pm$ 2.8	27.8 $\pm$ 2.8	28.3 $\pm$ 3.5
Gestational age (week)	27.67 $\pm$ 6.1	24.3 $\pm$ 13.6	
Body mass index (kg/m <sup>2</sup> )	33.4 $\pm$ 6.4*	25.8 $\pm$ 2.7	22.9 $\pm$ 2.5
Fasting blood glucose concentration (mmol/l)	4.70 $\pm$ 1.50	4.48 $\pm$ 0.37	4.48 $\pm$ 0.27
Fasting C-peptide (ng/ml)	6.00 $\pm$ 3.01	1.85 $\pm$ 0.97	1.11 $\pm$ 0.82
C-peptide/Blood glucose ratio	0.40 $\pm$ 0.15	1.27 $\pm$ 0.51	0.27 $\pm$ 0.13
HbA <sub>1c</sub> (%)	5.11 $\pm$ 0.71	5.30 $\pm$ 0.80	4.96 $\pm$ 0.45
Serum fructosamine concentration ( $\mu$ mol/l)	203.3 $\pm$ 15.4	191.0 $\pm$ 18.5	
Daily insulin dose (U)	33.30 $\pm$ 20.83		
Daily insulin dose/Body weight (U/kg)	0.41 $\pm$ 0.21		

n: number of cases, \*:  $p < 0.01$  as compared to values of healthy pregnant women, calculated by the Mann-Whitney test.

Table 2. Anthropometric parameters, absolute values and percentiles of the newborns (mean  $\pm$  SD)

	<b>NEWBORNS</b>	
	<b>of patients with gestational diabetes mellitus n = 30</b>	<b>of healthy pregnant women n = 23</b>
Boys/girls	13 / 17	11 / 12
Body weight (g)	3151 $\pm$ 672*	3575 $\pm$ 365
Percentile	55.87 $\pm$ 30.29	64.75 $\pm$ 16.10
Body length (cm)	51.90 $\pm$ 3.40*	55.80 $\pm$ 2.80
Percentile	86.4 $\pm$ 31.88	110.90 $\pm$ 16.24
Head circumference (cm)	33.80 $\pm$ 1.70*	34.75 $\pm$ 1.15
Percentile	65.50 $\pm$ 22.84	71.30 $\pm$ 32.14

n: number of cases, \*:  $p < 0.01$  as compared to neonates of healthy pregnant women calculated by the Mann-Whitney test.

Table 3. Serum tumor necrosis factor (TNF)- $\alpha$ , soluble TNF-receptor-1 and -2 (sTNFR-1 and sTNFR-2), leptin, fasting C-peptide concentrations and the C-peptide/Blood glucose ratio in patients with gestational diabetes mellitus (GDM), different trimesters of the healthy pregnant group and non-pregnant control females (mean  $\pm$  SD)

	GDM	Healthy pregnant women			Non-pregnant controls
		1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	
		trimester			
Serum TNF- $\alpha$ (pg/ml)	6.3 $\pm$ 0.6**	4.1 $\pm$ 0.4#+	4.4 $\pm$ 0.4#+	5.5 $\pm$ 0.7*	4.1 $\pm$ 0.4
Serum sTNFR-1 (ng/ml)	3.2 $\pm$ 0.5*	2.1 $\pm$ 0.5#+	2.3 $\pm$ 0.5#++	2.8 $\pm$ 0.9**	2.01 $\pm$ 0.1
Serum sTNFR-2 (ng/ml)	10.0 $\pm$ 6.9**	4.7 $\pm$ 2.1#+	5.3 $\pm$ 3.7#++	5.7 $\pm$ 2.6**	3.3 $\pm$ 0.2
Serum leptin (ng/ml)	40.4 $\pm$ 24.5**	11.4 $\pm$ 7.2#+	11.1 $\pm$ 5.2#+	33.5 $\pm$ 22.0**	12.0 $\pm$ 9.1
Serum fasting C-peptide	6.0 $\pm$ 3.0**	1.2 $\pm$ 0.8#+	1.3 $\pm$ 0.4#+	3.1 $\pm$ 1.7*	1.1 $\pm$ 0.8
Fasting C-peptide/Blood glucose ratio	1.4 $\pm$ 0.5**	0.3 $\pm$ 0.1#+	0.3 $\pm$ 0.1#+	0.7 $\pm$ 0.2*	0.2 $\pm$ 0.1

\*:  $p < 0.05$ , \*\*:  $p < 0.01$ , as compared to the non-pregnant controls, #:  $p < 0.01$ , as compared to patients with GDM, +:  $p < 0.05$ , ++:  $p < 0.01$ , as compared to pregnant women in the third trimester, calculated by the Mann-Whitney test.

Table 4. Correlation between fetuin/ $\alpha_2$ -HS-glycoprotein concentration and laboratory parameters of patients with gestational diabetes mellitus (GDM), healthy pregnant women, and non-pregnant controls

	<b>GDM</b>	<b>Healthy pregnant women</b>	<b>Non-pregnant controls</b>
	<b>n = 30</b>	<b>n = 38</b>	<b>n = 30</b>
Body mass index	0.5702 <sup>#</sup>	0.2180	-0.0021
	0.2537 – 0.7764	-0.1099 – 0.5030	-0.3884 – 0.3513
	(0.0010)	(0.1766)	(0.9104)
Tumor necrosis factor- $\alpha$	0.6614	0.5853	-0.0541
	0.3858 – 0.8286	0.3263 – 0.7625	-0.3224 – 0.4158
	(< 0.0001)	(< 0.0001)	(0.7766)
sTNFR-1	0.0308	0.3956	-0.0091
	-0.3432 – 0.3963	0.0836 – 0.6353	-0.3779 – 0.3621
	(0.8718)	(0.0115)	(0.9618)
sTNFR-2	0.1003	0.2123	0.0318
	-0.2802 – 0.4535	-0.1157 – 0.4986	-0.3422 – 0.3972
	(0.5981)	(0.1884)	(0.8673)
Leptin	0.5775	0.3528	-0.1212
	0.2639 – 0.7807	0.0368 – 0.6047	-0.4702 – 0.2605
	(0.0008)	(0.0255)	(0.5233)
C-peptide	0.6699	0.4020	0.0519
	0.3988 – 0.8334	0.0939 – 0.6398	-0.3244 – 0.4139
	(< 0.0001)	(0.0101)	(0.7855)

C-peptide/Blood	0.5116	0.3241	0.2151
glucose ratio	0.0939 - 0.6398	0.0044 – 0.5867	-0.1683 – 0.5420
	(0.0039)	(0.0413)	(0.2537)

#: calculated by the Spearman's rank correlation. First row: correlation coefficient, second row: 95% confidence interval. Significance values are in brackets in the third row. n: number of cases, sTNFR-1: soluble tumor necrosis factor receptor 1, sTNFR-2: soluble tumor necrosis factor receptor 2.

Table 5. Correlation between fetuin/ $\alpha_2$ -HS-glycoprotein concentration and anthropological parameters of the neonates of patients with gestational diabetes mellitus (GDM) and healthy pregnant women

	<b>GDM</b>	<b>Healthy pregnant women</b>
	<b>n = 30</b>	<b>n = 23</b>
Head circumference	-0.6256 <sup>#</sup>	-0.6417
	-0.8085 - -0.3325	-0.8374 - -0.3003
	(0.0002)	(0.0010)
Body length	-0.4068	-0.5016
	-0.6752 - -0.0433	-0.7628 - -0.0998
	(0.0257)	(0.0147)
Body weight	-0.3098	-0.4998
	-0.6099 - 0.0680	-0.7773 - -0.0594
	(0.0957)	(0.0248)

<sup>#</sup> Calculated by the Spearman's rank correlation. First row: correlation coefficient, second row: 95% confidence interval. Significance values are in brackets in the third row. n: number of cases.




Figure legend:

Figure 1. Serum fetuin/ $\alpha$ 2HS-glycoprotein concentrations in different trimesters (I, II, III) of normal pregnancy, third trimester of gestation diabetes (GDM) and non-pregnant women (NP). Horizontal lines within scattergram represent means. Statistical differences between individual groups, represented by horizontal lines between groups were calculated by the Mann-Whitney test.