

IL NUOVO CIMENTO **38 C** (2015) 4
DOI 10.1393/ncc/i2015-15004-7

COLLOQUIA: IF AE 2014

Composite Higgs searches at the LHC and beyond

D. BARDUCCI⁽¹⁾⁽²⁾, A. BELYAEV⁽¹⁾⁽²⁾, M. S. BROWN⁽¹⁾, S. DE CURTIS⁽²⁾,
S. MORETTI⁽¹⁾⁽²⁾ and G. M. PRUNA⁽³⁾

⁽¹⁾ *School of Physics and Astronomy, University of Southampton, Highfield, Southampton SO17 1BJ, UK*

⁽²⁾ *INFN, Sezione di Firenze, Via G. Sansone 1, 50019, Sesto Fiorentino (FI), Italy*

⁽³⁾ *Paul Scherrer Institute, CH-5232 Villigen PSI, Switzerland*

received 7 January 2015

Summary. — General Composite Higgs models provide an elegant solution to the hierarchy problem present in the Standard Model and give an alternative pattern leading to the mechanism of electroweak (EW) symmetry breaking. We present an analysis of a realistic realization of this general idea, namely the 4DCHM, analysing the Higgs production and decay modes, fitting them to the latest LHC showing the compatibility with the results of the CERN machine. We then present the prospects of a future electron positron collider of testing this model against the expected experimental accuracies in the various Higgs decay channels accessible herein.

PACS 12.60.Rc – Composite particle models.

PACS 12.60.Cn – Extensions of gauge sector.

PACS 14.80.Va – Nambu-Goldstone.

1. – Introduction

After the discovery of a Higgs like state at the mass of 125 GeV made at CERN from the ATLAS [1] and CMS [2] Collaborations, one of the primary questions that the physics community ought to answer is whether this particle is consistent with the one predicted by the Standard Model (SM). From an experimental point of view the properties of this particle show both agreement and tensions with the predictions of the SM, although the errors on these measurements do not allow yet to draw a final conclusion on its properties. Conversely, from a theoretical point of view there are many motivations to think that the SM is not the ultimate and complete theory of Nature, among which the naturalness argument plays a predominant role. The instability of the Higgs mass with respect to radiative corrections requires in fact an incredible high level of fine tuning in the precision of their cancellation in the SM in order to have an Higgs mass at the EW scale. Beside the supersymmetric solution to this problem, another possibility is to postulate the Higgs boson as a composite state arising as a bound state from a strongly interacting sector at




Fig. 1. – $\gamma\gamma$ signal strength as a function of the mass of the lightest extra quark with charge $2/3$ (a) and χ^2 fit for various model scales (b) with the 7 and 8 TeV LHC data.

the TeV scale [3]. Being composite the Higgs will be insensitive to radiative corrections above the composite scale and its lightness with respect to other resonances of the strong sector can be taken into account by postulating the Higgs as a pseudo Nambu Goldstone boson (pNGB), similarly to what happens for the pions in QCD. Among the various composite Higgs models present in the literature, we will show in this proceeding the compatibility of a recent proposed framework, the 4D Composite Higgs Model (4DCHM) of [4] based on a $SO(5)/SO(4)$ breaking pattern, with respect to the 7 and 8 TeV LHC data and the capabilities of a future proposed e^+e^- collider in unravelling the composite nature of the 125 GeV scalar boson. See refs. [5, 6] for further details.

2. – LHC analysis

The couplings of the 4DCHM Higgs to SM model gauge bosons and fermions are modified with respect to the SM ones through factors that depend on the model scale f , explicitly $g_{\bar{f}fH}/g_{\bar{f}fH}^{SM} = (1 - 2\xi)/\sqrt{1-\xi}$ and $g_{VVH}/g_{VVH}^{SM} = \sqrt{1-\xi}$, with $\xi = v^2/f^2$ and where v is the Higgs vacuum expectation value. Beside these modifications due to the pNGB nature, other source of modifications are due to mass mixing between SM fermions and gauge bosons with the extra ones present in the 4DCHM and, in case of loop induced processes, of extra particles that can run inside the loops. Taking into account all these aspects we plot in fig. 1(a) the prediction for the $\gamma\gamma$ signal strength, $\mu_{\gamma\gamma}$ (that is the ratio of the 4DCHM event rate with respect to the SM expectation), in function of the mass of the lightest extra top present in the model, for a model scale of 1 TeV and showing, with a vertical dashed red line, an approximate exclusion limit on the masses of the extra fermions obtained by recasting the available results for direct searches of top partners. The arrow indicates the expected 4DCHM signal strength in case of the decoupling of all the extra particle content of the model. Comparing then our predictions for the signal strength in the $b\bar{b}$, WW , ZZ and $\gamma\gamma$ channels we have performed a χ^2 fit, fig. 1(b), for various model scale choices, where we can see the compatibility of the 4DCHM with the latest LHC Higgs data, also with respect to the SM, represented by an horizontal black line.

3. – Future e^+e^- collider analysis

We then tested our framework against a future e^+e^- collider for which we have chosen as a benchmark the proposed International Linear Collider (ILC) [7]. We have analysed




Fig. 2. $-\mu_{bb}$ and μ_{WW} signal strengths for the Higgs-strahlung production process at the ILC with $\sqrt{s} = 250$ GeV for $f = 1$ TeV (blue) and $f = 0.8$ TeV (green). Red dashed lines represent the expected experimental accuracies in measuring these observables according to [7] while the circles represent the 4DCHM predictions in case of the decoupling of all the extra particle content.

the 4DCHM signal strengths and compared them with the predicted accuracies for this machines in measuring these observables. In case of Higgs-strahlung production at $\sqrt{s} = 250$ GeV we plot the values of μ_{bb} and μ_{WW} for two different model scales, $f = 1$ TeV and $f = 0.8$ TeV, in fig. 2, where we also show with dashed red lines the expected experimental accuracies for this observables with 250 fb^{-1} of integrated luminosity and with the circles the values of the signal strengths in the limit of the decoupling of all the extra particle content of the 4DCHM. From the plot we observe that the ILC has the potential, already at $\sqrt{s} = 250$ GeV, of disentangling the 4DCHM with respect to the SM hypothesis and also the importance of keeping the full particle spectrum of the model, which would indeed render the deviations from the SM manifest, in contrast to the decoupling limit, which would then be clearly inappropriate to adopt in this case.

4. – Conclusions

In conclusion in this proceeding we have shown that the 4DCHM shows compatibility with the LHC data pointing to the discovery of a Higgs boson at 125 GeV and that a future e^+e^- collider will be able to test with higher precision the properties of this state so as to understand its nature.

* * *

DB, AB and SM are financed in part through the NExT Institute.

REFERENCES

- [1] AAD G. *et al.* (ATLAS COLLABORATION), *Phys. Lett. B*, **716** (2012) 1 arXiv:1207.7214 [hep-ex].
- [2] CHATRCHYAN S. *et al.* (CMS COLLABORATION), *Phys. Lett. B*, **716** (2012) 30 arXiv:1207.7235 [hep-ex].
- [3] KAPLAN D. B. and GEORGI H., *Phys. Lett. B*, **136** (1984) 183.

- [4] DE CURTIS S., REDI M. and TESI A., *JHEP*, **04** (2012) 042 arXiv:1110.1613 [hep-ph].
- [5] BARDUCCI D., BELYAEV A., BROWN M. S., DE CURTIS S., MORETTI S. and PRUNA G. M., *JHEP*, **09** (2013) 047 arXiv:1302.2371 [hep-ph].
- [6] BARDUCCI D., DE CURTIS S., MORETTI S. and PRUNA G. M., *JHEP*, **02** (2014) 005 arXiv:1311.3305 [hep-ph].
- [7] BAER H., BARKLOW T., FUJII K., GAO Y., HOANG A., KANEMURA S., LIST J., LOGAN H. E. *et al.*, arXiv:1306.6352 [hep-ph].